

EXTREMAL PLANE QUASICONFORMAL MAPPINGS WITH GIVEN BOUNDARY VALUES

BY EDGAR REICH AND KURT STREBEL
 Communicated by F. W. Gehring, October 23, 1972

1. Introduction. Let Ω_i , $i = 1, 2$, be regions in the complex plane, $f(z)$ a quasiconformal mapping of Ω_1 onto Ω_2 . Let Q_f denote the class of all quasiconformal mappings of Ω_1 onto Ω_2 which have the same boundary values as f . A mapping $f^* \in Q_f$ will be called *extremal* for its boundary values if it is K^* -quasiconformal and if there exists no K -quasiconformal mapping in Q_f with $K < K^*$. The quantity $K^* = K^*(f)$ is the extremal dilatation for the class Q_f . (As is well known [3], there may be more than one K^* -quasiconformal mapping in the class Q_f .) In the present account, which is only an abstract, we restrict ourselves to the case $\Omega_1 = \Omega_2 = E = \{z \mid |z| < 1\}$. Generalizations to Riemann surfaces will be referred to in a detailed account, giving proofs, further results, and applications that is to appear elsewhere.

In what follows, the L^1 norm $\int \int_E |\varphi(z)| dx dy$ of functions $\varphi(z)$ holomorphic in E will be denoted by $\|\varphi\|$.

In 1969, R. S. Hamilton [1] proved the following: *If $f^* \in Q_f$ is an extremal mapping, $\kappa^*(z) = f^*/f_z^*$, then*

$$(1.1) \quad \sup_{\|\varphi\| \leq 1} \left| \int \int_E \kappa^*(z) \varphi(z) dx dy \right| = k^*(f) = \frac{K^*(f) - 1}{K^*(f) + 1}.$$

A central result of the present work is (§3) that condition (1.1) characterizes extremal mappings of E .

2. Estimates for $K^*(f)$. The following is a generalization of an inequality proved in [2] from the case $K^*(f) = 1$ to arbitrary $K^*(f)$.

THEOREM 2.1. *If $f(z)$ is a quasiconformal self-mapping of E , $\kappa(z) = f_z/f_z^*$, and if $\varphi(z)$ is holomorphic in E , then*

$$(2.1) \quad \left| \int \int_E \frac{\kappa(z)}{1 - |\kappa(z)|^2} \varphi(z) dx dy \right| \leq \frac{k^*(f)}{1 + k^*(f)} \|\varphi\| + \int \int_E \frac{|\kappa(z)|^2}{1 - |\kappa(z)|^2} |\varphi(z)| dx dy.$$

AMS (MOS) subject classifications (1970). Primary 30A60.
¹ Supported by GP 27275.

For certain special holomorphic functions $\varphi(z)$ the quantity on the left side of (2.1) can be estimated in a useful way from below. Let us select n equally spaced points on $\{|z| = 1\}$, $n \geq 4$. These are mapped by $w = f(z)$ onto n points of $\{|w| = 1\}$. The unique extremal quasiconformal mapping of E onto E , which preserves this correspondence of n -gons, is [4] a Teichmüller mapping with complex dilatation

$$k_n(\overline{\varphi_n(z)}/|\varphi_n(z)|),$$

where $\varphi_n(z)$ is holomorphic in E and has finite norm $\|\varphi_n\|$. Without loss of generality we can assume that $\|\varphi_n\| = 1$. It is then possible to prove the following:

THEOREM 2.2. *Let $f(z)$ be a quasiconformal self mapping of E , $\kappa(z) = f_{\bar{z}}/f_z$. Let $n \geq 4$ points be selected on ∂E and k_n and $\varphi_n(z)$ determined as above. Then*

$$\left| \iint_E \frac{\kappa(z)}{1 - |\kappa(z)|^2} \varphi_n(z) \, dx \, dy \right| \geq \frac{k_n}{1 - k_n} - \iint_E \frac{|\kappa(z)|^2}{1 - |\kappa(z)|^2} |\varphi_n(z)| \, dx \, dy.$$

Given f , let us introduce the quantities

$$I(f) = \sup_{\|\varphi\| \leq 1} \left| \iint_E \frac{\kappa(z)\varphi(z)}{1 - |\kappa(z)|^2} \, dx \, dy \right| \quad (\kappa(z) = f_{\bar{z}}/f_z),$$

$$\Delta(f) = \sup_{\|\varphi\| \leq 1} \iint_E \frac{|\kappa(z)|^2 |\varphi(z)|}{1 - |\kappa(z)|^2} \, dx \, dy.$$

The following then follows easily as a corollary of Theorems 2.1 and 2.2:

THEOREM 2.3. *The maximal dilatation $K^*(f)$ which is extremal for the class Q_f satisfies the estimate*

$$\frac{1}{1 - 2I(f) + 2\Delta(f)} \leq K^*(f) \leq 1 + 2I(f) + 2\Delta(f).$$

REMARK. When f is a Teichmüller mapping with finite norm both inequalities become equalities.

3. Characterization of complex dilatations of extremal mappings. Let

$$H(f) = \sup_{\|\varphi\| \leq 1} \left| \iint_E \kappa(z)\varphi(z) \, dx \, dy \right|.$$

By means of Theorem 2.2 one obtains a new proof of Hamilton’s necessary condition (1.1), with an explicit identification of an extremal sequence for $H(f)$:

THEOREM 3.1. *If $f^* \in Q_f$ is extremal, $\kappa^*(z) = f^*/f_z^*$, then*

$$\lim_{n \rightarrow \infty} \left| \iint_E \kappa^*(z) \varphi_n(z) dx dy \right| = H(f) = k^*(f).$$

Furthermore, as a deduction from Theorem 2.3, the converse also follows:

THEOREM 3.2. *A necessary and sufficient condition for the quasiconformal mapping f of E onto E to be extremal for its boundary values is that*

$$H(f) = \operatorname{ess\,sup}_{z \in E} |\kappa(z)|.$$

REFERENCES

1. Richard S. Hamilton, *Extremal quasiconformal mappings with prescribed boundary values*, Trans. Amer. Math. Soc. **138** (1969), 399–406.
2. Edgar Reich and Kurt Strebel, *On quasiconformal mappings which keep the boundary points fixed*, Trans. Amer. Math. Soc. **138** (1969), 211–222. MR **38** #6059.
3. Kurt Strebel, *Zur Frage der Eindeutigkeit extremaler quasikonformer Abbildungen des Einheitskreises*. I, II, Comment. Math. Helv. **36** (1962), 306–323; *ibid.* **39** (1964), 77–89. MR **31** #346.
4. ———, *Über quadratische Differentiale mit geschlossenen Trajektorien und extreme quasikonforme Abbildungen*, Festband zum 70. Geburtstag von R. Nevanlinna, Springer, Berlin, 1966, pp. 105–127.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455

MATHEMATISCHES INSTITUT, UNIVERSITÄT ZÜRICH, CH-8006 ZÜRICH/ZH, SWITZERLAND