

## PERIODIC AND HOMOGENEOUS STATES ON A VON NEUMANN ALGEBRA. III

BY MASAMICHI TAKESAKI<sup>1</sup>

Communicated by Jacob E. Feldman, September 12, 1972

In this paper, we will show with a fairly complete proof that most of the results in [10] hold for homogeneous periodic states on a factor without the assumption of *inner* homogeneity. As an application, we will see that nonisomorphic ergodic automorphisms  $\theta$  of  $\mathcal{L}_0$  give rise to nonisomorphic factors  $\mathcal{B}(\mathcal{M}_0, \theta)$  of type III. We keep most of the terminology and the notations in [9] and [10].

We consider an arbitrary pair of homogeneous periodic states  $\varphi$  and  $\psi$  on a factor  $\mathcal{M}$  of the same period, say  $T > 0$ . Let  $\kappa = e^{-2\pi/T}$ ,  $0 < \kappa < 1$ . We denote by  $\mathcal{M}_n^{\varphi, \psi}$  the set of all  $x \in \mathcal{M}$  such that  $\rho_t(x) = \kappa^{int}x$ ,  $t \in \mathbf{R}$ , which was denoted by  $\mathcal{V}_n$  in [10]. With this alternation of the notation, we first note that Lemmas 1 through 6 remain valid without the assumption of inner homogeneity. Since  $\mathcal{M}_0^\varphi$  and  $\mathcal{M}_0^\psi$  are no longer factors, we have to analyze more carefully the relation between  $\mathcal{M}_0^\varphi$ ,  $\mathcal{M}_0^{\varphi, \psi}$  and  $\mathcal{M}_0^\psi$ . We denote by  $\mathcal{Z}_0^\varphi$  and  $\mathcal{Z}_0^\psi$  the center of  $\mathcal{M}_0^\varphi$  and  $\mathcal{M}_0^\psi$  respectively, and by  $u_\varphi$  and  $u_\psi$  the isometries in  $\mathcal{M}_1^\varphi$  and  $\mathcal{M}_1^\psi$  respectively which give rise to isomorphisms  $\theta_\varphi$  and  $\theta_\psi$  of  $\mathcal{M}_0^\varphi$  and  $\mathcal{M}_0^\psi$  onto  $e_\varphi \mathcal{M}_0^\varphi e_\varphi$  and  $e_\psi \mathcal{M}_0^\psi e_\psi$  respectively, where  $e_\varphi = u_\varphi u_\varphi^*$  and  $e_\psi = u_\psi u_\psi^*$ . We also denote by  $\tilde{\theta}_\varphi$  and  $\tilde{\theta}_\psi$  the automorphisms of  $\mathcal{Z}_0^\varphi$  and  $\mathcal{Z}_0^\psi$  induced by  $\theta_\varphi$  and  $\theta_\psi$  respectively. Since  $\mathcal{M}$  is a factor, we know from [9, Proposition 9] that  $\tilde{\theta}_\varphi$  and  $\tilde{\theta}_\psi$  are both ergodic.

LEMMA 1. *For each  $n \in \mathbf{Z}$ , we have*

$$(1) \quad \mathcal{M}_{n-1}^{\varphi, \psi} = u_\varphi^* \mathcal{M}_n^{\varphi, \psi} \quad \text{and} \quad \mathcal{M}_{n+1}^{\varphi, \psi} = \mathcal{M}_n^{\varphi, \psi} u_\psi.$$

PROOF. From [10, Lemma 5], it follows that  $\mathcal{M}_{n-1}^{\varphi, \psi} \supset u_\varphi^* \mathcal{M}_n^{\varphi, \psi}$ ; so

$$\mathcal{M}_{n-1}^{\varphi, \psi} = u_\varphi^* u_\varphi \mathcal{M}_n^{\varphi, \psi} \subset u_\varphi^* \mathcal{M}_n^{\varphi, \psi} \subset \mathcal{M}_{n-1}^{\varphi, \psi}.$$

Hence we get  $\mathcal{M}_{n-1}^{\varphi, \psi} = u_\varphi^* \mathcal{M}_n^{\varphi, \psi}$ . By symmetry, the assertion for  $u_\psi$  follows. Q.E.D.

LEMMA 2. *For any nonzero projections  $p \in \mathcal{M}_0^\varphi$  and  $q \in \mathcal{M}_0^\psi$ , we have*

$$p \mathcal{M}_n^{\varphi, \psi} \neq \{0\} \quad \text{and} \quad \mathcal{M}_n^{\varphi, \psi} q \neq \{0\}.$$

---

AMS (MOS) subject classifications (1970). Primary 46L10.

Key words and phrases. von Neumann algebra, modular automorphism group, periodic state, homogeneous state.

<sup>1</sup>This work is supported in part by NSF Grant No. GP-33696X.

PROOF. Let  $\mathcal{I}_n$  be the set of all  $x \in \mathcal{M}_0^\psi$  with  $x\mathcal{M}_n^{\varphi,\psi} = \{0\}$ . By [10, Lemma 5],  $\mathcal{I}_n$  is a  $\sigma$ -weakly closed ideal of  $\mathcal{M}_0^\psi$ , so that there exists a projection  $z_n \in \mathcal{L}_0^\psi$  such that  $\mathcal{I}_n = \mathcal{M}_0^\psi z_n$ . If  $x \in \mathcal{I}_n$ , then  $x\mathcal{M}_{n+1}^{\varphi,\psi} = x\mathcal{M}_n^{\varphi,\psi}u_\psi = \{0\}$  by (1), which means that  $\mathcal{I}_n \subset \mathcal{I}_{n+1}$ ; so  $z_n \leq z_{n+1}$ . We have

$$\tilde{\theta}_\varphi(z_n)e_\varphi\mathcal{M}_{n+1}^{\varphi,\psi} = u_\varphi z_n u_\varphi^* \mathcal{M}_{n+1}^{\varphi,\psi} = u_\varphi z_n \mathcal{M}_n^{\varphi,\psi} = \{0\},$$

so that  $\tilde{\theta}_\varphi(z_n)e_\varphi \leq z_{n+1}$ ; hence  $\tilde{\theta}_\varphi(z_n) \leq z_{n+1}$ . Conversely, we have

$$\tilde{\theta}_\varphi^{-1}(z_{n+1})\mathcal{M}_n^{\varphi,\psi} = u_\varphi^* z_{n+1} u_\varphi \mathcal{M}_n^{\varphi,\psi} \subset u_\varphi^* z_{n+1} \mathcal{M}_{n+1}^{\varphi,\psi} \subset \{0\}.$$

Hence we get  $\tilde{\theta}_\varphi^{-1}(z_{n+1}) \leq z_n$ , so that  $z_{n+1} \leq \tilde{\theta}_\varphi(z_n)$ . Thus we have  $z_{n+1} = \tilde{\theta}_\varphi(z_n)$ . Hence  $z_n \leq \tilde{\theta}_\varphi(z_n)$ . The equality  $\varphi(z_n) = \varphi \circ \tilde{\theta}_\varphi(z_n)$  implies that  $z_n$  must be either 0 or 1. Since  $\mathcal{M}_n^{\varphi,\psi} \neq \{0\}$ , we have  $z_n = 0$ . Hence  $p\mathcal{M}_n^{\varphi,\psi} \neq \{0\}$ . By symmetry,  $\mathcal{M}_n^{\varphi,\psi}q \neq \{0\}$ . Q.E.D.

LEMMA 3. Let  $v_1$  and  $v_2$  be partial isometries in  $\mathcal{M}_n^{\varphi,\psi}$  with initial projections  $q_1, q_2$  and final projections  $p_1, p_2$  respectively. Then the following statements are equivalent:

- (i)  $p_1$  and  $p_2$  are centrally orthogonal in  $\mathcal{M}_0^\psi$ , i.e.  $p_1\mathcal{M}_0^\psi p_2 = \{0\}$ ;
- (ii)  $q_1$  and  $q_2$  are centrally orthogonal in  $\mathcal{M}_0^\psi$ , i.e.  $q_1\mathcal{M}_0^\psi q_2 = \{0\}$ .

PROOF. By symmetry, we have only to prove (i)  $\Rightarrow$  (ii). Suppose  $q_1\mathcal{M}_0^\psi q_2 \neq \{0\}$ . Let  $x$  be an element in  $\mathcal{M}_0^\psi$  with  $q_1 x q_2 \neq \{0\}$ . We have then  $v_1^* v_1 x v_2^* v_2 \neq 0$ , so that  $v_1 x v_2^* \neq 0$ . Hence  $p_1 v_1 x v_2^* p_2 = v_1 x v_2^* \neq 0$ . But this is impossible because  $v_1 x v_2^*$  is in  $\mathcal{M}_0^\psi$  by [10, Lemma 5]. Q.E.D.

Suppose  $\{v_i\}_{i \in I}$  is a maximal family of partial isometries in  $\mathcal{M}_n^{\varphi,\psi}$  such that the initial projections  $q_i = v_i^* v_i$  are centrally orthogonal in  $\mathcal{M}_0^\psi$ . Let  $p_i = v_i v_i^*$ . By Lemma 3,  $\{p_i\}_{i \in I}$  are centrally orthogonal in  $\mathcal{M}_0^\psi$ . Hence  $v = \sum_{i \in I} v_i$  is a partial isometry in  $\mathcal{M}_n^{\varphi,\psi}$ . Let  $p = vv^*$  and  $q = v^*v$ . By Lemma 3, we conclude that the central supports of  $p$  and  $q$  in  $\mathcal{M}_0^\psi$  and  $\mathcal{M}_0^\psi$  are both the identity. Therefore, there exists an isomorphism  $\sigma_v$  of  $\mathcal{L}_0^\psi$  onto  $\mathcal{L}_0^\psi$  such that

$$(2) \quad \sigma_v(a)p = vav^*, \quad a \in \mathcal{L}_0^\psi.$$

LEMMA 4. For every projection  $f \in \mathcal{L}_0^\psi$ ,  $\sigma_v(f)$  is characterized as the smallest projection  $e \in \mathcal{L}_0^\psi$  such that  $e\mathcal{M}_n^{\varphi,\psi}f = \mathcal{M}_n^{\varphi,\psi}f$ .

PROOF. Let  $e$  be the smallest projection in  $\mathcal{L}_0^\psi$  with  $e\mathcal{M}_n^{\varphi,\psi}f = \mathcal{M}_n^{\varphi,\psi}f$ . We have then  $evf = vf$ , so that

$$\sigma_v(f)p = vf v^* = evf v^* e = e\sigma_v(f)pe = \sigma_v(f)ep.$$

Hence  $(\sigma_v(f) - \sigma_v(f)e)p = 0$ . Since the central support of  $p$  is 1, we have  $\sigma_v(f) = \sigma_v(f)e$ ; that is,  $\sigma_v(f) \leq e$ . If  $e - \sigma_v(f) \neq 0$ , then there exists an  $x \in \mathcal{M}_n^{\varphi,\psi}$  with  $[e - \sigma_v(f)]xf = x \neq 0$ . Let  $x = wh = kw$  be the left and right polar decomposition of  $x$ . As in the arguments (8) in [19],  $w \in \mathcal{M}_n^{\varphi,\psi}$ .

By the choice of  $x$ , we have  $ww^* \leq e - \sigma_v(f)$  and  $w^*w \leq f$ . On the other hand, we have  $vf = \sigma_v(f)vf$ , so that  $(vf)(vf)^* \leq \sigma_v(f)$  and  $(vf)^*(vf) = fv^*vf = fq$ . Hence the central support of  $(vf)^*(vf)$  in  $\mathcal{M}_0^\psi$  is  $f$ . But this is impossible by Lemma 3 because the central supports of  $(vf)(vf)^*$  and  $ww^*$  in  $\mathcal{M}_0^\psi$  are orthogonal. Thus we get  $\sigma_v(f) = e$ . Q.E.D.

Therefore, the isomorphism  $\sigma_v$  does not depend on the choice of  $v$ , but only on  $n \in \mathbb{Z}$ ; so we denote it by  $\sigma_n$ .

LEMMA 5. For each  $n \in \mathbb{Z}$ , we have

$$(3) \quad \sigma_n \circ \tilde{\theta}_\psi = \sigma_{n+1} = \tilde{\theta}_\varphi \circ \sigma_n.$$

PROOF. Let  $f$  be an arbitrarily fixed projection in  $\mathcal{L}^\psi$ . Let  $e_n = \sigma_n(f) \in \mathcal{L}_0^\varphi$ . We have then

$$e_{n+1}u_\varphi \mathcal{M}_n^{\varphi, \psi} f = u_\varphi \mathcal{M}_n^{\varphi, \psi} f,$$

$$u_\varphi^* e_{n+1} u_\varphi \mathcal{M}_n^{\varphi, \psi} f = \mathcal{M}_n^{\varphi, \psi} f.$$

Hence we have  $\tilde{\theta}_\varphi^{-1}(e_{n+1}) \geq e_n$ ; equivalently,  $e_{n+1} \geq \tilde{\theta}_\varphi(e_n)$ .

On the other hand, putting  $z = 1 - \tilde{\theta}_\varphi(e_n)$ , we have

$$u_\varphi^* z u_\varphi \mathcal{M}_n^{\varphi, \psi} f = (1 - e_n) \mathcal{M}_n^{\varphi, \psi} f = \{0\};$$

$$z e_\varphi \mathcal{M}_{n+1}^{\varphi, \psi} f = z u_\varphi \mathcal{M}_n^{\varphi, \psi} f = u_\varphi u_\varphi^* z u_\varphi \mathcal{M}_n^{\varphi, \psi} f = \{0\}.$$

Hence we have  $z e_\varphi \leq (1 - e_{n+1})$ ; so  $z \leq 1 - e_{n+1}$ . Therefore we get  $\tilde{\theta}_\varphi(e_n) \geq e_{n+1}$ . Thus we have  $e_{n+1} = \tilde{\theta}_\varphi(e_n)$ ; that is,  $\tilde{\theta}_\varphi \circ \sigma_n(f) = \sigma_{n+1}(f)$  for every projection  $f \in \mathcal{L}_0^\psi$ , which means that  $\sigma_{n+1} = \tilde{\theta}_\varphi \circ \sigma_n$ .

By symmetry, the other half of our assertion follows. Q.E.D.

COROLLARY 6. The ergodic automorphisms  $\tilde{\theta}_\varphi$  of  $\mathcal{L}_0^\varphi$  and  $\tilde{\theta}_\psi$  of  $\mathcal{L}_0^\psi$  are isomorphic.

LEMMA 7. If  $v$  is a partial isometry in  $\mathcal{M}_n^{\varphi, \psi}$  such that the initial projection  $q = v^*v$  and the final projection  $p = vv^*$  have the central support 1, then we have

$$(4) \quad p^h = \alpha \kappa^n \sigma_n(q^h),$$

where  $\alpha$  is the real number defined in [10].

PROOF. Consider a faithful state  $\varphi \circ \sigma_n$  on  $\mathcal{L}_0^\psi$ . Then  $\varphi \circ \sigma_n \circ \tilde{\theta}_\psi = \varphi \circ \tilde{\theta}_\varphi \circ \sigma_n = \varphi \circ \sigma_n$ , so that  $\varphi \circ \sigma_n$  is  $\tilde{\theta}_\psi$ -invariant; hence  $\varphi \circ \sigma_n$  is a scalar multiple of  $\psi$  on  $\mathcal{L}_0^\psi$  by the ergodicity of  $\tilde{\theta}_\psi$ . But  $\varphi \circ \sigma_n$  and  $\psi$  are both states, so that  $\varphi \circ \sigma_n = \psi$  on  $\mathcal{L}_0^\psi$ .

We have next, for every  $a \in \mathcal{L}_0^\psi$ ,

$$\begin{aligned}
 \psi(a\sigma_n^{-1}(p^h)) &= \varphi(\sigma_n(a\sigma_n^{-1}(p^h))) = \varphi(\sigma_n(a)p^h) \\
 &= \varphi(\sigma_n(a)p) = \varphi(vav^*) \\
 &= \alpha\kappa^n\psi(av^*v) \quad \text{by [10, Lemma 4]} \\
 &= \alpha\kappa^n\psi(aq) = \alpha\kappa^n\psi(aq^h).
 \end{aligned}$$

Thus, we get  $\sigma_n^{-1}(p^h) = \alpha\kappa^n q^h = \alpha\kappa^n q^h$ , equivalently  $p^h = \alpha\kappa^n \sigma_n(q^h)$ .

Making use of the similar arguments as in Lemma 2, we conclude the following:

**LEMMA 8.** *If  $p \in \mathcal{M}_0^\phi$  and  $q \in \mathcal{M}_0^\psi$  are projections with central support  $e$  and  $f$  in  $\mathcal{M}_0^\phi$  and  $\mathcal{M}_0^\psi$  respectively, then  $p\mathcal{M}_n^{\phi,\psi}q = \{0\}$  if and only if  $e\sigma_n(f) = 0$ .*

Now, let  $\{v_i\}_{i \in I}$  be a maximal family of partial isometries in  $\mathcal{M}_n^{\phi,\psi}$  such that the initial projections  $q_i = v_i^*v_i$  and the final projections  $p_i = v_i v_i^*$  are orthogonal respectively. Let  $v = \sum_{i \in I} v_i$ ,  $p = \sum_{i \in I} p_i$  and  $q = \sum_{i \in I} q_i$ . By Lemma 3, the central supports of  $p$  in  $\mathcal{M}_0^\phi$  and  $q$  in  $\mathcal{M}_0^\psi$  are respectively the identity. By maximality, we have  $(1 - p)\mathcal{M}_n^{\phi,\psi}(1 - q) = \{0\}$ . Let  $e$  and  $f$  be the central supports of  $p$  in  $\mathcal{M}_0^\phi$  and  $q$  in  $\mathcal{M}_0^\psi$  respectively. By Lemma 8,  $e\sigma_n(f) = 0$ . On the other hand, we have  $p^h = \alpha\kappa^n \sigma_n(q^h)$  by Lemma 7. Hence  $p^h \leq \alpha\kappa^n \leq 1$  if  $n \geq 1$ . Hence we have  $e = 1$ , so that  $f = 0$ ; so  $q = 1$ . Hence  $v$  must be an isometry if  $n \geq 1$ . Similarly, if  $n \leq 0$ , then  $v$  is a co-isometry. For  $n = 0$ ,  $v$  is unitary if and only if  $\alpha = 1$ . Thus we reach the following conclusion:

**THEOREM 9.** *If  $\varphi$  and  $\psi$  are homogeneous periodic states on a factor  $\mathcal{M}$  with same period, then there exists isometries  $u$  and  $v$  in  $\mathcal{M}$  such that*

$$\begin{aligned}
 \psi(x) &= \varphi(uxu^*)/\varphi(uu^*), \\
 \varphi(x) &= \psi(vxv^*)/\psi(vv^*), \quad x \in \mathcal{M}; \\
 p &= uu^* \in \mathcal{M}_0^\phi \quad \text{and} \quad q = vv^* \in \mathcal{M}_0^\psi.
 \end{aligned}$$

From this theorem, we can conclude that Theorems 8 through 10 in [10] hold for homogeneous periodic states  $\varphi, \psi$  with the same period and/or for projections  $p$  and  $q$  with uniform relative dimensions.

Let  $\mathcal{F}$  be a hyperfinite  $\text{II}_1$ -factor and  $\mathcal{A} = L^\infty(0, 1)$ . Let  $\mathcal{M}_0 = \mathcal{F} \otimes \mathcal{A}$ . For  $0 < \kappa < 1$ , we choose a projection  $f \in \mathcal{F}$  with  $\tau(f) = \kappa$ , where  $\tau$  is the canonical trace of  $\mathcal{F}$ . Let  $\theta$  be a fixed isomorphism of  $\mathcal{F}$  onto  $f\mathcal{F}f$ . For each  $\sigma \in \text{Aut}(\mathcal{F})$ , let  $\theta_\sigma = \theta \circ \sigma$ . Let  $\tilde{\theta}$  be an ergodic automorphism of  $\mathcal{A}$  with invariant faithful normal state  $\mu$ . Changing  $\tilde{\theta}$  under an automorphism of  $\mathcal{A}$ , we may assume that  $\mu$  is given by the Lebesgue measure on  $(0, 1)$ . Let  $\varphi_0 = \tau \otimes \mu$ . We obtain then a factor  $\mathcal{R}(\mathcal{M}_0, \theta_\sigma \otimes \tilde{\theta}, \varphi_0)$  as described in [9]. We denote it by  $\mathcal{M}(\kappa, \sigma, \tilde{\theta})$ .

THEOREM 10. We choose  $\sigma_1, \sigma_2 \in \text{Aut}(\mathcal{F})$  and ergodic automorphisms  $\tilde{\theta}_1$  and  $\tilde{\theta}_2$  of  $\mathcal{A}$  and fix  $\kappa$ . A necessary and sufficient condition for  $\mathcal{M}(\kappa, \sigma_1, \tilde{\theta}_1) \cong \mathcal{M}(\kappa, \sigma_2, \tilde{\theta}_2)$  is that

- (i)  $\tilde{\theta}_1$  and  $\tilde{\theta}_2$  are isomorphic as ergodic automorphisms of  $\mathcal{A}$ ;
- (ii) there exist a projection  $p$  in  $\mathcal{F}$  with  $p \geq f$ , a partial isometry  $w$  in  $\mathcal{F}$  and an isomorphism  $\rho$  of  $\mathcal{F}$  onto  $p\mathcal{F}p$  such that

$$w\theta \circ \sigma_1 \circ \rho(x)w^* = \rho \circ \theta \circ \sigma_2(x);$$

$$\theta \circ \sigma_1 \circ \rho(x) = w^* \rho \circ \theta \circ \sigma_2(x)w, \quad x \in \mathcal{M}.$$

Furthermore, if  $\tilde{\theta}_1$  and  $\tilde{\theta}_2$  have no point spectrum other than 1, then the isomorphism  $\mathcal{M}(\kappa_1, \sigma_1, \tilde{\theta}_1) \cong \mathcal{M}(\kappa_2, \sigma_2, \tilde{\theta}_2)$  implies that  $\kappa_1 = \kappa_2$  as well as (i) and (ii).

ACKNOWLEDGEMENT. The author would like to thank Professor H. Araki for enjoyable discussions on the subject of this paper; Dr. A. Connes for his kindness informing the author of his new results on which this work depends; Professor H. A. Dye for his constant encouragement; and the members of the functional analysis seminar at Queen's University where the related topics were extensively discussed.

#### REFERENCES

1. A. Connes, *Un nouvel invariant pour les algèbres de von Neumann*, C. R. Acad. Sci. Paris Sér. A-B **273** (1971), A900–A903.
2. ———, *Calcul des deux invariants d'Araki et Woods par la théorie de Tomita et Takesaki*, C. R. Acad. Sci. Paris Sér. A-B **274** (1972), A175–A177.
3. ———, *Groupe modulaire d'une algèbre de von Neumann de genre dénombrable*, C.R. Acad. Sci. Paris Sér. A-B **274** (1972), A1923–A1926.
4. ———, *États presque périodiques sur une algèbre de von Neumann*, C. R. Acad. Sci. Paris Sér. A-B **274** (1972), A1402–A1405.
5. ———, A letter to the author, dated June 17, 1972.
6. J. Dixmier, *Les algèbres d'opérateurs dans l'espace hilbertien (Algèbre de von Neumann)*, Cahiers scientifiques, fasc. 25, Gauthier-Villars, Paris, 1957. MR **20** # 1234.
7. E. Størmer, *Spectra of states, and asymptotically abelian C\*-algebras* (to appear).
8. M. Takesaki, *Tomita's theory of modular Hilbert algebras and its applications*, Lecture Notes in Math., vol. 128, Springer-Verlag, Berlin and New York, 1970. MR **42** # 5061.
9. ———, *Periodic and homogeneous states on a von Neumann algebra*. I. Bull. Amer. Math. Soc. **79** (1973), 202–206.
10. ———, *Periodic and homogeneous states on a von Neumann algebra*. II, Bull. Amer. Math. Soc. **79** (1973), 416–420.
11. ———, *The structure of a von Neumann algebra with a homogeneous periodic state*, Acta Math. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024