

## INTERPOLATION OF OPERATORS FOR $\Lambda$ SPACES

BY ROBERT SHARPLEY

Communicated by Alberto Calderón, September 22, 1973

Lorentz and Shimogaki [2] have characterized those pairs of Lorentz  $\Lambda$  spaces which satisfy the interpolation property with respect to two other pairs of  $\Lambda$  spaces. Their proof is long and technical and does not easily admit to generalization. In this paper we present a short proof of this result whose spirit may be traced to Lemma 4.3 of [4] or perhaps more accurately to the theorem of Marcinkiewicz [5, p. 112]. The proof involves only elementary properties of these spaces and does allow for generalization to interpolation for  $n$  pairs and for  $M$  spaces, but these topics will be reported on elsewhere.

The Banach space  $\Lambda_\phi$  [1, p. 65] is the space of all Lebesgue measurable functions  $f$  on the interval  $(0, l)$  for which the norm

$$\|f\|_\phi = \int_0^l f^*(s)\phi(s) ds$$

is finite, where  $\phi$  is an integrable, positive, decreasing function on  $(0, l)$  and  $f^*$  (the decreasing rearrangement of  $|f|$ ) is the almost-everywhere unique, positive, decreasing function which is equimeasurable with  $|f|$ .

A pair of spaces  $(\Lambda_\phi, \Lambda_\psi)$  is called an interpolation pair for the two pairs  $(\Lambda_{\phi_1}, \Lambda_{\psi_1})$  and  $(\Lambda_{\phi_2}, \Lambda_{\psi_2})$  if each linear operator which is bounded from  $\Lambda_{\phi_i}$  to  $\Lambda_{\psi_i}$  (both  $i=1, 2$ ) has a unique extension to a bounded operator from  $\Lambda_\phi$  to  $\Lambda_\psi$ .

**THEOREM (LORENTZ-SHIMOGAKI).** *A necessary and sufficient condition that  $(\Lambda_\phi, \Lambda_\psi)$  be an interpolation pair for  $(\Lambda_{\phi_1}, \Lambda_{\psi_1})$  and  $(\Lambda_{\phi_2}, \Lambda_{\psi_2})$  is that there exist a constant  $A$  independent of  $s$  and  $t$  so that*

$$(*) \quad \Psi(t)/\Phi(s) \leq A \max_{i=1,2} (\Psi_i(t)/\Phi_i(s))$$

holds, where  $\Phi(s) = \int_0^s \phi(r) dr, \dots, \psi_2(t) = \int_0^t \Psi_2(r) dr$ .

**PROOF.** We only sketch the proof of the necessity since it is standard.

*AMS (MOS) subject classifications* (1970). Primary 46E30, 46E35, 47A30.

*Key words and phrases.* Lorentz  $\Lambda_\phi$  space, decreasing rearrangement, interpolation of operators.

Copyright © American Mathematical Society 1974

Suppose there are numbers  $s_n$  and  $t_n$  in  $(0, l)$  such that  $\Psi'(t_n)/\Phi(s_n) > n^3 \max_{i=1,2}(\Psi'_i(t_n)/\Phi_i(s_n))$ . Define the positive operator

$$T_n f(t) = \left( C_n \int_0^{s_n} f(s) ds/s_n \right) \chi_{(0,t_n)}(t),$$

where  $C_n = \min_{i=1,2}(\Phi_i(s_n)/\Psi'_i(t_n))$ .

For each  $f$  in  $\Lambda_{\phi_i}$ ,  $T_n f$  belongs to  $\Lambda_{\psi_i}$  and  $T_n$  has operator norm less than or equal to 1, but as an operator from  $\Lambda_{\phi}$  to  $\Lambda_{\psi}$ ,  $T_n$  has operator norm larger than  $n^3$ . Hence the operator  $T = \sum_1^\infty T_n/n^2$  is a bounded operator from  $\Lambda_{\phi_i}$  to  $\Lambda_{\psi_i}$  ( $i=1, 2$ ), but  $T$  is not a bounded operator from  $\Lambda_{\phi}$  to  $\Lambda_{\psi}$ .

To show that condition (\*) is sufficient, we prove that

$$(1) \quad \|Tf\|_{\psi} \leq 2AM \|f\|_{\phi}$$

where  $M$  is the maximum of the operator norms of  $T$  acting from  $\Lambda_{\phi}$  to  $\Lambda_{\psi_i}$  ( $i=1, 2$ ). We can assume that  $f$  is an arbitrary simple function with finite support since these functions are dense in  $\Lambda_{\phi}$ . We can also require  $f$  to be positive since  $\|f\|_{\phi} = \| |f| \|_{\phi}$ . Each function of this type can be written as  $f = \sum_1^n \alpha_i \chi_{E_i}$  where the  $\alpha_i$ 's are positive and  $E_n \subset \dots \subset E_1$ . Hence  $f^* = \sum_1^n \alpha_i \chi_{(0,a_i)}$  where  $a_i = mE_i$ . But then

$$(2) \quad \|T\chi_E\|_{\psi} \leq 2AM\Phi(mE), \quad \text{all measurable } E \subset (0, l)$$

is equivalent to relation (1), since

$$\|Tf\|_{\psi} \leq \sum_1^n \alpha_i \|T\chi_E\|_{\psi} \leq 2AM \sum_1^n \alpha_i \Phi(a_i) = 2AM \|f\|_{\phi}.$$

Hence, if we let  $g = (T\chi_E)^*$ , the proof is reduced to the following

LEMMA. *Suppose condition (\*) holds and  $g$  is a positive decreasing function that satisfies*

$$(3) \quad \|g\|_{\psi_i} \leq M\Phi_i(a) \quad (i = 1, 2),$$

then

$$(4) \quad \|g\|_{\psi} \leq 2AM\Phi(a).$$

PROOF. First assume  $g$  is a step function with finite support, i.e.,  $g = \sum_1^m \beta_j \chi_{(0,t_j)}$ . Set  $J = \{j | \max_{i=1,2}(\Psi'_i(t_j)/\Phi_i(a)) = \Psi'_1(t_j)/\Phi_1(a)\}$  and then let  $g_1 = \sum_{j \in J} \beta_j \chi_{(0,t_j)}$  and  $g_2 = g - g_1$ . Notice that both functions are positive, decreasing, step functions and

$$(5) \quad \|g_i\| \leq \|g\|$$

in any  $\Lambda$  space. Now using condition (\*), relations (5) and (3), we have

$$\begin{aligned} \|g_1\|_{\psi}/\Phi(a) &= \sum_{j \in J} \beta_j \Psi'(t_j)/\Phi(a) \\ &\leq A \sum_{j \in J} \beta_j \max_{i=1,2}(\Psi'_i(t_j)/\Phi_i(a)) \\ &= A \sum_{j \in J} \beta_j \Psi'_1(t_j)/\Phi_1(a) = A \|g_1\|_{\psi_1}/\Phi_1(a) \\ &\leq A \|g\|_{\psi_1}/\Phi_1(a) \leq AM. \end{aligned}$$

Similarly

$$\|g_2\|_{\psi}/\Phi(a) \leq A \|g_2\|_{\psi_2}/\Phi_2(a) \leq AM.$$

Hence, we obtain relation (4) for positive, decreasing, step functions.

Now suppose  $g$  is an arbitrary positive decreasing function and let  $\{g_n\}$  be a monotone increasing sequence of positive decreasing step functions converging pointwise to  $g$ . By (3) and (5)

$$\|g_n\|_{\psi_i} \leq M\Phi_i(a) \quad (i = 1, 2)$$

so

$$\|g_n\|_{\psi} \leq 2AM\Phi(a).$$

Applying the monotone convergence theorem to  $\{g_n\psi\}$ , we obtain relation (4).

The author wishes to thank Professor S. D. Riemenschneider for many helpful conversations regarding this work.

#### REFERENCES

1. G. G. Lorentz, *Bernstein polynomials*, Mathematical Expositions, no. 8, Univ. of Toronto Press, Toronto, 1953. MR 15, 217.
2. G. G. Lorentz and T. Shimogaki, *Interpolation theorems for operators in function spaces*, J. Functional Analysis 2 (1968), 31–51. MR 41 #2424.
3. ———, *Interpolation theorems for spaces  $\Lambda$* , Abstract Spaces and Approximation (Proc. Conf., Oberwolfach, 1968), Birkhäuser, Basel, 1969, pp. 94–98. MR 41 #2423.
4. R. C. Sharpley, *Spaces  $\Lambda_\alpha(X)$  and interpolation*, J. Functional Analysis 11 (1972), 479–513.
5. A. Zygmund, *Trigonometric series*. Vol. II, 2nd ed. reprinted with corrections and some additions, Cambridge Univ. Press, New York, 1968. MR 38 #4882.

DEPARTMENT OF MATHEMATICS, OAKLAND UNIVERSITY, ROCHESTER, MICHIGAN 48063