

RESEARCH ANNOUNCEMENTS

The purpose of this department is to provide early announcement of significant new results, with some indications of proof. Although ordinarily a research announcement should be a brief summary of a paper to be published in full elsewhere, papers giving complete proofs of results of exceptional interest are also solicited. Manuscripts more than eight typewritten double spaced pages long will not be considered as acceptable. All research announcements are communicated by members of the Council of the American Mathematical Society. An author should send his paper directly to a Council member for consideration as a research announcement. A list of members of the Council for 1974 is given at the end of this issue.

ON THE MAXIMUM N TH DIAMETER

BY B. VOLK

Communicated by Shlomo Sternberg, September 12, 1973

ABSTRACT. Herein is disproved, in some cases, a plausible conjecture on the maximum value of the N th diameter of a closed bounded connected planar set of logarithmic capacity one.

Introduction. Let N be a positive integer greater than one. Define the N th diameter of a closed bounded connected planar set to be the largest geometric mean of the distances between any N points of the set. This notion is a generalization of the usual diameter—which is just the case $N=2$. Then, if the above set is of logarithmic capacity one, how big can the N th diameter get?

Fix the set in question. Let d_N denote the N th diameter of the set. Then, as shown by M. Schiffer (in [4]),

$$d_N \leq 4^{1/N} N^{1/(N-1)} \quad \text{for } N = 2, 3.$$

This is the bound achieved only for the set consisting of the N -pronged slit forking at the origin and terminating at the N th roots of four, or translations of the set. This set is a contender for the set with maximum N th diameter, since the function of the form $az \downarrow$ power series in $(1/z)$ which maps the exterior of the unit disc onto the exterior of the set is $(z^N \downarrow 2 \downarrow z^{-N})^{1/N}$, whose coefficient of the z term is exactly one.

Thus the following conjecture seems plausible:

For all N , $d_N \leq 4^{1/N} N^{1/(N-1)}$. Equality is achieved iff the set, Γ , consists

AMS (MOS) subject classifications (1970). Primary 30A44.

Copyright © American Mathematical Society 1974

of N linear segments from the origin to the N th roots of four, or translations of the set.

This diameter conjecture is, as mentioned above, true for $N=2$ and for $N=3$. It was, however, stated by P. R. Garabedian and M. M. Schiffer (in [1]) that the conjecture is false for $N=4$. Here, it will be shown that the conjecture is in fact false for all N such that N is an even integer greater than two. The other cases, when N is an odd integer greater than three, remain open.

Let Σ' denote the class of functions

$$G(z) = z \perp b_1 z^{-1} \perp b_2 z^{-2} \perp \dots \perp b_n z^{-n} \perp \dots$$

which are analytic and univalent for all z such that $|z| > 1$. Let $Q(z) =$ that inverse function of $z^{-1}(1 \perp z^N)^{2/N}$ which is in Σ' . Let $\lambda =$ a positive real number greater than one. Then (G. Pólya and G. Szegő in [3]) a little reflection shows that, for any function $G(z)$ in Σ' , the function

$$H(z) = \frac{4^{1/N} \lambda}{(\lambda^N \perp 1)^{2/N}} G \left[Q \left(\frac{(\lambda^N \perp 1)^{2/N} (1 \perp z^N)^{2/N}}{4^{1/N} \lambda z} \right) \right]$$

is in Σ' and omits the N points

$$\frac{4^{1/N} \lambda}{(\lambda^N \perp 1)^{2/N}} G(\lambda \omega^k) \quad \text{with } \omega^N = 1, \text{ and } k = 1, 2, 3, \dots, N.$$

Hence, if D_N denotes the largest possible value of the N th diameter for any closed bounded connected planar set of logarithmic capacity one, then

$$D_N \geq \frac{4^{1/N} \lambda}{(1 \perp \lambda^N)^{2/N}} \prod_{i \neq j} |G(\lambda \omega^i) - G(\lambda \omega^j)|^{1/(N^2 - N)} \quad \text{where } \omega^N = 1.$$

Hence if the conjecture be true then

$$(1) \quad \prod_{j \neq k} \left| \frac{G(\lambda \omega^j) - G(\lambda \omega^k)}{\lambda \omega^j - \lambda \omega^k} \right| \leq (1 \perp 2\lambda^{-N} \perp \lambda^{-2N})^{N-1}.$$

Now define the double sequence of complex numbers $\{C_{P,Q}\}$ to be the Grunsky coefficients of $G(z)$ if

$$-\text{Log} \left\{ \frac{G(x) - G(y)}{x - y} \right\} = \sum_{P,Q \geq 1} C_{P,Q} x^{-P} y^{-Q}.$$

Then on comparing both sides of (1), supposedly,

$$|C_{1,N-1} \perp C_{2,N-2} \perp C_{3,N-3} \perp \dots \perp C_{N-1,1}| \leq 2 - 2/N.$$

The following lemma thus completes the disproof of the conjecture, for N an even integer greater than two.

LEMMA. $|C_{1,2M-1} \perp C_{2,2M-2} \perp \cdots \perp C_{2M-1,1}| > 2 - 1/M$ for some function $G_M(z)$ and for $M=2, 3, 4, \dots$.

PROOF. Let $M > 1$ be fixed. Let

$$F(z) = z \perp a_2 z^2 \perp \cdots \perp a_n z^n \perp \cdots$$

be univalent for z such that $|z| < 1$, and let

$$J(z) = F^{-1/M}(z^{-M}) \\ = z - M^{-1} a_2 z^{1-M} \perp (\frac{1}{2} M^{-1} a_2^2 \perp \frac{1}{2} M^{-2} a_2^2 - M^{-1} a_3) z^{1-2M} \perp \cdots .$$

Then some Grunsky coefficients of $J(z)$ are

$$C_{k,2M-k} = C_{2M-k,k} = \frac{1}{2} M^{-1} a_2^2 \perp \frac{1}{2} k M^{-2} a_2^2 - M^{-1} a_3$$

where $k = 1, 2, 3, \dots, M$

Therefore for $J(z)$

$$(2) \quad |C_{1,2M-1} \perp C_{2,2M-2} \perp \cdots \perp C_{2M-1,1}| = (2 - 1/M) \left| \left(a_3 - \frac{3M-1}{4M-2} a_2^2 \right) \right|.$$

Now select $F(z)$ to maximize the quantity

$$(3) \quad |a_3 - \alpha a_2^2| \quad \text{where } \alpha = \frac{3M-1}{4M-2}$$

over the set of functions with normalized power series which are univalent in the interior of the origin-centered unit disc. Then, as proven by G. M. Golusin (in [2]),

$$|a_3 - \alpha a_2^2| = 1 \perp 2 \exp(-2\alpha/(1 - \alpha)) \quad \text{for } \alpha \text{ in } (0, 1).$$

This, in conjunction with (2) and (3), proves the Lemma and the result.

ACKNOWLEDGEMENT. This paper is the core of a 1967 thesis at Yeshiva University, written under the direction of Professor H. E. Rauch. The education was graciously supported by the National Science Foundation.

REFERENCES

1. P. R. Garabedian and M. Schiffer, *A coefficient inequality for schlicht functions*, Ann. of Math. (2) **61** (1955), 116-136. MR **16**, 579.
2. G. M. Golusin, *Geometrische Funktionentheorie*, Hochschulbücher für Mathematik, Band 31, VEB Deutscher Verlag der Wissenschaften, Berlin, 1957. MR **19**, 735.
3. G. Pólya and G. Szegő, *Aufgaben und Lehrsätze aus der Analysis*, Springer, Berlin, 1925.
4. M. Schiffer, *Sur un probleme d'extremum de la représentation conforme*, Bull. Soc. Math. France **66** (1938), 432-449.

13-15 DICKENS STREET, FAR ROCKAWAY, NEW YORK 11691