

## FIXED POINTS OF DISK ACTIONS

BY ROBERT OLIVER

Communicated October 30, 1975

As a sequel to a previous announcement [3], the author can now give a complete classification up to homotopy type of which spaces can occur as fixed point sets of smooth actions of a given compact Lie group on disks. The result is contained in Theorems 1 to 3 below. For a group  $G$ ,  $G_0$  denotes its identity component.

**THEOREM 1.** *Let  $G$  be a compact Lie group, and  $F$  a finite CW complex. Then there exists a smooth action of  $G$  on a disk with fixed point set having the homotopy type of  $F$  if and only if:*

1.  $G \cong T^n$  ( $n \geq 1$ ):  $F$  is  $\mathbf{Z}$ -acyclic;
  2.  $G_0$  a torus and  $|G/G_0| = p^a$  ( $p$  prime,  $a \geq 1$ ):  $F$  is  $\mathbf{Z}_p$ -acyclic,
  3.  $G_0$  not a torus or  $G/G_0$  not of prime power order:  $\chi(F) \equiv 1 \pmod{n_G}$
- for some fixed integer  $n_G$ .

In order to describe the calculations of  $n_G$ , some classes of finite groups are defined, as in [3] and [4].  $G^1$  denotes the class of all  $G$  with normal subgroup  $P$  of prime power order, such that  $G/P$  is cyclic. For  $q$  prime,  $G^q$  denotes the class of all  $G$  with normal subgroup  $H \in G^1$  of  $q$ -power index. Then one gets

- THEOREM 2.**
1. *If  $G_0$  is not a torus, then  $n_G = 1$ .*
  2. *If  $G_0$  is a torus, then  $n_G = n_{G/G_0}$ .*
  3. *If  $G$  is finite, then  $n_G = 0$  if and only if  $G \in G^1$ ; if  $G \notin G^1$  then for any prime  $q$ ,  $q \mid n_G$  if and only if  $G \in G^q$ .*

In Theorem 1, the necessity of the conditions in (1) and (2) follow from standard Smith theory. Sufficiency follows in (2) from Jones [2], and in (1) is trivial ( $G * F$  is contractible and can be thickened up to a disk action by Theorem 6 of [4]).

For finite  $G$ , the existence of  $n_G$  and the calculations in Theorem 2, part 3, were proven in [4]. Furthermore, if  $G_0$  is a torus and  $G \supseteq G_0$ , then  $F$  clearly has the homotopy type of the fixed point set of a disk action of  $G$  if and only if it does the same for  $G/G_0$ , so  $n_G = n_{G/G_0}$ . The case where  $G_0$  is nontoral will be dealt with below; the above theorems say that *any* finite homotopy type can occur as fixed point set for such  $G$ .

The following result, completing the calculation of  $n_G$ , was obtained in

---

AMS (MOS) subject classifications (1970). Primary 57E25; Secondary 55C35.

Copyright © 1976, American Mathematical Society

Theorem 4 of [5] by studying the projective obstruction  $\gamma_G(F)$  first introduced in [4].

**THEOREM 3.** *For any finite group  $G$ ,  $n_G = 4$  if and only if:*

1.  $G$  is a semidirect product  $0 \rightarrow \mathbf{Z}_n \rightarrow G \rightarrow \mathbf{Z}_{2^k} \rightarrow 0$  ( $n$  odd) given by an automorphism  $\alpha \in \text{Aut}(\mathbf{Z}_n)$ .
2.  $G \notin G^1$ , but the subgroup of index 2 is in  $G^1$ .
3. Letting  $\alpha$  also denote the induced automorphism of  $\mathbf{Z}\xi_n$  (the ring generated by the  $n$ th roots of unity), there is no unit  $u \in (\mathbf{Z}\xi_n)^*$  such that  $\alpha(u) = -u$ . Otherwise,  $n_G$  equals 0, 1 or a product of distinct primes.

Groups fulfilling conditions 1–3 do actually exist, the smallest being given by  $\langle a, b : a^{15} = b^4 = e, bab^{-1} = a^2 \rangle$ .

It remains to describe the case of groups with nontoral identity component; by Bredon's construction [1, §I.8] it is enough to construct a fixed point free action of any such group on a disk. The following theorem provides some very specific examples of such actions. The concept of a *family* of subgroups is used, as defined by tom Dieck.

**THEOREM 4.** *Let  $G$  be a compact Lie group, and  $F$  a nonempty family of subgroups. Then there exists a smooth action of  $G$  on a disk  $D$  such that  $D^H$  is a disk for  $H \in F$  and empty for  $H \notin F$ , if and only if:*

1. For any pair of subgroups  $H \triangleleft K$  in  $G$ , for which  $K/H$  has prime order, either both  $H$  and  $K$  are in  $F$  or neither is.
2.  $F$  is closed in the space of closed subgroups of  $G$  with the Hausdorff topology.

In particular, the family of subgroups  $H$  such that  $H_0$  is a torus and  $H/H_0$  solvable meets these conditions.

#### REFERENCES

1. G. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972.
2. L. Jones, *The converse to the fixed point theorem of P. A. Smith*. I, *Ann. of Math.* (2) **94** (1971), 52–68. MR 45 #4427.
3. R. Oliver, *Fixed points of finite group actions on contractible complexes*, *Bull. Amer. Math. Soc.* **81** (1975), 482–484.
4. ———, *Fixed point sets of group actions on finite acyclic complexes*, *Comment Math. Helv.* **50** (1975), 155–177.
5. ———, *Projective obstructions to group actions on disks* (to appear)

MATEMATISK INSTITUT, ÅRHUS UNIVERSITET, ÅRHUS, DENMARK