

ON SURFACES OBTAINED FROM QUATERNION ALGEBRAS OVER REAL QUADRATIC FIELDS¹

BY IRA SHAVEL

Communicated by Hyman Bass, May 9, 1976

Let A be a totally indefinite division quaternion algebra with center $k = \mathbf{Q}(\sqrt{d})$, $d > 0$, \mathcal{O} a maximal order in A , and $\Gamma(1) = \{\alpha \in \mathcal{O} \mid \nu(\alpha) = 1\}$ where ν is the reduced norm from A to k . Fix an isomorphism λ such that $A \otimes_{\mathbf{Q}} \mathbf{R} \cong M_2(\mathbf{R}) \oplus M_2(\mathbf{R})$. Then $\lambda(\Gamma(1) \otimes_{\mathbf{Q}} 1) \subseteq \mathrm{SL}_2(\mathbf{R}) \times \mathrm{SL}_2(\mathbf{R})$, and $j(\Gamma(1)) = \Gamma(1)/(\text{center } \Gamma(1))$ acts holomorphically and properly discontinuously on $X = H \times H$, where H is the usual upper half plane. In general, if Γ is any group of holomorphic automorphisms of X acting properly discontinuously and without fixed points, then $\Gamma \backslash X$ is a complex manifold. Since A is division the quotient is compact, and it is known to be a projective algebraic variety. In this note we discuss the numerical invariants and second cohomology group of $U(\Gamma) = \Gamma \backslash H \times H$ where Γ is commensurable with $\Gamma(1)$.

(A) For any algebraic number field F , a quaternion algebra with center F is determined up to isomorphism by a finite set $S(A)$ of prime divisors of F . Denote this algebra by $A(F, S(A))$.

THEOREM 1. *Assume $h(k) = \text{class number of } k = 1$. Let $j(\Gamma(1)) = \Gamma(1)/\{\pm 1\}$, $A = A(k, S(A))$, and let*

$$\left(\frac{\cdot}{p} \right)$$

be the Kronecker symbol. $j(\Gamma(1))$ acts on X without fixed points \Leftrightarrow all of the following hold:

$$(1) \quad \left(\frac{-3}{p} \right) = 1 \quad \text{or} \quad \left(\frac{-D}{p} \right) = 1$$

for some $P \in S(A)$, where $p\mathbf{Z} = P \cap \mathbf{Z}$ and $-D'$ is the discriminant of the field $\mathbf{Q}(\sqrt{-3d})$.

$$(2) \quad \left(\frac{-1}{p} \right) = 1 \quad \text{or} \quad \left(\frac{-D'}{p} \right) = 1$$

for some $P \in S(A)$, where $p\mathbf{Z} = P \cap \mathbf{Z}$ and $-D'$ is the discriminant of the field $\mathbf{Q}(\sqrt{-d})$.

AMS (MOS) subject classifications (1970). Primary 14J20; Secondary 12A80, 22E40.

¹ Partial results of the author's dissertation [3] under M. Kuga.

(3) If $d = 5$, $\exists P \in S(A)$ such that $pZ = P \cap Z$ and $p \equiv 1 \pmod{5}$.

Let $A^{X^{++}} = \{\alpha \in A^X \mid \nu(\alpha) \text{ is totally positive}\}$ and call such α totally positive. Let $E^{++} = \mathcal{O}^X \cap A^{X^{++}}$. $|j(E^{++}): j(\Gamma(1))| = 2$ if ϵ_k , the fundamental unit of k greater than 1, is totally positive, and $|j(E^{++}): j(\Gamma(1))| = 1$ otherwise.

THEOREM 2. Assume $h(k) = 1$ and ϵ_k is totally positive. $j(E^{++})$ acts on X without fixed points \Leftrightarrow both of the following hold:

- (1) $j(\Gamma(1))$ has no elements of finite order.
- (2) $\exists P \in S(A)$ such that P splits in $k(\sqrt{-\epsilon_k})/k$.

Consider $B^{++} = \{\beta \in A^{X^{++}} \mid \beta\mathcal{O} = \mathcal{O}\beta\} = \text{normalizer of } \Gamma(1) \text{ in } A^{X^{++}}$. If $h(k) = 1$ then the class number of a maximal order in A is also 1. Therefore every 2-sided \mathcal{O} -ideal is principal. The set of all 2-sided maximal \mathcal{O} -ideals are in one-to-one correspondence with the prime ideals of \mathcal{O}_k . Let $P_i = \Pi_i\mathcal{O}$ correspond to $P_i = \pi_i\mathcal{O}_k$.

THEOREM 3. Assume $h(k) = 1$. Let ϵ be a fundamental unit of \mathcal{O}_k . Let $\{\pi_i\}_{i=1,2,\dots,n}$ correspond to $\{\Pi_i\mathcal{O}\}_{i=1,2,\dots,n} = S(A)$. For these π_i let $\eta(i_1, i_2, \dots, i_r) = \pi_{i_1}\pi_{i_2} \cdots \pi_{i_r}$ where $\pi_{i_s} \neq \pi_{i_t}$ for $s \neq t$. $j(B^{++})$ acts on X without fixed points if and only if both of the following hold:

- (1) $j(E^{++})$ has no elements of finite order.
- (2) For all totally positive $\eta(i_1, i_2, \dots, i_r)$, $\exists P \in S(A)$ such that P splits in $k(\sqrt{-\eta(i_1, i_2, \dots, i_r)})/k$, and for all totally positive $\eta(i_1, i_2, \dots, i_r)\epsilon$ (for some choice of ϵ), $\exists P \in S(A)$ such that P splits in $k(\sqrt{-\eta(i_1, i_2, \dots, i_r)\epsilon})/k$.

(B) Throughout this section Γ is a group commensurable with $j(\Gamma(1))$ acting on X without fixed points. Using a result of Matsushima and Shimura [2] we have

PROPOSITION 1. (1) The Euler characteristic E , the geometric genus p_g , and the arithmetic genus p_a of $\Gamma \backslash X$ have the following relationship: $E = 4(p_g + 1) = 4p_a$.

- (2) The irregularity q is 0.
- (3) Then m th plurigenus $P_m = (p_g + 1)(2m - 1)^2$, $m \geq 2$.

COROLLARY. $\Gamma \backslash X$ is a surface of general type.

Using the Riemann-Roch theorem we have

COROLLARY. $c_1^2 = 8p_g + 8$, where c_1 is the first Chern class of $\Gamma \backslash X$.

Using a formula of Shimizu [4] for the volume of a fundamental domain for the action of $j(\Gamma(1))$ on X , and the Gauss-Bonnet theorem we obtain

THEOREM 4. $E(U(1))$, the Euler characteristic of $j(\Gamma(1)) \backslash X$ is given by

$$E(U(1)) = \frac{B_d}{12} \prod_{P \in S(A)} (N_{k/Q} P - 1)$$

where B_d is the generalized Bernoulli number of the numerical character modulo d associated to the field $k = \mathbf{Q}(\sqrt{d})$.

For $d \neq 5$, B_d is an integer. With the aid of a computer, James Maiorana has calculated B_d for $d < 750$.

We have a complete list of surfaces with $p_g = 0$ and $p_g = 1$ which come from groups Γ , $j(\Gamma(1)) \subseteq \Gamma \subseteq j(B^{++})$.

(c) Let $U(1) = j(\Gamma(1)) \backslash X$ be an algebraic variety. $H_1(U(1), \mathbf{Z})$ is isomorphic to $H^2(U(1), \mathbf{Z})_{\text{torsion}}$ by Poincaré and Pontrjagin duality. Thus

$$H^2(U(1), \mathbf{Z})_{\text{tor}} \cong j(\Gamma(1)) / [j(\Gamma(1)), j(\Gamma(1))] \cong \Gamma(1) / \{\pm 1\} [\Gamma(1), \Gamma(1)].$$

By constructing a normal subgroup of $\Gamma(1)$ containing $[\Gamma(1), \Gamma(1)]$, we obtain

THEOREM 5. *Let $j(\Gamma(1))$ act on X without fixed points. Then $|H^2(U(1), \mathbf{Z})_{\text{tor}}|$ is divisible by $a \cdot b \cdot c \cdot \prod_{P \in S(A)} (N_{k/Q} P + 1)$ where*

$$a = \begin{cases} \frac{1}{2} & \text{if } N_{k/Q} P \equiv 1 \pmod{4} \text{ for some } P \in S(A), \\ 1 & \text{otherwise;} \end{cases}$$

$$b = \begin{cases} 4 & \text{if } \exists P, Q \text{ such that } P \neq Q, PQ = 2\mathbf{Z} \text{ and } P, Q \notin S(A), \\ 2 & \text{if } \exists P, Q \text{ such that } PQ = 2\mathbf{Z} \text{ and } P \notin S(A) \text{ but } Q \in S(A), \text{ or if } \exists P \text{ such} \\ & \text{that } P^2 = 2\mathbf{Z} \text{ and } P \notin S(A), \\ 1 & \text{otherwise;} \end{cases}$$

$$c = \begin{cases} 9 & \text{if } \exists P, Q \text{ such that } P \neq Q, PQ = 3\mathbf{Z} \text{ and } P, Q \notin S(A), \\ 3 & \text{if } \exists P, Q \text{ such that } PQ = 3\mathbf{Z} \text{ and } P \notin S(A) \text{ but } Q \in S(A), \text{ or if } \exists P \text{ such} \\ & \text{that } P^2 = 3\mathbf{Z} \text{ and } P \notin S(A), \\ 1 & \text{otherwise.} \end{cases}$$

EXAMPLE. Let $A = A(\mathbf{Q}(\sqrt{5}, \{P_5, P_{31}\}))$. We have $P_5^2 = 5\mathbf{Z}$, $N_{k/Q} P_5 = 5$, $P_{31} P'_{31} = 31\mathbf{Z}$, $N_{k/Q} P_{31} = 31$, $N_{k/Q} P_2 = 4$, $N_{k/Q} P_3 = 9$, $\epsilon_k = (1 + \sqrt{5})/2$, $N_{k/Q} \epsilon_k = -1$, and $B_5 = 4/5$. $U(1) = j(\Gamma(1)) \backslash X$ is smooth, $E(U(1)) = (1/12) \cdot (4/5)(5 - 1)(31 - 1) = 8$, so $p_g = 1$. $|H^2(U(1), \mathbf{Z})_{\text{tor}}|$ is divisible by $(1/2)(5 + 1)(31 + 1) = 96$. There are two subgroups between $j(\Gamma(1))$ and $j(B^{++})$ yielding $p_g = 0$ surfaces. For more examples see [3].

(D) Let K be the canonical line bundle of a surface of the above type. In conjunction with Gordon Jenkins, we have shown that in the case $P_g = 0$, $3K$ is very ample, that is, $3K$ determines a biholomorphic imbedding into some complex projective space.

Gordon Jenkins [1] has investigated cases where $[k : \mathbf{Q}] \geq 3$.

REFERENCES

1. G. Jenkins, Thesis, SUNY at Stony Brook, 1976.
2. Y. Matsushima and G. Shimura, *On the cohomology groups attached to certain vector valued differential forms on the product of the upper half planes*, Ann. of Math. (2) **78** (1963), 417–449. MR 27 #5274.
3. I. Shavel, Thesis, SUNY at Stony Brook, 1976.
4. H. Shimizu, *On discontinuous groups operating on the product of the upper half planes*, Ann. of Math. (2) **77** (1963), 33–71. MR 26 #2641.
For correction see: H. Shimizu, *On zeta functions of quaternion algebras*, Ann. of Math. (2) **81** (1965), 166–193. MR 30 #1998.

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK, CENTER AT STONY BROOK, STONY BROOK, NEW YORK 11794