

LOCAL GEVREY AND QUASI-ANALYTIC HYPOELLIPTICITY FOR \square_b

BY DAVID S. TARTAKOFF¹

Communicated by Daniel W. Stroock, May 27, 1976

Introduction. The $\bar{\partial}_b$ complex is well defined on any smooth CR manifold M , and once a metric is fixed, so is the complex Laplace-Beltrami operator \square_b on forms of type (p, q) . For compact M without boundary, the $\bar{\partial}_b$ cohomology of M may be studied via \square_b [6], and thus local smoothness of solutions to $\square_b u = f$ is important. In its own right, \square_b is a prototype of doubly characteristic operators. Under suitable convexity conditions on M , Kohn [7] established the following subelliptic estimate on (p, q) forms in $C_0^\infty(M)$:

$$\|\varphi\|_{\frac{1}{2}}^2 \leq C(\square_b \varphi, \varphi) + C'\|\varphi\|_0^2,$$

and in general such estimates imply C^∞ and Gevrey (G^s , $s \geq 2$) hypoellipticity locally [3], [8], [10] and no more [1]. In the special case of the Heisenberg group, Folland and Stein [5] found an explicit fundamental solution which gives local analytic hypoellipticity; while, in general, if M is compact, satisfies the convexity condition $Y(q)$ of Kohn, and has an invertible Levi form, the author proved \square_b is globally analytic hypoelliptic and so is the $\bar{\partial}$ -Neumann problem (joint work with M. Derridj, cf. [4], [9]).

In this note we assume $Y(q)$ and the invertibility of the Levi form and prove local regularity in all Gevrey classes G^s with $s > 1$ as well as in a quasi-analytic class. Full details will appear elsewhere.

Notations and definitions. The class $C^L(\Omega) \subset C^\infty(\Omega)$, Ω open in R^n , is defined by the condition that for all $K \subset\subset \Omega$ there exists a constant $C_{f,K}$ such that for any multi-index α ,

$$\sup_K |D^\alpha f| \leq C_{f,K}^{|\alpha|+1} L(|\alpha|)^{|\alpha|},$$

where we assume that the sequence $\{L(j)\}$ of positive numbers satisfies (1) $L(j)/j$ is nondecreasing and (2) $L(j+1)^{j+1} \leq C^j L(j)^j$ uniformly in j . The second condition implies that $C^L(\Omega)$ is closed under differentiation while the first implies that the class is preserved under composition. Thus one may speak of C^L manifolds. If, in addition, $\Sigma L(j)^{-1} < \infty$, the class is called non-quasi-analytic (NQA) and admits compactly supported functions. Common examples are the Gevrey classes $G^s(\Omega)$, obtained by taking $L(j) = j^s$. These are NQA if $s > 1$, while

AMS (MOS) subject classifications (1970). Primary 32F10, 35H05.

¹Partially supported by NSF MPS 72-05055 A03.

Copyright © 1976, American Mathematical Society

taking $s = 1$ gives the real analytic class. The class obtained by taking $L(j) = j \log j$ is quasi-analytic, and thus contains no compactly supported functions.

A smooth manifold M of dimension $2n - 1$ is called *CR* provided (a) $CT(M)_x = S_x \oplus \bar{S}_x \oplus F_x$ for all x , S a smooth subbundle of $CT(M)$ of complex dimension $n - 1$, orthogonal to \bar{S} under a smooth Hermitian inner product which induces a Riemannian metric on M , and F of complex dimension 1, and (b) if Y_1, Y_2 are local sections of S , so is their commutator $[Y_1, Y_2]$. $D^{p,q}(M)$, the space of smooth (p, q) forms on M , is defined to be the space of those smooth $p + q$ forms on M such that $h(t_1, \dots, t_{p+q}) = 0$ if p of the t 's and q of the \bar{t} 's have zero projection on S . For $v \in D^{p,q}(M)$ we define $\bar{\partial}_b v$ to be the projection on $D^{p,q+1}(M)$ of dv . The $\bar{\partial}_b$ form a complex and we denote by $\square_b: D^{p,q}(M) \rightarrow D^{p,q}(M)$ the operator $\bar{\partial}_b \bar{\partial}_b^* + \bar{\partial}_b^* \bar{\partial}_b$, where $\bar{\partial}_b^*$ denotes the formal L^2 adjoint of $\bar{\partial}_b$. When Y_1, \dots, Y_{n-1} forms a local frame for S and T denotes a local, nowhere zero, purely imaginary section of F , the matrix c_{ij} , given by $[Y_i, \bar{Y}_j] \equiv c_{ij}T$ modulo $S \oplus \bar{S}$, is the Levi form of M . The number of its nonzero eigenvalues and its signature in absolute value are independent of the choice of Y_j and of T . M satisfies $Y(q)$ if c_{ij} has $\max(q + 1, n - q)$ eigenvalues of the same sign or pairs of eigenvalues of opposite signs. M is strictly pseudoconvex if all eigenvalues are strictly of the same sign.

Results.

THEOREM. *Let M be a CR manifold of class C^L , L satisfying (1) and (2) above and non-quasi-analytic, of dimension $2n - 1$ with an invertible Levi form satisfying $Y(q)$ in an open set Ω . Then any $u \in D^{p,q}(\Omega)$ satisfying $\square_b u \in C^L(\Omega)$ is itself in $C^L(\Omega)$.*

PROPOSITION (BOMAN, cf. [2]). *The intersection of all $C^{L'}(\Omega)$, L satisfying (1) and (2) and non-quasi-analytic, is the class $C^{L'}$ where $L'(j) = j \log j$.*

COROLLARY. *Let M be a CR manifold of the quasi analytic class $C^{L'}$, $L'(j) = j \log j$, with invertible Levi form satisfying $Y(q)$. Then any $u \in D^{p,q}(\Omega)$, Ω open in M , with $\square_b u \in C^{L'}(\Omega)$, is itself in this class.*

Remarks. It is well known that one need only assume that $u \in D'(\Omega)$ for the above results to hold [6], [8]. Also, there is a direct proof of the Corollary which obviates the quasi-analyticity of $C^{L'}$ and obtains the local result by considering a family of compactly supported functions whose derivatives, up to a given order, grow uniformly as if the functions belonged to $C^{L'}$. Families of this sort were introduced by Ehrenpreis to localize some real analytic problems; while his families fail to satisfy an analogue of (1) above, one may approximate $C^{L'}$ by families which do.

REFERENCES

1. M. S. Baouendi and C. Goulaouic, *Analyticity for degenerate elliptic equations and*

applications, Proc. Sympos. Pure Math., vol. 23, Amer. Math. Soc., 1973, pp. 79–84. MR 50 #5167.

2. J. Boman, *On the intersection of classes of infinitely differentiable functions*, Ark. Mat. 5 (1963/65), 301–309. MR 36 #3116.

3. M. Derridj, *Sur la régularité G^2 au bord des solutions du problème de Neumann pour l'opérateur $\bar{\partial}$* (to appear).

4. M. Derridj and D. S. Tartakoff, *On the Global Real Analyticity of Solutions to the $\bar{\partial}$ -Neumann Problem*, Comm. in Partial Differential Equations.

5. G. B. Folland and E. Stein, *Estimates for the $\bar{\partial}_b$ -complex and analysis on the Heisenberg group*, Comm. Pure Appl. Math. 27 (1974), 429–522. MR 51 #3719.

6. J. J. Kohn, *Harmonic integrals on strongly pseudo-convex manifolds. I, II*, Ann. of Math. (2) 78 (1963), 112–148; *ibid.* 79 (1964), 450–472. MR 27 #2999; 34 #8010.

7. ———, *Boundaries of complex manifolds*, Proc. Conf. Complex Analysis, (Minneapolis, 1964), Springer-Verlag, Berlin, 1965, pp. 81–94. MR 30 #5334.

8. D. S. Tartakoff, *Gevrey hypoellipticity for subelliptic boundary value problems*, Comm. Pure Appl. Math. 26 (1973), 251–312. MR 49 #7586.

9. ———, *On the global real analyticity of solutions to \square_b on compact manifolds*, Comm. in Partial Differential Equations (to appear).

10. F. Trèves, *An invariant criterion of hypoellipticity*, Amer. J. Math. 83 (1961), 645–668. MR 24 #A2732.

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON,
NEW JERSEY 08540