

BORDISM OF DIFFEOMORPHISMS

BY M. KRECK

Communicated by P. T. Church, April 28, 1976

1. **Introduction.** In this note we determine the bordism groups Δ_n of orientation preserving diffeomorphisms of n -dimensional closed oriented smooth manifolds. These groups were introduced by W. Browder [1]. Winkelkemper showed that each diffeomorphism of the sphere S^n is nullbordant [7]. On the other hand, he showed that Δ_{4k+2} is not finitely generated. Medrano generalized this result to Δ_{4k} [5]. For this he introduced a powerful invariant in the Witt group $W_{\pm}(\mathbf{Z}, \mathbf{Z})$ (I_{\pm} in Medrano's notation) of isometries of free finite-dimensional \mathbf{Z} -modules with a symmetric (antisymmetric) unimodular bilinear form. The invariant is given by the middle homology modulo torsion, the intersection form and the isometry induced by the diffeomorphism. For a diffeomorphism $f: M \rightarrow M$ we denote this invariant by $I(M, f)$, the isometric structure of (M, f) . It is a bordism invariant and leads to a homomorphism $I: \Delta_{2k} \rightarrow W_{(-1)k}(\mathbf{Z}, \mathbf{Z})$.

Neumann has shown that the homomorphism I is surjective, that $W_{\pm}(\mathbf{Z}, \mathbf{Z}) \otimes \mathbf{Q} \cong \mathbf{Q}^{\infty}$ and that $W_{\pm}(\mathbf{Z}, \mathbf{Z})$ contains infinitely many summands of orders 2 and 4 [6]. On the other hand, $W_{\pm}(\mathbf{Z}, \mathbf{Z})$ is a subgroup of $W_{\pm}(\mathbf{Z}, \mathbf{Q})$, the Witt group of isometries of finite-dimensional \mathbf{Q} -vector spaces. This group plays an important role in the computation of bordism groups C_{2k-1} of odd-dimensional knots, which can be embedded in $W_{(-1)k}(\mathbf{Z}, \mathbf{Q})$. It is known that $W_{\pm}(\mathbf{Z}, \mathbf{Q}) \cong \mathbf{Z}^{\infty} \oplus \mathbf{Z}_2^{\infty} \oplus \mathbf{Z}_4^{\infty}$ [3]. Thus the group $W_{\pm}(\mathbf{Z}, \mathbf{Z})$ is also of the form $\mathbf{Z}^{\infty} \oplus \mathbf{Z}_2^{\infty} \oplus \mathbf{Z}_4^{\infty}$.

It turns out that the isometric structure is essentially the only invariant for bordism of diffeomorphisms.

2. **Bordism of odd-dimensional diffeomorphisms.** Two diffeomorphisms (M_1, f_1) and (M_2, f_2) are called bordant if there is a diffeomorphism (N, F) on an oriented manifold with boundary such that $\partial(N, F) = (M_1, f_1) + (-M_2, f_2)$. The bordism classes $[M^n, f]$ form a group under disjoint sum, called Δ_n .

The mapping torus of a diffeomorphism (M, f) is $M_f = I \times M / (0, x) \sim (1, f(x))$. This construction leads to a homomorphism $\Delta_n \rightarrow \Omega_{n+1}([M, f] \mapsto [M_f])$, where Ω_{n+1} is the ordinary bordism group of oriented manifolds.

In [4] we proved the following result.

AMS (MOS) subject classifications (1970). Primary 57D90.

Copyright © 1976, American Mathematical Society

THEOREM 1. *For $k > 2$ the map $[M, f] \mapsto ([M], [M_f])$ is an isomorphism $\Delta_{2k-1} \rightarrow \Omega_{2k-1} \oplus \hat{\Omega}_{2k}$, where $\hat{\Omega}_{2k}$ is the kernel of the signature homomorphism τ .*

3. The even-dimensional case. Consider triples (G, \langle, \rangle, h) , where G is a finite-dimensional free \mathbb{Z} -module, \langle, \rangle a symmetric (resp. antisymmetric) unimodular bilinear form on G and h an isometry of (G, \langle, \rangle) . (G, \langle, \rangle, h) is called hyperbolic if there exists an invariant subkernel, i.e. a subspace $U \subset G$ with $U \subset U^\perp$, $2 \dim U = \dim G$ and $h(U) \subset U$. (G, \langle, \rangle, h) and $(G', \langle', \rangle', h')$ are called bordant if $(G, \langle, \rangle, h) \oplus (G', -\langle', \rangle', h')$ is hyperbolic. This is an equivalence relation. The equivalence classes form a group under orthogonal sum, called $W_+(\mathbb{Z}, \mathbb{Z})$ (resp. $W_-(\mathbb{Z}, \mathbb{Z})$).

The isometric structure of a diffeomorphism (M^{2k}, f) is given by $(H_k(M; \mathbb{Z})/\text{Tor}, \circ, f_*)$, where \circ is the intersection form. If (M, f) bounds a diffeomorphism (N, F) the isometric structure is hyperbolic, an invariant subkernel being given by the kernel of $i_*: H_k(M; \mathbb{Z})/\text{Tor} \rightarrow H_k(N; \mathbb{Z})/\text{Tor}$, so we have a homomorphism $I: \Delta_{2k} \rightarrow W_{(-1)k}(\mathbb{Z}, \mathbb{Z})$. Neumann has shown that this homomorphism is surjective [6].

THEOREM 2. *For $k > 1$ the homomorphism*

$$\begin{aligned} \Delta_{2k} &\rightarrow W_{(-1)k}(\mathbb{Z}, \mathbb{Z}) \oplus \hat{\Omega}_{2k} \oplus \Omega_{2k+1}, \\ [M, f] &\mapsto (I(M, f), [M] - \tau(M)[P_k \mathbb{C}], [M_f]) \end{aligned}$$

is an isomorphism (k even), injective with cokernel \mathbb{Z}_2 (k odd).

4. Idea of the proof. Consider a diffeomorphism (M^n, f) such that M and M_f are nullbordant. This implies that there exists a differentiable map $g: N^{n+2} \rightarrow S^1$ with $\partial N = M_f$ and $g|_{\partial N}$ the canonical projection from M_f to S^1 . Let $x \in S^1$ be a regular value of g . $F := g^{-1}(x)$ is a 1-codimensional submanifold of N with trivial normal bundle meeting ∂N transversally along ∂F : Cut N along F to obtain a differentiable manifold N_F with corners. The boundary of N_F consists of two copies F_0 and F_1 of F with opposite orientations and a manifold $V = \partial N_{\partial F}$ fibred over the unit interval I . $\partial V = \partial F_0 + \partial F_1$. The corners of N_F are at ∂F_0 and ∂F_1 .

We now make the following strong assumption (compare [2, 2.3]): The components of F are simply connected and F_0 and F_1 are deformation retracts of N_F .

Then $(N_F; F_0, F_1)$ is a relative h -cobordism between $(F_0, \partial F_0)$ and $(F_1, \partial F_1)$. For $n > 5$ the h -cobordism theorem implies that the diffeomorphism on M can be extended to a diffeomorphism of F .

Our aim is, starting with an arbitrary N_F to obtain an N'_F , which satisfies the above assumption. For this we modify N_F by addition and subtraction of handles. If n is odd ($n \neq 3$) we have shown in [4] that this works. If n is even

($n > 2$) and $I(M, f) = 0$ we can do the same. The details of the proof will appear elsewhere.

REFERENCES

1. W. Browder, *Surgery and the theory of differentiable transformation groups*, Proc. Conf. on Transformation Groups (New Orleans, La., 1967), Springer-Verlag, New York, 1968, pp. 1–46. MR 41 #6242.
2. W. Browder and J. Levine, *Fibering manifolds over a circle*, Comment. Math. Helv. 40 (1966), 153–160. MR 33 #3309.
3. M. A. Kervaire, *Knot cobordism in codimension two*, Manifolds—Amsterdam 1970 (Proc. Nuffic Summer School), Lecture Notes in Math., vol. 197, Springer-Verlag, Berlin and New York, 1971, pp. 83–105. MR 44 #1016.
4. M. Kreck, *Cobordism of odd-dimensional diffeomorphisms*, Topology (to appear).
5. S. López de Medrano, *Cobordism of diffeomorphisms of $(k - 1)$ -connected $2k$ -manifolds*, Proc. Second Conf. on Compact Transformation Groups (Amherst, 1971), Part I., Lecture Notes in Math., vol. 298, Springer-Verlag, Berlin and New York, 1972, pp. 217–227. MR 51 #1847.
6. W. D. Neumann, *Signature related invariants of manifolds* (in preparation).
7. H. E. Winkelkemper, *On equators of manifolds and the actions of θ^n* , Thesis, Princeton Univ., 1970.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT, SONDERFORSCHUNGSBE-
REICH THEORETISCHE MATHEMATIK, WEGELERSTR. 10, 53 BONN, GERMANY