

FINITE RICKART C^* -ALGEBRAS

BY DAVID HANDELMAN AND JOHN LAWRENCE¹

Communicated by Barbara Osofsky, May 23, 1977

1. A C^* -algebra is Rickart [1] if the right annihilator of each element t is generated by a projection $1 - RP(t)$. If in addition, $xx^* = 1$ implies $x^*x = 1$, the ring is called a finite Rickart C^* -algebra. In this note we announce several new results on finite Rickart C^* -algebras. Detailed proofs will appear elsewhere.

Two projections e and f in a Rickart C^* -algebra are $*$ -equivalent ($e \overset{*}{\sim} f$) if there exists a w such that $ww^* = e$ and $w^*w = f$. Kaplansky asked whether left projections in a Rickart C^* -algebra were $*$ -equivalent to right projections, that is, whether $RP(t) \overset{*}{\sim} LP(t)$ [5]. We have the following partial answer.

THEOREM 1 [2]. *In a finite Rickart C^* -algebra, left projections are equivalent to right projections. In fact $RP(t)$ and $LP(t)$ are unitarily equivalent for each element t in the algebra.*

A consequence of this is the following.

COROLLARY 2 [2]. *A simple homomorphic image of a finite Rickart C^* -algebra is a finite AW^* -factor.*

THEOREM 3 [3]. *If T is a finite Rickart C^* -algebra that is either abelian or an $n \times n$ matrix ring over some ring for $n > 1$, then all matrix rings over T are also finite Rickart C^* -algebras.*

A finite Rickart C^* -algebra is a subdirect product of its simple homomorphic images.

THEOREM 4 [4]. *In a finite Rickart C^* -algebra, the intersection of the maximal (two-sided) ideals is zero.*

Applying Theorem 4 and Corollary 2, we have

COROLLARY 5 [4]. *A finite Rickart C^* -algebra can be embedded in a finite AW^* -algebra.*

2. **Outline of the method.** In [2] we show that a finite Rickart C^* -algebra T has an \aleph_0 -continuous (unit) regular quotient ring R to which the involution on T can be lifted. Certain questions concerning finite Rickart C^* -algebras can then be 'lifted' to this quotient ring. This is implemented by a close study

AMS (MOS) subject classifications (1970). Primary 16A30, 47L05.

¹Research supported by NRC Grant A4540 and University of Waterloo Research Grant 131-7052.

Copyright © 1978, American Mathematical Society

of the relations between the lattices of projections of R and T . By considering the rank functions on R , and by an examination of the Grothendieck group of an \aleph_0 -continuous ring as a directed abelian group, we have the following intermediate result.

THEOREM 6 [4]. *In an \aleph_0 -continuous ring, the intersection of the maximal (two-sided) ideals is zero.*

REFERENCES

1. S. Berberian, *Baer *-rings*, Die Grundlehren der math. Wissenschaften, Band 195, Springer-Verlag, New York (1972).
2. D. Handelman, *Rickart C^* -algebras and their properties*, Advances in Math. (to appear).
3. ———, *Rickart C^* -algebras. II* (to appear).
4. D. Handelman, D. Higgs and J. Lawrence, *Directed abelian groups, \aleph_0 -continuous rings and Rickart C^* -algebras* (to appear).
5. I. Kaplansky, *Projections in Banach algebras*, Ann. of Math. (2) 53 (1951), 235–249.
6. J. Lawrence, *\aleph_0 complete Archimedean antilattices and prime Rickart C^* -algebras* (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY,
UTAH 84112

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO
N2L 3G1, ONTARIO, CANADA

Current address (D. Handelman): Department of Mathematics, University of Ottawa,
Ottawa, Canada