

## A NONSTANDARD IDEAL OF A RADICAL BANACH ALGEBRA OF POWER SERIES

BY MARC P. THOMAS

**1. Introduction.** We will be concerned here with a result about certain radical Banach algebras of power series. Let  $\mathbf{C}[[z]]$  denote the algebra of formal power series over the complex field  $\mathbf{C}$ . We say that a sequence of positive reals  $\{w(n)\}$  is an *algebra weight* provided the following hold:

$$(1.1) \quad w(0) = 1 \quad \text{and} \quad 0 < w(n) \leq 1 \quad (n \in \mathbf{Z}^+),$$

$$(1.2) \quad w(m+n) \leq w(m)w(n) \quad (m, n \in \mathbf{Z}^+),$$

$$(1.3) \quad \lim_{n \rightarrow \infty} w(n)^{1/n} = 0.$$

If these conditions hold it is routine to check that

$$l^1(w(n)) \equiv \left\{ y = \sum_{n=0}^{\infty} y(n)z^n : \sum_{n=0}^{\infty} |y(n)|w(n) < \infty \right\}$$

is both a subalgebra of  $\mathbf{C}[[z]]$  and a radical Banach algebra with identity adjoined. The norm and multiplication are defined in the natural way (see [3, 4, and 8]). There are obvious closed ideals in  $l^1(w(n))$ :

$$M(n) \equiv \left\{ \sum_{k=0}^{\infty} y(k)z^k \in l^1(w(n)) : y(0) = y(1) = \dots = y(n-1) = 0 \right\}$$

and, of course, the zero ideal. Such ideals are referred to as *standard* ideals. Any other closed ideals are denoted *nonstandard* ideals. It has been an open question for some time whether there exists any algebra weight  $\{w(n)\}$  such that  $l^1(w(n))$  contains a nonstandard ideal [6, p. 189], and the problem seems to go back to Šilov (a proposed solution appearing in the literature [6, p. 205] is in error). Interest has also been focused upon the quotient algebras  $(l^1(w(n))/I)$ , where  $I$  is assumed to be nonstandard, since these algebras are representative of all radical Banach algebras with power series generators [1]. Our result is the following:

**THEOREM.** *There exists an algebra weight  $\{w(n)\}$  for which the algebra  $l^1(w(n))$  contains a nonstandard ideal.*

---

Received by the editors February 14, 1983.

1980 *Mathematics Subject Classification.* Primary 46J20; Secondary 47A15.

© 1983 American Mathematical Society  
 0273-0979/83 \$1.00 + \$.25 per page

In the next section we will sketch a procedure for constructing such an algebra weight  $\{w(n)\}$ . We will define

$$x = \sum_{n=1}^{\infty} x(n)z^n \in l^1(w(n)), \quad x(1) \neq 0,$$

such that the closed ideal generated by  $x$  is nonstandard. If  $T$  denotes the operator of right translation on  $l^1(w(n))$ , to prove that this ideal is nonstandard is equivalent to showing that  $z \notin \overline{\text{span}}\{T^k x\}_{k=0}^{\infty}$ . We note in passing that by weighting the operator rather than the space, one can rephrase the problem in terms of weighted shift operators which are quasinilpotent and strictly cyclic (see [7]).

**2. Semimultiplicative weights.** Our weight will belong to a special class of weights which are called *semimultiplicative*. Let  $\{n(k)\}$  be a strictly increasing sequence of positive integers with  $n(1) = 1$ . Let the values  $w(n(k))$  be assigned subject to the conditions:

$$w(n(k+1))^{1/(n(k+1)+n(k))} \leq w(n(k))^{1/n(k)} \quad (k \in \mathbf{N})$$

and

$$(2.1) \quad \lim_{k \rightarrow \infty} w(n(k))^{1/n(k)} = 0.$$

The weight is defined for other values by setting  $w(0) = 1$  and setting

$$w(tn(k) + j) = w(n(k))^t w(j) \quad \text{if } tn(k) + j < n(k+1).$$

One says that  $\{w(n)\}$  is the semimultiplicative weight *generated* by  $\{w(n(k))\}$ . It can be shown that such a weight is an algebra weight [10, Lemmas 2.4 and 2.5]. We will need to specify further conditions on the rate of increase of the sequence  $\{n(k)\}$  and require the sequence  $\{w(n(k))\}$  to approach zero sufficiently rapidly. When  $\{w(n)\}$  has been specified, the element  $x$  is chosen to be the lacunary series  $x = \sum_{k=1}^{\infty} x(n(k))z^{n(k)}$ , where  $x(n(1)) = 1$  and  $|x(n(k))|w(n(k)) = 2^{-k+1}$  for  $k \in \mathbf{Z}^+$ ,  $k \geq 2$ . We remark in passing that weights similar to semimultiplicative weights have been shown to possess pathological multiplier algebras [2].

**3. Remarks on the proof.** Let  $A = l^1(w(n))$ . To show that the closed principal ideal  $Ax^-$  is nonstandard, it suffices to prove that there exists  $\sigma > 0$  such that

$$(3.1) \quad \left\| z - \sum_{n=0}^N a(n)T^n x \right\| \geq \sigma$$

for all choices of finite sequences  $\{a(n)\}_{n=0}^N$ . Let  $\{c(n)\}$  be the *associated sequence* [9, Definition 2.1], which is the sequence of coefficients for which one has

$$x * \sum_{n=0}^N c(n)z^n = z$$

in the algebra  $\mathbf{C}[[z]]$  of formal power series. The series  $\sum_{n=0}^{\infty} c(n)z^n$  will not, in general, lie in  $A$ , since many of its coefficients are too large. However, sums

of the form

$$x * \sum_{n=0}^N c(n)z^n = \sum_{n=0}^N c(n)T^n x$$

would be natural choices for approximating  $z$ . It has been shown that, for the type of weights we are considering, an estimate of the form (3.1) does hold for these sums [5]. The key to the proof that (3.1) holds for all finite sums lies in the fact that, if there is a sequence  $\{\sum_{n=0}^{N_p} a^{(p)}(n)T^n x\}$  which converges to  $z$ , then for each  $k$ , the sequence  $\{a^{(p)}(k)\}$  converges to  $c(k)$ . A contradiction is then reached through a complicated recursive argument, based upon the lacunarity of the element  $x = \sum_{k=1}^{\infty} x(n(k))z^{n(k)}$  and the rate of decrease of the terms  $w(n(k))$ , which shows that any element  $\sum_{n=0}^{\infty} a(n)T^n x$  of  $A$  which is sufficiently close in norm to  $z$  must necessarily have infinitely many nonzero terms.

#### REFERENCES

1. G. R. Allan, *Commutative Banach algebras with power series generators*, Radical Banach Algebras and Automatic Continuity (Proc. Long Beach, 1981), edited by J. M. Bachar et al, vol. 975, Springer-Verlag, 1983.
2. W. G. Bade, H. G. Dales and K. B. Laursen, *Multipliers of radical Banach algebras of power series*, Mem. Amer. Math. Soc. (to appear).
3. S. Grabiner, *Weighted shifts and Banach algebras of power series*, Amer. J. Math. **97** (1975), 16–42.
4. K. B. Laursen, *Ideal structure in radical sequence algebras*, Radical Banach Algebras and Automatic Continuity (Proc. Long Beach, 1981), edited by J. M. Bachar et al, vol. 975, Springer-Verlag, 1983.
5. J. P. McClure, *Nonstandard ideals and approximations in primary weighted  $l^1$ -algebras*, preprint.
6. N. K. Nikolskiĭ, *Selected problems of weighted approximation and spectral analysis*, Trudy Mat. Inst. Steklov **120** (1974). Izdat. "Nauka" Leningrad Otdel, Leningrad, 1974; English transl., Proc. Steklov Inst. Math., No. 120 (1974), Amer. Math. Soc., Providence, R.I., 1976.
7. A. L. Shields, *Weighted shift operators and analytic function theory*, Topics in Operator Theory (C. Pearcy, ed.), Math. Surveys, No. 13, Amer. Math. Soc., Providence, R.I., 1974.
8. M. P. Thomas, *Closed ideals and bi-orthogonal systems in radical Banach algebras of power series*, Proc. Edinburgh Math. Soc. **25** (1982), 245–257.
9. —, *Approximation in the radical algebra  $l^1(w_n)$  when  $\{w_n\}$  is star-shaped*, Radical Banach Algebras and Automatic Continuity (Proc. Long Beach, 1981), edited by J. M. Bachar et al, vol. 975, Springer-Verlag, 1983.
10. —, *A non-standard closed subalgebra of a radical Banach algebra of power series*, J. London Math. Soc. (to appear).

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE COLLEGE, BAKERSFIELD, CALIFORNIA 93309

