

RESEARCH REPORTS

BULLETIN (New Series) OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 34, Number 3, July 1997, Pages 293–312
S 0273-0979(97)00723-4

HOMOLOGY OF ALGEBRAIC VARIETIES: AN INTRODUCTION TO THE WORKS OF SUSLIN AND VOEVODSKY

MARC LEVINE

ABSTRACT. We give an overview of the ideas Suslin and Voevodsky have introduced in their works on algebraic cycles and their relation to the mod- n homology of algebraic varieties.

1. INTRODUCTION

The recent series of papers by Suslin, Voevodsky, Suslin-Voevodsky and Friedlander-Voevodsky ([49], [50], [51], [52], [53] and [20]) has developed a remarkable new viewpoint in the study of algebraic cycles. A new “topology” defined by Voevodsky, the qfh-topology, relates algebraic cycles to certain representing sheaves for this topology, and thus allows a systematic application of the powerful methods of sheaf-theory in areas which heretofore have been approached by essentially algebro-geometric or homotopy theoretic means. We hope to give here, not a full overview, but rather a sample of these new techniques and the results they have made possible, concentrating on the applications to the mod- n theory.

2. HISTORICAL BACKGROUND

Let X be a CW-complex. The topological K_0 (see [2], [3]) of X , $K_0^{\text{top}}(X)$, is defined as an abelian group via generators and relations: the generators are the isomorphism classes $[E]$ of complex vector bundles $E \rightarrow X$, with relations given by *stable equivalence*:

$$[E] \equiv [E'] \iff E \oplus e^n \cong E' \oplus e^n \text{ for some } n,$$

Received by the editors March 4, 1996, and, in revised form, January 7, 1997.
1991 *Mathematics Subject Classification*. Primary 19-02, 19E15, 14C25; Secondary 19E08, 19E20, 14F20, 18F10.

Key words and phrases. Motives, cycles.

Research supported by the NSF.

©1997 American Mathematical Society

where e^n is the trivial rank n bundle on X . For X a finite CW complex, this group is also given as the homotopy classes of maps of X to the classifying space $BGL_{\mathbb{C}}$ of the topological group

$$GL_{\mathbb{C}} := \varinjlim_N GL_{N, \mathbb{C}},$$

where $GL_{N, \mathbb{C}}$ is the topological group of invertible n by n matrices over \mathbb{C} . The higher topological K -theory of a finite CW-complex X can then be defined as the homotopy groups of a function space:

$$K_n^{\text{top}}(X) := \pi_n(\text{Hom}(X, BGL_{\mathbb{C}})).$$

The filtration of X via its k -skeleta, together with Bott periodicity:

$$K_n^{\text{top}}(X) \cong K_{n+2}^{\text{top}}(X),$$

gives rise to the *Atiyah-Hirzebruch spectral sequence* relating singular cohomology and topological K -theory; this spectral sequence degenerates rationally, giving the isomorphisms

$$\begin{aligned} K_{\text{even}}^{\text{top}}(X) \otimes \mathbb{Q} &\cong \bigoplus_n H^{2n}(X, \mathbb{Q}) \\ K_{\text{odd}}^{\text{top}}(X) \otimes \mathbb{Q} &\cong \bigoplus_n H^{2n+1}(X, \mathbb{Q}). \end{aligned}$$

The algebraic K_0 of an algebraic variety X is defined as the abelian group with generators the isomorphism classes $[E]$ of algebraic vector bundles E on X , with relations given by setting

$$[E] = [E'] + [E'']$$

if there exists an exact sequence

$$0 \rightarrow E' \rightarrow E \rightarrow E'' \rightarrow 0.$$

If X is affine, all such exact sequences split, and one gets the same group by imposing the relation of stable equivalence as in the topological case; but in general, stable equivalence is a weaker relation (they agree, however, in the topological setting). In fact, the algebraic K_0 was defined (by Grothendieck) *before* the topological case, but the higher K -theory in the topological setting was defined before the algebraic case.

Pursuing the analogy with the topological situation, Karoubi and Villamayor [31] gave a definition of higher algebraic K -theory of a ring R by means of the *discrete* group $GL(R)$, where the topology of $GL_{\mathbb{C}}$ is replaced by a certain simplicial structure. As this idea is central to our whole discussion, we give a description in a somewhat more general setting.

The algebraic version of homotopy is gotten by replacing the unit interval with the affine line \mathbb{A}^1 . Following this further, one considers the cosimplicial variety Δ^* , with n -cosimplices Δ^n given as the hyperplane in $n+1$ -space defined by the linear equation

$$\sum_{i=0}^n t_i = 1.$$

If one takes the t_i to be real numbers, with $0 \leq t_i \leq 1$, this is the usual n -simplex; as the usual expressions for the co-face and co-degeneracy maps between the n -simplex and the $n-1$ -simplex are linear functions of the t_i , one obtains the structure of a cosimplicial variety on Δ^* by using the same formulas as in the real case.

Starting with the map

$$\begin{aligned} \mathbb{A}^1 \times \mathbb{A}^1 &\rightarrow \mathbb{A}^1 \\ (s, t) &\mapsto st \end{aligned}$$

one can construct a map

$$(2.1) \quad H: \mathbb{A}^1 \times \Delta^* \rightarrow (\mathbb{A}^1 \times \Delta^*)^{[0,1]},$$

having the formal properties of a “homotopy” between the identity map on $\mathbb{A}^1 \times \Delta^*$ and the map

$$(x, t) \mapsto (0, t).$$

Now, suppose we have a functor (here **Sch** is the category of schemes)

$$F: \mathbf{Sch}^{\text{op}} \rightarrow \mathbf{Sets}.$$

We may form the new functor

$$(2.2) \quad F_h: \mathbf{Sch}^{\text{op}} \rightarrow \mathbf{Simplicial Sets}$$

by

$$F_h(X) = F(X \times \Delta^*).$$

The simplicial set $F_h(X)$ then satisfies the *homotopy property*: by applying F to (2.1), one shows that the natural map

$$F_h(X) \rightarrow F_h(X \times \mathbb{A}^1)$$

is a homotopy equivalence on the geometric realization. Similarly, if we have a functor

$$F: \mathbf{Sch}^{\text{op}} \rightarrow \mathbf{Ab},$$

we may form the new functor with values in chain complexes

$$(2.3) \quad F_{h\mathbb{Z}}: \mathbf{Sch}^{\text{op}} \rightarrow \mathbf{C}(\mathbf{Ab})$$

by taking the chain complex associated to the simplicial abelian group F_h . The functor $F_{h\mathbb{Z}}$ is also homotopy invariant: the natural map

$$F_{h\mathbb{Z}}(X) \rightarrow F_{h\mathbb{Z}}(X \times \mathbb{A}^1)$$

is a quasi-isomorphism.

Now back to Karoubi-Villamayor K -theory. Suppose X is an affine variety with ring of functions R . The group $\text{GL}_N(R)$ is the group of algebraic maps

$$X \rightarrow \text{GL}_N,$$

where GL_N is the open subscheme of affine N^2 space defined as the locus where the determinant function is non-zero. Replacing X with $X \times \Delta^*$ gives the simplicial ring $\Delta_*(R)$ with

$$\Delta_n(R) = R[t_0, \dots, t_n] / \left(\sum_i t_i - 1 \right)$$

$$\text{Spec } \Delta_*(R) = X \times \Delta^*,$$

and the simplicial group $\text{GL}_N(\Delta_*(R))$:

$$\text{GL}_N(\Delta_n(R)) = \text{Hom}_{\text{alg}}(X \times \Delta^n, \text{GL}_N).$$

One then takes the geometric realization $|\mathrm{GL}_N(\Delta_*(R))|$ of this simplicial set, passes to the limit over N , $|\mathrm{GL}(\Delta_*(R))|$, and defines the *Karoubi-Villamayor K-theory of R* by

$$KV_n(R) = \pi_{n-1}(|\mathrm{GL}(\Delta_*(R))|).$$

One can extend this definition to an arbitrary scheme by a sheafification process.

Quillen gave another definition of higher algebraic K -theory, first for rings by using his plus-construction [38], then for arbitrary exact categories via his categorical Q -construction [39]; this definition gained wide acceptance as the “correct” one in the general setting. It turns out that, for a regular scheme X (the algebro-geometric version of a manifold), the Karoubi-Villamayor K -theory agrees with the Quillen K -theory, although for singular schemes, the two definitions are not in general the same.

In contrast with the topological case, the algebraic version of singular cohomology, the so-called “motivic cohomology”, was not completely defined until some fifteen years after the definition of algebraic K -theory. The first step towards the full definition of motivic cohomology was, however, taken quite a bit earlier, arising in the 50’s with the construction of the Chow ring of algebraic cycles modulo rational equivalence (see e.g. [56], [45], [8], [9], [7], [12] and [25]). What emerged from these constructions was an algebraic homology-like theory built out of the free abelian group on the algebraic subvarieties of X , the *algebraic cycles on X* . Replacing the unit interval with the affine line as above leads to the relation of *rational equivalence*, an algebraic version of homology. More precisely, a cycle is *rationally equivalent to zero* if it is of the form

$$\mathrm{pr}_X(W \cdot (X \times 0) - W \cdot (X \times 1))$$

for W a cycle on $X \times \mathbb{A}^1$, where pr_X is the projection on X , and \cdot is the intersection product (one requires that W contains no component of the form $W_0 \times t$, so that the intersection product is defined). One then defines the *Chow group of X* as the group of algebraic cycles modulo rational equivalence.

For smooth quasi-projective varieties, this homology-like theory has a cohomological flavor as well: the intersection of subvarieties extends to the *intersection product* on the group of cycles mod rational equivalence, defining the *Chow ring*

$$\mathrm{CH}^*(X) := \bigoplus_q \mathrm{CH}^q(X),$$

with the grading by codimension. In addition, the Chow ring admits functorial pull-back maps

$$f^*: \mathrm{CH}^*(X) \rightarrow \mathrm{CH}^*(Y)$$

for arbitrary maps $f: Y \rightarrow X$ between smooth, quasi-projective varieties. Grothendieck’s theory of Chern classes with values in the Chow ring [25], together with the Grothendieck-Riemann-Roch theorem [7], gives the isomorphism

$$\sum_q c_q: K_0(X) \otimes \mathbb{Q} \cong \bigoplus_q \mathrm{CH}^q(X) \otimes \mathbb{Q}.$$

The major inadequacy of both the algebraic K_0 and the Chow ring is the lack of good *localization sequence*, relating the theory for X , an open subscheme U , and the closed complement $Z := X \setminus U$. Both theories have the beginning (or end, depending on your point of view) of such a sequence; Quillen’s uniform definition

of the higher K -groups, together with his localization theorem (see [39]), filled this gap on the K -theory side, giving the long exact sequence

$$\dots \rightarrow K_p(Z) \rightarrow K_p(X) \rightarrow K_p(U) \rightarrow K_{p-1}(Z) \rightarrow \dots \rightarrow K_0(X) \rightarrow K_0(U) \rightarrow 0$$

(this is for X and Z smooth; there is a similar sequence in general).

The first successful definition of motivic cohomology was supplied by Bloch in 1985, with his construction of the *higher* Chow groups [4]. The idea is to make sense of the algebraic cycles on the co-simplicial variety $X \times \Delta^*$. The technical problem here is that an arbitrary codimension q subvariety of $X \times \Delta^p$ may not intersect a face $X \times \Delta^{p'}$ in codimension q , and thus the intersection product with this face is not defined; Bloch solves this by considering the subgroup generated by subvarieties which *do* intersect all faces in the correct codimension. This gives the homological complex $\mathcal{Z}^q(X, *)$, with $\mathcal{Z}^q(X, p)$ being the subgroup of the cycles on $X \times \Delta^p$ just described, and differential given by the pull-back via the coboundary maps in $X \times \Delta^*$ (i.e., intersection with the codimension one faces $X \times \Delta^{p-1}$ of $X \times \Delta^p$). The higher Chow groups are then defined as the homology

$$\text{CH}^q(X, p) := H_p(\mathcal{Z}^q(X, *)).$$

This extends the definition of the Chow ring $\text{CH}^q(X)$, via the identity

$$\text{CH}^q(X) = \text{CH}^q(X, 0).$$

It wasn't until around 1992 that the localization property for $\text{CH}^q(X, *)$ was proved (see [5]), and this only for varieties over a field of characteristic 0. We do know, however (see e.g. [32]), that there is a close relation between the higher algebraic K -theory and the higher Chow groups of a variety X which is smooth over a field, so the higher Chow groups do seem to be the "correct" groups for motivic cohomology.

3. THE QUILLEN-LICHTENBAUM CONJECTURE

Let X be a variety defined over \mathbb{C} , $X(\mathbb{C})$ the topological space of solutions to the equations defining X . Passing from algebraic maps to continuous maps defines the comparison map from algebraic to topological K -theory

$$K_p^{\text{alg}}(X) \rightarrow K_p^{\text{top}}(X(\mathbb{C})).$$

In general, this map is far from being an isomorphism. However, the situation seems to get quite a bit simpler if one takes K -theory with mod n coefficients (defined via mod n homotopy groups of the appropriate spaces):

$$K_p^{\text{alg}}(X; \mathbb{Z}/n) \rightarrow K_p^{\text{top}}(X(\mathbb{C}); \mathbb{Z}/n).$$

The so-called *Quillen-Lichtenbaum conjecture* asserts (in one of its forms) that this map is an isomorphism for $p \geq 2 \dim_{\mathbb{C}}(X)$.

One can extend this to varieties over more general fields as follows. Grothendieck *et al.* [26] have defined the *étale cohomology*, $H_{\text{ét}}^p(X, \mathbb{Z}/n)$, of a scheme X , which (for n invertible on X) has many of the formal properties of the mod n singular cohomology of a space (see §5 for more information). In fact, the *comparison theorem* of Artin gives a natural isomorphism between the étale cohomology $H_{\text{ét}}^p(X, \mathbb{Z}/n)$ of a scheme X defined over \mathbb{C} , and the singular cohomology $H^p(X(\mathbb{C}), \mathbb{Z}/n)$ of the analytic space of solutions $X(\mathbb{C})$. Building on the methods of Grothendieck, Friedlander [14], [15], and Dwyer-Friedlander [13] constructed an algebro-geometric version of mod n topological K -theory, the *étale K -theory* $K_*^{\text{ét}}(X; \mathbb{Z}/n)$. There is an

Atiyah-Hirzebruch spectral sequence in the étale case as well, relating $H_{\text{ét}}^*(X, \mathbb{Z}/n)$ and $K_*^{\text{ét}}(X; \mathbb{Z}/n)$.

The Quillen-Lichtenbaum conjecture then asserts that the natural map of mod- n algebraic K -theory to mod- n étale K -theory of a variety X over a field k :

$$K_p^{\text{alg}}(X; \mathbb{Z}/n) \rightarrow K_p^{\text{ét}}(X; \mathbb{Z}/n)$$

is an isomorphism for $p \geq 2\dim_k(X) + \text{c.d.}(k)$, where $\text{c.d.}(k)$ is the mod n cohomological dimension of the Galois group of \bar{k}/k . Via the theory of Chern classes to étale cohomology, this can be interpreted as stating that the Chern class maps induce isomorphisms (for F a field)

$$\sum_{2q-p=m} c_{q,p}: K_m(F; \mathbb{Z}/n) \rightarrow \oplus_{2q-p=m} H_{\text{ét}}^p(F, \mathbb{Z}/n(q)); \quad 2q \geq p,$$

for n prime to “small primes” (depending on m) and prime to the characteristic of F .

We should mention here that the original conjecture of Quillen and Lichtenbaum (see e.g. [34]) concerned itself only with the K -groups and étale cohomology of *number rings*, and that the conjecture stated above is a slight modification of Conjecture 3.9 in [15].

Thus, one hopes for natural isomorphisms,

$$(3.1) \quad H_{\mu}^p(X, \mathbb{Z}/n(q)) \rightarrow H_{\text{ét}}^p(X, \mathbb{Z}/n(q)); \quad 2q \geq p, \quad q \geq \dim_k(X),$$

where $H_{\mu}^p(X, \mathbb{Z}/n(q))$ is the mod- n motivic cohomology of a smooth k -scheme X ; at least provisionally, this can be defined via Bloch’s cycle complex as

$$H_{\mu}^p(X, \mathbb{Z}/n(q)) = H_{2q-p}(\mathcal{Z}^q(X, *) \otimes \mathbb{Z}/n).$$

One main result of Suslin and Voevodsky ([49] and [48]) is that, for varieties over an algebraically closed field, such isomorphisms exist.

In fact, the major player in the proof is not Bloch’s cycle complex, but a homological version introduced by Suslin (in a 1988 lecture at Luminy). The *Suslin homology* of a scheme X is defined in terms of families of zero cycles on X , parametrized by the co-simplicial scheme Δ^* . There was no serious progress in the study or application of Suslin homology until Voevodsky introduced his qfh-topology and h-topology in 1992; this provided the needed breakthrough by allowing an interpretation of the mod n Suslin cohomology (the dual of Suslin homology) as a cohomology theory arising from sheaves on a Grothendieck site.

4. RELATIVE 0-CYCLES AND SUSLIN HOMOLOGY OF SCHEMES

A pointed topological space $(X, *)$ freely generates a commutative monoid with $*$ acting as identity, the *pointed infinite symmetric product* $Sp^{\infty}X$, whose points are the finite formal sums $\sum_i x_i$ with $x_i \in X$, modulo the relation

$$n \cdot * + \sum_i x_i \sim \sum_i x_i.$$

If $(X, *)$ is a connected CW complex, the theorem of Dold and Thom [11] shows that $Sp^{\infty}X$ represents the homology of X , via a natural isomorphism

$$\pi_n(Sp^{\infty}X) \cong H_n(X; \mathbb{Z}).$$

One can mimick this in the algebraic setting. Rather than attempting to deal with the infinite dimensional scheme $Sp^{\infty}X$, Suslin considers a functor which is

in principle the one represented by the group completion of $Sp^\infty X$. As in §2, one replaces the topology on $Sp^\infty X$ induced by that of X with the simplicial structure induced by Δ^* .

DEFINITION 4.1. Let X and S be k -schemes, with S smooth and irreducible. Define the group $C_0(S; X)$ to be the free abelian group on the subvarieties W of $X \times S$ such that the projection $p_2: W \rightarrow S$ is finite and surjective (recall that a map $f: Y \rightarrow Z$ is finite if f is proper and each fiber of f is a finite set). Set

$$C_n(S; X) = C_0(S \times \Delta^n; X).$$

The group $C_0(S; X)$ is covariantly functorial in X and contravariant in S ; thus, we may form the complex (see (2.2))

$$C_*(S; X) := C_0(S \times \Delta^*; X) = C_0(S; X)_{h\mathbb{Z}},$$

and the homology

$$H_p^{\text{sing}}(S; X) := H_p(C_*(S; X)).$$

The groups $H_p^{\text{sing}}(S; X)$ are thus covariant in X and contravariant in S .

EXAMPLE 4.2. Take $S = \text{Spec } k$. Then $H_p^{\text{sing}}(k; X)$ is the Suslin homology of X . Sending X to $H_p^{\text{sing}}(X; \mathbb{Z}) := H_p^{\text{sing}}(k; X)$ defines the functor

$$H_p^{\text{sing}}(-; \mathbb{Z}): \mathbf{Sch}/k \rightarrow \mathbf{Ab}.$$

Let Y be a smooth irreducible k -scheme, and define $C_0(X)(Y)$ by

$$(4.1) \quad C_0(X)(Y) := C_0(Y; X).$$

This defines the (contravariant) functor $C_0(X)$ on the category \mathbf{Sm}/k of smooth, finite-type k -schemes.

5. GROTHENDIECK TOPOLOGIES

Grothendieck topologies play a central role in the arguments of Suslin-Voevodsky; we give here a brief overview. For more details see, e.g., [1], [35], [10], and [26].

Grothendieck introduced the notion of a Grothendieck topology, and the associated category of sheaves for the topology, in the construction of étale cohomology, the algebro-geometric replacement of singular cohomology. To form a topology on a set X , one selects a collection of subsets of X , the open subsets for the topology, subject to certain axioms. Grothendieck extended this notion by considering the inclusion of an open subset as a special case of a morphism in a category and considering, for a category \mathcal{C} , families of morphisms in \mathcal{C}

$$f_\alpha: U_\alpha \rightarrow X,$$

satisfying axioms which generalize the notion of an open cover.

More specifically, for each object U of \mathcal{C} , one must define when a family of morphisms

$$\{U_\alpha \rightarrow U \mid \alpha \in A\}$$

in \mathcal{C} is a cover of U ; one requires

1. *stability under base-change*: if

$$\{U_\alpha \rightarrow U \mid \alpha \in A\}$$

is a cover of U , and $V \rightarrow U$ is a morphism in \mathcal{C} , then the fiber product $U_\alpha \times_U V$ exists for each α , and the family

$$\{p_2: U_\alpha \times_U V \rightarrow V \mid \alpha \in A\}$$

is a cover of V .

2. *stability under composition*: if

$$\{U_\alpha \xrightarrow{f_\alpha} U \mid \alpha \in A\}$$

is a cover of U , and if

$$\{V_\beta \xrightarrow{g_{\alpha\beta}} U_\alpha \mid \beta \in B_\alpha\}$$

is a cover of U_α for each α , then

$$\{V_\beta \xrightarrow{f_\alpha \circ g_{\alpha\beta}} U \mid \beta \in B_\alpha, \alpha \in A\}$$

is a cover of U .

3. each isomorphism

$$U' \rightarrow U$$

forms a cover of U .

A category together with a Grothendieck topology is called a *Grothendieck site*.

We have actually defined above the notion of a Grothendieck *pre-topology* on \mathcal{C} ; this suffices to define the primary objects of interest: presheaves and sheaves. A Grothendieck pre-topology generates a Grothendieck topology, just as a basis for a topology generates a topology, so we will freely confuse the two notions in what follows.

One can view a topology in the usual sense on a set X as a Grothendieck topology, by defining \mathcal{C} to be the category formed by the inclusion maps $V \rightarrow U$ among the open subsets of X , and defining the covers of $U \subset X$ to be collections

$$\{U_\alpha \rightarrow U\},$$

with

$$\cup_\alpha U_\alpha = U,$$

i.e., open covers of U .

Conversely, if one has a Grothendieck topology on a category \mathcal{C} , one can consider the morphisms

$$f: V \rightarrow U$$

which occur in some cover $\{f_\alpha: U_\alpha \rightarrow U\}$ as an “open subset” of U . If the category \mathcal{C} has a final object X , and if each morphism $V \rightarrow U$ occurs in some cover of U , then one may consider the Grothendieck topology on \mathcal{C} as a “Grothendieck topology on X ”, and the objects of \mathcal{C} as the opens for the Grothendieck topology.

In the general case, a Grothendieck topology \mathfrak{T} on a category \mathcal{C} induces the Grothendieck topology \mathfrak{T}_X on each object X of \mathcal{C} by forming the category \mathcal{C}_X of maps $U \rightarrow X$ which occur in a cover of X , where a map $(U \rightarrow X) \rightarrow (V \rightarrow X)$ is a commutative triangle

$$\begin{array}{ccc} U & \rightarrow & V \\ & \searrow & \swarrow \\ & X & \end{array}$$

and using the covers in \mathfrak{T} to define the covers in \mathfrak{T}_X . The operation of restricting to an object of \mathcal{C} is like taking the induced topology on a subset of a topological

space. One can consider a Grothendieck topology on a category \mathcal{C} as being defined by giving a Grothendieck topology on X for each object X of \mathcal{C} , with this assignment being natural in X .

Once one has a Grothendieck topology \mathfrak{T} on a category \mathcal{C} , one defines a presheaf F (say of abelian groups) for the topology as a contravariant functor from \mathcal{C} to \mathbf{Ab} . A sheaf is a presheaf F that satisfies the sheaf axiom: if

$$\{i_\alpha: U_\alpha \rightarrow U\}$$

is a cover of U , then the sequence

$$0 \rightarrow F(U) \xrightarrow{\prod_\alpha F(i_\alpha)} \prod_\alpha F(U_\alpha) \xrightarrow{F(p_2)-F(p_1)} \prod_{\alpha,\beta} F(U_\alpha \times_V U_\beta)$$

is exact. The category $\text{Sh}_\mathfrak{T}$ of sheaves of abelian groups for the topology \mathfrak{T} is then an abelian category, so one can perform the usual operations of homological algebra in this setting. If X is in \mathcal{C} , using the topology \mathfrak{T}_X on \mathcal{C}_X , one has the notions of a pre-sheaf or a sheaf on X , giving the category of sheaves on X for the topology \mathfrak{T} , $\text{Sh}_\mathfrak{T}(X)$.

There is also the operation of sheafifying a presheaf, so one has in particular the “constant” sheaf A_X on X for each abelian group A . As one has the canonical isomorphism

$$\text{Hom}_{\text{Sh}_\mathfrak{T}(X)}(\mathbb{Z}_X, \mathcal{F}) \cong \mathcal{F}(X),$$

the groups $\text{Ext}^p(\mathbb{Z}_X, \mathcal{F})$ are the higher derived functors of the global sections functor; one can therefore define the cohomology of a sheaf \mathcal{F} on X for the topology \mathfrak{T} by

$$(5.1) \quad H_\mathfrak{T}^p(X; \mathcal{F}) = \text{Ext}_{\text{Sh}_\mathfrak{T}(X)}^p(\mathbb{Z}_X, \mathcal{F}).$$

EXAMPLE 5.1.

1. The *Zariski* topology on a scheme X is a topology in the classical sense, with opens being complements of algebraic subsets F of X . One makes the Zariski topology into a Grothendieck topology as described above.
2. For X a smooth scheme of finite type over an algebraically closed field k , an étale map is a morphism $i: U \rightarrow X$ whose differential di_u is an isomorphism for all k points $u \in U$. For example, if X is a smooth variety over \mathbb{C} , and $f: Y \rightarrow X$ is *proper*, then f is étale if and only if the map of complex manifolds $f(\mathbb{C}): Y(\mathbb{C}) \rightarrow X(\mathbb{C})$ is a covering space (necessarily finite). More generally, if X is still over \mathbb{C} , but not necessarily smooth, then a map $f: Y \rightarrow X$ is étale if and only if, for each point y of Y , there are neighborhoods U of $y \in Y(\mathbb{C})$ and V of $f(y) \in X(\mathbb{C})$ (in the \mathbb{C} -topology) such that f gives an isomorphism

$$f(\mathbb{C}): U \rightarrow V.$$

We omit the general definition of an étale morphism.

The étale topology on a scheme X is given by taking the category of “opens of X ” to be the étale maps $U \rightarrow X$ (of finite type), and saying that a cover of $U \rightarrow X$ is a collection of étale maps

$$\{f_\alpha: U_\alpha \rightarrow U\}$$

such that

$$U = \cup_\alpha f_\alpha(U_\alpha).$$

Using the methods discussed above, one has the étale cohomology of a scheme X with coefficients in an étale sheaf \mathcal{F} , $H_{\text{ét}}^*(X, \mathcal{F})$.

REMARK 5.2. Let x be a point of a variety X . The *henselization* X_x^h of X at x plays the role, for the étale topology, of a small neighborhood of x in X in the classical topology. The henselization is gotten by taking the inverse limit over the the collection of pointed étale map

$$(Y, y) \rightarrow (X, x).$$

For instance, if the base field k is algebraically closed, and x is a smooth k -point of a smooth k -variety X of dimension d , then X_x^h is isomorphic to the henselization $(\mathbb{A}^d)_0^h$ of affine d -space at the origin.

The examples in 5.1 describe the so-called *small* site: giving a Grothendieck topology for a single scheme X ; giving a Grothendieck topology on the category \mathbf{Sch}/k of schemes over a fixed base field k defines what is known as a *big* site.

For example, the étale topology on a scheme X is natural in X : if $U \rightarrow X$ is étale, and $f: Y \rightarrow X$ is a morphism of schemes, then $U \times_X Y \rightarrow Y$ is étale. This gives us the big étale site on \mathbf{Sch}/k . The big Zariski site is defined similarly.

Now, suppose we have a sheaf \mathcal{F} on \mathbf{Sch}/k for some topology \mathfrak{T} . For a k -scheme X , one can restrict \mathcal{F} to X to give the sheaf \mathcal{F}_X for the topology \mathfrak{T}_X , and the cohomology $H^*(X, \mathcal{F}_X)$, defined as in (5.1) as the Ext groups

$$H^*(X, \mathcal{F}_X) = \text{Ext}_{\text{Sh}_{\mathfrak{T}_X}(X)}^*(\mathbb{Z}_X, \mathcal{F}_X).$$

One can also define the cohomology on X entirely in the category $\text{Sh}_{\mathfrak{T}}/k$ of sheaves on \mathbf{Sch}/k in the following way: Let

$$\text{Hom}_k(-, X): \mathbf{Sch}/k^{\text{op}} \rightarrow \mathbf{Sets}$$

be the functor represented by X :

$$Y \mapsto \text{Hom}_k(Y, X).$$

Form the free abelian group on $\text{Hom}_k(-, X)$, giving the presheaf

$$\mathbb{Z}[\text{Hom}_k(-, X)]: \mathbf{Sch}/k^{\text{op}} \rightarrow \mathbf{Ab}.$$

Let $\mathbb{Z}_{\mathfrak{T}}(X)$ denote the sheafification of $\mathbb{Z}[\text{Hom}_k(-, X)]$ for the topology \mathfrak{T} . It follows from the Yoneda lemma that there is a natural isomorphism

$$(5.2) \quad H_{\mathfrak{T}_X}^*(X, \mathcal{F}_X) = \text{Ext}_{\text{Sh}_{\mathfrak{T}_X}(X)}^*(\mathbb{Z}_X, \mathcal{F}_X) \cong \text{Ext}_{\text{Sh}_{\mathfrak{T}}/k}^*(\mathbb{Z}_{\mathfrak{T}}(X), \mathcal{F}),$$

which is the interpretation of sheaf cohomology on X we wanted.

We shall see in the next section that two new topologies, both finer than the étale topology, provide via (5.2) the link between mod n étale cohomology and Suslin homology.

6. THE h-TOPOLOGY AND THE qfh-TOPOLOGY

Voevodsky's qfh-topology allows one to use the machinery of sheaf cohomology in the study of algebraic cycles; the h-topology is perhaps a more natural construction, but the resulting sheaf cohomology is not so obviously related to cycles. As we shall later see, the mod- n cohomologies in the h- and qfh-topologies agree, so the two points of view complement each other.

We fix a base field k , and write \mathbf{Sch}/k for the category of schemes of finite type over k .

DEFINITION 6.1.

1. Let $f: X \rightarrow Y$ be a map of k -schemes. f is a *topological epimorphism* if f is surjective on points and if Y has the quotient topology; i.e., a subset U of Y is open if and only if $f^{-1}(U)$ is open in X . f is a *universal topological epimorphism* if, for each map of schemes $Z \rightarrow Y$, the projection

$$Z \times_Y X \rightarrow Z$$

is a topological epimorphism.

2. The *h-topology* on \mathbf{Sch}/k is the Grothendieck topology for which an h-cover of Y is a universal topological epimorphism $X \rightarrow Y$.
3. The *qfh-topology* on \mathbf{Sch}/k is the Grothendieck topology for which a qfh-cover of Y is a universal topological epimorphism $X \rightarrow Y$ which is quasi-finite over Y (the inverse image of each point of Y is a finite set).

The following structure theorem gives a more concrete idea of the h-topology and qfh-topology.

Theorem 6.2. *i) Let $V \rightarrow Y$ be an h-cover. There is a refinement of V ,*

$$\begin{array}{ccc} U & \rightarrow & V \\ & \searrow & \swarrow \\ & Y & \end{array}$$

and a factorization of $U \rightarrow Y$ as

$$(6.1) \quad U = \coprod_i U_i \xrightarrow{j} X \xrightarrow{p} Z \xrightarrow{q} Y$$

where p is a finite morphism, q is the blow-up of a closed subscheme of Y , and j is a Zariski open cover. Conversely, each morphism $U \rightarrow Y$ which factors as in (6.1) is an h-cover.

ii) Let $V \rightarrow Y$ be a qfh-cover. There is a refinement of V ,

$$\begin{array}{ccc} U & \rightarrow & V \\ & \searrow & \swarrow \\ & Y & \end{array}$$

and a factorization of $U \rightarrow Y$ as

$$(6.2) \quad U = \coprod_i U_i \xrightarrow{j} X \xrightarrow{p} Y$$

where p is a finite morphism and j is a Zariski open cover. Conversely, each morphism $U \rightarrow Y$ which factors as in (6.2) is a qfh-cover.

REMARK 6.3. i) Clearly the h-topology is finer than the qfh-topology. The qfh-topology is finer than the étale topology.

ii) If the characteristic of k is zero, one may use Hironaka's resolution of singularities to show that each h cover of a k -scheme Y has a refinement $U \rightarrow Y$ with U smooth over k . In characteristic $p > 0$, the recent work of de Jong [29] gives the same result. This is certainly *not* the case for qfh-covers, even of a smooth k -scheme. This points out an important technical advantage of the h-topology over the qfh-topology: every k -scheme is locally smooth in the h-topology, while only 0 and 1-dimensional k -schemes are locally smooth in the qfh-topology.

7. FAMILIES OF 0-CYCLES AND THE qfh-TOPOLOGY

We have seen in (5.2) that the sheafification $\mathbb{Z}_{\mathfrak{X}}(X)$ of the abelian group generated by the representing presheaf $\text{Hom}(-, X)$ plays a central role in understanding sheaf cohomology. We have the “family of 0-cycles on X ” functor of example 4.1:

$$C_0(X): \mathbf{Sm}/k^{\text{op}} \rightarrow \mathbf{Ab}.$$

The qfh-topology links sheaf cohomology with algebraic cycles via

Theorem 7.1. *Let p be the exponential characteristic of the base field k ($p = \text{char}(k)$ if $\text{char}(k) > 0$, $p = 1$ if $\text{char}(k) = 0$). The functor $C_0(X)[1/p]$ on \mathbf{Sm}/k extends uniquely to a qfh-sheaf (denoted $c_0(X)$) on \mathbf{Sch}/k . Moreover, $c_0(X)$ is naturally isomorphic to the “representing” qfh-sheaf $\mathbb{Z}_{\text{qfh}}(X)[1/p]$.*

The map relating $\mathbb{Z}_{\text{qfh}}(X)[1/p]$ and $C_0(X)[1/p]$ is gotten by sending a morphism

$$f: Y \rightarrow X$$

to the transpose of the graph in $X \times Y$. The proof of the first part of Theorem 7.1 can be divided into two steps: the first step consists in extending $C_0(X)[1/p]$ to normal schemes. This is accomplished in [48] by Galois-theoretic methods, and in [50] by a type of limit process, reminiscent of the method of Weil [56] for defining intersection multiplicities. The extension from normal schemes to arbitrary schemes of finite type is then a fairly formal process, using the fact that a reduced scheme of finite type is a direct limit of normal schemes.

Another crucial property of qfh-sheaves is that they admit *transfers*. Recall that a map of schemes $f: X \rightarrow Y$ is *finite* if f is proper and quasi-finite ($f^{-1}(y)$ is a finite set for each $y \in Y$). For instance, a finite extension of field $K \rightarrow L$ defines a finite morphism $\text{Spec } L \rightarrow \text{Spec } K$. If P is a presheaf on \mathbf{Sch}/k , a transfer on P is gotten by giving a map

$$\text{Tr}_{X/S} = \text{Tr}_p: P(X) \rightarrow P(S),$$

for each finite morphism $p: X \rightarrow S$ with X and S reduced and irreducible, and S smooth. These maps should be compatible with pull-back in cartesian squares in a sense which we will leave vague. Now, in the qfh-topology, each finite map $X \rightarrow S$ is an open cover, and the sheaf axiom shows that, at least if $X \rightarrow S$ is Galois with group of automorphisms G , taking the trace

$$\begin{aligned} \text{Tr}: P(X) &\rightarrow P(X) \\ x &\mapsto \sum_{g \in G} x^g \end{aligned}$$

defines a map $P(X) \rightarrow P(S)$. The general case follows with a bit of additional work.

8. THE h-TOPOLOGY AND THE RIGIDITY THEOREM

As the h-topology is finer than the qfh-topology, an h-sheaf also has transfers. The crucial property of sheaves for the h-topology is expressed by the *rigidity theorem*.

The original precursor of the rigidity theorem may be found in Roitman’s work on zero-cycles. The work [42] considers the torsion subgroup of the group of zero-cycles

modulo rational equivalence on a smooth projective variety X over an algebraically closed field k , and the behavior of this group under the *Albanese mapping*:

$$\alpha_X : \text{CH}_0(X) \rightarrow \text{Alb}(X).$$

Here $\text{Alb}(X)$ is the Albanese variety of X , a finite dimensional projective algebraic group. Roitman [41], extending the work of Mumford [37] for surfaces, had shown that the map α_X has a “huge” kernel, assuming that X admits a global algebraic p -form with p at least 2 (and that $k = \mathbb{C}$). In contrast with this, Roitman shows in [42] that α_X induces an isomorphism on the torsion subgroups (prime to the characteristic of k).

The main point in his argument is the rigidity result: Suppose we have a family of zero-cycles on X , parametrized by an algebraic curve C , which are n -torsion mod rational equivalence, with n prime to the characteristic of k . Then the family is *constant* mod rational equivalence. The argument is quite simple: Let $C(k)$ be the set of k -points of C and let $\text{CH}_0(X)_n$ be the n -torsion subgroup of $\text{CH}_0(X)$. Sending $x \in C(k)$ to the corresponding 0-cycle gives the map

$$\rho : C(k) \rightarrow \text{CH}_0(X)_n;$$

one easily shows that ρ extends to a group homomorphism from the k -points of the Jacobian variety

$$J(\rho) : J(C)(k) \rightarrow \text{CH}_0(X)_n.$$

Since k is algebraically closed, the group $J(C)(k)$ is n -divisible; hence $J(\rho)$ is the zero map. Since $J(C)(k)$ is generated by the differences $x - y$, where x and y are k -points of C , this shows that $\rho(x) = \rho(y)$ for all $x, y \in C(k)$. The fundamental rigidity result, Theorem 8.2 below, is essentially a formal version of this result; various other extensions, with application to K -theory, had been given by Suslin [46], [47], Gillet-Thomason [27] and Gabber [22].

DEFINITION 8.1. A presheaf \mathcal{F} on \mathbf{Sch}/k is called *homotopy invariant* if the map

$$p_1^* : \mathcal{F}(X) \rightarrow \mathcal{F}(X \times \mathbb{A}^1)$$

is an isomorphism for all X .

Theorem 8.2. *Let \mathcal{F} be presheaf on \mathbf{Sch}/k which*

1. *is homotopy invariant,*
2. *has transfers,*
3. *is n -torsion: $n\mathcal{F}(T) = 0$ for all k -schemes T , where n is prime to the exponential characteristic of k .*

Let x be a smooth point on a k -variety X , X_x^h the henselization of X at x (see Remark 5.2) and let

$$i_x : \text{Spec } k \rightarrow X_x^h$$

be the inclusion. Then the map

$$i_0^* : \mathcal{F}(X_x^h) \rightarrow \mathcal{F}(\text{Spec } k)$$

is an isomorphism.

It follows from Theorem 8.2 and Remark 6.3 that the h-sheaf associated to a homotopy invariant presheaf with transfers \mathcal{F} and the constant h-sheaf with value $\mathcal{F}(\text{Spec } k)$ have the same n -torsion and n -cotorsion, assuming that k is algebraically closed. This implies the fundamental cohomological rigidity theorem

Theorem 8.3. *Let k be an algebraically closed field, and let \mathcal{F} be a homotopy invariant presheaf on \mathbf{Sch}/k which admits transfers. Denote by $\tilde{\mathcal{F}}_h$ the sheafification of \mathcal{F} for the h -topology. Then for all n prime to $\text{char}(k)$, there is a canonical isomorphism*

$$\text{Ext}_h^*(\tilde{\mathcal{F}}_h, \mathbb{Z}/n) \rightarrow \text{Ext}_{\mathbf{Ab}}^*(\mathcal{F}(\text{Spec } k), \mathbb{Z}/n).$$

REMARK 8.4. The proof in [49] assumes characteristic zero to allow the use of resolution of singularities, which enables one to show that each k -scheme of finite type admits an h -cover which is smooth over k ; the result of de Jong noted in Remark 6.3 allows the proof to go through in arbitrary characteristic.

9. CHANGE OF TOPOLOGY

Suppose we have two Grothendieck topologies on \mathbf{Sch}/k , \mathfrak{T} and \mathfrak{T}' , such that \mathfrak{T} is finer than \mathfrak{T}' ; i.e., each cover in \mathfrak{T}' is a cover in \mathfrak{T} .

We may take a sheaf for the topology \mathfrak{T}' and sheafify it for the topology \mathfrak{T} , defining the functor

$$i^* : \text{Sh}_{\mathfrak{T}'} \rightarrow \text{Sh}_{\mathfrak{T}}.$$

For example, we may compare the h -, qfh - and étale topologies,

$$\mathbf{Sch}/k_h \xrightarrow{\alpha} \mathbf{Sch}/k_{qfh} \xrightarrow{\beta} \mathbf{Sch}/k_{\text{ét}}.$$

The main comparison result is

Theorem 9.1. *Let \mathcal{F} be an étale sheaf and let \mathcal{G} be a qfh -sheaf on \mathbf{Sch}/k . Then*

$$\begin{aligned} \text{Ext}_{\text{ét}}^*(\mathcal{F}, \mathbb{Z}/n) &= \text{Ext}_{qfh}^*(\beta^* \mathcal{F}, \mathbb{Z}/n) \\ \text{Ext}_{qfh}^*(\mathcal{G}, \mathbb{Z}/n) &= \text{Ext}_h^*(\alpha^* \mathcal{G}, \mathbb{Z}/n). \end{aligned}$$

The proof is fairly straightforward: the comparison of the étale and qfh -cohomology relies on the structure of qfh -covers (Theorem 6.2(ii)) and some elementary facts on finite covers of strictly hensel schemes. One then compares the h -cohomology with étale cohomology, using the structure of h -covers (Theorem 6.2(i)) and the Künneth formula for étale cohomology.

10. SUSLIN COHOMOLOGY OF qfh -SHEAVES

We now have all the main ingredients needed to relate Suslin homology and étale cohomology: the representability theorem (Theorem 7.1) and the rigidity theorem (Theorem 8.3). Helped along by the comparison theorem (Theorem 9.1), a straightforward spectral sequence argument completes the proof. It is technically useful to work in a somewhat more general setting, working with an arbitrary presheaf rather than the constant presheaf \mathbb{Z} .

Let \mathcal{F} be a presheaf on \mathbf{Sch}/k , and let X be in \mathbf{Sch}/k . Applying \mathcal{F} to the cosimplicial scheme $X \times \Delta^*$ (as in §2) gives the simplicial abelian group $\mathcal{F}(X \times \Delta^*)$, and the associated (homological) complex $\mathcal{F}_*(X)$:

$$\mathcal{F}_n(X) = \mathcal{F}(X \times \Delta^n).$$

This forms the presheaves \mathcal{F}_n on \mathbf{Sch}/k , and the complex of presheaves \mathcal{F}_* . For an abelian group A , set

$$\begin{aligned} C_*(\mathcal{F}) &:= \mathcal{F}_*(\text{Spec } k) = \mathcal{F}(\Delta^*) \\ H_*^{\text{sing}}(\mathcal{F}, A) &:= H_*(C_*(\mathcal{F}) \otimes^L A) \\ H_{\text{sing}}^*(\mathcal{F}, A) &:= H^*(\text{RHom}(C_*(\mathcal{F}), A)). \end{aligned}$$

Let $\tilde{\mathcal{F}}_n$ denote the qfh-sheaf associated to \mathcal{F}_n and $\tilde{\mathcal{F}}_{\text{qfh}}$ the qfh-sheaf associated to \mathcal{F} . We also write $C_*(\mathcal{F})$ for the complex of constant sheaves.

Theorem 10.1. *Let k be an algebraically closed field, and let $n > 0$ be prime to $\text{char}(k)$. Let \mathcal{F} be a presheaf on \mathbf{Sch}/k which admits transfers. Then there is a canonical isomorphism*

$$H_{\text{sing}}^*(\mathcal{F}, \mathbb{Z}/n) \cong \text{Ext}_{\text{qfh}}^*(\tilde{\mathcal{F}}_{\text{qfh}}, \mathbb{Z}/n).$$

Using Theorem 9.1, and the fact that qfh-sheaves admit transfers (§7), Theorem 10.1 implies

Corollary 10.2. *Let \mathcal{F} be a qfh-sheaf on \mathbf{Sch}/k , with k algebraically closed. Then there is a canonical isomorphism*

$$H_{\text{sing}}^*(\mathcal{F}, \mathbb{Z}/n) \cong \text{Ext}_{\text{ét}}^*(\mathcal{F}, \mathbb{Z}/n)$$

for all n prime to $\text{char}(k)$.

Taking $\mathcal{F} = \mathbb{Z}_{\text{qfh}}(X)$ and using Theorem 7.1 and formula (5.2) gives the main result:

Corollary 10.3. *Let X be a scheme of finite type over an algebraically closed field k , and let n be prime to $\text{char}(k)$. Then there is a natural isomorphism*

$$H_{\text{sing}}^*(X, \mathbb{Z}/n) \cong H_{\text{ét}}^*(X, \mathbb{Z}/n).$$

The proof of Theorem 10.1 starts by considering the two spectral sequences

$$I_1^{p,q} = \text{Ext}_{\text{qfh}}^p((\tilde{\mathcal{F}}_q)_{\text{qfh}}, \mathbb{Z}/n) \implies \text{Ext}^{p-q}((\tilde{\mathcal{F}}_*)_{\text{qfh}}, \mathbb{Z}/n),$$

$$II_2^{p,q} = \text{Ext}_{\text{qfh}}^p(H_q((\tilde{\mathcal{F}}_*)_{\text{qfh}}), \mathbb{Z}/n) \implies \text{Ext}^{p-q}((\tilde{\mathcal{F}}_*)_{\text{qfh}}, \mathbb{Z}/n).$$

The comparison of qfh-cohomology and étale cohomology (Theorem 9.1), and the homotopy invariance of étale cohomology

$$H_{\text{ét}}^*(X, \mathbb{Z}/n) \cong H_{\text{ét}}^*(X \times \mathbb{A}^1, \mathbb{Z}/n)$$

lead to a proof that the projections $X \times \Delta^p \rightarrow X$ induce isomorphisms

$$\text{Ext}_{\text{qfh}}^p(\tilde{\mathcal{F}}_{\text{qfh}}, \mathbb{Z}/n) \cong \text{Ext}_{\text{qfh}}^p((\tilde{\mathcal{F}}_q)_{\text{qfh}}, \mathbb{Z}/n).$$

This implies the degeneration of the spectral sequence I at E_1 , giving the isomorphism

$$\text{Ext}_{\text{qfh}}^*(\tilde{\mathcal{F}}_{\text{qfh}}, \mathbb{Z}/n) \cong \text{Ext}_{\text{qfh}}^*((\tilde{\mathcal{F}}_*)_{\text{qfh}}, \mathbb{Z}/n).$$

By the homotopy invariance of the functors (2.3), the homology presheaf

$$H_q((\tilde{\mathcal{F}}_*)_{\text{qfh}})$$

forms a homotopy invariant presheaf on \mathbf{Sch}/k ; the transfers for the qfh-sheaves $(\tilde{\mathcal{F}}_q)_{\text{qfh}}$ give transfers for $H_q((\tilde{\mathcal{F}}_*)_{\text{qfh}})$. This allows one to use the comparison with h-cohomology (Theorem 9.1) and the rigidity theorem (Theorem 8.3) to compare the spectral sequence II with the spectral sequence

$$III_2^{p,q} = \text{Ext}_{\mathbf{Ab}}^p(H_q(\mathcal{F}_*(\text{Spec } k)), \mathbb{Z}/n) \implies \text{Ext}_{\mathbf{Ab}}^{p-q}(\mathcal{F}_*(\text{Spec } k), \mathbb{Z}/n),$$

giving the isomorphism

$$\text{Ext}_{\mathbf{Ab}}^*(\mathcal{F}_*(\text{Spec } k), \mathbb{Z}/n) \cong \text{Ext}_{\text{qfh}}^*((\tilde{\mathcal{F}}_*)_{\text{qfh}}, \mathbb{Z}/n)$$

and completing the proof.

11. SUSLIN HOMOLOGY AND BLOCH'S HIGHER CHOW GROUPS

To complete the discussion, it is necessary to relate Suslin homology and Bloch's higher Chow groups.

For X in \mathbf{Sch}/k , and Y normal, let $Z_0(X)(Y)$ denote the free abelian group on the subvarieties W of $X \times Y$ which are *quasi-finite* over Y . As in §4, sending a smooth Y to $Z_0(X)(Y)$ defines a contravariant functor on \mathbf{Sm}/k . Taking Y to be the cosimplicial scheme Δ^* and taking the associated homological complex define the complex $Z_*(X)$, with

$$Z_n(X) = Z_0(X)(\Delta^n).$$

Suppose X has pure dimension d over k . As a subvariety W of $X \times \Delta^p$ which is quasi-finite over Δ^p obviously has the proper intersection with all faces, there is the inclusion of complexes

$$(11.1) \quad Z_*(X) \rightarrow \mathcal{Z}^d(X, *).$$

Suslin [48] has shown

Theorem 11.1. *Suppose that X is affine. Then the inclusion (11.1) is a quasi-isomorphism.*

Let $\mathrm{CH}^q(X, p; \mathbb{Z}/n)$ denote the homology

$$\mathrm{CH}^q(X, p; \mathbb{Z}/n) = H_p(\mathcal{Z}^q(X, *) \otimes \mathbb{Z}/n) = H_p((\mathcal{Z}^q(X, *) \otimes^L \mathbb{Z}/n)).$$

There is an analog of Theorem 7.1:

Theorem 11.2. *There is a unique extension of the functor $Z_0(X)[1/p]$ on \mathbf{Sm}/k to a qfh-sheaf $z_0(X)$ on \mathbf{Sch}/k .*

Applying Theorem 10.1 gives

Proposition 11.3. *Let n be prime to $\mathrm{char}(k)$ and suppose X is affine of dimension d over k . There is a natural isomorphism*

$$\mathrm{CH}^d(X, *; \mathbb{Z}/n)^\vee = H_*^{\mathrm{sing}}(z_0(X), \mathbb{Z}/n)^\vee \cong \mathrm{Ext}_{\mathrm{qfh}}^*(z_0(X), \mathbb{Z}/n).$$

Here M^\vee is the dual

$$M^\vee := \mathrm{Hom}_{\mathbb{Z}/n}(M, \mathbb{Z}/n)$$

of a \mathbb{Z}/n -module M .

The qfh-sheaves $z_0(X)$ are contravariantly functorial in X for open immersions, and covariantly functorial for proper maps. If Y is a closed subscheme of X with complement U , the following sequence

$$(11.2) \quad 0 \rightarrow z_0(Y) \rightarrow z_0(X) \rightarrow z_0(U)$$

is easily seen to be exact. Letting $\tilde{z}_0(X)_h$ denote the h-sheaf associated to $z_0(X)$, there is the

Lemma 11.4. *The sequence*

$$(11.3) \quad 0 \rightarrow \tilde{z}_0(Y)_h \rightarrow \tilde{z}_0(X)_h \rightarrow \tilde{z}_0(U)_h \rightarrow 0$$

induced from the sequence (11.2) is an exact sequence of h-sheaves on \mathbf{Sch}/k .

REMARK 11.5. As mentioned in §4, the Suslin homology of the functor $C_0(X)$ is an algebraic version of homology; the Suslin homology of the functor $Z_0(X)$ can similarly be viewed as an algebraic version of *Borel-Moore homology*. In fact, recall from Theorem 7.1 that the functor $C_0(X)[1/p]$ extends to the qfh-sheaf $c_0(X)$ on \mathbf{Sch}/k . The h-sheaf $\tilde{c}_0(-)_h$ associated to the qfh-sheaf $c_0(-)$ satisfies a Mayer-Vietoris property, giving the exact sequence

$$(11.4) \quad 0 \rightarrow \tilde{c}_0(U \cap V)_h \rightarrow \tilde{c}_0(U)_h \oplus \tilde{c}_0(V)_h \rightarrow \tilde{c}_0(U \cup V)_h \rightarrow 0.$$

By Theorem 10.1, the sequences (11.3) and (11.4) give rise to a localization sequence for the mod n Suslin homology $H_*^{\text{sing}}(Z_0(-), \mathbb{Z}/n)$ and a Mayer-Vietoris sequence for the mod n Suslin homology $H_*^{\text{sing}}(-, \mathbb{Z}/n)$ (assuming n prime to the characteristic). In [50], modifications of the sequences (11.3) and (11.4) give rise to a localization sequence for $H_*^{\text{sing}}(Z_0(-), \mathbb{Z})$ of $Z_0(-)$ and a Mayer-Vietoris sequence for $H_*^{\text{sing}}(-, \mathbb{Z})$ (assuming characteristic zero).

If X is proper over k , we have

$$z_0(X) = c_0(X).$$

By Theorem 9.1, formula (5.2) and Corollary 10.2, we have the isomorphisms

$$(11.5) \quad \text{Ext}_h^*(\tilde{z}_0(X)_h, \mathbb{Z}/n) \cong H_{\text{ét}}^*(X, \mathbb{Z}/n)$$

for X proper. Using the long exact Ext-sequence arising from the exact sequence of Lemma 11.4, and the corresponding long exact Gysin sequence for étale cohomology with compact supports, one extends the isomorphism (11.6) to the isomorphism

$$(11.6) \quad \text{Ext}_h^*(\tilde{z}_0(X)_h, \mathbb{Z}/n) \cong H_c^*(X, \mathbb{Z}/n)$$

to arbitrary X ; here H_c^* is the étale cohomology with compact supports.

With the help of (11.6), one arrives at the following comparison of the mod- n higher Chow groups, and étale cohomology for affine X :

Theorem 11.6. *Let X be an affine variety over an algebraically closed field k , and let $n > 0$ be an integer prime to $\text{char}(k)$. Assume that $q > d := \dim(X)$. Then there is a natural isomorphism*

$$CH^q(X, p; \mathbb{Z}/n) \cong H_c^{2(d-q)+p}(X, \mathbb{Z}/n(d-q))^\vee.$$

The proof uses the homotopy property to replace X with $X \times \mathbb{A}^{q-d}$, which reduces to the case $q = d$; one then applies Proposition 11.3 and the isomorphism (11.6). Using Poincaré duality, Theorem 8.3 implies

Corollary 11.7. *Let X be a smooth affine variety over an algebraically closed field k , and let $n > 0$ be an integer prime to $\text{char}(k)$. Assume that $q \geq d := \dim(X)$. Then there is a natural isomorphism*

$$CH^q(X, p; \mathbb{Z}/n) \cong H_{\text{ét}}^{2d-p}(X, \mathbb{Z}/n(q)).$$

Theorem 11.6 (resp. Corollary 11.7) extends to quasi-projective X (resp. smooth quasi-projective X) if $\text{char}(k) = 0$, using the localization property for the higher Chow groups [5] and the analogous Gysin sequence for étale cohomology with compact supports.

12. EPILOGUE

We have not described the beautiful constructions and results in the fundamental papers [50], [52] and [20], which form the foundations for many of the constructions we have described. We omitted as well mention of the works on Lawson homology and related topics ([17], [18], and [16]), which formed a large part of the motivation for many of the constructions of [50] and [20]. We have also not described the various constructions of triangulated motivic categories and motivic complexes, in [23], [28], [34], [36], [53] and [33]. Presumably, the categorical constructions all lead to equivalent categories and the complexes all yield equivalent cohomologies, but this is at present not known.

More recently, Suslin and Voevodsky [51] have reduced the Quillen-Lichtenbaum conjectures for motivic cohomology to the *Bloch-Kato conjecture*, which asserts that the *Galois symbol* on Milnor K -theory of a field F defines an isomorphism

$$K_q^M(F)/n \rightarrow H_{\text{ét}}^q(F, \mu_n^{\otimes q})$$

for all n prime to the characteristic of F . Relying on results of Rost [43], [44], Voevodsky has given a proof of the mod 2 Bloch-Kato conjecture (see [54]), which thus completely solves the mod 2 Quillen-Lichtenbaum conjecture for motivic cohomology. Using the Bloch-Lichtenbaum spectral sequence [6] relating K -theory and motivic cohomology, Weibel [55] has shown how to compute the mod 2 K -theory of \mathbb{Z} ; this has recently been extended by Kahn [30] to a computation of the mod 2 K -theory of all number rings, verifying that mod 2 algebraic K -theory agrees with mod 2 étale K -theory for all number rings. There is a whole host of unanswered questions in the area of motivic cohomology and algebraic K -theory; the new viewpoint offered by Voevodsky, Suslin and others will surely lead to further breakthroughs.

REFERENCES

- [1] M. Artin, **Grothendieck topologies**, Seminar notes, Harvard Univ. Dept of Math., 1962.
- [2] M. Atiyah, **K-theory**, Benjamin 1967. MR **36**:7130
- [3] M. Atiyah and F. Hirzebruch, *Vector bundles and homogeneous spaces*, Amer. Math. Soc. Symp. in Pure Math. **III**(1961) 7-38. MR **25**:2617
- [4] S. Bloch, *Algebraic cycles and higher K-theory*, Adv. in Math. **61** No. 3(1986) 267-304. MR **88f**:18010
- [5] S. Bloch, *The moving lemma for the higher Chow groups*, J. Alg. Geom. **3**(1994) 537-568. MR **96c**:14007
- [6] S. Bloch and S. Lichtenbaum, *A spectral sequence for motivic cohomology*, preprint (1995).
- [7] A. Borel and J.-P. Serre, *Le théorème de Riemann-Roch*, Bull. Soc. Math. France **86**(1958) 97-136. MR **22**:6817
- [8] C. Chevalley, *Anneaux de Chow et Applications*, Sémin. C. Chevalley, **2** Paris, 1958.
- [9] W.-L. Chow, *On the equivalence classes of cycles in an algebraic variety*, Ann. of Math. **64**(1956) 450-479. MR **18**:509a
- [10] P. Deligne *et al.*, *Cohomologie Etale*, Lecture Notes in Math. **569**, Springer Verlag 1977. MR **57**:3132
- [11] A. Dold and R. Thom, *Quasifaserungen und unendliche symmetrische Produkte*, Ann. of Math. **67**(1958) 239-281. MR **20**:3542
- [12] A. Duady and J.-L. Verdier, *Séminaire de Géométrie Analytique de l'Ecole Normal Sup.* Astérisque **36-7**, 1976.
- [13] W. Dwyer and E. Friedlander, *Algebraic and étale K-theory*, Trans. Amer. Math. Soc. **292**(1985) 247-280. MR **87h**:18013
- [14] E. Friedlander, *Etale K-theory I: Connections with Etale Cohomology and Algebraic Vector Bundles*, Invent. Math. **60**(1980) 105-134. MR **82e**:14029

- [15] E. Friedlander, *Etale K-theory II: Connections with algebraic K-theory*, Ann. Sci. Ec. Norm. Sup. 4^e sér. **15** no. 2(1982) 231-256. MR **85c**:14014
- [16] E. Friedlander and O. Gabber, *Cycle spaces and intersection theory*, in **Topological methods in modern mathematics**, Publish or Perish, Houston, TX (1993) 325-370. MR **94j**:14010
- [17] E. Friedlander and B. Lawson, *Duality relating spaces of algebraic cocycles and cycles*, Topology **36** (1997) 533-565. MR **1**:415 605
- [18] E. Friedlander and B. Lawson, *Moving algebraic cycles of bounded degree*, preprint (1994).
- [19] E. Friedlander and B. Mazur, *Correspondence homomorphisms for singular varieties*, Ann. Inst. Fourier (Grenoble) **44** (1994) 703-727. MR **95j**:14009
- [20] E. Friedlander and V. Voevodsky, *Bivariant cycle cohomology*, preprint (1995).
- [21] W. Fulton, **Intersection Theory**, Springer-Verlag, New York 1984. MR **85k**:14004
- [22] O. Gabber, *K-theory of henselian local rings and henselian pairs*, in **Algebraic K-theory, Commutative Algebra, and Algebraic Geometry**, Contemp. Math. **126**, Amer. Math Soc., Providence, RI (1992) 59-70. MR **93c**:19005
- [23] D. Grayson, *Weight filtrations via commuting automorphisms*, K-Theory **9** (1995) 139-172. MR **96h**:19001
- [24] A. Grothendieck, *Classes de faisceaux et théorème de Riemann-Roch*, Exposé 0, **SGA 6**, Lecture Notes in Math. 225(1971) 297-364.
- [25] A. Grothendieck, *La théorie des classes des Chern*, Bull. Soc. Math. France **86**(1958) 137-154. MR **22**:6818
- [26] A. Grothendieck *et al.*, *Théorie de Topos et Cohomologie Etale des Schémas*, I, Lecture Notes in Math. **269**, Springer Verlag 1972. MR **50**:7130
- [27] H. Gillet and R. Thomason, *On the K-theory of strict hensel local rings and a theorem of Suslin*, JPAA **34**(1984) 241-254. MR **86e**:18014
- [28] M. Hanamura, *Mixed motives and algebraic cycles*, I, Math. Res. Lett. **2** (1995) 811-821; II, preprint (1995). MR **1**:362 972
- [29] A. J. de Jong, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math. **83** (1996) 51-93. MR **1**:423 020
- [30] B. Kahn, *The Quillen-Lichtenbaum conjecture at the prime 2*, preprint (1997).
- [31] M. Karoubi and O. Villamayor, *K-théorie algébrique et K-théorie topologique*, I. Math. Scand. **28**(1971) 265-307. MR **47**:1915
- [32] M. Levine, *Bloch's higher Chow groups revisited*, in **K-theory, Strasbourg, 1992**, Asterisque **226**(1994)235-320. MR **96c**:14008
- [33] M. Levine, *Motivic cohomology and algebraic cycles*, preprint(1995).
- [34] S. Lichtenbaum, *Values of zeta-functions at non-negative integers*, in **Number Theory**, Lecture Notes in Math. 1068, Springer, Berlin (1984) 127-138. MR **756**:089
- [35] J.S. Milne, **Etale Cohomology**, Princeton Math. Ser. **33**, Princeton Univ. Press 1980. MR **81j**:14002
- [36] F. Morel, *Théorie de l'homotopie et motifs*, I, preprint(1995).
- [37] D. Mumford, *Rational equivalence of zero-cycles on surfaces*, J. Math. Kyoto Univ. **9** (1968), 195-204. MR **40**:2673
- [38] D. Quillen, *Cohomology of groups*, International Congress of Mathematics, Nice 1970. MR **58**:7627a
- [39] D. Quillen, *Higher algebraic K-theory I*, in **Algebraic K-theory I: Higher K-theories**, Lecture Notes in Math. **341**, Springer Verlag(1973) 85-147. MR **49**:2895
- [40] J. Roberts, *Chow's moving lemma*, in **Algebraic Geometry**, Oslo 1979, F. Oort editor. MR **52**:3154
- [41] A.A. Roitman, *Rational equivalence of zero-dimensional cycles*, Math. of the USSR Sbornik **18**(1972) 571-588. MR **48**:6109
- [42] A.A. Roitman, *The torsion of the group of 0-cycles modulo rational equivalence*, Ann. of Math. **111**(1980) 553-569. MR **81g**:14003
- [43] M. Rost, *On the spinor norm and $A_0(X, \mathcal{K}_1)$ for quadrics*, preprint (1988).
- [44] M. Rost, *Some new results on the Chow groups of quadrics*, preprint (1990).
- [45] F. Severi, *Problèmes résolus et problèmes nouveaux dans la théories des systèmes d'équivalence*, Proc. Intern. Cong. Math. **3**(1954), 529-541. MR **20**:5206
- [46] A. Suslin, *On the K-theory of algebraically closed fields*, Invent. Math. **73**(1983) 241-245. MR **85h**:18008a

- [47] A. Suslin, *On the K-theory of local fields*, JPAA **34**(1984) 301-318. MR **86d**:18010
- [48] A. Suslin, *Higher Chow groups of affine varieties and étale cohomology*, preprint (1994).
- [49] A. Suslin and V. Voevodsky, *Singular homology of abstract algebraic varieties*, Inv. Math. **123** Fasc. 1(1996) 61-94. MR **1**:376 246
- [50] A. Suslin and V. Voevodsky, *Relative cycles and Chow sheaves*, preprint (1994).
- [51] A. Suslin and V. Voevodsky, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, preprint(1995).
- [52] V. Voevodsky, *Homology of schemes II*, preprint (1994).
- [53] V. Voevodsky, *Triangulated categories of motives over a field*, preprint (1995).
- [54] V. Voevodsky, *The Milnor conjecture*, preprint (1996).
- [55] C. Weibel, *The two-torsion in the K-theory of the integers*, preprint (1996).
- [56] A. Weil, **Foundations of Algebraic Geometry**, AMS Colloquium publ. **29**, AMS 1962. MR **26**:2439

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MASSACHUSETTS 02115
E-mail address: `marc@neu.edu`