

SELF-BÄCKLUND CURVES IN CENTROAFFINE GEOMETRY AND LAMÉ'S EQUATION

MISHA BIALY, GIL BOR, AND SERGE TABACHNIKOV

ABSTRACT. Twenty five years ago U. Pinkall discovered that the Korteweg-de Vries equation can be realized as an evolution of curves in centroaffine geometry. Since then, a number of authors interpreted various properties of KdV and its generalizations in terms of centroaffine geometry. In particular, the Bäcklund transformation of the Korteweg-de Vries equation can be viewed as a relation between centroaffine curves.

Our paper concerns self-Bäcklund centroaffine curves. We describe general properties of these curves and provide a detailed description of them in terms of elliptic functions. Our work is a centroaffine counterpart to the study done by F. Wegner of a similar problem in Euclidean geometry, related to Ulam's problem of describing the (2-dimensional) bodies that float in equilibrium in all positions and to bicycle kinematics.

We also consider a discretization of the problem where curves are replaced by polygons. This is related to discretization of KdV and the cross-ratio dynamics on ideal polygons.

CONTENTS

1. Introduction	232
2. Bäcklund transformations of centroaffine curves	237
3. Self-Bäcklund curves: First study	246
4. Self-Bäcklund curves and the Lamé equation	250
5. Self-Bäcklund polygons	272
6. Appendix: From the centroaffine plane to the hyperbolic plane	278
Acknowledgments	280
References	280

1. INTRODUCTION

The motivation for this work is the interpretation of the Korteweg-de Vries equation in terms of centroaffine geometry. This growing body of work started with U. Pinkall's paper [38], see [15, 25, 26, 46] for a sampler.

In [44], the Bäcklund transformation of the KdV equation is interpreted as a relation between centroaffine curves. We start with a very brief description of this approach to KdV.

Received by the editors December 7, 2020, and, in revised form, May 9, 2022, and July 16, 2022.
2020 *Mathematics Subject Classification.* Primary 53B99.

The first author was supported by ISF grant 580/20. The second author was supported by Conacyt grant #A1-S-45886. The third author was supported by NSF grants DMS-1510055 and DMS-2005444.

Let $\gamma(t)$ be a parametrized smooth curve in the affine plane with a fixed area form. The curve is *centroaffine* if the Wronski determinant is constant: $[\gamma(t), \gamma'(t)] = 1$ for all $t \in \mathbb{R}$. The group $SL_2(\mathbb{R})$ acts on centroaffine curves, and we shall also consider the moduli space of such curves.

Unless specified otherwise, we assume that the curves are π -anti-periodic: $\gamma(t + \pi) = -\gamma(t)$ for all t . That is, the curve is closed, centrally symmetric and 2π -periodic (the last condition, if not satisfied by a centrally symmetric centroaffine curve, can be arranged by an appropriate rescaling.)

The rationale for assuming that the curves are centrally symmetric is as follows. An orientation preserving diffeomorphism of $\mathbb{R}P^1$ admits a unique area preserving and homogeneous of degree 1 lifting to a diffeomorphism of the punctured plane. The image of the unit circle under such a diffeomorphism is a centrally symmetric star-shaped curve, and projectively equivalent diffeomorphisms correspond to $SL_2(\mathbb{R})$ -equivalent curves. See [36] for details.

Our results can be extended to non-centrally symmetric curves, but we do not dwell on it in this paper.

Given a centroaffine curve, one has $\gamma''(t) = p(t)\gamma(t)$ where p is a π -periodic potential function of the Hill operator $-d^2/dt^2 + p(t)$. In the language of centroaffine geometry, p is the *centroaffine curvature* of the curve γ (alternatively, some authors call $-p$ the centroaffine curvature, but we shall adopt the plus sign convention).

For example, $\gamma(t) = (\cos t, \sin t)$ has $p(t) = -1$. This unit circle and its $SL_2(\mathbb{R})$ images are trivial examples of centroaffine curves. We refer to these curves as *centroaffine conics*.

A tangent vector to a centroaffine curve $\gamma(t)$, in the space of π -anti-periodic centroaffine curves, is given by a vector field along it of the form $g(t)\gamma(t) + f(t)\gamma'(t)$, where f, g are π -periodic. Taking the derivative of the centroaffine condition $[\gamma, \gamma'] = 1$ with respect to this vector field we obtain $f' + 2g = 0$. Thus such a vector field has the form

$$(1) \quad V_f := -\frac{1}{2}f'(t)\gamma(t) + f(t)\gamma'(t),$$

where f is a π -periodic function. Pinkall observed in [38] that the evolution of the curves $\gamma(t)$ with the potential function $p(t)$ under the vector field V_p is a centroaffine version of the Korteweg-de Vries equation: the potential evolves according to the equation

$$\dot{p} = -\frac{1}{2}p''' + 3p'p$$

(where dot is the time derivative).

We say that two centroaffine curves, $\gamma(t)$ and $\delta(t)$, are *c-related* if $[\gamma(t), \delta(t)] = c$ for all t . See Figure 1. It is shown in [44] that this relation is a geometric realization of the Bäcklund transformation for the KdV equation.

In this paper we are mostly interested in *self-Bäcklund* centroaffine curves, the curves $\gamma(t)$ for which there exist $\alpha \in (0, \pi)$ and a constant c such that

$$(2) \quad [\gamma(t), \gamma(t + \alpha)] = c \text{ for all } t.$$

For example, the centroaffine conics are self-Bäcklund for every choice of α with $c = \sin \alpha$. To exclude trivial cases, we assume that $c \neq 0$. We call α in equation (2) the




FIGURE 1. Bäcklund transformation: As the end points of the line segment AB trace the two curves, OA and OB sweep area with the same rate and the area of the shaded triangle OAB remains constant

rotation number of a self-Bäcklund curve. See Figure 2 for examples of self-Bäcklund curves.




FIGURE 2. Self-Bäcklund curves (blue), with winding numbers 1 (left) and 3 (right). A line segment (green) moves with its end points sliding along the curve, forming a constant area triangle with the origin, while the midpoint of the line segment traces a curve (red), always tangent to the line segment at its midpoint. The two curves depicted here are members of an infinite family of self-Bäcklund curves described explicitly in Section 4 in terms of the Weierstrass \wp -function. For more images and animations, see [12].

An analogous problem in Euclidean geometry was thoroughly studied relatively recently. The problem is to describe the closed smooth arc length parametrized curves $\gamma(t) \subset \mathbb{R}^2$ for which there exist constants s and ℓ such that $|\gamma(t+s) - \gamma(t)| = \ell$ for all t . For example, a circle is a trivial solution to this problem.

Although the full solution of this problem is not available yet, there is a wealth of results, including many non-trivial examples of such curves. See [11, 41, 43, 47–49] for a sampler.

This problem originated in two seemingly unrelated theories. First, such curves are the boundaries of 2-dimensional bodies that float in equilibrium in all positions – to

describe such bodies (in all dimensions) is S. Ulam's problem in flotation theory, see [34, problem 19].

Second, an interesting problem in the study of bicycle kinematics is to describe the pairs of front and rear bicycle tracks for which one cannot determine the direction of the bicycle motion. The above mentioned curves appear in this problem as the front tracks in such ambiguous pairs; they are referred to as *bicycle curves*. See [23] for a survey of this approach to bicycle kinematics.

This geometric problem is intimately related to another completely integrable equation of soliton type, the filament – or binormal, or smoke ring, or local induction – equation; more precisely, to the planar filament equation

$$\dot{\gamma} = \frac{1}{2}k^2T + \frac{dk}{ds}N,$$

where $\gamma(s)$ is an arc length parameterized plane curve, dot means the time evolution, k is the curvature, and T and N is a Frenet frame along γ .

Two arc length parametrized curves, $\gamma(t)$ and $\delta(t)$, are in bicycle correspondence if the length of the segment $\gamma(t)\delta(t)$ is constant and the velocity of its midpoint is aligned with the segment for all t . This correspondence is a geometric realization of the Bäcklund transformation of the planar filament equation, and in this sense, bicycle curves are self-Bäcklund.

We must say more about the work of Franz Wegner, cited above. He discovered a large variety of bicycle curves (or solutions to the 2-dimensional Ulam's problem), explicitly described in terms of elliptic functions. Wegner made his discovery by assuming that the desired solutions have a certain geometrical property, resulting in a differential equation on their curvature that was solved in elliptic functions. Then he proved that indeed, for a proper choice of parameters, these curves solved the problem.

It is shown in [11] that Wegner's curves are solutions to a variational problem: they are buckled rings (the relative extrema of the elastic – or bending – energy, subject to the length and area constraints), and they are solitons: under the planar filament flow, they evolve by isometries.

Our main goal in this paper is to obtain centroaffine analogs of these results.

In the spirit of discrete differential geometry, we also consider centroaffine polygons, a discretization of centroaffine curves. These are centrally symmetric $2n$ -gons P_1, \dots, P_{2n} such that $[P_i, P_{i+1}] = 1$ and $P_{i+n} = -P_i$ for all i (the index is understood cyclically). A centroaffine $2n$ -gon is a *self-Bäcklund* (n, k) -gon if there exists a constant c such that $[P_i, P_{i+k}] = c$ for all i . A trivial example is an affine-regular $2n$ -gon which is a self-Bäcklund (n, k) -gon for all k . The problem is to describe non-trivial self-Bäcklund (n, k) -gons.

These polygons are centroaffine analogs of the discretization of the bicycle curves, the bicycle polygons, studied in [41, 45]. Some of our results on self-Bäcklund (n, k) -gons were included in Section 7.3 of the original (but not the final) version of [6], and were motivated by the study of the cross-ratio dynamics on ideal polygons in the hyperbolic plane and hyperbolic space therein.

Centroaffine polygons are closely related to linear second-order difference equations with periodic solutions and with Coxeter's frieze patterns, see [35]. In particular, given a simple centroaffine $2n$ -gon, the determinants $[P_i, P_j]$ with $|i - j| < n$ form the entries, all positive, of a frieze pattern of width $n - 3$. In these terms, we are interested in

frieze patterns that have a row consisting of the same numbers, but not every row being constant.

A word about the terminology that we use. We call a closed smooth curve *star-shaped* if every ray emanating from the origin intersects the curve transversely and only once. A curve is *locally star-shaped* if the above property holds locally, near every point. Equivalently, $[\gamma(t), \gamma'(t)] \neq 0$ for all t . Star-shaped curves have winding number 1, but locally star-shaped curves can go around the origin several times.

The contents of this paper are as follows.

Section 2 concerns Bäcklund transformations of centroaffine curves. We describe a centroaffine analog of the rear track curve (in the above mentioned bicycle setting). We also interpret the Miura transformation in terms of centroaffine geometry.

Section 2.4 is devoted to the following problem: given a centroaffine curve γ , for which c do c -related curves exist? We provide a complete answer to this question. This result is a centroaffine analog of Menzin's conjecture – now a theorem, originally formulated for hatchet planimeters, but it also applies to the bicycle model, see [30] or [23].

Section 3 comprises several results on self-Bäcklund curves. In Theorem 3 we prove that a non-trivial infinitesimal deformation of a central conic as a self-Bäcklund curve exists if and only if either $\alpha = \pi/2$ or α satisfies the equation

$$\tan(k\alpha) = k \tan \alpha$$

for some integer $k \geq 4$. A similar result is known for bicycle curves, see [41].

We show that if $\alpha = \pi/3$ or $\alpha = \pi/4$ then only the central conics are self-Bäcklund. In contrast, if $\alpha = \pi/2$, one has a family of self-Bäcklund centroaffine curves with functional parameters. Example 4.11 provides families of analytic curves with rotation number $\pi/2$ and, at the same time, examples of analytic Radon curves.

Section 4 is the core part of the paper. We start by developing a centroaffine analog of Wegner's ansatz, that is, we guess what geometric properties self-Bäcklund curves may possess. This leads to the assumption that these curves correspond to the traveling wave solutions of the KdV equation, that is, their centroaffine curvature is an elliptic function.

Thus we assume that the coordinates of our self-Bäcklund curves satisfy the Lamé equation, the Hill equation whose potential is an elliptic function. In Section 4.2 we construct these curves and describe the conditions on the parameters for which the curves are self-Bäcklund. This work is analogous to the one done by F. Wegner. In Section 4.3 we show that central conics indeed admit a deformation into self-Bäcklund centroaffine curves for each α appearing in Theorem 3.

Section 5 concerns self-Bäcklund centroaffine polygons. We start by showing that the c -relations on centroaffine curves satisfy the Bianchi permutability property (Theorem 9).

We describe a discrete version of Bäcklund transformation on centroaffine polygons (this transformation is studied in detail in [2]). Theorem 10 presents some pairs (n, k) for which non-trivial self-Bäcklund polygons do not exist, and some pairs for which they do. We also describe necessary and sufficient conditions for the existence of non-trivial infinitesimal deformations of regular centroaffine n -gons in the class of self-Bäcklund polygons. Similar results were known for bicycle polygons, see [41].

In the appendix we connect centroaffine geometry with another geometry associated with the group $SL_2(\mathbb{R})$, two-dimensional hyperbolic geometry. We assign to a centroaffine curve a curve in the hyperbolic plane, its dual. The centroaffine curvature p of a curve and the curvature κ of its dual in H^2 are related by the equation $(1+p)(1+\kappa) = 2$.

We make extensive use of the formulas involving the Weierstrass elliptic function. We refer to [39] for a compendium of such formulas. The arXiv preprint version of this paper contains an appendix listing these formulas.

It was pointed out by a referee that it is more common to use Jacobi elliptic functions in KdV theory, whereas we use the Weierstrass elliptic functions. In this regard, we quote from [1]:

The fact that the integral in Jacobi form or Riemann form contains only one parameter, and not two like the Weierstrass integral, is very convenient for various computations. The Weierstrass form is almost always preferable for theoretical considerations.

2. BÄCKLUND TRANSFORMATIONS OF CENTROAFFINE CURVES

2.1. The middle curve. Let $\gamma(t)$ be a centroaffine curve satisfying $\gamma''(t) = p(t)\gamma(t)$. Construct a new centroaffine curve $\delta(t) = f(t)\gamma(t) + g(t)\gamma'(t)$, where $f(t)$ and $g(t)$ are π -periodic functions. Lemma 2.1 repeats Lemma 1.2 of [44].

Lemma 2.1. *The curves γ and δ are c -related if and only if $g(t) = c$ and*

$$(3) \quad cf'(t) - f^2(t) + c^2p(t) + 1 = 0.$$

Proof. One has

$$c = [\gamma(t), \delta(t)] = g(t)[\gamma(t), \gamma'(t)] = g(t),$$

and therefore $g'(t) = 0$. Next,

$$\delta'(t) = (f'(t) + p(t)g(t))\gamma(t) + (f(t) + g'(t))\gamma'(t),$$

hence

$$1 = [\delta(t), \delta'(t)] = f^2(t) - c(f'(t) + cp(t)).$$

This implies equation (3). □

Note that equation (3) is a Riccati equation on the unknown function $f(t)$.

Lemma 2.2. *Let γ and δ be c -related and let $\Gamma(t)$ be the midpoint of the segment $\gamma(t)\delta(t)$. Then the velocity of Γ is aligned with this segment:*

$$\Gamma'(t) \sim \delta(t) - \gamma(t)$$

for all t . In addition, Γ is locally star-shaped, that is, $[\Gamma(t), \Gamma'(t)] \neq 0$ for all t .

Proof. Since $[\gamma, \gamma'] = [\delta, \delta'] = 1$ and $[\gamma, \delta] = c$, one has

$$[\gamma' + \delta', \delta - \gamma] = [\gamma', \delta] - [\delta', \gamma] = [\gamma, \delta]' = 0,$$

as needed.

For the second statement, if $[\Gamma(t), \Gamma'(t)] = 0$ then the line connecting $\gamma(t)$ and $\delta(t)$ passes through the origin, and then $c = 0$. □

Remark 2.3. The locus of midpoints in Lemma 2.2 plays the role of the rear bicycle track in the analogous problem mentioned in Section 1. This middle curve may have cusps.

We describe a method of constructing pairs of c -related curves. Start with a locally star shaped curve Γ , with a centroaffine parameter s and curvature $p(s)$, so that $[\Gamma, \Gamma_s] = 1$, $\Gamma_{ss} = p\Gamma$. Let $\gamma_{\pm} := \Gamma \pm (c/2)\Gamma_s$. The condition $[\gamma_-, \gamma_+] = c$ is immediate; however, in general, s is not a centroaffine parameter for γ_{\pm} .

Proposition 2.4. *If $c^2 p \neq 4$ along Γ (for example, if Γ is locally convex, that is, $p < 0$), then γ_{\pm} can be simultaneously reparametrized by a centroaffine parameter t , so that $[\gamma_{\pm}, (\gamma_{\pm})_t] = 1$.*

Proof. We calculate that $[\gamma_{\pm}, (\gamma_{\pm})_s] = 1 - (c^2/4)p$. If this does not vanish, then the desired parameter t is defined by

$$\frac{dt}{ds} = 1 - \frac{c^2 p(s)}{4}.$$

With this new parameter one has $[\gamma_{\pm}, (\gamma_{\pm})_t] = 1$, as needed. □

Remark 2.5. As we mentioned, and as is seen from illustrations in this paper, the middle curve Γ may have cusps. The above construction of the curves γ_{\pm} from Γ extends to the case when Γ has cusps and the curves γ_{\pm} remain smooth. Without going into details, we illustrate this with an example.

Let $\Gamma(x) = (x^2, x^3 + 1)$ be a cusp, and let s be a centroaffine parameter. Then $\Gamma_x = (2x, 3x^2)$ and

$$\frac{ds}{dx} = [\Gamma, \Gamma_x] = x^4 - 2x.$$

It follows that

$$\gamma_{\pm} = \Gamma \pm \frac{c}{2}\Gamma_s = \left(\mp\frac{c}{2}, 1\right) + \left(0, \mp\frac{3c}{4}\right)x + O(x^2),$$

which, for $c \neq 0$ and x close to zero, are smooth curves.

Remark 2.6. Consider an oriented smooth closed strictly convex plane curve Γ . The outer billiard transformation T is a map of its exterior, defined as follows: given a point x , draw the oriented tangent line from x to Γ , and reflect x in the tangency point to obtain the point $T(x)$. See [20] for a survey.

The relation of our topic to outer billiards is as follows: if γ is a self-Bäcklund curve and the respective middle curve Γ is convex, then γ is an invariant curve of the outer billiard map about Γ .

2.2. Curves c -related to centroaffine conics. In this section we consider the curves that are c -related to centroaffine conics and identify self-Bäcklund curves among them. These curves will have points at infinity.

Let $\gamma(t) = (\cos t, \sin t)$, and let us construct a c -related curve as in Lemma 2.1: $\delta(t) = f(t)\gamma(t) + c\gamma'(t)$. The respective Riccati equation for the function f is

$$(4) \quad cf'(t) = f^2(t) + c^2 - 1.$$

Assume that $c > 1$. This differential equation is easily solved:

$$(5) \quad f(t) = a \tan\left(\frac{at}{c}\right), \quad \text{where } a = \sqrt{c^2 - 1}$$

and a choice of the constant of integration has been made so that $f(0) = 0$ (any other solution is obtained by a parameter shift).

The function f has poles (the same is true for the solutions with $c < 1$ and $c = 1$), and the respective centroaffine curve goes to infinity, having there an inflection point.

For example, let $c = 5/3, a = 4/3$, see Figure 3. This curve is periodic with period 10π .




FIGURE 3. The curve $\delta(t) = \left(\frac{4}{3} \tan\left(\frac{4t}{5}\right) \cos t - \frac{5}{3} \sin t, \frac{4}{3} \tan\left(\frac{4t}{5}\right) \sin t + \frac{5}{3} \cos t\right)$

Let us look for self-Bäcklund curves among the above curves δ .

Lemma 2.7. *Let δ be the centroaffine curve c -related to the unit circle $\gamma(t) = (\cos t, \sin t)$, where $c > 1$. Then δ is self-Bäcklund with rotation number α , that is, $[\delta(t), \delta(t + \alpha)] = \text{const}$, if and only if α satisfies*

$$(6) \quad \tan(u\alpha) = u \tan \alpha, \quad \text{where } u = \frac{\sqrt{c^2 - 1}}{c}.$$

Furthermore, given such an α , one has $[\delta(t), \delta(t + \alpha)] = \sin \alpha$.

Proof. The statement is invariant under parameter shift so it is enough to consider $\delta = f\gamma + c\gamma'$, where f is given by formula (5). Next, by a straightforward calculation, the derivative of $[\delta(t), \delta(t + \alpha)]$ with respect to t is some non-zero function times $\tan(u\alpha) - u \tan \alpha$. It follows that $[\delta(t), \delta(t + \alpha)]$ is constant if and only if $\tan(u\alpha) = u \tan \alpha$. Using this equation for α , we calculate that $[\delta(t), \delta(t + \alpha)] = \sin \alpha$. \square

In general, for a fixed $u \in (0, 1)$, equation (6) has infinitely many solutions. See Figure 4. If u is rational then δ is periodic and there are finitely many solutions α within a period.

A solution of equation (4) for $c < 1$ is similar:

$$f(t) = -a \tanh\left(\frac{at}{c}\right),$$

where $a^2 = 1 - c^2$. The associated c -related curve $\delta = f\gamma + c\gamma'$ is non-periodic and stays bounded; it is self-Bäcklund with a parameter shift α satisfying

$$\tanh(u\alpha) = u \tan \alpha, \quad \text{where } u = \frac{\sqrt{1 - c^2}}{c},$$




FIGURE 4. Solutions to equation (6), $u \tan \alpha = \tan(u\alpha)$, $u \in (0, 1)$, are given by the intersection points of the (red) graph of the π -periodic function $y = \tan^{-1}(u \tan \alpha) - \alpha + \pi n$, $\pi n - \frac{\pi}{2} \leq \alpha \leq \pi n + \frac{\pi}{2}$, $n \in \mathbb{Z}$, and any of the (blue) lines $y = (u - 1)\alpha + n\pi$, $n \in \mathbb{Z}$. If u is rational then $f = a \tan(ut)$ is periodic and δ is closed, self-Bäcklund with rotation numbers α given by the intersection points within a period of f . In the figure above, $u = 2/7$, f is 7π -periodic, δ is 14π -periodic, and there are 8 solutions $\alpha \in (0, 14\pi)$ with $\sin \alpha \neq 0$.




FIGURE 5. The curve $\delta(t) = \left(-\frac{4}{5} \tanh\left(\frac{4t}{3}\right) \cos t - \frac{3}{5} \sin t, -\frac{4}{5} \tanh\left(\frac{4t}{3}\right) \sin t + \frac{3}{5} \cos t\right)$

and the constant determinant is $\sin \alpha$. This equation admits infinitely many solutions $\pm\alpha_1, \pm\alpha_2, \dots$, with $\alpha_n \in (n\pi, n\pi + \pi/2)$. For $t \rightarrow \pm\infty$, the curve approaches the unit circle, see Figure 5.

Another solution of (4) for $c < 1$ is

$$f(t) = -a \operatorname{coth}\left(\frac{at}{c}\right),$$

with the respective value of α given by

$$\operatorname{coth}(u\alpha) = u \tan \alpha, \quad \text{where } u = \frac{\sqrt{1 - c^2}}{c}$$

and the constant determinant is $\sin \alpha$. There are infinitely many solutions here as well, $\pm\alpha_0, \pm\alpha_1, \dots$, with $\alpha_n \in (n\pi, n\pi + \pi/2)$. This curve approaches the unit circle as $t \rightarrow \pm\infty$ and goes to infinity as $t \rightarrow 0$. See Figure 6.

If $c = 1$, a solution of equation (4) is $f(t) = -1/t$. This curve is self-Bäcklund with a parameter shift α satisfying $\tan \alpha = \alpha$ and the constant determinant is $\sin \alpha$. There are infinitely many solutions $\pm\alpha_1, \pm\alpha_2, \dots$, with $\alpha_n \in (n\pi, n\pi + \pi/2)$. Its asymptotic behavior is the same as in the previous example, see Figure 7.

For completeness, consider the case of a straight line $\gamma(t) = (t, -1)$. This centroaffine curve is self-Bäcklund for an arbitrary parameter shift. A c -related curve




FIGURE 6. The curve $\delta(t) = \left(-\frac{4}{5} \coth\left(\frac{4t}{3}\right) \cos t - \frac{3}{5} \sin t, -\frac{4}{5} \coth\left(\frac{4t}{3}\right) \sin t + \frac{3}{5} \cos t\right)$




FIGURE 7. The curve $\delta(t) = \left(-\frac{1}{t} \cos t - \sin t, -\frac{1}{t} \sin t + \cos t\right)$

$f\gamma + c\gamma'$ has $f(t) = -\tanh(t/c)$, see Figure 8. This curve is not self-Bäcklund: the respective equation on the parameter shifts b is

$$\tanh\left(\frac{b}{c}\right) = \frac{b}{c},$$

and the only solution is $b = 0$.




FIGURE 8. The curve $\delta(t) = (1 - t \tanh t, \tanh t)$ (red), a Bäcklund transform of the line $y = -1$ (black)

2.3. c -Related curves and Miura transformation. The Miura transformation connects the Korteweg-de Vries equation $\dot{u} = u''' + 6uu'$ and the modified Korteweg-de Vries equation $\dot{v} = v''' - 6v^2v'$: if v satisfies mKdV then $u = -v' - v^2$ satisfies KdV. More generally, if

$$(7) \quad u = -v' - v^2 + \lambda,$$

and v satisfies

$$(8) \quad \dot{v} = v''' - 6v^2v' - 6\lambda v',$$

then u satisfies KdV. See [24].

Given u , equation (7) is a Riccati equation on v , just like equation (3) on the function $f(t)$ that describes the curves, c -related to a centroaffine curve with curvature $p(t)$. This provides a geometrical interpretation of the Miura transformation in centroaffine geometry.

The details are described by Theorem 1.

Theorem 1. *Let γ be a centroaffine curve, and $\delta = f\gamma + c\gamma'$ be a c -related curve. Let the curves γ and δ evolve by the KdV flow. Then they remain c -related, and the function f evolves according to a version of mKdV:*

$$\dot{f} = -\frac{1}{2}f''' + \frac{3}{c^2}(f^2 - 1)f'.$$

Proof. Let q be the centroaffine curvature of δ , that is, $\delta''(t) = q(t)\delta(t)$. Then $\dot{\gamma} = V_p$, $\dot{\delta} = V_q$, where we use the notation as in equation (1).

We start with the observation that $\gamma = f\delta - c\delta'$, and then we express the curvatures p and q from equation (3) as follows

$$(9) \quad p = \frac{1}{c^2}(f^2 - 1 - cf'), \quad q = \frac{1}{c^2}(f^2 - 1 + cf')$$

(compare with Lemma 3.1 in [44]). It follows that

$$(10) \quad q - p = \frac{2}{c}f', \quad p' + q' = \frac{4}{c^2}ff'.$$

That γ and δ remain c -related under the KdV flow follows from the fact the c -relation commutes with the KdV flow, see [44]. Here is an independent verification.

We have: $\delta' = (f' + cp)\gamma + f\gamma'$, and

$$\begin{aligned} [\gamma, \delta]' &= [V_p, \delta] + [\gamma, V_q] = [-\frac{1}{2}p'\gamma + p\gamma', \delta] + [\gamma, -\frac{1}{2}q'\delta + q\delta'] = \\ &= -\frac{1}{2}c(p' + q') + f(q - p) = 0, \end{aligned}$$

the last equality due to equation (10).

To calculate \dot{f} , note that $f = [\delta, \gamma']$. Then

$$\dot{f} = [\dot{\delta}, \gamma'] + [\delta, \dot{\gamma}'] = [V_q, \gamma'] + [\delta, V_p'] = [-\frac{1}{2}q'\delta + q\delta', \gamma'] + [\delta, (-\frac{1}{2}p'\gamma + p\gamma')].$$

After substituting the values of p and q and their derivatives in terms of f from equation (9) and collecting terms we obtain the stated equality. \square

One can expand a periodic solution of equation (3) in a power series in c :

$$\begin{aligned} f &= 1 + \frac{c^2}{2}p + \frac{c^3}{4}p' + \frac{c^4}{8}(p'' - p^2) + \frac{c^5}{16}(p''' - 8pp') \\ &\quad + \frac{c^6}{32}[p'''' - 10pp'' - 9(p')^2 + 2p^3] + \dots \end{aligned}$$

Given the relation of f with the Miura transformation, one has the next statement; see Section 1.1 of [24].

Corollary 2.8. *The integrals of the odd terms of this series vanish, and the integrals of the even terms are integrals of the KdV equation:*

$$\int_0^\pi p \, dt, \int_0^\pi p^2 \, dt, \int_0^\pi \left(p^3 + \frac{1}{2}(p')^2 \right) dt, \dots$$

See [11, Section 3.3] for a similar statement about the bicycle transformation and the filament equation.

2.4. Range of the parameter c . The aim of this section is to describe, for a given centroaffine closed π -anti-periodic curve $\gamma(t)$, the range of the parameter c for which γ admits closed centroaffine c -related curves. The main result is Theorem 2, describing this range (a closed interval) in terms of the lowest eigenvalue of a Hill equation associated with γ . For a convex γ we obtain as a corollary an upper bound on c in terms of the area enclosed by its dual curve γ^* . This result can be viewed as a centroaffine analog of Menzin's conjecture for hatchet planimeters (equivalently, bicycle monodromy), discussed and proved in [30].

As we saw in Lemma 2.1, finding a centroaffine curve c -related to a given curve γ amounts to finding a solution $f(t)$ to the Riccati equation

$$(11) \quad cf' - f^2 + c^2p(t) + 1 = 0,$$

where $p = [\gamma'', \gamma']$ (the centroaffine curvature of γ). The corresponding c -related centroaffine curve is $\delta = f\gamma + c\gamma'$. If γ is π -anti-periodic then p in equation (11) is π -periodic and we are looking for the values of the parameter c for which the equation admits a π -periodic solution, so that δ is π -anti-periodic as well. Note that for $c = 0$ the equation admits the trivial solution $f \equiv 1$.

Our study of the Riccati equation (11) is based on its relation with the Hill equation

$$(12) \quad y'' + (\lambda - p(t))y = 0.$$

To state this relation we recall first that a solution $y(t)$ of (12) is called π -quasi-periodic if $y(t + \pi) = \mu y(t)$ for all t and some $\mu \in \mathbb{R}, \mu \neq 0$, called the Floquet multiplier of $y(t)$. If $\mu = 1$ then the solution is π -periodic and if $\mu = -1$ it is π -anti-periodic.

Proposition 2.9. *The Riccati equation (11) with a π -periodic $p(t)$ admits a π -periodic solution $f(t)$ for a parameter value $c \neq 0$ if and only if the Hill equation (12) admits a positive π -quasi-periodic solution $y(t)$ for $\lambda = -1/c^2$.*

Proof. Indeed, if there exists such $y(t)$, then $f := -cy'/y$ is a periodic solution of equation (11). In the opposite direction: if f is a periodic solution of equation (11) and F is a primitive of f then $y := e^{-F/c}$ is the required solution of equation (12). □

We now borrow a well-known result from the general theory of the Hill equation, due to Lyapunov and Haupt (ca. 1910, see Theorem 2.1 on page 11 of [31]).

Theorem (Spectrum of the Hill operator). *Consider equation (12),*

$$y'' + (\lambda - p(t))y = 0,$$

where $y(t)$ is an unknown real function, $p(t)$ is a real π -periodic function and λ a real parameter. Then there exist two unbounded sequences of real numbers

$$\begin{aligned} \lambda_0 < \lambda_1 \leq \lambda_2 < \lambda_3 \leq \lambda_4 < \dots, \\ \mu_0 \leq \mu_1 < \mu_2 \leq \mu_3 < \mu_4 \leq \dots, \end{aligned}$$

satisfying

$$(13) \quad \lambda_0 < \mu_0 \leq \mu_1 < \lambda_1 \leq \lambda_2 < \mu_2 \leq \mu_3 < \lambda_3 \leq \lambda_4 < \dots,$$

such that equation (12) has a non-trivial π -periodic solution if and only if $\lambda = \lambda_k$, and a π -anti-periodic non-trivial solution if and only if $\lambda = \mu_k$, $k = 0, 1, \dots$. The number of zeros on $[0, \pi)$ of a solution corresponding to λ_{2k-1} or λ_{2k} is $2k$. In particular, if a π -periodic solution has no zeros, then $\lambda = \lambda_0$. Similarly, the number of zeros on $[0, \pi)$ of a non-trivial solution corresponding to μ_{2k} or μ_{2k+1} is $2k + 1$. Moreover, a solution to equation (12) is unstable (that is, unbounded) if and only if λ belongs to one of the intervals $(-\infty, \lambda_0)$, (μ_0, μ_1) , (λ_1, λ_2) , \dots (called instability intervals, or ‘gaps’). See Figure 9.




FIGURE 9. The spectrum of Hill’s equation (12), stability and instability intervals

Concerning the lowest eigenvalue λ_0 , we have the following.

Lemma 2.10. *Let λ_0 be the first eigenvalue of the spectrum (13) of the Hill equation (12) associated with a π -anti-periodic centroaffine curve γ . Then*

$$\lambda_0 < 0, \quad \lambda_0 \leq -P,$$

where

$$(14) \quad P := -\frac{1}{\pi} \int_0^\pi p(t) dt.$$

Proof. Each of the two coordinate components of γ is a non-trivial π -anti-periodic solution of equation (12) for $\lambda = 0$. This implies that $\mu_k = 0$ for some $k \geq 1$, hence $\lambda_0 < 0$.

The inequality $\lambda_0 \leq -P$ is due to Borg (see Theorem 3.3.1 of [22]). The following argument is due to Ungar: Take a positive periodic solution $y(t)$ of equation (12) corresponding to λ_0 . Then $h(t) = y'(t)/y(t)$ is a periodic solution of the Riccati equation $h' + h^2 + (\lambda_0 - p(t)) = 0$. Integrating this equation over the period gives:

$$\int_0^\pi (\lambda_0 - p(t))dt \leq 0.$$

This yields the result. □

Remark 2.11. If γ is locally convex, so that $p(t)$ is strictly negative, then $P > 0$ and we have $\lambda_0 \leq -P < 0$. The geometric meaning of P is the area bounded by the dual curve γ^* (we refer to [28] and [42] for this and related facts).

Theorem 2. *Let γ be a centroaffine π -anti-periodic curve and $\lambda_0 < 0$ the lowest π -periodic eigenvalue of the associated Hill equation (12). Then γ admits a c -related closed curve if and only if $|c| \leq 1/\sqrt{-\lambda_0}$.*

An immediate consequence of Theorem 2 and Lemma 2.10 is the following.

Corollary 2.12. *Suppose $P > 0$ (for example γ is locally convex) and γ admits a c -related π -anti-periodic closed curve. Then $|c| \leq 1/\sqrt{P}$.*

Proof of Theorem 2. By Proposition 2.9, we need to show that equation (12) admits a π -quasi-periodic positive solution if and only if $\lambda \leq \lambda_0$.

Consider first the “if” part. If $\lambda = \lambda_0$ then equation (12) has a positive periodic solution, hence quasi-periodic. So we shall assume now that $\lambda < \lambda_0$. In this case equation (12) has no conjugate points, that is, a non-trivial solution vanishing at two distinct points t_1, t_2 because, by the Sturm Comparison Theorem, any solution for every larger λ must have a zero between t_1, t_2 . However for λ_0 there is a positive periodic solution. To complete the proof of the “if” part we make use of Lemma 2.13.

Lemma 2.13. *The equation $y'' + q(t)y = 0$, where $q(t + \pi) = q(t)$, has no conjugate points if and only if it admits a positive π -quasi-periodic solution.*

As far as we know, Lemma 2.13 is due to E. Hopf [29]. For completeness, we give its proof below.

Now we prove Theorem 2 in the opposite direction. We need to show that equation (12) admits no positive π -quasi-periodic solution for $\lambda > \lambda_0$. Assume $y(t)$ is such a solution, $y(t + \pi) = \mu y(t)$, where $\mu > 0$. There are two cases:

- If $\mu = 1$ then $y(t)$ is a positive periodic solution. But this is possible only for $\lambda = \lambda_0$, a contradiction.
- If $\mu \neq 1$ then the solution $y(t)$ is unbounded, and hence λ belongs to one of the instability zones. In particular, $\lambda > \mu_0$. But then, by the Sturm Comparison Theorem, $y(t)$ cannot be positive since solutions for μ_0 have zeroes.

This completes the proof of Theorem 2. □

Proof of Lemma 2.13 (after E. Hopf). If a Hill equation $y'' + q(t)y = 0$ has no conjugate points then for every two distinct $a, b \in \mathbb{R}$ there exists a unique solution $y(t; a, b)$ satisfying

$$y(a; a, b) = 1, \quad y(b; a, b) = 0.$$

By uniqueness, one has the relation for distinct a, a' :

$$(15) \quad y(t; a, b) = y(a'; a, b)y(t; a', b).$$

Using disconjugacy, one can show that a limiting solution exists and is positive everywhere:

$$y(t; a) := \lim_{b \rightarrow +\infty} y(t; a, b).$$

These positive solutions are π -quasi-periodic. Indeed, setting $a' \mapsto a + \pi, t \mapsto t + \pi$ in equation (15) and passing to the limit $b \rightarrow +\infty$, we get

$$(16) \quad y(t + \pi; a) = y(a + \pi; a)y(t + \pi; a + \pi) = y(a + \pi; a)y(t; a),$$

where the last equality is due to the π -periodicity of $q(t)$. Thus, $y(t; a)$ is π -quasi-periodic with multiplier $\mu = y(a + \pi; a)$, as needed.

In the opposite direction the claim is obvious: if $y'' + q(t)y = 0$ admits a positive solution then, by the Sturm Oscillation Theorem, any non-trivial solution has no conjugate points. □

3. SELF-BÄCKLUND CURVES: FIRST STUDY

3.1. Infinitesimal deformations of centroaffine conics. In this section we study infinitesimal deformations of centroaffine conics in the class of self-Bäcklund centroaffine curves. (This includes, as we recall from Section 1, the requirement for π -anti-periodicity). We describe the values of the parameter α for which centroaffine conics admit non-trivial infinitesimal deformations. Later, in Section 4.3, we shall show that these values of α are realized by actual deformations, see Corollary 4.20.

Here is a brief reminder about deformations. Let $\gamma(t)$ be a self-Bäcklund centroaffine curve, satisfying

$$(17) \quad [\gamma, \gamma'] = 1, \quad [\gamma(t), \gamma(t + \alpha)] = c,$$

for some constants α, c . A *deformation* of such a curve, within the class of self-Bäcklund centroaffine curves, is a function $\tilde{\gamma}(t, \varepsilon)$ defined on $\mathbb{R} \times (-\varepsilon_0, \varepsilon_0)$ for some $\varepsilon_0 > 0$, and functions $\tilde{\alpha}(\varepsilon), \tilde{c}(\varepsilon)$ defined on $(-\varepsilon_0, \varepsilon_0)$, satisfying equation (17) for each fixed ε , namely

$$(18) \quad \left[\tilde{\gamma}, \frac{\partial}{\partial t} \tilde{\gamma} \right] = 1, \quad [\tilde{\gamma}(t, \varepsilon), \tilde{\gamma}(t + \tilde{\alpha}(\varepsilon), \varepsilon)] = \tilde{c}(\varepsilon),$$

and such that $\gamma = \tilde{\gamma}(\cdot, 0)$, $\alpha = \tilde{\alpha}(0)$ and $c = \tilde{c}(0)$.

An *infinitesimal deformation* of γ is a formal expression $\tilde{\gamma} = \gamma(t) + \varepsilon\gamma_1(t)$, satisfying equation (18) for each ε , modulo ε^2 , for some $\tilde{\alpha} = \alpha + \varepsilon\alpha_1$, $\tilde{c} = c + \varepsilon c_1$. Clearly, if $\tilde{\gamma}$ is a deformation of γ , then its first jet, $\gamma + \varepsilon \frac{\partial}{\partial \varepsilon} \Big|_{\varepsilon=0} \tilde{\gamma}$, is an infinitesimal deformation of γ . However, the converse is not necessarily true, that is, given an infinitesimal deformation $\gamma + \varepsilon\gamma_1$, it is not clear *a priori* that there exists an ‘actual’ deformation $\tilde{\gamma}$ of γ such that $\gamma_1 = \frac{\partial}{\partial \varepsilon} \Big|_{\varepsilon=0} \tilde{\gamma}$.

An infinitesimal deformation is *trivial* if it is induced by a shift of the argument, $\tilde{\gamma}(t, \varepsilon) = \gamma(t + \varepsilon\alpha)$, or by the action of $SL_2(\mathbb{R})$, $\tilde{\gamma}(t, \varepsilon) = e^{\varepsilon A} \gamma(t)$, $A \in \mathfrak{sl}_2(\mathbb{R})$.

Theorem 3. *Let $\gamma(t) = (\cos t, \sin t)$. Then*

- (1) *A non-trivial infinitesimal deformation of γ within the class of self-Bäcklund π -anti-periodic centroaffine curves exists if and only if $\tilde{\alpha} = \alpha + \varepsilon\alpha_1$ where $\alpha = \pi/2$, or $\alpha \neq \pi/2$ and α satisfies the equation*

$$(19) \quad \tan(k\alpha) = k \tan \alpha$$

for some integer $k \geq 4$.

- (2) *For $k \geq 2$, there are exactly $k - 2$ solutions of equation (19) in the interval $(0, \pi)$, counting also $\alpha = \pi/2$ as a solution for k odd.*

Proof. (1) We make calculations mod ε^2 . The first equation of (18) means that γ_1 is a vector field along γ , hence $\gamma_1 = -(1/2)f'\gamma + f\gamma'$ for a π -periodic function $f(t)$, see equation (1). The second equation of (18) implies

$$(20) \quad [\gamma_1(t), \gamma(t + \alpha)] + [\gamma(t), \gamma_1(t + \alpha)] + \alpha_1[\gamma(t), \gamma'(t + \alpha)] = c_1.$$

For $\gamma(t) = (\cos t, \sin t)$ we have

$$(21) \quad \begin{aligned} [\gamma(t), \gamma(t + \alpha)] &= \sin \alpha, & [\gamma'(t), \gamma(t + \alpha)] &= -\cos \alpha, \\ [\gamma(t), \gamma'(t + \alpha)] &= \cos \alpha, \end{aligned}$$

hence (20) becomes

$$[\gamma_1(t), \gamma(t + \alpha)] + [\gamma(t), \gamma_1(t + \alpha)] = c_1 + \alpha_1 \cos \alpha = \text{const.}$$

It follows that

$$\begin{aligned} &[-\frac{1}{2}f'(t)\gamma(t) + f(t)\gamma'(t), \gamma(t + \alpha)] \\ &+ [\gamma(t), -\frac{1}{2}f'(t + \alpha)\gamma(t + \alpha) + f(t + \alpha)\gamma'(t + \alpha)] = \text{const.} \end{aligned}$$

In view of equation (21), this implies

$$(22) \quad \frac{1}{2} [f'(t) + f'(t + \alpha)] \sin \alpha - [f(t + \alpha) - f(t)] \cos \alpha = \text{const.}$$

Since the integral of the left hand side over the period is zero, the constant on the right hand side is also zero.

Recall that f is a π -periodic function and let

$$f(t) = \sum_{k=-\infty}^{\infty} a_k e^{2ikt}$$

be its Fourier expansion, with $a_{-k} = \bar{a}_k$. Then

$$\begin{aligned} f'(t) &= 2i \sum_{k=-\infty}^{\infty} k a_k e^{2ikt}, & f(t + \alpha) &= \sum_{k=-\infty}^{\infty} a_k e^{2ik\alpha} e^{2ikt}, \\ f'(t + \alpha) &= 2i \sum_{k=-\infty}^{\infty} k a_k e^{2ik\alpha} e^{2ikt}. \end{aligned}$$

Substitute this in equation (22) to conclude that

$$a_k [ik(1 + e^{2ik\alpha}) \sin \alpha - (e^{2ik\alpha} - 1) \cos \alpha] = 0$$

for each k . Hence $a_k = 0$, unless

$$ik(1 + e^{2ik\alpha}) \sin \alpha = (e^{2ik\alpha} - 1) \cos \alpha,$$

or

$$k \frac{e^{ik\alpha} + e^{-ik\alpha}}{2} \sin \alpha = \frac{e^{ik\alpha} - e^{-ik\alpha}}{2i} \cos \alpha,$$

that is, $k \tan \alpha = \tan(k\alpha)$.

Conversely, if equation (19) holds, then one can choose $f(t)$ to be a pure harmonic of order $2k$, and then equation (18) holds modulo ε^2 . Likewise, if $\alpha = \pi/2$, one can choose $g(t)$ to be a pure harmonic of order $2k$ with odd $k \geq 3$ or a linear combination of such harmonics.

Note that equation (19) holds trivially for $k = 0$ and $k = 1$. The former case corresponds to $f(t)$ being constant, a shift of the argument of $\gamma(t)$. The latter case corresponds to the action of $\mathfrak{sl}(2, \mathbb{R})$, a stretching of the unit circle to an ellipse bounding area π .

For $k = 2$ there are no solutions $\alpha \in (0, \pi)$ to equation (19) and for $k = 3$ the only solution is $\alpha = \pi/2$ (see next item).

(2) See Proposition 2 of [27], or Lemma 4.8 of [11].

□

Remark 3.1. Equation (19) appeared in the context of bicycle kinematics in [11,41] and in the papers by Wegner, summarized in [47]. It also appeared in [27] in the context of billiards and flotation problems, and in [8], [9], [10] for magnetic, outer and wire billiards. This ubiquitous equation has a countable number of solutions but, except for $\pi/2$, there are no π -rational solutions [18].

3.2. **Periods 3 and 4.**

Theorem 4. *Let $\gamma(t)$ be a π -anti-periodic self-Bäcklund centroaffine curve, that is, $[\gamma(t), \gamma(t + \alpha)] = c \neq 0$. If $\alpha = \pi/3$ or $\alpha = \pi/4$ then γ is a centroaffine ellipse.*

Proof. Consider the case of $\alpha = \pi/3$. Let us use the shorthand notation

$$\gamma(t) = \gamma_0, \quad \gamma\left(t + \frac{\pi}{3}\right) = \gamma_1, \quad \gamma\left(t + \frac{2\pi}{3}\right) = \gamma_2.$$

Then

$$[\gamma_0, \gamma_1] = [\gamma_1, \gamma_2] = [\gamma_2, -\gamma_0] = c,$$

hence $[\gamma_0, \gamma_2] = [\gamma_0, \gamma_1]$, and the vector $\gamma_1 - \gamma_2$ is collinear with γ_0 . Likewise, $\gamma_2 + \gamma_0$ is collinear with γ_1 , and $\gamma_1 - \gamma_0$ with γ_2 . We write

$$\gamma_1 - \gamma_2 = \varphi_0 \gamma_0, \gamma_2 + \gamma_0 = \varphi_1 \gamma_1, \gamma_1 - \gamma_0 = \varphi_2 \gamma_2.$$

Since $[\gamma_0, \gamma_1] \neq 0$, the linear map $\mathbb{R}^3 \rightarrow \mathbb{R}^2, (x_0, x_1, x_2) \mapsto \sum x_i \gamma_i$, has rank 2, hence nullity 1. It follows that the matrix

$$\begin{bmatrix} -\varphi_0 & 1 & -1 \\ 1 & -\varphi_1 & 1 \\ -1 & 1 & -\varphi_2 \end{bmatrix}$$

has rank 1, hence $\varphi_0 = \varphi_1 = \varphi_2 = 1$. Thus $\gamma_2 = \gamma_1 - \gamma_0$.

It follows that $\gamma'_2 = \gamma'_1 - \gamma'_0$, and hence

$$1 = [\gamma_2, \gamma'_2] = [\gamma_1 - \gamma_0, \gamma'_1 - \gamma'_0] = 2 - [\gamma_0, \gamma'_1] + [\gamma'_0, \gamma_1].$$

Since $[\gamma_0, \gamma_1] = c$, one has $[\gamma'_0, \gamma_1] + [\gamma_0, \gamma'_1] = 0$. This implies that

$$[\gamma_0, \gamma'_1] = \frac{1}{2}, \quad [\gamma'_0, \gamma_1] = -\frac{1}{2},$$

and hence $\gamma_1 = (1/2)\gamma_0 + c\gamma'_0$.

It follows that in equation (3) one has $f = 1/2$, and hence, by Lemma 2.1, $c^2 p = -3/4$. That is, p is constant, which implies $p = -1$ and $c = \sqrt{3}/2$, and therefore the curve is a centroaffine conic.

The case $\alpha = \pi/4$ is similar. In analogous notations, one has

$$[\gamma_0, \gamma_1] = [\gamma_1, \gamma_2] = [\gamma_2, \gamma_3] = [\gamma_3, -\gamma_0] = c,$$

hence

$$\gamma_0 \sim \gamma_1 - \gamma_3, \gamma_1 \sim \gamma_0 + \gamma_2, \gamma_2 \sim \gamma_1 + \gamma_3, \gamma_3 \sim -\gamma_0 + \gamma_2.$$

This implies

$$(23) \quad \gamma_1 = g(\gamma_0 + \gamma_2), \quad \gamma_3 = g(\gamma_2 - \gamma_0)$$

for some function $g(t)$.

Since $[\gamma_1, \gamma'_1] = [\gamma_3, \gamma'_3] = 1$, equation (23) implies

$$2g^2 = 1, \quad [\gamma_0, \gamma'_2] + [\gamma_2, \gamma'_0] = 0.$$

But $[\gamma_0, \gamma_2] = c$, hence $[\gamma'_0, \gamma_2] + [\gamma_0, \gamma'_2] = 0$, and therefore $[\gamma'_0, \gamma_2] = [\gamma_0, \gamma'_2] = 0$. In particular, $\gamma_2 \sim \gamma'_0$.

It follows that $\gamma_1 = (1/\sqrt{2})\gamma_0 + c\gamma'_0$. Then, in equation (3), one has $f = 1/\sqrt{2}$, and hence, by Lemma 2.1, $c^2 p = -1/2$. Thus $p = -1, c = 1/\sqrt{2}$, and the curve is a centroaffine conic. □

Remark 3.2. An analogous result, rigidity for periods 3 and 4, holds for bicycle curves, see [13, 14, 41].

3.3. Period two: Flexibility and Radon curves. In this section we show that self-Bäcklund curves of period two, that is, $\alpha = \pi/2$, exhibit a substantial flexibility. A similar result, for the value of the density $1/2$, was known for a long time for Ulam's flotation in equilibrium problem [7, 50].

Let us construct a self-Bäcklund curve of period two as a closed trajectory of a vector field V on the space of origin-centered parallelograms. Let the vertices be $P_1, P_2, -P_1, -P_2$, and let the vector field have the values $V_1, V_2, -V_1, -V_2$ at these vertices, respectively.

We want the trajectories of the points $P_1, P_2, -P_1, -P_2$ to coincide and to form a self-Bäcklund curve with $\alpha = \pi/2$. Let $(R_1(t), P_2(t))$ be an integral curve of such a vector field. Then $P_2(t) = R_1(t + \pi/2)$. The centroaffine conditions $[P_i, P_i'] = 1$ and the c -relation $[R_1, P_2] = c$ amount to

$$(24) \quad [P_1, V_1] = [P_2, V_2] = 1, \quad [V_1, P_2] + [R_1, V_2] = 0.$$

Note that the area of the parallelogram $(P_1, P_2, -P_1, -P_2)$ remains constant.

Lemma 3.3. *Equation (24) is satisfied if and only if*

$$V_1 = fP_1 + \frac{1}{c}P_2, \quad V_2 = -\frac{1}{c}P_1 - fP_2,$$

where $f(P_1, P_2)$ is an odd function, in the sense that $f(P_2, -P_1) = -f(P_1, P_2)$.

Proof. Write $V_1 = fP_1 + gP_2, V_2 = \bar{f}P_1 + \bar{g}P_2$ and substitute into equation (24), using $[P_1, P_2] = c$, to obtain $f + \bar{g} = 0, g = -\bar{f} = 1/c$. That f is odd follows from the central symmetry of the parallelogram. □

Thus one has a functional parameter f to play with. The boundary conditions

$$(25) \quad R_1(0) = (1, 0), \quad R_1\left(\frac{\pi}{2}\right) = P_2(0) = (0, c), \quad P_2\left(\frac{\pi}{2}\right) = -R_1(0) = (-1, 0)$$

impose a finite-dimensional restriction on the function f . As a result, we obtain a functional space of self-Bäcklund curves of period two.

For example, if f is identically zero and $c = 1$, then $P_1'' = P_2' = -P_1$, and the curve is a centroaffine ellipse. See Figure 10 for a non-trivial example. In Example 4.11 (Figure 16) we construct explicitly many analytic curves.

Remark 3.4. The space of origin-centered parallelograms of a fixed area is identified with $SL_2(\mathbb{R})$. If $P = (p_1, p_2), Q = (q_1, q_2)$, then the first equation (24), $[P, U] = [Q, V]$, means that the curve under consideration is tangent to the kernel of the 1-form $p_1 dp_2 - p_2 dp_1 + q_2 dq_1 - q_1 dq_2$. This form defines a contact structure on $SL_2(\mathbb{R})$, and the curve is Legendrian.

Let Γ be a smooth closed convex curve, symmetric with respect to the origin. Let $x, y \in \Gamma$. One says that y is Birkhoff orthogonal to x if y is parallel to the tangent line to Γ at x . This relation is not necessarily symmetric; if it is symmetric, then Γ is called a *Radon curve*. Radon curves comprise a functional space, with ellipses providing a trivial example.

Radon curves have been thoroughly studied since their introduction more than 100 years ago; see [32] for a modern treatment.




FIGURE 10. A self-Bäcklund curve with rotation angle $\alpha = \pi/2$ and $c = 1$, using Lemma 3.3 and equation (25), where $f(P_1, P_2) = u(P_1)u(P_2)$ and $u(x, y) = 1.2x - 4x^3 - 4x^5$ (approximately)

Let Γ be a Radon curve, $x \in \Gamma$ be a point, and $y \in \Gamma$ be its Birkhoff orthogonal. Then the tangent lines at points $x, y, -x, -y$ form a parallelogram circumscribed about Γ . As x traverses Γ , the vertices of the parallelogram describe a curve γ . The latter curve is an invariant curve of the outer billiard transformation about Γ , see Remark 2.6.

The relation of self-Bäcklund curves with Radon curves is as follows. Let γ be a self-Bäcklund curve with rotation number $\pi/2$, then the points $\gamma(t), \gamma(t + \pi/2), \gamma(t + \pi), \gamma(t + 3\pi/2)$ form a parallelogram. Therefore the middle curve Γ is a Radon curve. Example 4.11 provides analytic families of Radon curves.

4. SELF-BÄCKLUND CURVES AND THE LAMÉ EQUATION

4.1. Traveling wave solutions of KdV and Wegner’s ansatz. The first two in the hierarchy of integrals of the Korteweg-de Vries equation are the functionals

$$(26) \quad \int p(t) dt, \int p^2(t) dt$$

on centroaffine curves. In particular, KdV is the Hamiltonian flow of the former functional with respect to the symplectic form $\int [V_f, V_g] dt$, where we use formula (1) for tangent vector fields [38].

Consider a centroaffine curve that is a relative extremum of the second functional (26) subject to the constraint given by the first one. Lemma 4.1 is well known and we do not present its proof, see [21].

Lemma 4.1. *These relative extrema are characterized by the differential equation on the centroaffine curvature*

$$(27) \quad p''' = 6pp' + ap',$$

where a is a Lagrange multiplier.

Equation (27) describes traveling wave solutions of KdV, see [21]. For the centroaffine curves satisfying equation (27), the KdV evolution is described by the equation $\dot{p} = ap'$, that is, by a parameter shift of the curvature $p(t)$. Two centroaffine curves with the same curvature function differ by an element of $SL_2(\mathbb{R})$. Therefore these curves evolve in time by special linear transformations.

Equation (27) can be integrated to

$$(28) \quad (p')^2 = 2p^3 + ap^2 + 2bp + c,$$

where a, b, c are constants.

Lemma 4.2. *The curves described in Section 2.2 satisfy equation (27).*

Proof. Let $q(t)$ be the centroaffine curvature of the curve $f\gamma + c\gamma'$ where γ is a unit circle and f satisfies equation (4). Then

$$q = \frac{2}{c^2}(f^2 - 1) - 1,$$

see Lemma 3.1 in [44] for this calculation. Hence

$$q' = \frac{4ff'}{c^2} = \frac{4f}{c^2} \left(\frac{f^2}{c} + c - \frac{1}{c} \right).$$

One needs to check that $(q')^2 = 2q^3 + aq^2 + 2bq + c$ for some constants a, b, c . One has

$$(q')^2 = \frac{16f^2}{c^4} \left(\frac{f^2}{c} + c - \frac{1}{c} \right)^2$$

a cubic polynomial in f^2 with the leading coefficient $16/c^6$. The same holds for $2q^3 + aq^2 + 2bq + c$, so one can choose the coefficients a, b, c as needed. \square

Now we develop a centroaffine analog of F. Wegner's approach to 2-dimensional bodies that float in equilibrium in all positions (or bicycle curves) [47–49].

Consider a centroaffine curve $\gamma(t) = (r(t) \cos \alpha(t), r(t) \sin \alpha(t))$. The centroaffine condition $[\gamma, \gamma'] = 1$ is satisfied if $\alpha' = r^{-2}$. We use prime to denote the derivative with respect to t ; the derivative with respect to α is denoted as r_α .

Emulating Wegner's approach and using material of Section 2.1, fix a small ε and consider the curves $\Gamma_\pm = \gamma \pm \varepsilon\gamma'$. These curves are 2ε -related. We want them to be obtained from the same curve, Γ , by rotating it through small angles $\pm\delta$. The assumption is that δ is of order ε^3 ; all the calculations below are mod ε^4 . We use the notations in Figure 11.

Lemma 4.3. *One has:*

$$\varphi = \tan^{-1} \left(\frac{\varepsilon}{r^2 + \varepsilon r r'} \right), \quad \rho = \sqrt{r^2 + 2\varepsilon r r' + \varepsilon^2(r^{-2} + r'^2)}.$$

Proof. One has $|\gamma'| = r^{-1}\sqrt{1 + r^2 r'^2}$, hence $|AB_+| = \varepsilon r^{-1}\sqrt{1 + r^2 r'^2}$. Next, $1 = [\gamma, \gamma'] = |\gamma||\gamma'| \sin \psi$, hence

$$\sin \psi = \frac{1}{\sqrt{1 + r^2 r'^2}}, \quad \cos \psi = -\frac{r r'}{\sqrt{1 + r^2 r'^2}}.$$

Then

$$\tan \varphi = \frac{|AB_+| \sin \psi}{|OA| - |AB_+| \cos \psi} = \frac{\varepsilon}{r^2 + \varepsilon r r'}.$$

Finally, by the cosine rule,

$$|OB_+|^2 = |OA|^2 + |AB_+|^2 - 2|OA||AB_+| \cos \psi = r^2 + 2\varepsilon r r' + \varepsilon^2(r^{-2} + r'^2),$$

as claimed. \square




FIGURE 11. Notation for Lemma 4.3: $r = |OA|$, $\rho = |OB_-| = |OB| = |OB_+|$, $\varphi = \angle AOB_+$, $\psi = \angle OAB_+$, $\delta = \angle BOB_+ = \angle B_-OB$. γ and Γ are given in polar coordinates by $r(\alpha)$ and $\rho(\beta)$ (respectively).

Thus we have an equation for Γ in polar coordinates:

$$(29) \quad \rho(\beta) = \rho(\alpha + \varphi - \delta) = \sqrt{r^2 + 2\varepsilon r r' + \varepsilon^2(r^{-2} + r'^2)},$$

where φ is given in Lemma 4.3, and where $\delta = c\varepsilon^3$ with c being a constant.

To solve equation (29), consider the cubic Taylor polynomials of both sides and equate the even and odd parts separately (since the equation holds for $\pm\varepsilon$). One has

$$\begin{aligned} \varphi &= \varepsilon r^{-2} - \varepsilon^2 r^{-3} r' + \varepsilon^3 \left(r^{-4} r'^2 - \frac{1}{3} r^{-6} \right), \\ \varphi^2 &= \varepsilon^2 r^{-4} - 2\varepsilon^3 r^{-5} r', \quad \varphi^3 = \varepsilon^3 r^{-6}, \\ \sqrt{r^2 + 2\varepsilon r r' + \varepsilon^2(r^{-2} + r'^2)} &= r + \varepsilon r' + \frac{\varepsilon^2}{2} r^{-3} - \frac{\varepsilon^3}{2} r^{-4} r'. \end{aligned}$$

To expand the left hand side of equation (29), we calculate ρ_α , $\rho_{\alpha\alpha}$ and $\rho_{\alpha\alpha\alpha}$, using $\alpha' = r^{-2}$:

$$\rho_\alpha = r^2 \rho', \quad \rho_{\alpha\alpha} = 2r^3 r' \rho' + r^4 \rho'', \quad \rho_{\alpha\alpha\alpha} = 6r^4 r'^2 \rho' + 2r^5 r'' \rho' + 6r^5 r' \rho'' + r^6 \rho''''.$$

Now we have for the left hand side of equation (29)

$$\begin{aligned} \rho(\alpha + \varphi - \delta) &= \rho + \varphi \rho_\alpha + \frac{1}{2} \varphi^2 \rho_{\alpha\alpha} + \frac{1}{6} \varphi^3 \rho_{\alpha\alpha\alpha} - \delta \rho_\alpha \\ &= \rho + \varepsilon \rho' + \frac{1}{2} \varepsilon^2 \rho'' + \frac{1}{6} \varepsilon^3 (r^{-2} \rho'''' + 2r^{-1} r'' \rho' - 2r^{-4} \rho') - c\varepsilon^3 r^2 \rho'. \end{aligned}$$

Thus

$$\begin{aligned} \rho + \frac{1}{2} \varepsilon^2 \rho'' &= r + \frac{1}{2} \varepsilon^2 r^{-3}, \\ \rho' + \frac{1}{6} \varepsilon^2 (\rho'''' + 2r^{-1} r'' \rho' - 2r^{-4} \rho' - 6c r^2 \rho') &= r' - \frac{1}{2} \varepsilon^2 r^{-4} r'. \end{aligned}$$

Differentiate the first equation and subtract from the second one, setting, following Wegner, $\rho = r$ (since ε is infinitesimal), to obtain

$$r'''' - r^{-1} r' r'' + 4r^{-4} r' + 3c r^2 r' = 0.$$

Multiply this by r^{-1} and write it as

$$\left(r^{-1}r'' - r^{-4} - \frac{3}{2}cr^2\right)' = 0,$$

or

$$r'' - r^{-3} + \frac{3}{2}cr^3 - br = 0,$$

where b is a constant. Multiply this by $2r'$ and write it as

$$\left(r'^2 + r^{-2} + \frac{3}{4}cr^4 - br^2\right)' = 0.$$

Hence

$$r'^2 = -r^{-2} - \frac{3}{4}cr^4 + br^2 + a,$$

where a is another constant. Multiply by $4r^2$ to obtain

$$4r^2r'^2 = -4 - 3cr^6 + 4br^4 + ar^2.$$

Finally, setting $R = r^2$ and renaming the constants, we obtain the differential equation

$$(30) \quad R'^2 = aR^3 + bR^2 + cR - 4.$$

Thus $R(t)$ is an elliptic function. The curve is given by a parametric equation

$$(31) \quad \Gamma(t) = (R(t)^{1/2} \cos \alpha(t), R(t)^{1/2} \sin \alpha(t))$$

with R as in equation (30) and $\alpha' = R^{-1}$.

Remark 4.4. If the curve is a centroaffine ellipse, one has $a = 0$ in equation (30).

Concerning the centroaffine curvature of this curve, it is also an elliptic function.

Lemma 4.5. *One has*

$$p(t) = \frac{1}{2}aR(t) + \frac{1}{4}b.$$

Proof. Differentiating equation (31) twice, we find that

$$p = -\frac{1}{4}R^{-2}(R'^2 + 4) + \frac{1}{2}R^{-1}R''.$$

Differentiating equation (30), we obtain

$$R'' = \frac{3}{2}aR^2 + bR + \frac{1}{2}c.$$

Substitute this and equation (30) in the above formula for p to obtain the result. □

Renaming the constants again, we obtain from equation (30)

$$p'^2 = 2p^3 + ap^2 + bp + c,$$

which coincides with equation (28).

Let us also calculate the (Euclidean) curvature k of a curve satisfying equation (30).

Lemma 4.6. *One has*

$$k = -\frac{4aR + 2b}{(aR^2 + bR + c)^{\frac{3}{2}}}.$$

Proof. Since t is the centroaffine parameter, we have for the curvature

$$k = \frac{[\gamma', \gamma'']}{|\gamma'|^3} = \frac{-p(t)}{|\gamma'|^3}.$$

We have

$$|\gamma'| = \sqrt{r'^2 + r^2\alpha'^2} = \sqrt{\frac{R'^2}{4R} + \frac{1}{R}} = \sqrt{\frac{R'^2 + 4}{4R}} = \frac{\sqrt{aR^2 + bR + c}}{2}.$$

Hence

$$k = \frac{-8p(t)}{\sqrt{aR^2 + bR + c}^3} = -\frac{4aR + 2b}{(aR^2 + bR + c)^{\frac{3}{2}}}.$$

□

Thus the curvature is a function of the distance from the origin. This is a special class of curves, studied in [16, 40]. One can think of these curves as the trajectories of a charge in a rotationally symmetric magnetic field whose strength is a function of the distance from the origin. Note that Wegner's curves also have this property: their curvature satisfies $k = ar^2 + b$, where a, b are constants.

Likewise one can interpret equation $\gamma'' = p\gamma$ as Newton's Second Law, that is, $\gamma(t)$ is the trajectory of a point-mass in a central force field whose potential V is rotationally symmetric. By Lemma 4.5, and renaming the constants, one has $V(r) = ar^4 + br^2 + c$. Using conservation of energy and momentum, one can solve the equation of motion in quadratures.

Remark 4.7. Consider a particular case when V is a pure 4th power of the distance, that is, the force is proportional to r^3 . According to a corollary of the Bohlin theorem, see Theorem 5, Appendix 1 in [4], some trajectories in this field are the images of straight lines under the conformal transformation $w = z^{1/3}$. These are cubic curves, see Figure 12.




FIGURE 12. The curve $2(x^3 - 3xy^2) - 5(3x^2y - y^3) + 1 = 0$, the image of the line $2a - 5b + 1 = 0$ under the conformal transformation $w = z^{1/3}$

4.2. Self-Bäcklund curves as solutions of the Lamé equation. In this section we give an explicit construction of a large family of self-Bäcklund curves, given by the Wegner ansatz of Section 4.1. We shall make frequent use of standard facts about the Weierstrass elliptic functions \wp, ζ, σ , such as: the addition formulas [1, pages 40-41], quasi-periodicity properties [1, pages 35-37], reality conditions [37, pages 29-32], degenerate cases of Weierstrass functions [1, pages 201]. We shall also use applications of elliptic functions to the Lamé equation which can be found in [37, pages 48-54].

4.2.1. *Constructing the curves.* Our starting point is equation (28),

$$(p')^2 = 2p^3 + ap^2 + 2bp + c,$$

for the curvature $p(t)$ of the self-Bäcklund curves suggested by the Wegner's ansatz. Comparing this equation to the equation satisfied by the Weierstrass \wp function,

$$(32) \quad (\wp')^2 = 4\wp^3 - g_2\wp - g_3,$$

we conclude that $p(t)$ is given, in terms of \wp , by

$$(33) \quad p(t) = 2\wp(t + \omega') + C.$$

Here \wp is the Weierstrass function with half periods ω, ω' , where the first one is real and the second one is pure imaginary, see Figure 13. Since $p(t)$ needs to be periodic, we are in the case of three real roots $e_1 > e_2 > e_3$ of the right hand side of equation (32). In formula (33) the shift of the argument by ω' is performed in order to get a real, smooth, 2ω -periodic potential $p(t)$.




FIGURE 13. The Weierstrass function $\wp(z)$ with real invariants and fundamental half periods $\omega \in \mathbb{R}, \omega' \in i\mathbb{R}$. (a) The fundamental rectangle in the z plane. The boundary of the rectangle $(0, \omega', \omega + \omega', \omega)$ is mapped by \wp onto the extended real axis $\mathbb{R} \cup \{\infty\}$. (b) The phase plane of $(\wp')^2 = 4(\wp - e_1)(\wp - e_2)(\wp - e_3)$. (c) The line $\{t + \omega' | t \in \mathbb{R}\}$ is mapped, 2ω -periodically, onto the segment $[e_3, e_2]$.

The constant C can be written as $C = \wp(a)$ for some $a \in \mathbb{C}$. Thus

$$(34) \quad p(t) = 2\wp(t + \omega') + \wp(a).$$

We write our curve in complex form $X(t) = x(t) + iy(t)$, satisfying

$$(35) \quad X'' + (-\wp(a) - 2\wp(t + \omega'))X = 0,$$

which is precisely the Lamé equation (equation (6) of [1, page 186]).

In order to construct a centroaffine π -anti-periodic curve, we shall require the following:

- (1) The Wronskian $[X, X'] = 1$. This can be achieved by rescaling of any solution of equation (35) satisfying $[X, X'] = \text{const} > 0$ (see item (4) of Proposition 4.8).
- (2) $\omega = \pi/2k$ for some integer $k \geq 2$, so that p is π/k -periodic.
- (3) The solution X is rotated over the period 2ω by $\pi n/k$, where $0 < n < k$ is odd and co-prime to k , so that after k periods we have $X(t + \pi) = -X(t)$. In other words, we require $X(t)$ to be a complex 2ω -quasi-periodic solution of equation (35), with Floquet multiplier $\mu = e^{i\pi n/k}$:

$$X(t + 2\omega) = X(t)e^{i\pi n/k}.$$

A basis X_+, X_- for the solutions of the Lamé equation (35) can be written in the following form (see [1, page 37]):

$$(36) \quad X_{\pm}(t) = e^{-t\zeta(\pm a)} \frac{\sigma(\pm a + t + \omega')\sigma(\omega')}{\sigma(\pm a + \omega')\sigma(t + \omega')},$$

where ζ, σ are the Weierstrass zeta and sigma functions, respectively.

The construction of the self-Bäcklund curves in this section boils down to a careful choice of the parameter a in equation (35).

Proposition 4.8. *For every $a \in (0, \omega') \cup (\omega, \omega + \omega')$,*

- (1) $\wp(a)$ is real, hence the potential $2\wp(t + \omega') + \wp(a)$ in the Lamé equation (35) is real as well.
- (2) $X_+(t)$ is a regular curve, that is, $X'_+(t) \neq 0$ for all t .
- (3) $X_+(0) = 1$ and $X'_+(0) = ib$ for some $b \in \mathbb{R}, b > 0$.
- (4) $X_+(t)$ is locally star-shaped and positively oriented:

$$[X_+(t), X'_+(t)] = \text{const} > 0.$$

- (5) $X_+(t + 2\omega) = X_+(t)e^{2f(a)}$, where

$$(37) \quad f(a) := a\zeta(\omega) - \omega\zeta(a).$$

That is, $X_+(t)$ is a 2ω -quasi-periodic solution of equation (35) with a Floquet multiplier $\mu = e^{2f(a)}$.

- (6) The function f of the previous item satisfies the identities

$$f(-a) = -f(a), \quad f(a + 2\omega) = f(a), \quad f(a + 2\omega') = f(a) + i\pi.$$

Proof. (1) See pages 31-32 of [37].

(2) Differentiating equation (36), and using $\zeta = \sigma'/\sigma$ and the addition formula for ζ , we compute:

$$X'_+(t) = X_+(t) [\zeta(a + t + \omega') - \zeta(a) - \zeta(t + \omega')] = X_+(t) \frac{\wp'(a) - \wp'(t + \omega')}{2[\wp(a) - \wp(t + \omega')]}.$$

Notice that the numerator in the last fraction cannot vanish, since $\wp'(t + \omega')$ is real and $\wp'(a)$ is purely imaginary, both non-vanishing (\wp' vanishes in the fundamental rectangle only at $0, \omega, \omega', \omega + \omega'$). It follows that $X'_+(t)$ does not vanish.

(3) Substituting $t = 0$ into equation (36) gives $X_+(0) = 1$. From the previous item we have

$$X'_+(0) = \frac{\wp'(a)}{2(\wp(a) - e_3)}.$$

For $a \in (0, \omega') \cup (\omega, \omega + \omega')$ the numerator $\wp'(a)$ is purely imaginary and the denominator is real, both non-vanishing. Hence we can write $X'_+(0) = ib$, $b \in \mathbb{R}$, $b \neq 0$. Moreover, $\wp(a) < e_3$ and $\text{Im}[\wp'(a)] < 0$ for $a \in (0, \omega')$. When $a \in (\omega, \omega + \omega')$ we have that $\wp(a) > e_3$ is positive and $\text{Im}[\wp'(a)] > 0$. (All this is evident in Figure 13.) Hence, in both cases, $b > 0$.

(4) Since X_+ is a solution of the Lamé equation (35), which has no X' term, one has

$$\text{Wronskian} = [X_+(t), X'_+(t)] = \text{const.}$$

The constant must be positive, due to item (2).

(5) See [37, page 52].

(6) See [37, page 86].

□

Remark 4.9. Following Proposition 4.8 (item (4)) and the proof of item (3), we can normalize the solutions of the Lamé equation (35) given by formula (36) by the constant factor

$$N := \sqrt{|X'_\pm(0)|} = \sqrt{\frac{\wp'(a)}{2i(\wp(a) - e_3)}},$$

so that the normalized solutions $Y_\pm(t) := \frac{1}{N}X_\pm(t)$ satisfy the centroaffine condition $[Y(t), Y'(t)] = 1$.

Next, due to requirement (3) and Proposition 4.8 (item (5)), we need to solve $2f(a) \equiv i\pi n/k \pmod{2\pi i}$, or

$$(38) \quad f(a) = \frac{i\pi n}{2k} + i\pi m,$$

for some integers $m, n \in \mathbb{Z}$, where n is odd, relatively prime to k , and $0 < n < k$.

To solve equation (38), it is enough to restrict a to the fundamental rectangle. Indeed, if a_1 and a_2 are two congruent solutions of equation (38), then the corresponding potentials (34) of the Lamé equation are equal, and the curves constructed by formula (36) are equivalent under the action of $\text{SL}_2(\mathbb{R})$.

One may further restrict to solutions of equation (38) where a belongs to one of the segments $(0, \omega')$ or $(\omega, \omega + \omega')$, and $m \geq 0$. This follows from the properties of f listed in Proposition 4.8 and the monotonicity property of f on the segments $[0, 2\omega']$ and $[\omega, \omega + 2\omega']$. On the segment $[0, 2\omega']$ the function f varies monotonically from $+\infty$ to $-\infty$. On the segment $[\omega, \omega + 2\omega']$ it varies from 0 to $i\pi$.

Theorem 5. Consider equation (38) for fixed integers k, n , where $k \geq 2$ and n is odd, relative prime to k , and $0 < n < k$. Then

- (1) For each integer $m \geq 0$ there is a unique solution $a_m \in (0, \omega') \cup (\omega, \omega + \omega')$.
- (2) For $m > 0$, $a_m \in (0, \omega')$.
- (3) For $m = 0$, $a_0 \in (\omega, \omega + \omega')$.

- (4) The sequence $\lambda_m(\mu) := -\wp(a_m)$ is strictly monotone increasing and, in particular, the value $\lambda_0(\mu) = -\wp(a_0)$ is the smallest one.

Proof. The proof of items (1)–(3) uses the behavior of the function f . Since $\frac{\pi n}{2k} < \frac{\pi}{2}$, for $m = 0$ there is a unique solution a_0 in the segment $[\omega, \omega + \omega']$, because f is pure imaginary on $[\omega, \omega + \omega']$ and varies monotonically from 0 at ω to $i\pi/2$ at $\omega + \omega'$.

For $m > 0$, one can find a unique a_m in the segment $[0, \omega']$ since there f is pure imaginary, varying monotonically from $+i\infty$ at 0 to $i\pi/2$ at ω' . Moreover, the sequence a_m is monotone decreasing on $[0, \omega']$.

In order to prove (4), notice that on the segment $[0, \omega']$ the function \wp is real-valued and monotone increasing from $-\infty$ to e_3 . Hence $-\wp(a_m)$ is monotone increasing for $m \geq 1$. Moreover, $-\wp(a_m) > -e_3$ for every $m \geq 1$. As for $m = 0$,

$$-\wp(a_0) \in (-e_1, -e_2),$$

because on the interval $[\omega, \omega + \omega']$ the function \wp is monotonically decreasing and takes the values e_1, e_2 at the end points, respectively. Since $e_3 < e_2 < e_1$, this proves item (4) (see Figure 13). □

Moreover we have the following result.

Theorem 6. For each k, m, n as in Theorem 5, consider the curve X_+ determined by the value a_m .

- (1) X_+ is locally star-shaped π -anti-periodic curve, with the winding number

$$w = 2k \left\lfloor \frac{m}{2} \right\rfloor + n.$$

- (2) X_+ is embedded (simple) if and only if $m = 0, n = 1$.

Proof. It follows from Theorem 5 that the sequence $\lambda_m(\mu) := -\wp(a_m)$ is the sequence of Floquet eigenvalues for the problem

$$X'' + (\lambda - 2\wp(t + \omega'))X = 0, \quad X(t + 2\omega) = \mu X(t), \quad \mu := e^{i\pi n/k},$$

and that $\lambda_m(\mu)$ is monotone increasing.

It follows from Proposition 4.8 that the curve is locally star-shaped and positively oriented.

In order to compute the winding number of the curve, we need first to see what happens over one period $[0, 2\omega]$. Denote by $y_m(t)$ the imaginary part of the solution X_+ corresponding to a_m . We know by Proposition 4.8 (claim (2)) that at the end points of the period one has

$$y_m(0) = 0, \quad y'_m(0) > 0, \quad y_m(2\omega) = \sin\left(\frac{\pi n}{k}\right) > 0.$$

This implies that the number of zeroes of y_m on $(0, 2\omega]$ is even for every m .

In order to find the number of zeroes of y_m on the interval $(0, 2\omega)$ we use Sturm theory, comparing y_m with the Dirichlet eigenfunctions of the Lamé equation, as follows.

Let us denote by $\Lambda_m, \Psi_m, m \geq 0$, the eigenvalues and eigenfunctions corresponding to Dirichlet boundary conditions of the equation

$$(39) \quad \Psi'' + (\lambda - 2\wp(t + \omega'))\Psi = 0.$$

Thus the eigenfunctions Ψ_m vanish at the end points of the interval $[0, 2\omega]$ and have exactly m zeros in $(0, 2\omega)$.

We claim that the number of zeroes of y_m in $(0, 2\omega)$ is given by the formula:

$$(40) \quad \#\{t \in (0, 2\omega) : y_m(t) = 0\} = 2 \left\lceil \frac{m}{2} \right\rceil.$$

To prove this, we shall consider two cases (see Figure 14):

(1) If $m = 2l$ then $\Lambda_{2l-1} < \lambda_{2l}(\mu) < \Lambda_{2l}$. In this case, the zeroes of Ψ_{2l-1} divide the interval into $2l$ subsegments. In each of them, y_{2l} must vanish somewhere (by Sturm theory). Hence there are at least $2l$ zeroes. In fact, this number must be exactly $2l$, because otherwise it would be at least $2l + 2$ zeros (y_m has an even number of zeroes). But then Ψ_{2l} would have more than $2l$ zeroes.




FIGURE 14. Graph of the function $\Delta(\lambda) := y_1(\lambda, 2\omega) + y'_2(\lambda, 2\omega)$, where $y_1(\lambda, t), y_2(\lambda, t)$ are the basic solutions of equation (39) with $y_1(\lambda, 0) = y'_2(\lambda, 0) = 1, y'_1(\lambda, 0) = y_2(\lambda, 0) = 0$; the positions of the periodic (λ_n), anti-periodic (μ_n), Dirichlet (Λ_n), and Floquet ($\lambda_n(\mu)$) eigenvalues are indicated

(2) If $m = 2l + 1$ then $\Lambda_{2l} < \lambda_{2l+1}(\mu) < \Lambda_{2l+1}$. The zeroes of Ψ_{2l} divide the interval into $2l + 1$ subintervals, in each of which y_{2l+1} must vanish somewhere (by Sturm theory), implying that y_{2l+1} has at least $2l + 1$ zeroes. But then this number is at least $2l + 2$, because it is even. Hence, the number of zeroes of y_{2l+1} is exactly $2l + 2$, because otherwise Ψ_{2l+1} would have more than $2l + 1$ zeroes. This completes the proof of the claim.

As a consequence of formula (40), we see that for $a = a_m$ the solution X_+ makes $\lceil \frac{m}{2} \rceil$ full turns over the period $[0, 2\omega]$, plus an angle of $\frac{\pi n}{k}$, which is a $\frac{n}{2k}$ fraction of a full turn. Altogether, after $2k$ periods, the number of turns is

$$w = 2k \left(\left\lceil \frac{m}{2} \right\rceil + \frac{n}{2k} \right) = 2k \left\lceil \frac{m}{2} \right\rceil + n.$$

This proves the first claim of Theorem 6.

The last formula implies that the curve is simple, that is, $w = 1$, if and only if $m = 0, n = 1$, proving the second claim. This completes the proof. □

4.2.2. *Establishing the self-Bäcklund property.*

Proposition 4.10. *The curve X_+ of equation (36) satisfies the self-Bäcklund property $[X_+(t), X_+(t + \alpha)] = \text{const}$ for a value of the parameter $\alpha \in (0, \pi)$ if and only if*

$$(41) \quad \sigma(a + \alpha) = e^{2\alpha\zeta(a)}\sigma(a - \alpha).$$

Proof. Set $\beta = \alpha/2$. Then equation (2) can be rewritten as

$$\text{Im} \left(X_+(t + \beta)\overline{X_+(t - \beta)} \right) = c,$$

where overline denotes the complex conjugation. We can rewrite this equation as

$$X_+(t + \beta)X_-(t - \beta) - X_-(t + \beta)X_+(t - \beta) = 2c.$$

Next we substitute in the last equation the expressions for X_{\pm} from equation (36):

$$\begin{aligned} 2c = & e^{-(t+\beta)\zeta(a)} \frac{\sigma(a + t + \beta + \omega')\sigma(\omega')}{\sigma(a + \omega')\sigma(t + \beta + \omega')} e^{(t-\beta)\zeta(a)} \frac{\sigma(-a + t - \beta + \omega')\sigma(\omega')}{\sigma(-a + \omega')\sigma(t - \beta + \omega')} \\ & - e^{(t+\beta)\zeta(a)} \frac{\sigma(-a + t + \beta + \omega')\sigma(\omega')}{\sigma(-a + \omega')\sigma(t + \beta + \omega')} e^{-(t-\beta)\zeta(a)} \frac{\sigma(a + t - \beta + \omega')\sigma(\omega')}{\sigma(a + \omega')\sigma(t - \beta + \omega')}. \end{aligned}$$

This can be simplified, using the identity

$$(42) \quad \wp(z) - \wp(w) = -\frac{\sigma(z - w)\sigma(z + w)}{\sigma^2(z)\sigma^2(w)}$$

(see [37, page 25]). We get

$$\begin{aligned} 2c = & e^{-2\beta\zeta(a)} \frac{[\wp(t + \omega') - \wp(a + \beta)]\sigma^2(a + \beta)\sigma^2(\omega')}{[\wp(t + \omega') - \wp(\beta)]\sigma^2(\beta)\sigma(a + \omega')\sigma(-a + \omega')} \\ & - e^{2\beta\zeta(a)} \frac{(\wp(t + \omega') - \wp(a - \beta))\sigma^2(a - \beta)\sigma^2(\omega')}{[\wp(t + \omega') - \wp(\beta)]\sigma^2(\beta)\sigma(a + \omega')\sigma(-a + \omega')}. \end{aligned}$$

Multiplying by the common denominator and renaming the constant,

$$\tilde{c} := 2c\sigma^2(\beta)\sigma(a + \omega')\sigma(-a + \omega')/\sigma^2(\omega'),$$

we get

$$\begin{aligned} \tilde{c} [\wp(t + \omega') - \wp(\beta)] = & e^{-2\beta\zeta(a)} [\wp(t + \omega') - \wp(a + \beta)]\sigma^2(a + \beta) \\ & - e^{2\beta\zeta(a)} [\wp(t + \omega') - \wp(a - \beta)]\sigma^2(a - \beta). \end{aligned}$$

Thus we must have

$$\begin{aligned} \tilde{c} &= e^{-2\beta\zeta(a)}\sigma^2(a + \beta) - e^{2\beta\zeta(a)}\sigma^2(a - \beta), \\ \wp(\beta)\tilde{c} &= e^{-2\beta\zeta(a)}\wp(a + \beta)\sigma^2(a + \beta) - e^{2\beta\zeta(a)}\wp(a - \beta)\sigma^2(a - \beta). \end{aligned}$$

Substituting \tilde{c} from the first identity into the second and simplifying, we get

$$\sigma^2(a + \beta) [\wp(a + \beta) - \wp(\beta)] = e^{4\beta\zeta(a)}\sigma^2(a - \beta) [\wp(a - \beta) - \wp(\beta)].$$

Now, using equation (42) again, we obtain $\sigma(a + \alpha) = e^{2\alpha\zeta(a)}\sigma(a - \alpha)$, as needed. □

Theorem 7 states the self-Bäcklund property of the curves X_+ .

Theorem 7. For each k, m, n as in Theorem 5, the associated curve X_+ satisfies the self-Bäcklund property $[X_+(t), X_+(t + \alpha)] = \text{const}$ for $k - 2$ values of $\alpha \in (0, \pi)$.

Example 4.11. Let us look for solutions of equation (41) of the form $\alpha = l\omega$, where l is an integer. Using the quasi-periodicity property of σ (see [1, page 37], [37, page 20]), we write

$$\begin{aligned} \sigma(a + \alpha) &= \sigma(a + l\omega) = \sigma(a - \alpha + 2l\omega) = (-1)^l e^{2l\zeta(\omega)(a - \alpha + l\omega)} \sigma(a - \alpha) \\ &= (-1)^l e^{2la\zeta(\omega)} \sigma(a - \alpha). \end{aligned}$$

Comparing with equation (41), we require $(-1)^l e^{2la\zeta(\omega)} = e^{2\alpha\zeta(a)}$. We choose l to be odd and require

$$2\alpha\zeta(a) = 2l\omega\zeta(a) = 2la\zeta(\omega) - i\pi.$$

Hence $f(a) = a\zeta(\omega) - \omega\zeta(a) = i\pi/2l$. But, according to equation (38), $f(a) = i\pi n/2k + i\pi m$. Therefore, choosing $m = 0, n = 1$ implies $l = k$, and so $\alpha = l\omega = k\pi/2k = \pi/2$. In this way, we construct an infinite family of self-Bäcklund simple closed curves with rotation number $\alpha = \pi/2$, as discussed in Section 3.3, but now we have an analytical example. See Figure 15.




FIGURE 15. Example 4.11. Self-Bäcklund centroaffine simple curves $X_+(t)$ of Wegner type (blue) with $2k$ -fold symmetry, $k = 3, 5, 7$, with rotation number $\alpha = \pi/2$ (one quarter of a turn). The red curve is traced by the midpoint of the line segment $X_+(t)X_+(t + \pi/2)$ (black) and is tangent to it. For large enough ω' , the midpoint curve is smooth and convex (top); as ω' becomes smaller, cusps appear (bottom).

4.2.3. Proof of the self-Bäcklund property (Theorem 7). We shall distinguish between two cases. In both cases we shall rewrite equation (41) in a more tractable form.

Case 1. Let us start with the most important case $m = 0$ (the curve is simple if and only if $n = 1$). For $m = 0$ we have from equation (38) that $f(a) = \frac{i\pi n}{2k}$, where

$$a = \omega + ib \in [\omega, \omega + \omega'], \quad b \in \mathbb{R}.$$

We have from equation (41) that

$$(43) \quad -\frac{\sigma(\alpha + \omega + ib)}{\sigma(\alpha - \omega - ib)} = e^{2\alpha\zeta(\omega+ib)}.$$

Using the quasi-periodicity of σ , one has

$$-\sigma(\alpha + \omega + ib) = \sigma(\alpha - \omega + ib)e^{2\zeta(\omega)(\alpha+ib)}.$$

Substituting into equation (43), we get

$$\frac{\sigma(\alpha - \omega + ib)}{\sigma(\alpha - \omega - ib)} = e^{2\alpha\zeta(\omega+ib)-2\zeta(\omega)(\alpha+ib)} = e^{2\alpha[\zeta(\omega+ib)-\zeta(\omega)]-2i\zeta(\omega)b},$$

or, equivalently,

$$-\frac{\sigma(\alpha - \omega + ib)}{\sigma(-\alpha + \omega + ib)} = e^{2\alpha[\zeta(\omega+ib)-\zeta(\omega)]-2i\zeta(\omega)b}.$$

Taking log, we obtain

$$i2\pi l + \int_{-\alpha+\omega}^{\alpha-\omega} \zeta(ib+t)dt = i\pi + 2\alpha[\zeta(\omega+ib) - \zeta(\omega)] - 2i\zeta(\omega)b.$$

Hence

$$(44) \quad \pi l + \operatorname{Im} \left(\int_0^{\alpha-\omega} \zeta(ib+t)dt \right) = \frac{\pi}{2} + \frac{\alpha}{i} [\zeta(\omega+ib) - \zeta(\omega)] - \zeta(\omega)b.$$

Let us denote

$$g(\alpha) := \operatorname{Im} \left(\int_0^{\alpha-\omega} \zeta(ib+t)dt \right).$$

Lemma 4.12. *For any $r \in \mathbb{N} \cup \{0\}$, we have*

$$\operatorname{Im} \left(\int_0^{2\omega r-\omega} \zeta(ib+t)dt \right) = (2r-1)b\zeta(\omega) - \pi r + \frac{\pi}{2}.$$

Proof. Apply the Cauchy residue formula to the rectangular path

$-\omega(2r-1) + ib \rightarrow \omega(2r-1) + ib \rightarrow \omega(2r-1) - ib \rightarrow -\omega(2r-1) - ib \rightarrow -\omega(2r-1) + ib$
to obtain the result. \square

Using the quasi-periodicity of ζ and Lemma 4.12, we have

$$\begin{aligned} g(\alpha + 2\omega) &= \operatorname{Im} \left(\int_0^{2\omega r+\omega} \zeta(ib+t)dt \right) \\ &= \operatorname{Im} \left(\int_0^{2\omega r-\omega} \zeta(ib+t)dt \right) + \operatorname{Im} \left(\int_{-\omega}^{\omega} \zeta(ib+t)dt \right) \\ &= \operatorname{Im} \left(\int_0^{2\omega r-\omega} \zeta(ib+t)dt \right) + 2\operatorname{Im} \left(\int_0^{\omega} \zeta(ib+t)dt \right) \\ &= \operatorname{Im} \left(\int_0^{2\omega r-\omega} \zeta(ib+t)dt \right) + 2b\zeta(\omega) - \pi = g(\alpha) + 2b\zeta(\omega) - \pi. \end{aligned}$$

Therefore we can write g in the form

$$(45) \quad g(\alpha) = \left(\frac{2b\zeta(\omega) - \pi}{2\omega} \right) \alpha + h(\alpha),$$

where h is a 2ω -periodic function. Moreover, by Lemma 4.12 (with $r = 0$),

$$h(0) = g(0) = -b\zeta(\omega) + \frac{\pi}{2}.$$

It is convenient to use h_0 instead of h :

$$h_0(\alpha) := h(\alpha) - h(0) = h(\alpha) + b\zeta(\omega) - \frac{\pi}{2},$$

so that h_0 is 2ω -periodic with $h_0(0) = 0$. Thus

$$(46) \quad g(\alpha) = \left(\frac{2b\zeta(\omega) - \pi}{2\omega} \right) \alpha + h_0(\alpha) - b\zeta(\omega) + \frac{\pi}{2}.$$

Substituting equation (46) into equation (44), we obtain the equation:

$$\begin{aligned} \pi l + \left(\frac{2b\zeta(\omega) - \pi}{2\omega} \right) \alpha + h_0(\alpha) - b\zeta(\omega) + \frac{\pi}{2} \\ = \frac{\pi}{2} + \frac{\alpha}{i} [\zeta(\omega + i\alpha) - \zeta(\omega)] - \zeta(\omega)b. \end{aligned}$$

This is the same as

$$(47) \quad \begin{aligned} \pi l + h_0(\alpha) &= \alpha \left(\frac{-2b\zeta(\omega) + \pi}{2\omega} + \frac{(\zeta(\omega + i\alpha) - \zeta(\omega))}{i} \right) \\ &= \alpha \left(\frac{\pi}{2\omega} + \frac{2\omega\zeta(\omega + i\alpha) - 2\omega\zeta(\omega) - 2ib\zeta(\omega)}{2i\omega} \right) \\ &= \alpha \left(\frac{\pi}{2\omega} - \frac{2f(\omega + i\alpha)}{2i\omega} \right) = \alpha \left(\frac{\pi}{2\omega} - \frac{2f(a)}{2i\omega} \right). \end{aligned}$$

Taking into account that $f(a) = \frac{i\pi n}{2k}$ and $2\omega k = \pi$, we come to the final form of the equation:

$$(48) \quad \pi l + h_0(\alpha) = \alpha(k - n).$$

We claim that equation (48) has at least $k - n - 1$ solutions for α in the open interval $(0, \pi)$.

Indeed, since $h_0(0) = h_0(\pi) = 0$, the end points $\alpha = 0, \alpha = \pi$ of the open interval are solutions of equation (48) for $l = 0$ and $l = k - n$, respectively. (These two solutions are geometrically trivial, corresponding to $\alpha = 2\beta = 0$ and $\alpha = 2\beta = \pi$ for the initial geometric problem.) Therefore, for all intermediate levels of l , that is, for $l \in [1, k - n - 1]$, there exists a solution of equation (48). This proves the claim.

We shall prove now that the number of solutions of equation (48) in the interval $(0, \pi)$ is exactly equal to $(k - n - 1)$. For equation (44), it suffices to show that the function

$$\operatorname{Im} \left(\int_0^{\alpha - \omega} \zeta(ib + t) dt \right) - \frac{\alpha}{i} [\zeta(\omega + i\alpha) - \zeta(\omega)]$$

has non-vanishing derivative with respect to α . Arguing by contradiction, suppose that

$$\operatorname{Im} (\zeta(ib + \alpha - \omega) - [\zeta(\omega + i\alpha) - \zeta(\omega)]) = 0.$$

Notice that $\zeta(\omega)$ is real, and $\zeta(\omega + \alpha + ib)$ and $\zeta(-\omega + \alpha + ib)$ have the same imaginary part. Hence

$$(49) \quad \text{Im}(\zeta(ib + \alpha + \omega) - \zeta(\omega + ib)) = 0.$$

Using the addition formula, we have

$$\zeta(ib + \omega + \alpha) = \zeta(ib + \omega) + \zeta(\alpha) + \frac{\wp'(ib + \omega) - \wp'(\alpha)}{2(\wp(ib + \omega) - \wp(\alpha))}.$$

It then follows from equation (49) that

$$\zeta(\alpha) + \frac{\wp'(ib + \omega) - \wp'(\alpha)}{2(\wp(ib + \omega) - \wp(\alpha))} \in \mathbb{R}.$$

Moreover, the values $\zeta(\alpha)$, $\wp(ib + \omega)$, $\wp(\alpha)$, $\wp'(\alpha)$ are all real. We conclude that $\wp'(ib + \omega) \in \mathbb{R}$.

On the other hand,

$$ib + \omega \in (\omega, \omega') \Rightarrow e_2 < \wp(ib + \omega) < e_1.$$

Thus the equation $(\wp')^2 = 4(\wp - e_1)(\wp - e_2)(\wp - e_3)$ implies that $\wp'(ib + \omega) \in i\mathbb{R}$, a contradiction. This completes the proof of Theorem 7 in Case 1.

Case 2. In this case $m > 0$, $a = ib \in [0, \omega']$, $b \in \mathbb{R}$. Using $\frac{\sigma'}{\sigma} = \zeta$, we write

$$\sigma(z) = \sigma(z_0) \exp\left(\int_{z_0}^z \zeta(t) dt\right).$$

Taking log, we rewrite equation (41) in the form

$$\int_{-\alpha}^{\alpha} \zeta(ib + t) dt + 2\pi il = 2\alpha\zeta(ib), \quad l \in \mathbb{Z}.$$

Using that ζ is odd, rewrite this as

$$2\pi il + \int_0^{\alpha} [\zeta(ib + t) - \zeta(-ib + t)] dt = 2\alpha\zeta(ib).$$

Notice that both sides of this equation are purely imaginary, and hence

$$(50) \quad \pi l + \text{Im}\left(\int_0^{\alpha} \zeta(ib + t) dt\right) = \frac{1}{i}\alpha\zeta(ib).$$

On the right hand side we have a linear function of α . Let us denote the integral on the left hand side of equation (50) by

$$g(\alpha) := \text{Im}\left(\int_0^{\alpha} \zeta(ib + t) dt\right).$$

Lemma 4.13. *For any $r \in \mathbb{N}$, we have*

$$\text{Im}\left(\int_0^{2\omega r} [\zeta(ib + t) dt]\right) = -\pi r + 2r\zeta(\omega)b.$$

Proof. This follows from the residue formula for the rectangular path

$$ib \rightarrow 2\omega r + ib \rightarrow 2\omega r - ib \rightarrow -ib \rightarrow ib,$$

avoiding the singular points of ζ at 0 and $2\omega r$ by small half circles. □

In particular, using Lemma 4.13 for $r = 1$ and the quasi-periodicity of ζ , we compute

$$g(\alpha + 2\omega) = g(\alpha) + \frac{1}{i} \int_0^{2\omega} \zeta(ib + t)dt = g(\alpha) - \pi + 2\zeta(\omega)b.$$

Using this, one can express g as the sum of a linear and a 2ω -periodic function as follows:

$$g(\alpha) = \left(\frac{-\pi + 2\zeta(\omega)b}{2\omega} \right) \alpha + h(\alpha), \quad g(0) = h(0) = 0,$$

where h is 2ω -periodic. Therefore, equation (50) takes the form

$$\pi l + h(\alpha) = - \left(\frac{-\pi + 2\zeta(\omega)b}{2\omega} \right) \alpha + \frac{1}{i} \alpha \zeta(ib),$$

hence

$$\pi l + h(\alpha) = \alpha \left(\frac{1}{i} \zeta(ib) - \frac{-\pi + 2\zeta(\omega)b}{2\omega} \right).$$

Thus we arrive at the following equation

$$\pi l + h(\alpha) = \alpha \left(\frac{\pi}{2\omega} + \frac{2\omega\zeta(ib) - 2\zeta(\omega)ib}{2\omega i} \right) = \alpha \left(\frac{\pi}{2\omega} - \frac{2f(ib)}{2\omega i} \right).$$

Next, taking into account that $f(ib) = f(a) = \frac{i\pi n}{2k}$ and $2\omega k = \pi$, we obtain the simplest possible form:

$$(51) \quad \pi l + h(\alpha) = \alpha(k - n).$$

Also in this case we claim that equation (51) has at least $k - n - 1$ solutions for α in the open interval $(0, \pi)$.

Indeed, since $h(0) = h(\pi) = 0$, the end points $\alpha = 0, \alpha = \pi$ of the open interval are solutions of equation (51) for $l = 0$ and $l = k - n$, respectively. Therefore, for all intermediate levels of l , that is, for $l \in [1, k - n - 1]$, there exists a solution of equation (51). This proves the claim.

We shall prove now that the number of solutions of equation (51) in the interval $(0, \pi)$ equals exactly $k - n - 1$. Consider equation (50). We shall check that the function

$$\text{Im} \left(\int_0^\alpha \zeta(ib + t)dt - \alpha \zeta(ib) \right)$$

has everywhere non-vanishing derivative with respect to α when $ib \in (0, \omega')$.

Suppose, on the contrary, that the derivative vanishes for some α :

$$(52) \quad \text{Im} (\zeta(ib + \alpha) - \zeta(ib)) = 0.$$

Using the addition formula for ζ , we have

$$\zeta(ib + \alpha) = \zeta(ib) + \zeta(\alpha) + \frac{\wp'(ib) - \wp'(\alpha)}{2(\wp(ib) - \wp(\alpha))}.$$

Taking the imaginary part and using equation (52), we obtain

$$\zeta(\alpha) + \frac{\wp'(ib) - \wp'(\alpha)}{2(\wp(ib) - \wp(\alpha))} \in \mathbb{R}.$$

Also we know that $\zeta(\alpha), \wp(ib), \wp(\alpha), \wp'(\alpha)$ are all real. Therefore we conclude that $\wp'(ib) \in \mathbb{R}$. But, on the other hand, \wp satisfies the equation $(\wp')^2 = 4(\wp - e_1)(\wp - e_2)(\wp - e_3)$. Moreover,

$$ib \in (0, \omega') \Rightarrow \wp(ib) < e_3 \Rightarrow \wp'(ib) \in i\mathbb{R}.$$

This contradiction completes the proof in Case 2.

□

Theorem 7 has Corollary 4.14.

Corollary 4.14. *All the solutions of equation (41) are transversal and hence change smoothly as one varies the parameter ω' of the elliptic functions involved.*

4.3. Self-Bäcklund curves as deformations of conics. In this section we use the self-Bäcklund curves of Section 4.2 in order to construct genuine non-trivial self-Bäcklund deformations of a central conic, as was promised in Section 3.1, see Corollary 4.20.

To state the result, we recall briefly from Section 4.2 our construction of *simple* self-Bäcklund centroaffine π -anti-periodic curves. For every integer $k \geq 3$ and $\omega' \in i\mathbb{R}_+$ one considers the Weierstrass \wp -function with half periods $\omega = \pi/2k$, ω' , the associated σ - and ζ -functions and the (unique) solution $a \in (\omega, \omega')$ to

$$(53) \quad a\zeta(\omega) - \omega\zeta(a) = i\omega,$$

then set

$$(54) \quad Y(t) := X(t)/N,$$

where

$$(55) \quad X(t) := \frac{\sigma(a + t + \omega')\sigma(\omega')}{\sigma(a + \omega')\sigma(t + \omega')} e^{-t\zeta(a)}, \quad N := \sqrt{|X'(0)|}.$$

Remark 4.15. The normalization factor $N = \sqrt{|X'(0)|}$ in equations (54)-(55) is introduced so as to render the normalized curve Y centroaffine and π -anti-periodic (enclosing area π). See Remark 4.9 for an explicit expression for N .

The deformations of the unit circle we are seeking are obtained by fixing k and letting $\omega' \rightarrow \infty$ in the above construction. To examine this limit we let $\omega' = i/s$, $s \in (0, 1]$, and use henceforth the subscript s to denote all associated objects, such as $\wp_s, \sigma_s, \zeta_s, a_s, X_s, N_s$ and Y_s (suppressing the dependence on k , which is fixed throughout the section). Our goal in this section is to prove Theorem 8, illustrated in Figure 16.

Theorem 8. *For each integer $k \geq 3$,*

- (1) *The family of curves $Y_s(t)$, $s \in (0, 1]$, given by equations (53)-(55) with $\omega = \pi/2k$, $\omega' = i/s$, extends smoothly to $s \in [0, 1]$ by setting $Y_0(t) := e^{it}$.*
- (2) *Each curve $Y_s(t)$ is a centroaffine π -anti-periodic simple curve with $2k$ -fold symmetry, $Y_s(t + \pi/k) = Y_s(t)e^{i\pi/k}$, self-Bäcklund for $s > 0$ with respect to $k - 2$ rotation numbers $\alpha \in (0, \pi)$, varying smoothly in $s \in [0, 1]$ and converging as $s \rightarrow 0$ to the $k - 2$ solutions of equation (19), $\tan(k\alpha) = k \tan \alpha$.*
- (3) *The deformation Y_s , $s \in [0, 1]$, is analytic away from $s = 0$ but not at $s = 0$. In fact, one has $(\partial_s)^n|_{s=0} Y_s(t) = 0$, $n \geq 1$, so the associated infinitesimal deformation of the unit circle vanishes to all orders, yet the deformation itself is non-trivial.*

(4) *The change of parameter,*

$$(56) \quad \varepsilon := \begin{cases} e^{-2k/s}, & s > 0, \\ 0, & s = 0, \end{cases}$$

gives a deformation Y_ε of the unit circle Y_0 , analytic in $\varepsilon \in [0, e^{-2k}]$.

(5) *The infinitesimal deformation associated with the analytic deformation Y_ε is non-trivial. That is,*

$$Y_\varepsilon(t) = e^{it} + Y_1(t)\varepsilon + O(\varepsilon^2),$$

where Y_1 is non-vanishing.




FIGURE 16. Theorem 8. Three families of deformations of the circle (black) through a 1-parameter family of centroaffine self-Bäcklund curves Y_s (blue) with $2k$ -fold symmetry, $k = 3, 4, 5$.

Proof. The main idea of the proof of this theorem is to write the functions $X_s, s \in [0, 1]$, as suitably normalized Floquet eigenfunctions of a Hill operator depending smoothly on s , and use a general argument of smooth dependence of the eigenfunctions of a Hill operator depending on the smooth parameter. Similarly, when replacing s with ε the Hill operator depends analytically on ε and so do its eigenfunctions.

In more detail, we recall from Section 4.2 that $X_s, s \in (0, 1]$, is precisely the eigenfunction corresponding to the smallest eigenvalue $\lambda_{0,s}$ for the Floquet problem

$$(57) \quad X'' + (\lambda - 2q_s(t))X = 0, \quad X(t + \pi/k) = \mu X(t), \quad \mu = e^{i\pi/k},$$

where $q_s(t) = \wp(i/s + t)$ and X_s satisfy the normalization condition $X_s(0) = 1$. Moreover, we showed that $\lambda_{0,s} = -\wp_s(a_s)$, where $a_s \in (\omega, \omega')$ is the (unique) solution to equation (53).

Following this idea, we begin by extending q_s smoothly to $s = 0$.

Lemma 4.16.

(A) *The function*

$$q_s(t) := \begin{cases} \wp_s(t + i/s), & s \neq 0, \\ -k^2/3, & s = 0 \end{cases}$$

depends smoothly on $(s, t) \in [0, 1] \times \mathbb{R}$.

(B) *The change of parameter $s \mapsto \varepsilon$ of equation (56) transforms the deformation q_s to q_ε which is real analytic in $\varepsilon \in [0, e^{-2k}]$, with Taylor series*

$$(58) \quad q_\varepsilon = -\frac{k^2}{3} - 8k^2 \cos(2kt)\varepsilon + O(\varepsilon^2).$$

We postpone the proof of Lemma 4.16, as well as Lemmas 4.17–4.19, to the end of this section.

Lemma 4.17. *The eigenfunctions $X_s(t)$, $s \in [0, 1]$, corresponding to the first eigenvalue $\lambda_{0,s}$ of the Floquet problem (57), are uniquely determined by the condition $X_s(0) = 1$ and are smooth (analytic) in s if the potential q_s is smooth (analytic) in s .*

Lemma 4.18. *For every $s \in [0, 1]$ the curves Y_s are self-Bäcklund for $k - 2$ values of $\alpha_s \in (0, \pi)$, satisfying*

$$(59) \quad \frac{\sigma_s(a_s + \alpha_s)}{\sigma_s(a_s - \alpha_s)} = e^{2\alpha_s \zeta_s(a_s)}.$$

All $k - 2$ solutions α_s depend smoothly on $s \in [0, 1]$. For $s = 0$ this equation reduces to equation (19) of Theorem 3, $k \tan(\alpha) = \tan(k\alpha)$. Moreover, with respect to the parameter ε of equation (56) the $k - 2$ families α_ε are analytic in $\varepsilon \in [0, e^{-2k}]$.

Lemma 4.19. *X_ε has a Taylor series in ε ,*

$$X_\varepsilon(t) = e^{it} + X_1(t)\varepsilon + O(\varepsilon^2),$$

where X_1 is non-vanishing.

With Lemmas 4.16–4.19 the proof of the 5 items of Theorem 8 is straightforward: by Lemma 4.16, the Hill operator of equation (57) is smooth in $s \in [0, 1]$ and analytic in $\varepsilon \in [0, e^{-2k}]$. This implies, by Lemma 4.17, that X_s is smooth in s and X_ε is analytic in ε , therefore the same holds for Y_s and Y_ε . This proves items (1) and (4) of Theorem 8. Lemma 4.18 proves item (2). Item (3) follows from the well-known fact that $\varepsilon(s)$ of formula (56) is “flat” at $s = 0$ (all derivatives exist and vanish). Lemma 4.19 gives item (5). □

Corollary 4.20. *For every value of $\alpha \in (0, \pi)$ for which the unit circle admits a non-trivial infinitesimal self-Bäcklund deformation (solution of $\tan(k\alpha) = k \tan \alpha$ for some $k \geq 3$) there is a genuine analytic self-Bäcklund deformation realizing it.*

We now proceed to the promised proofs of Lemmas 4.16–4.19 appearing in the above proof of Theorem 8.

4.3.1. *Proof of Lemma 4.16.* By the definition of \wp , we have the following series representing q_s for $s > 0$:

$$(60) \quad q_s(t) = \wp_s(t + i/s) = (t + i/s)^{-2} + \sum_{(m,n) \neq (0,0)} \left[\left(t + \frac{\pi n}{k} + i \frac{2m+1}{s} \right)^{-2} - \left(\frac{\pi n}{k} + i \frac{2m}{s} \right)^{-2} \right].$$

Let $z := t + i/s$, $\Omega_{nm} := \pi n/k + 2mi/s$, $m, n \in \mathbb{Z}$, $s > 0$. We break the double sum in the series (60) as a sum $\sum_m Q_m$, where each Q_m is a series in n :

$$Q_m = \begin{cases} \sum_{n \in \mathbb{Z}} [(z - \Omega_{nm})^{-2} - (\Omega_{nm})^{-2}], & m \neq 0, \\ \sum_{n \in \mathbb{Z}, n \neq 0} [(z - \Omega_{n0})^{-2} - (\Omega_{n0})^{-2}], & m = 0. \end{cases}$$

We have for Q_m the exact expressions (see [1], page 197, Table I):

$$Q_m = \begin{cases} k^2 \left[\sin^{-2} \left(k \left(z - i \frac{2m}{s} \right) \right) - \sin^{-2} \left(i \frac{2km}{s} \right) \right], & m \neq 0, \\ k^2 \left[-\frac{1}{3} + \sin^{-2}(kz) \right], & m = 0. \end{cases}$$

Substituting into these formulas $z = t + i/s$, we get

$$Q_m = \begin{cases} k^2 \left[\sin^{-2} \left(k \left(t - i \frac{2m-1}{s} \right) \right) - \sin^{-2} \left(i \frac{2km}{s} \right) \right], & m \neq 0, \\ k^2 \left[-\frac{1}{3} + \sin^{-2} \left(k \left(t + \frac{i}{s} \right) \right) \right], & m = 0. \end{cases}$$

Thus we have

$$Q_m = \begin{cases} k^2 \left[\left(\sin(kt) \cosh \left(k \frac{2m-1}{s} \right) - i \cos(kt) \sinh \left(k \frac{2m-1}{s} \right) \right)^{-2} - \sinh^{-2} \left(\frac{2km}{s} \right) \right], & m \neq 0, \\ k^2 \left[-\frac{1}{3} + \left(\sin(kt) \cosh \left(\frac{k}{s} \right) + i \cos(kt) \sinh \left(\frac{k}{s} \right) \right)^{-2} \right], & m = 0. \end{cases}$$

Next introduce the change of parameter, $s \mapsto \tau = e^{-k/s}$, $0 \leq \tau \leq \tau_0 = e^{-k}$, ie $\varepsilon = \tau^2$. In terms of τ , we have

$$(61) \quad Q_m = \begin{cases} 4k^2 \left[(\sin(kt)(\tau^{1-2m} + \tau^{2m-1}) - i \cos(kt)(\tau^{1-2m} - \tau^{2m-1}))^{-2} - (\tau^{-2m} - \tau^{2m})^{-2} \right], & \tau > 0, m \neq 0, \\ 0, & \tau = 0, m \neq 0, \end{cases}$$

$$Q_0 = \begin{cases} k^2 \left[-\frac{1}{3} + 4 \left(\sin(kt)(\tau^{-1} + \tau) + i \cos(kt)(\tau^{-1} - \tau) \right)^{-2} \right], & \tau > 0, \\ -\frac{1}{3}k^2, & \tau = 0. \end{cases}$$

From formulas (61) one can conclude the following facts:

- (1) The series

$$q = \sum_{m \in \mathbb{Z}} Q_m$$

converges as $\tau \rightarrow 0$, uniformly in $(\tau, t) \in [0, \tau_0] \times \mathbb{R}$, to the constant function $q_0 = -\frac{1}{3}k^2$. This follows from the estimate

$$|Q_m| \leq C(\tau_0)(\tau_0^{4|m|} + \tau_0^{|4m-2|})$$

for some constant $C(\tau_0) > 0$. Since $\tau_0 = e^{-k} < 1$ this implies uniform convergence in $[0, \tau_0] \times \mathbb{R}$.

- (2) Every term Q_m in equation (61) is analytic in τ at $\tau = 0$ with radius of convergence $R_m = 1 > \tau_0$. To see this, one represents each term in the square brackets of (61) as a rational function of τ and finds that its poles all lie on the unit circle in the complex τ plane. Hence $R_m = 1$.
- (3) It follows from items (1) and (2), by Weierstrass theorem, that the sum of the series $\sum Q_m$, which equals exactly $q_\tau(t)$, is analytic in $\tau \in [0, \tau_0]$.
- (4) Each Q_m in equation (61) is clearly even in τ , hence so is q . Thus, with the change of variable $\varepsilon = \tau^2$, q_ε becomes analytic in ε .
- (5) The following 1st order Taylor expansions at $\tau = 0$ hold:

$$(62) \quad Q_0 = -\frac{k^2}{3} - 4k^2 e^{2ikt} \tau^2 + \dots, \quad Q_1 = -4k^2 e^{-2ikt} \tau^2 + \dots,$$

and Q_m is of order τ^{4m-2} for $m > 0$, which implies equation (58).

4.3.2. *Proof of Lemma 4.17.* Notice that, for a given periodic potential $q(t)$, the problem (57) of Floquet eigenvalues has the following properties (see [22, page 32]):

- (1) The eigenvalues $\lambda_m(\mu)$ are solutions of the equation

$$(63) \quad \Delta(\lambda) = 2 \cos\left(\frac{\pi}{k}\right).$$

Here and below, $\Delta(\lambda) = \text{tr}M(\lambda)$ is the trace of the monodromy matrix of equation (57). It is defined as follows. Fix a basis of solutions $\{y_1(\lambda, t), y_2(\lambda, t)\}$ of the second order differential equation

$$X'' + (\lambda - 2q(t))X = 0,$$

such that

$$y_1(\lambda, 0) = y_2'(\lambda, 0) = 1, \quad y_1'(\lambda, 0) = y_2(\lambda, 0) = 0.$$

Then the monodromy matrix is

$$M(t, \lambda) = \begin{pmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{pmatrix} = \begin{pmatrix} y_1(2\omega, \lambda) & y_2(2\omega, \lambda) \\ y_1'(2\omega, \lambda) & y_2'(2\omega, \lambda) \end{pmatrix}, \quad \det(M) = 1.$$

- (2) The graph of the function $\Delta(\lambda)$ (see Figure 14) is such that all the solutions of equation (63) are transversal. Hence all eigenvalues $\lambda_{m,s}(\mu)$ of equation (57), and, in particular, $\lambda_{0,s}(\mu)$, depend smoothly on the parameter s .
- (3) All Floquet eigenvalues $\lambda_{0,s}(\mu)$ of equation (57) have multiplicity 1, because if X is an eigenfunction for some non-real Floquet exponent μ , then \bar{X} is not.

In our situation, we have a smooth family of potentials $q_s(t)$. So we have the standard basis $\{y_1(\lambda, s, t), y_2(\lambda, s, t)\}$, where

$$y_1(\lambda, s, 0) = y_2'(\lambda, s, 0) = 1, \quad y_1'(\lambda, s, 0) = y_2(\lambda, s, 0) = 0,$$

and the monodromy matrix $M(\lambda, s)$ which is smooth in λ, s . We can write the eigenfunction corresponding to $\lambda_{m,s}(\mu)$ in the form

$$X = Ay_1 + By_2,$$

for some complex A, B . Then the Floquet boundary conditions in terms of A, B reads

$$(64) \quad (M(\lambda, s) - \mu Id) \cdot \begin{pmatrix} A \\ B \end{pmatrix} = 0.$$

Moreover, it follows from properties (2) and (3) above that, for $\lambda = \lambda_{m,s}$, the matrix $(M - \mu Id)$ has rank 1 and that $M(\lambda_{m,s}, s)$ depends smoothly on s . The normalization $X(0) = 1$ implies that $A = 1$ and hence B can be found uniquely from (64),

$$B = -(m_{11} - \mu)/m_{12}.$$

It is important that the denominator m_{12} in this formula cannot vanish, because otherwise the matrix M would be triangular having real eigenvalues, which is not the case, since μ is not real. Thus we conclude that the solution $\begin{pmatrix} A \\ B \end{pmatrix}$ of equation (64) is smooth in $s \in [0, 1]$. An analogous proof applies when the potential q_ε depends analytically on $\varepsilon \in [0, e^{-2k}]$. This completes the proof of our Lemma.

4.3.3. *Proof of Lemma 4.18.* The functions \wp_s, σ_s, ζ_s depend analytically on $s \in (0, 1]$ and can be shown to converge, as $s \rightarrow 0$, to the limiting functions (see [1, page 201])

$$(65) \quad \begin{aligned} \wp_0(z) &= -\frac{k^2}{3} + k^2 \sin^{-2}(kz), & \zeta_0(z) &= \frac{k^2}{3}z + k \cot(kz), \\ \sigma_0(z) &= \frac{1}{k} e^{k^2 z^2/6} \sin(kz). \end{aligned}$$

Using the above formula for ζ_0 , we compute that equation (53) for $s = 0$ is equivalent to

$$(66) \quad a = \frac{\pi}{2k} + ib, \quad \tanh\left(\frac{\pi b}{2}\right) = \frac{1}{k}.$$

Consider equation (59) on α for $s = 0$:

$$\frac{\sigma_0(a + \alpha)}{\sigma_0(a - \alpha)} = e^{2\alpha\zeta_0(a)},$$

where a is the solution of equation (53) for $s = 0$. Set

$$F(\alpha) := \frac{\sigma_0(a + \alpha)}{\sigma_0(a - \alpha)} e^{-2\alpha\zeta_0(a)}.$$

Using the explicit formulas (65)-(66), we have:

$$F(\alpha) = \frac{\sin(k(a + \alpha))}{\sin(k(a - \alpha))} e^{i2\alpha} = \frac{1 - i\frac{1}{k} \tan(k\alpha)}{1 + i\frac{1}{k} \tan(k\alpha)} e^{i2\alpha}.$$

This immediately implies that the equation $F = 1$ is equivalent to the familiar equation (19):

$$k \tan(\alpha) = \tan(k\alpha).$$

This means that, for $s = 0$, equation (59) has precisely $k - 2$ solutions for $\alpha \in (0, \pi)$.

Moreover, differentiating F at a point α where $F(\alpha) = 1$ we have:

$$F'(\alpha) = 2i \frac{(1 - k^2) \tan^2 k\alpha}{k^2 + \tan^2 k\alpha} \neq 0.$$

Applying the implicit function theorem, we conclude that all $k - 2$ solutions of equation (59) can be smoothly extended from $s = 0$ to $s > 0$. This, together with Theorem 7 and

Corollary 4.14, implies the existence of $k - 2$ solutions for every $s \in [0, 1]$, smoothly depending on s . An analogous proof applies for analytic dependence on $\varepsilon \in [0, e^{-2k}]$.

4.3.4. *Proof of Lemma 4.19.* We calculate mod ε^2 . Use the Taylor expansion (58),

$$q_\varepsilon = -\frac{k^2}{3} - 8k^2 \cos(2kt)\varepsilon + \dots,$$

and let $X_\varepsilon = e^{it} + X_1\varepsilon + \dots$, $\lambda_{0,\varepsilon} = \lambda_0 + \lambda_1\varepsilon + \dots$. Substitute these into $X'' + (\lambda - 2q)X = 0$ and solve for successive powers of ε . The ε^0 term gives

$$\lambda_0 = 1 - 2k^2/3$$

and the ε^1 term gives

$$X_1'' + X_1 + 8k^2 (e^{i(1+2k)t} + e^{i(1-2k)t}) + \lambda_1 e^{it} = 0.$$

The general solution is

$$X_1 = A_+ e^{i(1+2k)t} + A_- e^{i(1-2k)t} + B_+ e^{it} + B_- e^{-it} + \frac{\lambda_1}{2i} t e^{it},$$

where $A_\pm = 2k/(k \pm 1) \neq 0$ and $B_\pm \in \mathbb{C}$ are arbitrary. Since X_1 is periodic we must have $\lambda_1 = 0$ and what remains is non-vanishing.

5. SELF-BÄCKLUND POLYGONS

5.1. **Centroaffine butterflies, Bianchi permutability.** The central projection $\mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{R}P^1$ takes a centroaffine curve to a curve in the projective line. Conversely, a projective curve admits a unique lift to a centroaffine curve. Bianchi permutability for c -relation was established for projective curves, in [44]. Here we do it for centroaffine curves.

Let us say that a quadrilateral $P_1P_2P_3P_4$ forms a *centroaffine butterfly* if

$$(67) \quad [P_1, P_2] = [P_4, P_3] \text{ and } [P_2, P_3] = [P_1, P_4].$$

Note that a centroaffine butterfly is not necessarily a centroaffine polygon.

Lemma 5.1. *A generic quadrilateral $P_1P_2P_3P_4$ is a centroaffine butterfly if and only if any of the following equivalent conditions are satisfied:*

- (1) *There is a linear involution $I \in GL_2(\mathbb{R})$ interchanging P_1P_2 and P_3P_4 . That is, $I(P_1) = P_3, I(P_2) = P_4, I(P_3) = P_1, I(P_4) = P_2$.*
- (2) *The line segments P_1P_3, P_2P_4 are parallel and their midpoints are collinear. See Figure 17.*
- (3) *$P_aP_bP_cP_d$ is a centroaffine butterfly, where $abcd$ is any of the 8 permutations of 1234 generated by (1234), (24), (12)(34).*

Proof. (1) By applying a linear transformation, we can assume that $P_1 = (1, 0), P_3 = (0, 1)$. Let $P_2 = (c, d)$. Then equation (67) implies $P_4 = (d, c)$. Thus $I : (x, y) \mapsto (y, x)$ is the required symmetry.

(2) Note that the said segments are parallel and their midpoints are collinear if and only if $[P_1 \pm P_3, P_2 \pm P_4] = 0$ (‘-’ for the 1st statement, ‘+’ for the 2nd). By expanding these expressions we see that they are equivalent to $[P_1, P_2] = [P_4, P_3], [P_2, P_3] = [P_1, P_4]$.

(3) This is a simple verification (omitted).

□




FIGURE 17. A centroaffine butterfly

It follows from Lemma 5.1 that, given a generic triple of points P_1, P_2, P_3 , there is a unique fourth point P_4 such that $P_1P_2P_3P_4$ form a centroaffine butterfly. Namely, by property (1), $P_4 = IP_2$ where I is defined by $IP_1 = P_3, IP_3 = P_1$. More geometrically, by property (2), one constructs the line ℓ through P_2 and parallel to P_1P_3 , intersects ℓ with the line through the origin O and the midpoint of P_1P_3 , then finds the unique point P_4 on ℓ such that this intersection point is the midpoint of P_2P_4 .

Theorem 9 (Bianchi permutability). *Consider three centroaffine curves γ, δ , and Γ such that Γ and δ are b - and c -related to γ (respectively). Then there exists a fourth centroaffine curve Δ that is b -related to δ and c -related to Γ . In fact, $\Delta(t)$ is the unique point such that $\delta(t)\gamma(t)\Gamma(t)\Delta(t)$ form a centroaffine butterfly.*

Proof. The idea of the proof is that if $\gamma(t), \delta(t)$ and $\Gamma(t)$ are considered as three vertices of time-evolving centroaffine butterfly, then $\Delta(t)$ is its fourth vertex.

Specifically, we have

$$[\gamma, \delta] = [\Gamma, \Delta] = c, [\gamma, \Gamma] = [\delta, \Delta] = b,$$

and need to check that $\Delta(t)$ is a centroaffine curve, that is, $[\Delta, \Delta'] = 1$.

Using the above relations, one can write Δ as a linear combination of δ and Γ ,

$$\Delta = \frac{[\gamma, \delta]}{[\Gamma, \delta]} \delta - \frac{[\gamma, \Gamma]}{[\Gamma, \delta]} \Gamma = \frac{c\delta - b\Gamma}{[\Gamma, \delta]}.$$

Then

$$[\Delta, \Delta'] = \frac{[c\delta - b\Gamma, c\delta' - b\Gamma']}{[\Gamma, \delta]^2} = \frac{b^2 + c^2 - bc([\delta, \Gamma'] + [\Gamma, \delta'])}{[\Gamma, \delta]^2}.$$

Thus we want to show that

$$(68) \quad b^2 + c^2 - bc([\delta, \Gamma'] + [\Gamma, \delta']) = [\Gamma, \delta]^2.$$

We have

$$\delta = f\gamma + c\gamma', \quad \Gamma = g\gamma + b\gamma',$$

hence

$$\delta' = (f' + cp)\gamma + f\gamma', \quad \Gamma' = (g' + bp)\gamma + g\gamma'.$$

It follows that

$$[\Gamma, \delta] = cg - bf, [\delta, \Gamma'] = fg - cg' - bcp, [\Gamma, \delta'] = fg - bf' - bcp.$$

In addition, one has by equation (3):

$$cf' = f^2 - c^2p - 1, bg' = g^2 - b^2p - 1.$$

Substitute these formulas into equation (68) to obtain a true identity. □

5.2. Rigidity results and flexible examples of self-Bäcklund polygons. Bäcklund transformation can be defined on centroaffine polygons. Similarly to its continuous version, it is a completely integrable dynamical system. We refer to [2] for a detailed study; see also [33].

For the purpose of this paper, we recall, from Section 1, that an origin-symmetric $2n$ -gon \mathbf{P} in \mathbb{R}^2 with vertices $P_i, i = 1, \dots, 2n$, is called a *self-Bäcklund* (n, k) -gon if

$$[P_i, P_{i+1}] = 1, [P_i, P_{i+k}] = c$$

for all i and $2 \leq k \leq n - 2$. Such polygons are acted upon by $SL_2(\mathbb{R})$. Since $P_{i+n} = -P_i$, we can assume, without loss of generality, that $k \leq n/2$.

A regular $2n$ -gon is a self-Bäcklund (n, k) -gon for all $2 \leq k \leq n/2$. We call these self-Bäcklund (n, k) -gons and their $SL_2(\mathbb{R})$ images trivial. The problem is to find non-trivial self-Bäcklund (n, k) -gons.

The next result is analogous to Theorem 9 of [41].

Theorem 10. *In the following cases every self-Bäcklund (n, k) -gon is trivial:*

- (1) n is arbitrary, $k = 2$;
- (2) n is odd, $k = 3$;
- (3) k is arbitrary, $n = 2k + 1$.
- (4) $n = 3k$.

On the other hand, there exist non-trivial self-Bäcklund (n, k) -gons in the following cases:

- (1) n is even and k is odd;
- (2) $n = 2k$.

Proof. Each next vertex is a linear combination of the preceding two: $P_{i+2} = a_i P_{i+1} - P_i$.

Let $k = 2$. Then $[P_i, P_{i+2}] = c$, hence $a_i = c$ for all i . Let A be the linear map defined by

$$A(P_1) = P_2, A(P_2) = P_3.$$

We claim that A is area preserving and $A(P_i) = P_{i+1}$ for all i . This would imply that the polygon \mathbf{P} is centroaffine regular, that is, trivial.

That A is area preserving follows from $[P_1, P_2] = [P_2, P_3]$. Next,

$$P_3 = -P_1 + cP_2, \text{ hence } A(P_3) = -P_2 + cP_3 = P_4.$$

Repeating this argument, we obtain $A(P_i) = P_{i+1}$ for all i .

Now let n be odd and $k = 3$. Consider four consecutive vertices of \mathbf{P} ; they satisfy the Ptolemy-Plücker relation

$$[P_i, P_{i+1}][P_{i+2}, P_{i+3}] + [P_{i+1}, P_{i+2}][P_i, P_{i+3}] = [P_i, P_{i+2}][P_{i+1}, P_{i+3}].$$

Therefore

$$1 + c = [P_i, P_{i+2}][P_{i+1}, P_{i+3}].$$

It follows that $[P_i, P_{i+2}] = [P_{i+2}, P_{i+4}]$ for all i .

Recall that n is odd and that $P_{i+n} = -P_i$ for all i . This implies that

$$[P_i, P_{i+2}] = [P_{n+i}, P_{n+i+2}] = [P_{i+1}, P_{i+3}],$$

and hence $[P_i, P_{i+2}]$ has the same value for all i . Thus \mathbf{P} is a self-Bäcklund $(n, 2)$ -gon, the already considered case.

Next, let $n = 2k + 1$. First we notice that $[P_i, P_{i+k+1}] = c$. Indeed,

$$[P_i, P_{i+k}] = [P_{i+k+1}, P_{i+n}] = [P_i, P_{i+k+1}].$$

Now consider the quadruple of vertices $P_i, P_{i+1}, P_{i+k}, P_{i+k+1}$. The Ptolemy-Plücker relation implies that

$$[P_{i+1}, P_{i+k}] = \frac{c^2 - 1}{c}$$

for all i . That is, $[P_i, P_{i+k-1}]$ is independent of i .

Continuing in the same way, we reduce k until we get to the case $k = 2$, considered above, and we conclude that \mathbf{P} is centroaffine regular.

Now let $n = 3k$. Let us scale the polygon so that $[P_i, P_{i+k}] = \sqrt{3}/2$ for all i (as for a regular $6k$ -gon inscribed in a unit circle). Then $[P_i, P_{i+1}] = t$, a constant.

Each hexagon $\mathbf{P}_i := (P_i, P_{i+k}, P_{i+2k}, P_{i+3k}, P_{i+4k}, P_{i+5k})$ is affine-regular, and they are all equivalent under $SL_2(\mathbb{R})$. Hence we assume, without loss of generality, that the vertices of \mathbf{P}_0 are the sixth roots of unity. Let $A \in SL_2(\mathbb{R})$ take \mathbf{P}_0 to \mathbf{P}_1 . A quick calculation, using the equations

$$[P_0, P_1] = [P_k, P_{k+1}] = [P_{2k}, P_{2k+1}] = [P_{3k}, P_{3k+1}] = [P_{4k}, P_{4k+1}] = [P_{5k}, P_{5k+1}] = t,$$

reveals that A is a rotation

$$A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}, \quad t = \sin \alpha.$$

The same argument, applied to the linear map that takes \mathbf{P}_1 to \mathbf{P}_2 , shows that this map is the same rotation, A . And so on, showing that the polygon is regular.

Let us construct non-trivial self-Bäcklund (n, k) -gons for even n and odd k . Start with a regular $2n$ -gon, and consider the midpoints of its sides. These points are the vertices of another regular $2n$ -gon. Dilate the latter $2n$ -gon with the center of dilation at its center. We obtain a centrally symmetric $4n$ -gon having a dihedral symmetry, and this symmetry implies $[P_i, P_{i+k}] = [P_{i+1}, P_{i+k+1}]$. See Figure 18 on the left. (The projection of this polygon to $\mathbb{R}P^1$ is a regular n -gon therein.)

The construction of a non-trivial self-Bäcklund $(2k + 4, k + 2)$ -gon is presented in Figure 18 on the right (where $k = 2$).¹ This polygon has two axes of symmetry. In the general case, one has points $(a, 1), (a + 1, 1), \dots, (a + k, 1)$ on a horizontal line with

$$a = \frac{\sqrt{k^2 + 8} - k}{4}, \quad c = \frac{\sqrt{k^2 + 8} + k}{2}.$$

One checks that $[P_i, P_{i+1}] = 1$ and $[P_i, P_{i+k+2}] = c$ for all i . □

¹We are grateful to Michael Cuntz for suggesting this construction.




FIGURE 18. Left: a self-Bäcklund $(8, 3)$ -gon. Right: a self-Bäcklund $(8, 4)$ -gon.

5.3. Infinitesimal deformations of regular polygons. Here we consider the linearized problem, that is, infinitesimal deformations of regular polygons as self-Bäcklund (n, k) -gons; this is a discrete analog of the material in Section 3.1.

Call a regular polygon *infinitesimally rigid* as a self-Bäcklund (n, k) -gon if each of its infinitesimal deformations in the class of self-Bäcklund (n, k) -gons is induced by the action of $\mathfrak{sl}(2, \mathbb{R})$.

Theorem 11. *A regular $2n$ -gon is infinitesimally rigid as a self-Bäcklund (n, k) -gon unless one of the following holds:*

- (1) n is even and k is odd;
- (2) $n = 2k$ with even $k > 2$;
- (3) there exists an integer j with $2 \leq j \leq n - 2$ such that $n = 2(k + j)$ and n divides $(k - 1)(j - 1)$.

Corollary 5.2. *A regular $2n$ -gon is infinitesimally rigid as a self-Bäcklund (n, k) -gon if n is odd, or if both n and k are even, $k < n/2$, and $\gcd(n, k) > 2$.*

Proof. The first statement of the corollary follows immediately from Theorem 11.

For the second statement, assume that a non-trivial infinitesimal deformation exists. We claim that k and j are coprime. Indeed, if $(j, k) = p$, then $n = 2(j + k) \equiv 0 \pmod p$, but $(j - 1)(k - 1) \equiv 1 \pmod p$. This contradicts the fact that n divides $(j - 1)(k - 1)$. It follows that

$$(n, k) = (2(j + k), k) = 2(j, k) = 2,$$

proving the second statement. □

Now we prove Theorem 11.

Proof. Let

$$P_j = \left(\cos\left(\frac{\pi j}{n}\right), \sin\left(\frac{\pi j}{n}\right) \right), \quad j = 1, \dots, 2n,$$

be the vertices of a regular $2n$ -gon. We have

$$[P_j, P_{j+1}] = \sin\left(\frac{\pi}{n}\right) = a, \quad [P_j, P_{j+k}] = \sin\left(\frac{\pi k}{n}\right) = b.$$

(One can rescale to have $a = 1$, but it is not really needed for the argument.)

We also have the respective second-order linear recurrence

$$(69) \quad P_{j+1} = 2 \cos\left(\frac{\pi}{n}\right) P_j - P_{j-1}.$$

Consider an infinitesimal deformation $P_j + \varepsilon V_j$, where V_j is an n -anti-periodic sequence of vectors, that is, $V_{j+n} = -V_j$ for all j , and assume that the resulting polygon is a self-Bäcklund (n, k) -gon. By applying a dilation, we may assume that the constant a does not change. Then, calculating modulo ε^2 , we obtain two systems of equations

$$(70) \quad [P_j, V_{j+1}] + [V_j, P_{j+1}] = 0, \quad j = 1, \dots, n,$$

and

$$(71) \quad [P_j, V_{j+k}] + [V_j, P_{j+k}] = C, \quad j = 1, \dots, n,$$

where C is a constant.

Consider the system (70). Let

$$V_j = a_j P_j + b_j P_{j+1} = c_j P_j + d_j P_{j-1}.$$

Then the recurrence (69) implies that

$$\frac{c_j - a_j}{b_j} = 2 \cos\left(\frac{\pi}{n}\right), \quad \frac{d_j}{b_j} = -1.$$

Substitute vectors V_j into equation (70) to obtain

$$(72) \quad a_j = -c_{j+1}, \quad b_j = \frac{c_j + c_{j+1}}{2 \cos(\pi/n)}, \quad d_j = -\frac{c_j + c_{j+1}}{2 \cos(\pi/n)},$$

where c_j is an n -periodic sequence to be determined.

Now consider the system (71). Substituting vectors V_j , using equation (72), and collecting terms yields the linear system

$$(73) \quad \mu_{k-1} c_j - \mu_{k+1} c_{j+1} + \mu_{k+1} c_{j+k} - \mu_{k-1} c_{j+k+1} = C, \quad j = 1, \dots, n,$$

where $\mu_k = \sin(\pi k/n)$.

First, we note that C must be zero. Indeed, add equation (73): the left hand side vanishes, and so must the right hand side.

Second, system (73) has a 3-dimensional space of trivial solutions that correspond to the action of the Lie algebra $\mathfrak{sl}_2(\mathbb{R})$. These solutions are given by the formulas

$$c_j = 1; \quad c_j = \cos\left(\frac{\pi(2j-1)}{n}\right); \quad c_j = \sin\left(\frac{\pi(2j-1)}{n}\right).$$

We need to find out when there are no other solutions.

To this end, consider the eigenvalues of the matrix defining the system (73). This is a circulant matrix, and its eigenvalues are given by the formula

$$\lambda_j = \mu_{k-1} - \mu_{k+1} \omega_j + \mu_{k+1} \omega_j^k - \mu_{k-1} \omega_j^{k+1}, \quad j = 0, \dots, n-1,$$

where $\omega_j = e^{i \frac{2\pi j}{n}}$, see [19].

We are interested in zero eigenvalues. One has $\lambda_j = 0$ if and only if

$$\omega_j^{k+1} = \frac{\mu_{k-1} - \mu_{k+1} \omega_j}{\mu_{k-1} - \mu_{k+1} \bar{\omega}_j}.$$

Let 2α be the argument of the unit complex number on the right. A direct calculation yields

$$\tan \alpha = -\frac{\sin\left(\frac{\pi(k+1)}{n}\right)\sin\left(\frac{2\pi j}{n}\right)}{\sin\left(\frac{\pi(k-1)}{n}\right) - \sin\left(\frac{\pi(k+1)}{n}\right)\cos\left(\frac{2\pi j}{n}\right)}.$$

The argument of ω_j^{k+1} is $2\pi j(k+1)/n$, hence (after cleaning up the formulas)

$$\sin\left(\frac{\pi j(k+1)}{n}\right)\sin\left(\frac{\pi(k-1)}{n}\right) = \sin\left(\frac{\pi j(k-1)}{n}\right)\sin\left(\frac{\pi(k+1)}{n}\right),$$

or, equivalently,

$$(74) \quad \tan\left(\frac{\pi j}{n}\right)\tan\left(\frac{\pi k}{n}\right) = \tan\left(\frac{\pi jk}{n}\right)\tan\left(\frac{\pi}{n}\right).$$

Note the trivial solutions $j = 0, 1, n - 1$, corresponding to the action of $\mathfrak{sl}_2(\mathbb{R})$. Let us assume that $2 \leq j \leq n - 2$.

One also has other trivial solutions, when both sides of equation (74) are infinite: $n = 2j$ and k odd, and $n = 2k$ and j odd. Note that, in the latter case, $k > 2$. Indeed, if $k = 2$, then $n = 4$, and since $2 \leq j \leq n - 2$, we have $j = 2$, contradicting that j is odd.

Equation (74) appeared in [41] and in [3], and it was solved in [17]. This equation has non-trivial solutions if and only if $n = 2(j+k)$ and n divides $(j-1)(k-1)$. This completes the proof. \square

Remark 5.3. As we know from Theorem 10, if n is even and k is odd, or if $n = 2k$, non-trivial self-Bäcklund (n, k) -gons indeed exist. The smallest values in case 3) of Theorem 11 are $k = 4, n = 30$. Does there exist a non-trivial self-Bäcklund $(30, 4)$ -gon?

Remark 5.4. One wonders whether the symmetry between k and j in the formulation of Theorem 11 corresponds to some kind of duality between self-Bäcklund (n, k) - and (n, j) -gons.

6. APPENDIX: FROM THE CENTROAFFINE PLANE TO THE HYPERBOLIC PLANE

In this appendix we connect two geometries associated with the group $SL_2(\mathbb{R})$, the centroaffine and the hyperbolic ones.

Consider the 3-dimensional space of quadratic forms $ax^2 + 2bxy + cy^2$ with the pseudo-Euclidean metric given by quadratic form $b^2 - ac$, the negative of the determinant of the quadratic form. The projectivization of the subspace of the positive-definite forms is the hyperbolic plane H^2 ; the degenerate forms comprise the circle at infinity. In the modern literature, this approach to hyperbolic geometry was developed in [5].

In the coordinates (u, v, w) , such that

$$a = u + v, \quad b = w, \quad c = u - v,$$

one has the standard Minkowski metric $v^2 + w^2 - u^2$. The unit-determinant quadratic forms comprise the hyperboloid of two sheets, and the condition $a + c > 0$ describes its upper half, the pseudo-sphere.

A “unit” central ellipse of area π is an $SL_2(\mathbb{R})$ image of the unit circle, given by an equation of the form $ax^2 + 2bxy + cy^2 = 1$ with $ac - b^2 = 1$ and $a + c > 0$. This defines a point of the hyperbolic plane H^2 in the pseudo-sphere model.

Likewise, a central hyperbola, which is an $SL_2(\mathbb{R})$ image of the “unit” hyperbola $xy = 1$, is given by an equation of the form $ax^2 + 2bxy + cy^2 = 1$ with $ac - b^2 = -1$. It defines a point of the hyperboloid of one sheet.

Lemma 6.1. *Let a unit central ellipse $ax^2 + 2bxy + cy^2 = 1$ and a unit central hyperbola $a'x^2 + 2b'xy + c'y^2 = 1$ be tangent at point (x, y) . Then the vectors (a, b, c) and (a', b', c') are orthogonal.*

Proof. The group $SL_2(\mathbb{R})$ acts transitively on the space of contact elements of the punctured plane whose line does not pass through the origin. And it acts by isometries on the space of quadratic forms. Therefore it suffices to consider the point $(1, 0)$ and the vertical direction. In this case the two conics are $x^2 + y^2 = 1$ and $x^2 - y^2 = 1$, and the vectors $(1, 0, 1)$ and $(1, 0, -1)$ are indeed orthogonal. \square

To a point (x, y) of the punctured plane there corresponds the affine plane $ax^2 + 2bxy + cy^2 = 1$ in the 3-dimensional space of quadratic forms. The normal vector of this plane is isotropic, and this plane lies above the origin. Hence its intersection with the pseudo-sphere is a horocycle in H^2 . The symmetric point $(-x, -y)$ yields the same horocycle.

To summarize, a point of the centroaffine plane is a horocycle in H^2 , and a unit central ellipse is a point of H^2 .

Let $\gamma(t)$ be a centroaffine curve. The *osculating ellipse* at a point $(x, y) = \gamma(t)$ is a unit central ellipse tangent to γ at this point. As t varies, one obtains a curve $\gamma^*(t) \subset H^2$, the dual curve of γ . Due to the central symmetry of γ , this curve closes up after t is increased by π . Equivalently, the curve γ^* is the envelope of the horocycles corresponding to the points of the curve γ .

Lemma 6.2. *If $[\gamma(t), \gamma'(t)] = 1$, then $|\gamma^*(t)'| = |1 + p(t)|$.*

Proof. Let $\gamma(t) = (x(t), y(t))$. Then $xy' - x'y = 1$.

The osculating ellipse at a point (x, y) satisfies the equations

$$ax^2 + 2bxy + cy^2 = 1, (ax + by, bx + cy) \cdot (x', y') = 0.$$

Taking $ac - b^2 = 1$ into account, one solves these equations to obtain

$$a = y^2 + y'^2, b = -(xy + x'y'), c = x^2 + x'^2.$$

This is the equation of γ^* .

Next, $x'' = px, y'' = py$. Then

$$(\gamma^*)' = (1 + p)(2yy', -(x'y + xy'), 2xx'),$$

and $|(\gamma^*)'| = |1 + p|$, as claimed. \square

Let k be curvature of the curve γ^* .

Lemma 6.3. *One has*

$$k = \frac{1 - p}{1 + p} \text{ or } (1 + p)(1 + k) = 2.$$

For example, when γ is a unit central ellipse with $p = -1$, the dual curve is a point, and the formula accordingly gives $k = \infty$. If γ is a unit central hyperbola with $p = 1$, then the formula gives $k = 0$. Indeed, Lemma 6.1 implies that γ^* is a straight line, the intersection of the pseudo-sphere with the 2-dimensional subspace orthogonal to the vector corresponding to this hyperbola.

Proof. Let τ be the arc length parameter on γ^* . Then $dt/d\tau = 1/(1+p)$.

The curvature is the magnitude of the projection of the vector $d^2\gamma^*/d\tau^2$ on the pseudo-sphere. If u is a position vector of a point of the pseudo-sphere and v is a vector with foot point u , then the projection of u is given by $u + (u \cdot v)v$.

From Lemma 6.3, we know that

$$\frac{d\gamma^*}{d\tau} = (2yy', -(x'y + xy'), 2xx'),$$

hence

$$\frac{d^2\gamma^*}{d\tau^2} = \frac{1}{1+p}(2yy', -(x'y + xy'), 2xx')' = \frac{2}{1+p}(py^2 + y'^2, -pxy - x'y', px^2 + x'^2).$$

Next,

$$\frac{d\gamma^*}{d\tau} \cdot \gamma^* = 0 \Rightarrow \frac{d^2\gamma^*}{d\tau^2} \cdot \gamma^* + \frac{d\gamma^*}{d\tau} \cdot \frac{d\gamma^*}{d\tau} = 0 \Rightarrow \frac{d^2\gamma^*}{d\tau^2} \cdot \gamma^* = -1,$$

therefore the projection of $d^2\gamma^*/d\tau^2$ on the pseudo-sphere is

$$\begin{aligned} \frac{d^2\gamma^*}{d\tau^2} - \gamma^* &= \frac{2}{1+p}(py^2 + y'^2, -pxy - x'y', px^2 + x'^2) - \\ &(y^2 + y'^2, -(xy + x'y'), x^2 + x'^2) = \frac{1-p}{1+p}(y'^2 - y^2, xy - x'y', x'^2 - x^2), \end{aligned}$$

and it remains to notice that the vector in the parentheses is unit. \square

Remark 6.4. According to a theorem of E. Ghys, see [36], the potential $p(t)$ of the curve γ assumes the value -1 at least four times on the period $[0, \pi)$. It follows that the curve γ^* has at least four cusps; in particular, it cannot be smooth.

ACKNOWLEDGMENTS

We are grateful to L. Buhovsky, D. Fuchs, E. Gluskin, A. Izosimov, M. Cuntz, A. Mironov, A. Sodin, and F. Wegner for fruitful discussions. We thank the referees for their constructive criticisms and suggestions.

REFERENCES

- [1] N. I. Akhiezer, *Elements of the theory of elliptic functions*, Translations of Mathematical Monographs, vol. 79, American Mathematical Society, Providence, RI, 1990. Translated from the second Russian edition by H. H. McFaden, DOI 10.1090/mmono/079. MR1054205
- [2] M. Arnold, D. Fuchs, and S. Tabachnikov, *A family of integrable transformations of centroaffine polygons: geometrical aspects*, arXiv:2112.08124, 2021.
- [3] Tarik Aougab, Xidian Sun, Serge Tabachnikov, and Yuwen Wang, *On curves and polygons with the equiangular chord property*, Pacific J. Math. **274** (2015), no. 2, 305–324, DOI 10.2140/pjm.2015.274.305. MR3332906
- [4] V. I. Arnol'd, *Huygens and Barrow, Newton and Hooke*, Birkhäuser Verlag, Basel, 1990. Pioneers in mathematical analysis and catastrophe theory from evolvents to quasicrystals; Translated from the Russian by Eric J. F. Primrose, DOI 10.1007/978-3-0348-9129-5. MR1078625
- [5] V. Arnold, *Lobachevsky triangle altitudes theorem as the Jacobi identity in the Lie algebra of quadratic forms on symplectic plane*, J. Geom. Phys. **53** (2005), no. 4, 421–427, DOI 10.1016/j.geomphys.2004.07.008. MR2125401
- [6] Maxim Arnold, Dmitry Fuchs, Ivan Izmestiev, and Serge Tabachnikov, *Cross-ratio dynamics on ideal polygons*, Int. Math. Res. Not. IMRN **9** (2022), 6770–6853, DOI 10.1093/imrn/rnaa289. MR4411469
- [7] H. Auerbach, *Sur un problème de M. Ulam concernant l'équilibre des corps flottants*, Studia Math. **7** (1938), 121–142.

- [8] Misha Bialy, Andrey E. Mironov, and Lior Shalom, *Magnetic billiards: non-integrability for strong magnetic field; Gutkin type examples*, J. Geom. Phys. **154** (2020), 103716, 14, DOI 10.1016/j.geomphys.2020.103716. MR4099906
- [9] Misha Bialy, Andrey E. Mironov, and Lior Shalom, *Outer billiards with the dynamics of a standard shift on a finite number of invariant curves*, Exp. Math. **30** (2021), no. 4, 469–474, DOI 10.1080/10586458.2018.1563514. MR4346647
- [10] Misha Bialy, Andrey E. Mironov, and Serge Tabachnikov, *Wire billiards, the first steps*, Adv. Math. **368** (2020), 107154, 27, DOI 10.1016/j.aim.2020.107154. MR4085879
- [11] Gil Bor, Mark Levi, Ron Perline, and Sergei Tabachnikov, *Tire tracks and integrable curve evolution*, Int. Math. Res. Not. IMRN **9** (2020), 2698–2768, DOI 10.1093/imrn/rny087. MR4095423
- [12] G. Bor, *Centroaffine curves*, <https://www.cimat.mx/~gil/centro-affine/>, 2017.
- [13] J. Bracho, L. Montejano, and D. Oliveros, *A classification theorem for Zindler carrousel*, J. Dynam. Control Systems **7** (2001), no. 3, 367–384, DOI 10.1023/A:1013099830164. MR1848363
- [14] J. Bracho, L. Montejano, and D. Oliveros, *Carousels, Zindler curves and the floating body problem*, Period. Math. Hungar. **49** (2004), no. 2, 9–23, DOI 10.1007/s10998-004-0519-6. MR2106462
- [15] Annalisa Calini, Thomas Ivey, and Gloria Mari-Beffa, *Remarks on KdV-type flows on star-shaped curves*, Phys. D **238** (2009), no. 8, 788–797, DOI 10.1016/j.physd.2009.01.007. MR2522973
- [16] Ildelfonso Castro, Ildelfonso Castro-Infantes, and Jesús Castro-Infantes, *New plane curves with curvature depending on distance from the origin*, Mediterr. J. Math. **14** (2017), no. 3, Paper No. 108, 19, DOI 10.1007/s00009-017-0912-z. MR3633368
- [17] Robert Connelly and Balázs Csikós, *Classification of first-order flexible regular bicycle polygons*, Studia Sci. Math. Hungar. **46** (2009), no. 1, 37–46, DOI 10.1556/SScMath.2008.1074. MR2656480
- [18] Van Cyr, *A number theoretic question arising in the geometry of plane curves and in billiard dynamics*, Proc. Amer. Math. Soc. **140** (2012), no. 9, 3035–3040, DOI 10.1090/S0002-9939-2012-11258-4. MR2917076
- [19] Philip J. Davis, *Circulant matrices*, A Wiley-Interscience Publication, John Wiley & Sons, New York-Chichester-Brisbane, 1979. MR543191
- [20] Serge Tabachnikov and Filiz Dogru, *Dual billiards*, Math. Intelligencer **27** (2005), no. 4, 18–25, DOI 10.1007/BF02985854. MR2183864
- [21] P. G. Drazin and R. S. Johnson, *Solitons: an introduction*, Cambridge Texts in Applied Mathematics, Cambridge University Press, Cambridge, 1989, DOI 10.1017/CBO9781139172059. MR985322
- [22] M. S. P. Eastham, *The spectral theory of periodic differential equations*, Texts in Mathematics (Edinburgh), Scottish Academic Press, Edinburgh; Hafner Press, New York, 1973. MR3075381
- [23] Robert Foote, Mark Levi, and Serge Tabachnikov, *Tractrices, bicycle tire tracks, hatchet planimeters, and a 100-year-old conjecture*, Amer. Math. Monthly **120** (2013), no. 3, 199–216, DOI 10.4169/amer.math.monthly.120.03.199. MR3030293
- [24] A. P. Fordy, *A historical introduction to solitons and Bäcklund transformations*, Harmonic maps and integrable systems, Aspects Math., E23, Friedr. Vieweg, Braunschweig, 1994, pp. 7–28, DOI 10.1007/978-3-663-14092-4_2. MR1264180
- [25] Atsushi Fujioka and Takashi Kurose, *Hamiltonian formalism for the higher KdV flows on the space of closed complex equicentroaffine curves*, Int. J. Geom. Methods Mod. Phys. **7** (2010), no. 1, 165–175, DOI 10.1142/S0219887810003987. MR2647776
- [26] Atsushi Fujioka and Takashi Kurose, *Multi-Hamiltonian structures on spaces of closed equicentroaffine plane curves associated to higher KdV flows*, SIGMA Symmetry Integrability Geom. Methods Appl. **10** (2014), Paper 048, 11, DOI 10.3842/SIGMA.2014.048. MR3210587
- [27] Eugene Gutkin, *Capillary floating and the billiard ball problem*, J. Math. Fluid Mech. **14** (2012), no. 2, 363–382, DOI 10.1007/s00021-011-0071-0. MR2925114
- [28] H. Guggenheimer, *Hill equations with coexisting periodic solutions*, J. Differential Equations **5** (1969), 159–166, DOI 10.1016/0022-0396(69)90109-0. MR239193
- [29] Eberhard Hopf, *Closed surfaces without conjugate points*, Proc. Nat. Acad. Sci. U.S.A. **34** (1948), 47–51, DOI 10.1073/pnas.34.2.47. MR23591
- [30] Mark Levi and Serge Tabachnikov, *On bicycle tire tracks geometry, hatchet planimeter, Menzin's conjecture, and oscillation of unicycle tracks*, Experiment. Math. **18** (2009), no. 2, 173–186. MR2549686
- [31] Wilhelm Magnus and Stanley Winkler, *Hill's equation*, Interscience Tracts in Pure and Applied Mathematics, No. 20, Interscience Publishers John Wiley & Sons, New York-London-Sydney, 1966. MR0197830
- [32] Horst Martini and Konrad J. Swanepoel, *Antinorms and Radon curves*, Aequationes Math. **72** (2006), no. 1-2, 110–138, DOI 10.1007/s00010-006-2825-y. MR2258811

- [33] Nozomu Matsuura, *Discrete KdV and discrete modified KdV equations arising from motions of planar discrete curves*, Int. Math. Res. Not. IMRN **8** (2012), 1681–1698, DOI 10.1093/imrn/rnr080. MR2920827
- [34] R. Daniel Mauldin (ed.), *The Scottish Book*, 2nd ed., Birkhäuser/Springer, Cham, 2015. Mathematics from the Scottish Café with selected problems from the new Scottish Book; Including selected papers presented at the Scottish Book Conference held at North Texas University, Denton, TX, May 1979, DOI 10.1007/978-3-319-22897-6. MR3242261
- [35] Sophie Morier-Genoud, *Coxeter’s frieze patterns at the crossroads of algebra, geometry and combinatorics*, Bull. Lond. Math. Soc. **47** (2015), no. 6, 895–938, DOI 10.1112/blms/bdv070. MR3431573
- [36] V. Ovsienko and S. Tabachnikov, *Sturm theory, Ghys theorem on zeroes of the Schwarzian derivative and flattening of Legendrian curves*, Selecta Math. (N.S.) **2** (1996), no. 2, 297–307, DOI 10.1007/BF01587937. MR1414890
- [37] G. Pastras, *Four lectures on Weierstrass elliptic function and applications in classical and quantum mechanics*, arXiv:1706.07371.
- [38] Ulrich Pinkall, *Hamiltonian flows on the space of star-shaped curves*, Results Math. **27** (1995), no. 3-4, 328–332, DOI 10.1007/BF03322836. MR1331105
- [39] P. Reinhardt and P. L. Walker, *Weierstrass elliptic and modular functions*, NIST Handbook Math. Functions, chap. 23, Cambridge Univ. Press, Cambridge, 2010, <https://dlmf.nist.gov/23>.
- [40] David A. Singer, *Curves whose curvature depends on distance from the origin*, Amer. Math. Monthly **106** (1999), no. 9, 835–841, DOI 10.2307/2589616. MR1732664
- [41] Serge Tabachnikov, *Tire track geometry: variations on a theme*, Israel J. Math. **151** (2006), 1–28, DOI 10.1007/BF02777353. MR2214115
- [42] Serge Tabachnikov, *Variations on R. Schwartz’s inequality for the Schwarzian derivative*, Discrete Comput. Geom. **46** (2011), no. 4, 724–742, DOI 10.1007/s00454-011-9371-7. MR2846176
- [43] Serge Tabachnikov, *On the bicycle transformation and the filament equation: results and conjectures*, J. Geom. Phys. **115** (2017), 116–123, DOI 10.1016/j.geomphys.2016.05.013. MR3623617
- [44] Serge Tabachnikov, *On centro-affine curves and Bäcklund transformations of the KdV equation*, Arnold Math. J. **4** (2018), no. 3-4, 445–458, DOI 10.1007/s40598-019-00105-y. MR3949812
- [45] S. Tabachnikov and E. Tsukerman, *On the discrete bicycle transformation*, Publ. Mat. Urug. **14** (2013), 201–219. MR3235356
- [46] Chuu-Lian Terng and Zhiwei Wu, *Central affine curve flow on the plane*, J. Fixed Point Theory Appl. **14** (2013), no. 2, 375–396, DOI 10.1007/s11784-014-0161-8. MR3248564
- [47] F. Wegner, *Floating bodies of equilibrium in 2D, the tire track problem and electrons in a parabolic magnetic field*, arXiv:physics/0701241.
- [48] Franz Wegner, *Floating bodies of equilibrium*, Stud. Appl. Math. **111** (2003), no. 2, 167–183, DOI 10.1111/1467-9590.t01-1-00231. MR1990237
- [49] F. Wegner, *Three problems – one solution*, <http://www.tphys.uni-heidelberg.de/~wegner/F12mvs/Movies.html#float>.
- [50] Konrad Von Zindler, *Über konvexe Gebilde* (German), Monatsh. Math. Phys. **31** (1921), no. 1, 25–56, DOI 10.1007/BF01702711. MR1549095

SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, ISRAEL
 Email address: bialy@post.tau.ac.il

CIMAT, A.P. 402, GUANAJUATO, GTO. 36000, MEXICO
 Email address: gil@cimat.mx

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY
 Email address: tabachni@math.psu.edu