

PARUSIŃSKI'S “KEY LEMMA” VIA ALGEBRAIC GEOMETRY

Z. REICHSTEIN AND B. YOUSIN

(Communicated by David Kazhdan)

ABSTRACT. The following “Key Lemma” plays an important role in the work by Parusiński on the existence of Lipschitz stratifications in the class of semi-analytic sets: For any positive integer n , there is a finite set of homogeneous symmetric polynomials W_1, \dots, W_N in $\mathbb{Z}[x_1, \dots, x_n]$ and a constant $M > 0$ such that

$$|dx_i/x_i| \leq M \max_{j=1, \dots, N} |dW_j/W_j|,$$

as densely defined functions on the tangent bundle of \mathbb{C}^n . We give a new algebro-geometric proof of this result.

1. INTRODUCTION

Parusiński's fundamental work on the existence of Lipschitz stratifications in the class of semianalytic sets relies on the following result.

Theorem 1.1 (Parusiński [P, pp. 202–203]). *For any positive integer n , there is a finite set of homogeneous symmetric polynomials $W_1, \dots, W_N \in \mathbb{Z}[x_1, \dots, x_n]$ and a constant $M > 0$ such that*

$$(1.1) \quad \left| \frac{dx_i}{x_i}(p, v) \right| \leq M \max_{j=1, \dots, N} \left| \frac{dW_j}{W_j}(p, v) \right|$$

for all $p \in \mathbb{C}^n$ and $v \in T_p\mathbb{C}^n$ for which both sides are defined. Here for any $P \in \mathbb{C}[x_1, \dots, x_n]$ we view the meromorphic differential form $\frac{dP}{P}$ on \mathbb{C}^n as a densely defined function on the total space of the tangent bundle $T\mathbb{C}^n$.

Parusiński refers to Theorem 1.1 as the “Key Lemma”; the proof of this result in [P, Section 6] is quite difficult, being apparently the hardest part of [P]. The purpose of this paper is to show that Theorem 1.1, in spite of its analytic appearance, has a natural proof in the framework of algebraic geometry. Our argument is an application of the results of [RY] about group actions on algebraic varieties; these results, in turn, rely on canonical resolution of singularities.

We remark that Parusiński proves the inequality (1.1) under the additional assumption that $dV(p, v) = 0$ if $V(p) = 0$ for every V belonging to a finite set \mathcal{V} of polynomials. Since this additional requirement does not affect a dense Zariski open subset of $T\mathbb{C}^m$ (given by $V(p) \neq 0$ for every $V \in \mathcal{V}$), it can be dropped. We also note

Received by the editors October 16, 1999.

1991 *Mathematics Subject Classification*. Primary 14E15, 14F10, 14L30; Secondary 16S35, 32B10, 58A40.

Z. Reichstein was partially supported by NSF grant DMS-9801675 and (during his stay at MSRI) by NSF grant DMS-9701755.

that the statement of the Key Lemma in [P] only asserts the existence of polynomials W_1, \dots, W_N with real coefficients; however, the construction of W_1, \dots, W_N given there, produces polynomials over \mathbb{Z} . Thus, while Theorem 1.1 appears to be stronger than the "Key Lemma" in [P], the two are, in fact, equivalent.

We are grateful to R. Guralnick and S. Montgomery for informative conversations on the subject of group algebras, and to D. Kazhdan for helpful comments on an earlier draft of this paper.

2. PRELIMINARIES

Notational conventions. All algebraic varieties considered in this paper, are assumed to be irreducible and defined over a field k of characteristic 0. The base field k is not assumed to be algebraically closed; the two cases of interest to us are $k = \mathbb{Q}$ and $k = \mathbb{C}$. By a point of a variety we shall always understand a closed point. Given an embedding $k \subset \mathbb{C}$ and a rational function f on X , we shall denote the corresponding rational function on $X_{\mathbb{C}} = X \otimes_k \mathbb{C}$ by f as well.

Throughout this paper G will be a finite group. A G -variety X is a variety with a regular action of G , $G \times X \rightarrow X$, where $G \times X$ is understood as the disjoint union of $|G|$ copies of X . We will always assume that the G -action is faithful, i.e., every nonidentity element of G acts nontrivially. By a morphism (respectively, rational map, birational isomorphism) of G -varieties we shall mean a G -equivariant morphism (respectively, rational map, birational isomorphism).

Stabilizers. For a point x in a G -variety X , we define its "naive" stabilizer $\text{NStab}(x)$ as the set of all $g \in G$ which preserve x . If k is not algebraically closed, the residue field $k(x)$ may be a nontrivial finite extension of k , and G may act on it nontrivially. We define the "honest" stabilizer $\text{Stab}(x)$ as the set of all $g \in \text{NStab}(x)$ that act on $k(x)$ trivially; cf. [MFK, Definition 0.4]. The subgroups $\text{NStab}(x)$ and $\text{Stab}(x)$ of G are sometimes called the *decomposition group* and the *inertia group* respectively.

If \bar{k} is the algebraic closure of k , then x is represented by a set of "conjugate" points of the variety $X_{\bar{k}} = X \otimes_k \bar{k}$ ("geometric points" of X), one for each embedding $k(x) \hookrightarrow \bar{k}$; $\text{Stab}(x)$ fixes each of these points while $\text{NStab}(x)$ permutes them. As an example, consider the action of $G = \mathbb{Z}/2\mathbb{Z}$ on the affine line $\mathbb{A}_{\mathbb{Q}}^1 = \text{Spec } \mathbb{Q}[t]$: the nontrivial element of this group acts by $t \mapsto -t$. Here $\text{Stab}(x) = \{1\}$ for any $x \in \mathbb{A}_{\mathbb{Q}}^1 - \{0\}$. On the other hand, $\text{NStab}(x) = G$ iff x corresponds to the ideal in $\mathbb{Q}[t]$ generated by an irreducible polynomial of the form $q(t^2)$ (e.g., $t^2 + 1$).

This phenomenon is entirely arithmetic; we are concerned with it here because the symmetric polynomials W_1, \dots, W_N in Theorem 1.1 are asserted to have integer coefficients. A reader who is only interested in the existence of such polynomials in $\mathbb{C}[x_1, \dots, x_n]$ may skip the rest of this section and assume that "naive" stabilizers always coincide with "honest" ones in the sequel.

Semilinear representations and skew group rings. Let X be a G -variety and let $x \in X$. The "naive stabilizer" $\text{NStab}(x)$ acts upon $T_x(X)^* = \mathfrak{m}_x/\mathfrak{m}_x^2$. However, if $\text{NStab}(x)$ is strictly larger than $\text{Stab}(x)$, then this action is not linear over $k(x)$ but rather "semilinear" in the following sense.

Definition 2.1. Suppose a finite group H acts by automorphisms on a field K . A *semilinear representation* of H over K is a K -vector space V with a K^H -linear

action of H on V having the property $g(\lambda v) = g(\lambda)g(v)$ for any $g \in H$, $\lambda \in K$ and $v \in V$.

For the rest of this section we shall assume that K is a field, K^* is the multiplicative group of K , H is a finite group acting on K by automorphisms, and H' is the kernel of this action. In the subsequent applications we will take $K = k(x)$, $H = \text{NStab}(x)$ and $H' = \text{Stab}(x)$.

Recall that the skew group algebra $K * H$ is defined as the set of formal sums $\sum_{h \in H} a_h h$ (where $a_h \in K$), with componentwise addition and with multiplication given, distributively, by $(a_1 h_1)(a_2 h_2) = a_1 h_1(a_2) h_1 h_2$. A semilinear representation of H is the same thing as a $(K * H)$ -module. (All modules in this paper are understood to be left modules.)

Remark 2.2. Note that $V = K$ has a natural structure of a $(K * H)$ -module. This module contains a vector $1 \in K$ which is fixed by H .

Recall that by Wedderburn's Theorem every semisimple ring R is a direct product of simple rings, called *the simple components* of R ; see [B, Theorem VIII.5.1].

Lemma 2.3. *$K * H$ is a semisimple K^H -algebra with at most $|H'|$ simple components. (Here H' is the kernel of the H -action on K , as above.)*

Proof. Semisimplicity of $K * H$ is proved by the same averaging argument as the usual Maschke's theorem; for details see, e.g., [Mo, Theorem 0.1 and Corollary 0.2].

Denote the simple components of $K * H$ by S_1, \dots, S_m . Then $Z(K * H) = Z(S_1) \times \dots \times Z(S_m)$, where $Z(A)$ denotes the center of A . It is easy to see directly that $\dim_{K^H}(Z(K * H)) \leq |H'|$. Hence, $m \leq |H'|$, as claimed. \square

The following proposition describes the particular kind of skew group rings we shall encounter in the sequel.

Proposition 2.4. *Suppose that H' is an abelian group of exponent e , K contains a primitive e th root of unity and χ_1, \dots, χ_m is a set of generators for the dual group $(H')^* = \text{Hom}(H', K^*)$. Assume further that for each i there is a one-dimensional semilinear representation V_i of H over K such that $h'(v) = \chi_i(h')v$ for every $v \in V_i$ and every $h' \in H'$. Then:*

- (a) *For every $\chi \in (H')^*$, there exists a unique semilinear representation V_χ of H such that $\dim_K(V_\chi) = 1$ and $h'(v) = \chi(h')v$ for every $v \in V_\chi$ and every $h' \in H'$.*
- (b) *Every simple $(K * H)$ -module is isomorphic to V_χ for some $\chi \in (H')^*$.*

Proof. Note that if V_1 and V_2 are semilinear representations of H over K , then so is $V_1 \otimes_K V_2$. Indeed,

$$h(\lambda v_1) \otimes h(v_2) = h(\lambda) \cdot (h(v_1) \otimes h(v_2)) = h(v_1) \otimes h(\lambda v_2).$$

To construct V_χ , write $\chi \in (H')^*$ as $\chi = \chi_1^{l_1} \dots \chi_m^{l_m}$ for some nonnegative integers l_1, \dots, l_m , and set $V_\chi = V_1^{\otimes l_1} \otimes_K \dots \otimes_K V_m^{\otimes l_m}$. The subgroup H' acts on V_χ by the character χ , as desired. As $\dim_K(V_\chi) = 1$, V_χ is a simple $(K * H)$ -module. Note that the $(K * H)$ -modules V_χ are pairwise nonisomorphic because H' acts on them by different characters.

The isomorphism classes of simple $(K * H)$ -modules are in 1–1 correspondence with the simple components of $K * H$; see [B, Proposition VIII.5.11]. Thus Lemma 2.3 implies that $K * H$ has $\leq |H'|$ nonisomorphic simple modules. On the other

hand, we have constructed $|H'|$ nonisomorphic simple modules V_χ . This proves (b) and the uniqueness of V_χ in (a). \square

Remark 2.5. One can show that, under the assumptions of Proposition 2.4, H is a semidirect product of H' and H/H' , where the action of H/H' on H' is given by embedding H' into $(K^*)^m$ via $h' \mapsto (\chi_1(h'), \dots, \chi_m(h'))$.

3. REDUCTION TO AN ALGEBRO-GEOMETRIC PROBLEM

We begin by restating (1.1) as an inequality involving densely defined functions on the tangent bundle of $\mathbb{P}_{\mathbb{C}}^{n-1}$ rather than the tangent bundle of \mathbb{C}^n . Since $\mathbb{P}_{\mathbb{C}}^{n-1}$ is compact in the metric topology, this will allow us to pass from local to global estimates.

Proposition 3.1. *Let X be a projective G -variety over a field $k \subset \mathbb{C}$, and let f be a rational function on X . Then there exist G -invariant rational functions β_1, \dots, β_m on X and a constant $K > 0$ such that*

$$(3.1) \quad \left| \frac{df}{f}(p, v) \right| \leq K \max_{j=1, \dots, m} \left| \frac{d\beta_j}{\beta_j}(p, v) \right|$$

for any $(p, v) \in T(X_{\mathbb{C}})$ such that p is a smooth point of $X_{\mathbb{C}}$ and does not lie on the divisors of $f, \beta_1, \dots, \beta_m$.

Reduction 3.2. *Proposition 3.1 \implies Theorem 1.1.*

Indeed, apply Proposition 3.1 with $k = \mathbb{Q}$, $X = \mathbb{P}_{\mathbb{Q}}^{n-1}$, $G = S_n$ and $f = x_1/s_1$; here S_n acts on $\mathbb{P}_{\mathbb{Q}}^{n-1}$ by permuting the homogeneous coordinates x_1, \dots, x_n , and $s_1 = x_1 + \dots + x_n$. Write each β_j as a quotient of two homogeneous polynomials (of the same degree, with integer coefficients) in x_1, \dots, x_n :

$$\beta_1 = W_1/W_2, \dots, \beta_m = W_{2m-1}/W_{2m}.$$

We claim that the polynomials $W_1, \dots, W_{2m}, W_{2m+1} \stackrel{\text{def}}{=} s_1$ have the property asserted in Theorem 1.1. Indeed, since $df/f = dx_1/x_1 - ds_1/s_1$ and $d\beta_i/\beta_i = dW_{2i-1}/W_{2i-1} - dW_{2i}/W_{2i}$, inequality (3.1) translates into

$$\begin{aligned} \left| \left(\frac{dx_1}{x_1} - \frac{ds_1}{s_1} \right) (p, v) \right| &\leq K \max_{i=1, \dots, m} \left| \left(\frac{dW_{2i-1}}{W_{2i-1}} - \frac{dW_{2i}}{W_{2i}} \right) (p, v) \right| \\ &\leq 2K \max_{j=1, \dots, 2m} \left| \frac{dW_j}{W_j}(p, v) \right|. \end{aligned}$$

Consequently,

$$\begin{aligned} \left| \frac{dx_1}{x_1}(p, v) \right| &\leq 2K \max_{j=1, \dots, 2m} \left| \frac{dW_j}{W_j}(p, v) \right| + \left| \frac{ds_1}{s_1}(p, v) \right| \\ &\leq (2K + 1) \max_{j=1, \dots, 2m+1} \left| \frac{dW_j}{W_j}(p, v) \right|. \end{aligned}$$

This means that (1.1) holds for $i = 1$, with $M = 2K + 1$ and $N = 2m + 1$. By symmetry, (1.1) holds for all i . \square

Definition 3.3. Let X be a projective G -variety and f be a rational function on X . We shall say that the pair (X, f) has property (*) if there is a Zariski open

covering $X = \bigcup_i U_i$ and, for each i , rational functions $\beta_{i1}, \dots, \beta_{i,q_i} \in k(X)^G$ and regular functions $\gamma_{i1}, \dots, \gamma_{i,q_i} \in \mathcal{O}_X(U_i)$ such that

$$(3.2) \quad \frac{df}{f} = \gamma_{i1} \frac{d\beta_{i1}}{\beta_{i1}} + \dots + \gamma_{i,q_i} \frac{d\beta_{i,q_i}}{\beta_{i,q_i}}.$$

In other words, the pair (X, f) has property (*) if df/f is a global section of the sheaf of differentials on X generated over \mathcal{O}_X by $d\beta/\beta$, as β ranges over some finite subset of $k(X)^G$ (or, equivalently, as β ranges over all of $k(X)^G$).

Reduction 3.4. *Proposition 3.1 holds, assuming the pair (X, f) that appears there, has property (*).*

Indeed, the Zariski open cover $\bigcup_i U_i$ of X , as in Definition 3.3, gives rise to a Zariski open cover $\bigcup_i U_{i,\mathbb{C}}$ of $X_{\mathbb{C}}$. The functions γ_{ij} are continuous on $U_{i,\mathbb{C}}$ with respect to the metric topology. Thus any point $x \in X$ has an open neighborhood U_x (in the metric topology) such that $U_x \subset U_{i_x,\mathbb{C}}$ for some i_x , and

$$\left| \frac{df}{f}(p, v) \right| \leq K_x \max_{j=1, \dots, q_{i_x}} \left| \frac{d\beta_{i_x,j}}{\beta_{i_x,j}}(p, v) \right|$$

whenever $v \in T_p(X)$ and p is a smooth point of U_x which does not lie on the divisors of $f, \beta_{i_x,1}, \dots, \beta_{i_x,q_{i_x}}$. The open sets U_x form a cover of X ; since X is compact in the metric topology, we can choose a finite subcover U_{x_1}, \dots, U_{x_r} . Now if $K > K_{x_1}, \dots, K_{x_r}$, then

$$\left| \frac{df}{f}(p, v) \right| \leq K \max_{i,j} \left| \frac{d\beta_{ij}}{\beta_{ij}}(p, v) \right|.$$

This shows that Proposition 3.1 holds. \square

Reduction 3.5. *Suppose X and X' are birationally isomorphic G -varieties over $k \subset \mathbb{C}$. If Proposition 3.1 holds for X and $f \in k(X)$, then it holds for X' and the same $f \in k(X') = k(X)$.*

Indeed, X and X' have isomorphic Zariski-open subsets U and U' . After passing to smaller subsets if necessary, we may assume that U and U' are smooth and do not intersect the divisors of $f, \beta_1, \dots, \beta_n$ on X and X' respectively. Thus if inequality (3.1) holds for every (p, v) such that $p \in U_{\mathbb{C}}$ and $v \in T_p(X_{\mathbb{C}})$, then it holds for every (p, v) such that $p \in U'_{\mathbb{C}}$ and $v \in T_p(X'_{\mathbb{C}})$. The subset $U'_{\mathbb{C}}$ is dense in $X'_{\mathbb{C}}$ with respect to the metric topology; hence, by continuity the same inequality (with the same β_j and the same K) holds for every $(p, v) \in T(X'_{\mathbb{C}})$ such that p is a smooth point of $X'_{\mathbb{C}}$ and does not lie in the union of divisors of $f, \beta_1, \dots, \beta_m$. This means that Proposition 3.1 holds for X' as claimed. \square

We have thus shown that Theorem 1.1 is a consequence of the following, purely algebraic statement (see Reductions 3.2, 3.4 and 3.5).

Proposition 3.6. *Let G be a finite group, X a projective G -variety, and $f \in k(X)$. Then there exists a birational morphism $\pi: X' \rightarrow X$ of G -varieties such that the pair $(X', \pi^*(f))$ has property (*) (see Definition 3.3).*

A proof of Proposition 3.6 (and thus of Theorem 1.1) will be given in the next section. The idea is to construct $\pi: X' \rightarrow X$ by resolving the G -action on X to “standard form” with respect to a divisor containing the divisor of f ; see below. The simplest (affine) example of such X' is $X' = \mathbb{A}^1 = \text{Spec } k[t]$, where k contains a primitive m th root of unity, $G = \mathbb{Z}/m\mathbb{Z}$ acts on \mathbb{A}^1 linearly by a faithful character,

and $f = t$. In this case we can take $\beta = t^m \in k(X)^G$; the equality $df/f = \frac{1}{m}d\beta/\beta$ shows that (X', f) has property (*).

4. CONCLUSION OF THE PROOF

***G*-varieties in standard form.**

Definition 4.1 ([RY, Definition 3.1]). We say that a generically free *G*-variety X is in standard form with respect to a divisor Y if

- (a) X is smooth and Y is a normal crossing divisor on X ,
- (b) the G -action on $X - Y$ is free, and
- (c) for every $g \in G$ and for every irreducible component Y_0 of Y either $g(Y_0) = Y_0$ or $g(Y_0) \cap Y_0 = \emptyset$.

Theorem 4.2 ([RY, Corollary 3.6]). *Let X be a G -variety and $Y \subset X$ be a Zariski closed G -invariant subvariety such that the action of G on $X - Y$ is free. Then there is a sequence of blowups*

$$(4.1) \quad \pi: X_n \xrightarrow{\pi_n} X_{n-1} \xrightarrow{\pi_{n-1}} \dots \xrightarrow{\pi_2} X_1 \xrightarrow{\pi_1} X_0 = X$$

with smooth G -invariant centers $C_i \subset X_i$ such that X_n is in standard form with respect to a divisor \tilde{Y} containing $\pi^{-1}(Y)$.

Theorem 4.3. *Let X be a G -variety in standard form with respect to a divisor Y , let x be a point of X , let Y_1, \dots, Y_m be the irreducible components of Y passing through x , and let $W = Y_1 \cap \dots \cap Y_m$. Then*

- (a) [RY, Theorem 4.1] $\text{Stab}(x)$ is commutative.
- (b) [RY, Remark 4.4] *The action of $\text{Stab}(x)$ on the normal space to W at x is faithful and decomposes into the sum of one-dimensional representations as follows:*

$$(4.2) \quad T_x(X)/T_x(W) = \bigoplus_{i=1}^m \frac{T_x(Y_1) \cap \dots \cap \widehat{T_x(Y_i)} \cap \dots \cap T_x(Y_m)}{T_x(W)}.$$

- (c) [RY, Remark 4.5] *Let e be the exponent of $\text{Stab}(x)$. Then the residue field $k(x)$ of x contains a primitive e th root of unity.*

Remark 4.4. Under the assumptions of Theorem 4.3, set $H = \text{NStab}(x)$ and $H' = \text{Stab}(x)$. Recall that H acts on $T_x(X)$ semilinearly; see Definition 2.1. Property (c) of Definition 4.1 implies that Y_i is preserved by the action of H , and hence, the subspace $T_x(Y_i)$ is H -invariant for each i . It follows that all spaces appearing in (4.2), are $(k(x) * H)$ -modules.

The conormal space $(T_x(X)/T_x(Y_i))^*$ is dual to the i th summand in (4.2); it is a $(k(x) * H)$ -module of dimension 1 over $k(x)$, and H' acts on it by a character. Denote this character by ξ_i . Theorem 4.3(b) implies that the characters ξ_1, \dots, ξ_m generate the dual group $(H')^*$. Combining this observation with Theorem 4.3(c), we conclude that Proposition 2.4 applies in this setting.

A local coordinate system. Suppose that X is an algebraic variety and x is a point of X . Recall that $u_1, \dots, u_n \in \mathfrak{m}_x$ are said to form a local coordinate system on X at x if their classes modulo \mathfrak{m}_x^2 form a basis of $\mathfrak{m}_x/\mathfrak{m}_x^2$ as a $k(x)$ -vector space.

Proposition 4.5. *Let X be a quasiprojective G -variety in standard form with respect to a divisor Y . Suppose Y_1, \dots, Y_m are the irreducible components of Y passing through a point x of X . Then there exists a local coordinate system $u_1, \dots, u_m, v_1, \dots, v_l$ at x with the following properties:*

- (a) *Let $\bar{u}_i = u_i \bmod \mathfrak{m}_x^2$ and $\bar{v}_j = v_j \bmod \mathfrak{m}_x^2$. Then each of $\bar{u}_1, \dots, \bar{u}_m, \bar{v}_1, \dots, \bar{v}_l$ generates a one-dimensional H -invariant $k(x)$ -subspace of $\mathfrak{m}_x/\mathfrak{m}_x^2$.*
- (b) *For every $i = 1, \dots, m$ and every $g \in G$, u_i is a local equation of $g(Y_i)$ at gx .*
- (c) *For every $j = 1, \dots, l$ there are integers $e_{j1}, \dots, e_{jm} \geq 0$ such that $h_j = u_1^{e_{j1}} \cdots u_m^{e_{jm}} v_j$ is a G -invariant rational function on X .*

Proof. We begin by constructing u_1, \dots, u_m . Consider the divisor $D_i = \sum g(Y_i)$, where each summand of the form $g(Y_i)$ (for some $g \in G$) appears in this sum exactly once. Since X is quasiprojective, the divisor D_i can be “moved off” the finite set Gx ; see [Sh, Theorem III.1.1]. In other words, for every $i = 1, \dots, m$ there is a rational function u_i on X such that the support of the divisor $D_i - (u_i)$ does not intersect Gx . It is now easy to see that u_1, \dots, u_m satisfy (a) and (b).

Next we turn to the construction of v_1, \dots, v_l . Each \bar{u}_i generates a one-dimensional $k(x)$ -subspace $\langle \bar{u}_i \rangle = (T_x(X)/T_x(Y_i))^* \subset \mathfrak{m}_x/\mathfrak{m}_x^2$. We have seen in Remark 4.4 that $\langle \bar{u}_i \rangle$ is H -invariant; H' acts on it by the character ξ_i . In view of Lemma 2.3 and Proposition 2.4, we can write

$$\mathfrak{m}_x/\mathfrak{m}_x^2 = \langle \bar{u}_1 \rangle \oplus \cdots \oplus \langle \bar{u}_m \rangle \oplus V_1 \oplus \cdots \oplus V_l,$$

where each V_i is a simple $(k(x)*H)$ -module and $\dim_{k(x)}(V_i) = 1$. Choose $\bar{v}_1, \dots, \bar{v}_l \in \mathfrak{m}_x/\mathfrak{m}_x^2$ so that \bar{v}_i generates V_i as a $k(x)$ -vector space. Denote the character of H' associated to \bar{v}_j by η_j . By Remark 4.4, ξ_1, \dots, ξ_m generate the dual group $(H')^*$; consequently, each η_j can be written in the form

$$\eta_j = \xi_1^{-e_{j1}} \cdots \xi_m^{-e_{jm}}$$

for some integers $e_{j1}, \dots, e_{jm} \geq 0$. Note that $\bar{h}_j = \bar{u}_1^{e_{j1}} \cdots \bar{u}_m^{e_{jm}} \bar{v}_j$ is an H' -invariant element of $\mathfrak{m}_x^{d_j}/\mathfrak{m}_x^{d_j+1}$, where $d_j = e_{j1} + \cdots + e_{jm} + 1$. Clearly, \bar{h}_j generates an H -invariant one-dimensional $k(x)$ -subspace $\langle \bar{h}_j \rangle \subset \mathfrak{m}_x^{d_j}/\mathfrak{m}_x^{d_j+1}$ on which H' acts trivially. By the uniqueness statement in Proposition 2.4(a), $\langle \bar{h}_j \rangle \cong k(x)$ as $(k(x)*H)$ -modules, and by Remark 2.2, after replacing \bar{v}_j by $\lambda \bar{v}_j$ for some $\lambda \in k(x)$, we may assume

$$(4.3) \quad \bar{h}_j = \bar{u}_1^{e_{j1}} \cdots \bar{u}_m^{e_{jm}} \bar{v}_j \text{ is an } H\text{-invariant element of } \mathfrak{m}_x^{d_j}/\mathfrak{m}_x^{d_j+1}.$$

We claim that we can choose $v_1, \dots, v_l \in \mathfrak{m}_x$ so that $\bar{v}_j = v_j \bmod \mathfrak{m}_x^2$ and each $h_j = u_1^{e_{j1}} \cdots u_m^{e_{jm}} v_j$ is G -invariant. If we can do this, then $u_1, \dots, u_m, v_1, \dots, v_l$ will clearly satisfy the requirements of the proposition.

To prove the claim, let $R = \bigcap_{y \in Gx} \mathcal{O}_{y,X}$ be the ring of rational functions on X that are well defined on Gx , and let $\mathcal{I}_{Gx} = \bigcap_{y \in Gx} \mathfrak{m}_y$ be a G -invariant ideal in R consisting of all elements that vanish on Gx . By our choice of u_1, \dots, u_m ,

$$(4.4) \quad g^*(u_i)/u_i \text{ is defined and invertible at every point of } Gx$$

for every $g \in G$, and every $i = 1, \dots, m$. Consequently, $u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathcal{I}_{Gx}$ is a G -invariant ideal of R .

For each $y \in Gx$, the functions u_1, \dots, u_m vanish at y , and hence,

$$u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathcal{I}_{Gx} \subset \mathfrak{m}_y^{d_j}.$$

Consider the G -equivariant projection map

$$\psi: u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathcal{I}_{Gx} \longrightarrow \bigoplus_{y \in Gx} \mathfrak{m}_y^{d_j} / \mathfrak{m}_y^{d_j+1}.$$

If $v \in \mathcal{I}_{Gx}$, then $\psi(u_1^{e_{j1}} \cdots u_m^{e_{jm}} v)$ depends only on the image of v in $\bigoplus_{y \in Gx} \mathfrak{m}_y / \mathfrak{m}_y^2$. Moreover, since X is quasiprojective, the finite set Gx lies in an affine open subset of X and hence, by the Chinese Remainder Theorem, the projection map $\mathcal{I}_{Gx} \longrightarrow \bigoplus_{y \in Gx} \mathfrak{m}_y / \mathfrak{m}_y^2$ is surjective. Thus

$$\text{Im}(\psi) = \bigoplus_{y \in Gx} (u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathfrak{m}_y) \bmod \mathfrak{m}_y^{d_j+1} \subset \bigoplus_{y \in Gx} \mathfrak{m}_y^{d_j} / \mathfrak{m}_y^{d_j+1}.$$

We shall denote elements of $\text{Im}(\psi)$ by $a = (a_y \mid y \in Gx)$, where

$$a_y \in (u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathfrak{m}_y) \bmod \mathfrak{m}_y^{d_j+1}.$$

Recall that by (4.3), \bar{h}_j is a nonzero H -invariant element of $(u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathfrak{m}_x) \bmod \mathfrak{m}_x^{d_j+1}$. Let a_j be the element of $\text{Im}(\psi)$ such that $(a_j)_y = (g^{-1})^*(\bar{h}_j)$, where $y = gx$; in view of (4.4),

$$(a_j)_y \in (g^{-1})^*[(u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathfrak{m}_x) \bmod \mathfrak{m}_x^{d_j+1}] = (u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathfrak{m}_y) \bmod \mathfrak{m}_y^{d_j+1}.$$

Note that since \bar{h}_j is H -invariant, $(a_j)_y$ is independent of the choice of g . By our construction a_j is G -invariant and $(a_j)_x = \bar{h}_j \in \mathfrak{m}_x^{d_j} / \mathfrak{m}_x^{d_j+1}$.

The homomorphism ψ has a G -equivariant k -linear splitting and consequently, there exists a G -invariant element $h_j = u_1^{e_{j1}} \cdots u_m^{e_{jm}} v_j \in u_1^{e_{j1}} \cdots u_m^{e_{jm}} \mathcal{I}_{Gx}$ such that $\psi(h_j) = a_j$. In particular, $\bar{h}_j = \bar{u}_1^{e_{j1}} \cdots \bar{u}_m^{e_{jm}} \bar{v}_j = h_j \bmod \mathfrak{m}_x^{d_j+1}$ and hence $\bar{v}_j = v_j \bmod \mathfrak{m}_x^2$. This proves the claim and thus shows that $u_1, \dots, u_m, v_1, \dots, v_l$ have the required properties. \square

Property (*). We are now ready to revisit property (*) of Definition 3.3.

Lemma 4.6. *Suppose X is a quasiprojective G -variety in standard form with respect to a divisor Y , $x \in X$, $u_1, \dots, u_m, v_1, \dots, v_l$, and h_1, \dots, h_l , are as in Proposition 4.5, and $w_i = \prod_{g \in G} g^*(u_i)$. Let f be a rational function on X whose divisor is supported on Y . Then*

$$df/f \in \mathcal{M},$$

where \mathcal{M} is the $\mathcal{O}_{x,X}$ -module generated by dw_i/w_i and dh_j/h_j with $i = 1, \dots, m$ and $j = 1, \dots, l$.

Note that w_i and h_j are G -invariant rational functions on X for every $i = 1, \dots, m$ and $j = 1, \dots, l$.

Proof. Let $(\Omega_X^1)_x$ be the $\mathcal{O}_{x,X}$ -module of germs at x of regular differential forms on X . Since X is smooth, $(\Omega_X^1)_x$ is a free $\mathcal{O}_{x,X}$ -module generated by $du_1, \dots, du_m, dv_1, \dots, dv_l$. Let \mathcal{M}' be the $\mathcal{O}_{x,X}$ -module generated by du_i/u_i , and dv_j/v_j , where $i = 1, \dots, m$ and $j = 1, \dots, l$. Clearly, $(\Omega_X^1)_x \subset \mathfrak{m}_x \mathcal{M}'$.

We claim that $\mathcal{M} = \mathcal{M}'$. It is clear that $\mathcal{M} \subset \mathcal{M}'$; to prove the opposite inclusion, we shall show that $dw_1/w_1, \dots, dw_m/w_m, dh_1/h_1, \dots, dh_l/h_l$ generate \mathcal{M}' as an $\mathcal{O}_{x,X}$ -module.

Note that by our choice of u_1, \dots, u_m , we can write $w_i = a_i u_i^{|G|}$ for some $a_i \in \mathcal{O}_{X,x} - \mathfrak{m}_x$; see (4.4). Thus

$$\frac{dw_i}{w_i} = \frac{da_i}{a_i} + |G| \frac{du_i}{u_i} \equiv |G| \frac{du_i}{u_i} \pmod{(\Omega_X^1)_x}.$$

In particular, since $(\Omega_X^1)_x \subset \mathfrak{m}_x \mathcal{M}'$, we conclude that

$$(4.5) \quad \frac{dw_i}{w_i} \equiv |G| \frac{du_i}{u_i} \pmod{\mathfrak{m}_x \mathcal{M}'}.$$

On the other hand, since $h_j = u_1^{e_{j1}} \cdots u_m^{e_{jm}} v_j$, we have

$$(4.6) \quad \frac{dh_j}{h_j} = \frac{dv_j}{v_j} + \sum_i e_{ji} \frac{du_i}{u_i}.$$

Examining (4.5) and (4.6), we see that dw_i/w_i ($i = 1, \dots, m$), and dh_j/h_j ($j = 1, \dots, l$) generate $\mathcal{M}'/\mathfrak{m}_x \mathcal{M}'$ as a $k(x)$ -vector space. Consequently, by Nakayama's lemma these elements generate \mathcal{M}' as an $\mathcal{O}_{x,X}$ -module. Thus $\mathcal{M}' = \mathcal{M}$, as claimed.

Since the divisor of f is supported on Y , locally near x it is a union of smooth hypersurfaces of the form $\{u_i = 0\}$. This means that $f = au_1^{e_1} \cdots u_m^{e_m}$ for some $a \in \mathcal{O}_{x,X} - \mathfrak{m}_x$ and $e_1, \dots, e_m \geq 0$; hence,

$$\frac{df}{f} = \frac{da}{a} + e_1 \frac{du_1}{u_1} + \cdots + e_m \frac{du_m}{u_m} \in \mathcal{M}' = \mathcal{M}.$$

□

Proposition 4.7. *Let X be a projective variety in standard form with respect to a divisor Y and let f be a rational function on X whose divisor is supported on Y . Then the pair (X, f) has property (*); see Definition 3.3.*

Proof. By Zariski compactness, it is enough to show that for any $x \in X$, there exist $\beta_1, \dots, \beta_q \in k(X)^G$ and $\gamma_1, \dots, \gamma_q \in \mathcal{O}_{x,X}$ such that

$$\frac{df}{f} = \gamma_1 \frac{d\beta_1}{\beta_1} + \cdots + \gamma_q \frac{d\beta_q}{\beta_q}.$$

The last assertion is immediate from Lemma 4.6. □

Proof of Theorem 1.1. As we showed in Section 3, it is enough to prove Proposition 3.6.

Assume X is a projective G -variety and $f \in k(X)$. Let X_0 be the subvariety of all points in X with nontrivial stabilizers. Let Y be the union of X_0 and (the supports of) the divisors of $g^*(f)$ for every $g \in G$; it is a G -invariant Zariski closed subvariety of X . By Theorem 4.2 there exists a birational morphism $\pi: X' \rightarrow X$ and a divisor $Y' \subset Y$, such that X' is in standard form with respect to Y' and $\pi^{-1}(Y) \subset Y'$. Note that the divisor of $\pi^*(f)$ is contained in $\pi^{-1}(Y)$ and hence in Y' . Proposition 3.6 (and thus Theorem 1.1) now follows from Proposition 4.7, which asserts that the pair $(X', \pi^*(f))$ has property (*). □

REFERENCES

- [B] N. Bourbaki, *Algèbre*, Hermann, Paris. MR **20**:4576
- [Mo] S. Montgomery, *Fixed points of finite automorphism groups of associative rings*, Lect. Notes in Math. **818**, Springer-Verlag, 1980. MR **81j**:16041
- [MFK] D. Mumford, J. Fogarty and F. Kirwan, *Geometric invariant theory*. Third enlarged edition, Springer, 1994. MR **95m**:14012
- [P] A. Parusiński, *Lipschitz properties of semianalytic sets*, Ann. Inst. Fourier, Grenoble **38** (1988), 189–213. MR **90e**:32016
- [RY] Z. Reichstein and B. Youssin, *Essential dimensions of algebraic groups and a resolution theorem for G -varieties*, with an appendix by J. Kollár and E. Szabó, preprint. Available at <http://ucs.orst.edu/~reichstz/pub.html>.

- [Sh] I. R. Shafarevich, Basic algebraic geometry, Springer-Verlag, Heidelberg, 1974. MR
51:3163

DEPARTMENT OF MATHEMATICS, OREGON STATE UNIVERSITY, CORVALLIS, OR 97331
E-mail address: `zinovy@math.orst.edu`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF THE NEGEV, BE'ER
SHEVA', ISRAEL
Current address: Hashofar 26/3, Ma'ale Adumim, Israel
E-mail address: `youssin@math.bgu.ac.il`