

AN ESTIMATE FOR CHARACTER SUMS

NICHOLAS M. KATZ

In this note, we give estimates for a class of character sums that occur as eigenvalues of adjacency matrices of certain graphs constructed by F. R. K. Chung. Her situation is as follows. We are given a finite field F , an integer $n \geq 1$, an extension field E of F of degree n , and an element x in E that generates E over F , i.e., an element x such that E is $F(x)$.

Theorem 1. *Let χ be any nontrivial complex-valued multiplicative character of E^\times (extended by zero to all of E), and x in E any element that generates E over F . Then*

$$\left\| \sum_{t \in F} \chi(t - x) \right\| \leq (n - 1) \sqrt{\#(F)}.$$

It turns out to be easier to consider the following more general situation. F is a finite field, $n \geq 1$ is an integer, and B is a finite etale F -algebra of dimension n over F (i.e., over a finite extension K of F , there exists an isomorphism of K -algebras $B \otimes_F K \simeq K \times K \times \cdots \times K$). We assume given an element x in B that is regular in the sense that its characteristic polynomial $\det_F(T - x | B)$ in the regular representation of B on itself has n distinct eigenvalues. (In terms of the above isomorphism $B \otimes_F K \simeq K \times K \times \cdots \times K$, x is regular if and only if $x \otimes 1 \simeq (x_1, \dots, x_n)$ with all distinct components x_i . Or equivalently, x is regular if and only if B is equal to the F -subalgebra $F[x]$ generated by x . In the special case when B is a field F , the element x is regular if and only if $F(x) = E$.)

Theorem 2. *Let χ be any nontrivial complex-valued multiplicative character of B^\times (extended by zero to all of B), and x in B any regular element. Then*

$$\left\| \sum_{t \in F} \chi(t - x) \right\| \leq (n - 1) \sqrt{\#(F)}.$$

Proof. The basic idea is that the theorem is an immediate consequence of Weil's estimates for one-variable character sums in the case when the F -algebra B is completely split, and that one can reduce to this case by thinking geometrically about suitable Lang torsors.

Received by the editors February 10, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 11L40, 11T21.

We begin by explaining how to view the problem geometrically. Given any finite-dimensional commutative F -algebra A , we denote by \mathbb{A} the smooth affine scheme over F given by “ A as algebraic group over F ”; concretely, for any F -algebra R , the group $\mathbb{A}(R)$ of R -valued points of \mathbb{A} is $A \otimes_F R$. We denote by \mathbb{A}^\times the open subscheme of \mathbb{A} given by “ A^\times as algebraic group over F ”; concretely, for any F -algebra R , the group $\mathbb{A}^\times(R)$ of R -valued points of \mathbb{A} is $(A \otimes_F R)^\times$. These concepts will be applied to the cases $A = B$ and $A = F$. It will be important in what follows to think of \mathbb{A}^\times as a smooth commutative group scheme over F , but to think of \mathbb{A} only as an ambient scheme (not as a group scheme) containing \mathbb{A}^\times as an open subscheme.

Because \mathbb{B}^\times is a smooth, geometrically connected commutative group scheme over the finite field F , the Lang isogeny $1 - \text{Frob}_F: \mathbb{B}^\times \rightarrow \mathbb{B}^\times$ makes \mathbb{B}^\times into a \mathbb{B}^\times -torsor over itself, the “Lang torsor” \mathbb{L} . Let us now fix a prime number $l \neq \text{char}(F)$, an algebraic closure \overline{Q}_l of Q_l , and an isomorphism of fields $C \simeq \overline{Q}_l$. This isomorphism allows us to view χ as a \overline{Q}_l -valued character of \mathbb{B}^\times , by which it makes sense to push out the Lang torsor \mathbb{L} to obtain a lisse rank one \overline{Q}_l -sheaf \mathbb{L}_χ on \mathbb{B}^\times which is pure of weight zero. If we denote by $j: \mathbb{B}^\times \rightarrow \mathbb{B}$ the inclusion, we may form the extension by zero $j_! \mathbb{L}_\chi$ on \mathbb{B} . Now consider the morphism of F -schemes of $f: \mathbb{F} \rightarrow \mathbb{B}$ defined by $f(t) := t - x$, and the pullback sheaf $\mathcal{F} := f^*(j_! \mathbb{L}_\chi)$ on \mathbb{F} . The sheaf \mathcal{F} is lisse of rank one and pure of weight zero on the open set $f^{-1}(\mathbb{B}^\times)$, and zero outside. The sheaf \mathcal{F} is everywhere tamely ramified, simply because on $f^{-1}(\mathbb{B}^\times)$ it is lisse of order dividing that of χ , hence of order prime to the characteristic of F .

In terms of this data, the character sum in question is given by

$$\sum_{t \in F} \chi(t - x) = \sum_{t \in f^{-1}(\mathbb{B}^\times)(F)} \text{Trace}(\text{Frob}_{t,F} | \mathcal{F}),$$

and by the Lefschetz Trace Formula this last sum is equal to

$$\sum_i (-1)^i \text{Trace}(\text{Frob}_F | H_{\text{comp}}^i(f^{-1}(\mathbb{B}^\times) \otimes_F \overline{F}, \mathcal{F})).$$

By Weil (but expressed in the language of Deligne’s paper [De]) we know that the above cohomology groups H_{comp}^i are mixed of weight $\leq i$. For dimension reasons, H_{comp}^i vanishes for $i > 2$, and H_{comp}^0 vanishes because \mathcal{F} is lisse on the incomplete curve $f^{-1}(\mathbb{B}^\times) \otimes_F \overline{F}$. It thus remains only to establish the following two facts:

- (a) $H_{\text{comp}}^2(f^{-1}(\mathbb{B}^\times) \otimes_F \overline{F}, \mathcal{F}) = 0$,
- (b) $\dim H_{\text{comp}}^1(f^{-1}(\mathbb{B}^\times) \otimes_F \overline{F}, \mathcal{F}) = n - 1$.

Both of these facts are geometric, i.e., they concern the situation over the algebraic closure of F , and hence it suffices to verify them universally in the case when the F -algebra B is completely split. (The key point here is that our hypothesis that χ is nontrivial is stable under finite extension of scalars.

Indeed, after extension of scalars from F to any finite extension field K , the pullback to $(\mathbf{B}^\times) \otimes_F K$ of \mathbb{L}_χ is $\mathbb{L}_{\tilde{\chi}}$, where $\tilde{\chi}$ is the character of $(B \otimes_F K)^\times$ obtained from χ by composition with the norm homomorphism $\text{Norm}_{K/F}$ from $(B \otimes_F K)^\times$ to B^\times . Because this norm map is surjective, the character $\tilde{\chi}$ is nontrivial provided that χ is nontrivial.)

Suppose now that B is simply the n -fold self product of F with itself. Then a nontrivial character χ of B^\times is simply an n -tuple (χ_1, \dots, χ_n) of characters of F^\times , not all of which are trivial, the regular element x is just an n -tuple (x_1, \dots, x_n) with all distinct components x_i , the open set $f^{-1}(\mathbf{B}^\times)$ is just the complement $\mathbb{F} - \{x_1, \dots, x_n\}$ of the n distinct points x_i in \mathbb{F} , the sheaf \mathcal{F} is just the tensor product of the sheaves $[t \mapsto t - x_i]^* \mathbb{L}_{\chi_i} |_{\mathbb{F} - \{x_1, \dots, x_n\}}$, and the sum in question is

$$\sum_{t \in \mathbb{F} - \{x_1, \dots, x_n\}} \chi_1(t - x_1) \chi_2(t - x_2) \cdots \chi_n(t - x_n).$$

By assumption, at least one of the χ_i is nontrivial. For such an index i , the sheaf $[t \mapsto t - x_i]^* \mathbb{L}_{\chi_i}$ is tamely but nontrivially ramified at x_i , while all the other factors $[t \mapsto t - x_j]^* \mathbb{L}_{\chi_j}$ with $j \neq i$ are lisse at x_i (by the hypothesis that all the x_j are distinct). Therefore, the sheaf \mathcal{F} is nontrivially ramified at the point x_i . Because \mathcal{F} is lisse of rank one on $\mathbb{F} - \{x_1, \dots, x_n\}$, its coinvariants under the inertia group I_{x_i} must vanish, and a fortiori its covariants under the entire π_1^{geom} of $\mathbb{F} - \{x_1, \dots, x_n\}$ must also vanish, i.e., its H_{comp}^2 vanishes. Once we have the vanishing of all the H_{comp}^i save for $i = 1$, the asserted dimension formula $\dim H_{\text{comp}}^1 = n - 1$ is then equivalent to the Euler characteristic formula

$$\sum_i (-1)^i \dim H_{\text{comp}}^i ((\mathbb{F} - \{x_1, \dots, x_n\}) \otimes_F \overline{\mathbb{F}}, \mathcal{F}) = 1 - n,$$

which holds because \mathcal{F} is lisse of rank one and everywhere tame on the open curve $(\mathbb{F} - \{x_1, \dots, x_n\}) \otimes_F \overline{\mathbb{F}}$, whose Euler characteristic is $1 - n$. Q.E.D.

Remarks and Questions. (1) If we drop the hypothesis that the element x be regular, then Theorem 2 remains valid for characters χ of B^\times whose restriction to F^\times is nontrivial. The proof proceeds along the same lines as above, reducing to the completely split case in which χ is simply an n -tuple (χ_1, \dots, χ_n) of characters of F^\times , with the property that their product $\prod_i \chi_i$ is nontrivial on F^\times . Now one gets the vanishing of H_{comp}^2 by observing that the sheaf \mathcal{F} is nontrivially ramified at ∞ (as an I_∞ -representation, \mathcal{F} is isomorphic to $\mathbb{L}_{\prod_i \chi_i}$), and the constant “ $n - 1$ ” actually improves to “(the number of distinct x_i) $- 1$.” Indeed, in the case of the choice $x := 0$, the character sum in question is exactly $\sum_{t \in F^\times} \chi(t)$. (Alternately, one could apply Theorem 2 directly to the (automatically finite etale) subalgebra $B_0 := F[x]$ of B generated by x over F , to the regular element x of B_0 , and to the nontrivial (because nontrivial on F^\times) character $\chi |_{(B_0)^\times}$.)

(2) What happens if we also drop the hypothesis that B be étale? Suppose that we are given an arbitrary n -dimensional commutative F -algebra A , a multiplicative character χ of A^\times (extended by zero to all of A) whose restriction to F^\times is nontrivial, and an element x in A . It seems plausible that the estimate

$$\left\| \sum_{t \in F} \chi(t - x) \right\| \leq (n - 1) \sqrt{\#(F)}$$

should still hold. For example, in the case when A is the algebra of dual numbers $F[x]/(x^2)$, the character sums in question are none other than the usual Gauss sums attached to the field F .

REFERENCES

- [Ch] F. R. K. Chung, *Diameters and eigenvalues*, J. Amer. Math. Soc. **2** (1989), 187–196.
 [De] P. Deligne, *La conjecture de Weil II*, Inst. Hautes Études Sci. Publ. Math., no. 52, 1981, pp. 313–428.
 [We] A. Weil, *On some exponential sums*, Proc. Nat. Acad. Sci. U.S.A. **34** (1948), 204–207.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08544