

ASYMPTOTIC PROPERTIES OF BANACH SPACES UNDER RENORMINGS

E. ODELL AND TH. SCHLUMPRECHT

§1. INTRODUCTION

A classical problem in functional analysis has been to give a geometric characterization of reflexivity for a Banach space. The first result of this type was D.P. Milman's [Mil] and B.J. Pettis' [P] theorem that a uniformly convex space is reflexive. While perhaps considered elementary today it illustrated how a geometric property can be responsible for a topological property. Of course a Banach space can be reflexive without being uniformly convex, even under renormings, as shown by M.M. Day [D2]. The problem considered for years by functional analysts was whether there exists a weaker property of a geometric nature which is equivalent to reflexivity. In this paper we give an affirmative solution by demonstrating that such a property exists. The property was suggested in 1971 by V.D. Milman [Mi] (see also [DGZ], problem IV, p.177). We prove that a separable Banach space X is reflexive (if and) only if there exists an equivalent norm $\|\cdot\|$ on X so that

$$(*) \quad \begin{aligned} &\text{whenever a sequence } (x_n) \subseteq X \text{ satisfies} \\ &\lim_n \lim_m \|x_n + x_m\| = 2 \lim_n \|x_n\| \\ &\text{then } (x_n) \text{ must converge in norm.} \end{aligned}$$

The "if" part of the characterization follows easily from James' famous characterization of reflexivity in terms of the sup of linear functionals [J1]. Indeed given $x^* \in X^*$ with $\|x^*\| = 1$ choose $(x_n) \subseteq X$ with $x^*(x_n) \rightarrow 1$ and $\|x_n\| = 1$ for all n . Then $\lim_m \lim_n \|x_n + x_m\| = 2$ and so $x_n \rightarrow x$ with $\|x\| = 1$. Thus $x^*(x) = 1$ so x^* attains its norm. Hence by [J1] X is reflexive.

The investigation of spaces having property $(*)$ (also called property $(2R)$ in [D1]) goes back to the 1950's. In [FG], for example, the relation of $(*)$ to other smoothness and rotundity properties was studied. For a more complete survey of these notions we refer the reader to [DGZ].

More recently (over the past 30 years) functional analysts have considered the question as to what sort of nice infinite dimensional subspaces one can find in an arbitrary infinite dimensional Banach space X . One can assume X has a basis and ask: What kinds of block bases does it have? Must one be unconditional? Is some block subspace either reflexive or isomorphic to c_0 or ℓ_1 ? These problems are related. James [J3] showed that if (x_i) is an unconditional basis for X , then

Received by the editors May 12, 1997 and, in revised form, September 15, 1997.

1991 *Mathematics Subject Classification*. Primary 46B03, 46B45.

Key words and phrases. Spreading model, Ramsey theory, ℓ_1 , c_0 , reflexive Banach space.

Research of both authors was supported by NSF and TARP.

either X is reflexive or some block basis is equivalent to the unit vector basis for c_0 or ℓ_1 . W.T. Gowers [G1] proved the following remarkable dichotomy theorem: X contains a subspace Y which either has an unconditional basis or is H.I. (hereditarily indecomposable; i.e., if $Z \subseteq Y$ and $Z = V \oplus W$, then V or W must be finite dimensional). Gowers and Maurey [GM] proved that both alternatives are possible. Then Gowers [G2] proved that a space need not contain c_0 , ℓ_1 or a reflexive subspace.

The search for an answer to this last problem led to much research into both characterizations of reflexivity and to the characterization as to when X contains isomorphs of c_0 or ℓ_1 (e.g., [J1], [J2], [J3], [R1], [R2], [R3], [BP], [M]). The proof of our characterization of reflexivity led to additional characterizations as to when X contains c_0 or ℓ_1 in terms of the asymptotic behavior of sequences in X .

There are two main notions of asymptotic properties in Banach spaces. The first is that of a spreading model. If (x_n) is bounded in X , then by using Ramsey theory (cf. [B], [BS], [O], [BL]) one can extract a subsequence (y_n) so that for all k and $(a_i)_1^k \subseteq \mathbb{R}$, we have the existence of the iterated limit

$$\lim_{n_1 \rightarrow \infty} \cdots \lim_{n_k \rightarrow \infty} \left\| \sum_{i=1}^k a_i y_{n_i} \right\| \equiv f(a_1, \dots, a_k).$$

If (y_n) does not converge in the norm topology, then $f(\cdot)$ is a norm on c_{00} , the linear space of all finitely supported real valued sequences. Let (e_i) be the unit vector basis of c_{00} . If (y_n) does not converge weakly to a nonzero element of X , then (e_i) is a basis for $E = [(e_i)]$, the completion of c_{00} under $f(\cdot)$. In this case we call (e_i) or E the *spreading model* of (y_n) . If (x_i) is weakly null, then the spreading model (e_i) is unconditional. In any event the spreading model is subsymmetric ($\|\sum a_i e_i\| = \|\sum a_i e_{n_i}\|$ if $(a_i) \subseteq \mathbb{R}$ and $n_1 < n_2 < \cdots$) and $(e_1 - e_2, e_3 - e_4, \dots)$ is unconditional.

The second notion of asymptotic structure is due to Maurey, Milman and Tomczak-Jaegermann (see [MT], [MMT]). Let X have a basis (x_i) . For $x, y \in X$ we write $x < y$ if $\max \text{supp } x < \min \text{supp } y$ where if $x = \sum a_i x_i$, then $\text{supp } x = \{i : a_i \neq 0\}$. $\langle x_i \rangle_{i \in I}$ denotes the linear span of $\{x_i : i \in I\}$ and $S_{\langle x_i \rangle_{i \in I}}$ denotes the unit sphere of this span. Let $n \in \mathbb{N}$ and let $(w_i)_1^n$ be a normalized basis for some n dimensional space. We say $(w_i)_1^n \in \{X\}_n$ if

$$\begin{aligned} &\forall k_1 \in \mathbb{N} \exists y_1 \in S_{\langle x_i \rangle_{k_1}}^\infty \forall k_2 \in \mathbb{N} \\ &\exists y_2 \in S_{\langle x_i \rangle_{k_2}}^\infty \cdots \forall k_n \in \mathbb{N} \exists y_n \in S_{\langle x_i \rangle_{k_n}}^\infty \end{aligned}$$

so that $(y_i)_1^n$ is $1 + \varepsilon$ -equivalent to $(w_i)_1^n$. This means that there exist A, B with $AB \leq 1 + \varepsilon$ so that for all $(a_i)_1^n \subseteq \mathbb{R}$

$$A^{-1} \left\| \sum_1^n a_i y_i \right\| \leq \left\| \sum_1^n a_i w_i \right\| \leq B \left\| \sum_1^n a_i y_i \right\|.$$

Note that if (e_i) is a spreading model of a normalized block basis of (x_i) , then $(e_i)_1^n \in \{X\}_n$ for all n .

Both notions give a more regular structure in general than that possessed by the original space X . They are a joining of the finite and infinite dimensional structures of the space. Generally only finite dimensional information can be gleaned about X from knowledge of its asymptotic structure.

For example, note that the Schreier space S ([CS], p.1) has a basis having a spreading model isometric to ℓ_1 and yet S is c_0 saturated (all infinite dimensional subspaces of S contain c_0). Tsirelson's space T (the dual of Tsirelson's original space [T] as described in [FJ]; see also [CS]) has a basis with the property that all spreading models are isomorphic to ℓ_1 and in addition every infinite dimensional subspace contains a sequence whose spreading model is isometric to ℓ_1 [OS]. Yet T is reflexive. We do have the following result which requires a very strong assumption on the class of spreading models.

Theorem ([OS]). *If (x_i) is a basis for X and if every spreading model (e_i) of any normalized block basis of (x_i) is 1-equivalent to the unit vector basis of ℓ_1 (respectively, c_0), then X contains an isomorph of ℓ_1 (respectively, c_0).*

In this paper we deduce information about the infinite dimensional structure of X from knowledge about its asymptotic structure under equivalent norms.

We shall show that a separable space $(X, \|\cdot\|)$ can be given a special renorming $\|\cdot\|$ so that certain information about a given spreading model E yields information about the infinite dimensional structure of X . For example if $\|e_i \pm e_2\| = 2$ (respectively, $\|e_1 + e_2\| = 1$) for some spreading model (e_i) of a normalized (and respectively, weakly null) sequence in X , then X contains ℓ_1 (respectively, c_0). Furthermore we show that a subspace Y of X is reflexive iff Y satisfies (*).

Our main result is the following theorem.

Main Theorem. *Every separable Banach space X admits an equivalent strictly convex norm $\|\cdot\|$ with the following properties.*

a) *If $(x_n) \subseteq X$ is relatively weakly compact and if*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 2 \lim_{n \rightarrow \infty} \|x_n\|,$$

then (x_n) is norm convergent.

b) *If $(x_n) \subseteq X$ satisfies*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m \pm x_n\| = 2 \lim_{n \rightarrow \infty} \|x_n\| > 0,$$

then some subsequence of (x_n) is equivalent to the unit vector basis of ℓ_1 .

c) *If $(x_n) \subseteq X$ is weakly null and satisfies*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = \lim_{n \rightarrow \infty} \|x_n\| > 0,$$

then some subsequence of (x_n) is equivalent to the unit vector basis of c_0 .

This theorem is proved in §2.

As a corollary we deduce Milman's suggested characterization of reflexivity. In addition we obtain that X contains ℓ_1 if under all equivalent norms, Y admits a normalized basic sequence having a spreading model (e_i) satisfying $\|e_1 \pm e_2\| = 2$. In particular if under all equivalent norms X admits a spreading model (e_i) which is 1-equivalent to the unit vector basis of ℓ_1 , then X contains an isomorph of ℓ_1 . If under all equivalent norms X admits a weakly null sequence having spreading model (e_i) with $\|e_1 + e_2\| = 1$ (e.g., if (e_i) is 1-equivalent to the unit vector basis of c_0), then X contains an isomorph of c_0 . From James' proof that ℓ_1 and c_0 are not distortable [J2] one obtains that both implications can be reversed.

In §3 we present some corollaries discussed briefly in this introduction. Our notation is standard as may be found in [LT].

§2. PROOF OF THE MAIN THEOREM

We first recall the following results of Maurey and Rosenthal.

Theorem ([M], [R1]). *Let X be a separable Banach space.*

- a) X is not reflexive if and only if there exists a normalized basic sequence $(x_n) \subseteq X$ satisfying for all $x \in X$ and $\alpha, \beta \geq 0$ with $\alpha + \beta = 1$,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x + \alpha x_n + \beta x_n\| = \lim_{m \rightarrow \infty} \|x + x_m\| .$$

- b) X contains an isomorph of ℓ_1 if and only if there exists a normalized basic sequence $(x_n) \subseteq X$ such that for all $x \in X$ and $\alpha, \beta \in \mathbb{R}$ with $|\alpha| + |\beta| = 1$,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x + \alpha x_m + \beta x_n\| = \lim_{m \rightarrow \infty} \|x + x_m\| .$$

- c) X contains an isomorph of c_0 iff there exists a normalized basic sequence $(x_n) \subseteq X$ such that for all $x \in X$ and $\alpha, \beta \in \mathbb{R}$ with $|\alpha| \vee |\beta| = 1$,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x + \alpha x_m + \beta x_n\| = \lim_{m \rightarrow \infty} \|x + x_m\| .$$

The intuition behind these results and the techniques employed to prove them had their origin in [KM] where *types* were introduced (and further developed in [HM]). A type $\tau_{(x_n)}$ on X is a function on X defined by a bounded sequence $(x_n) \subseteq X$, $\tau_{(x_n)}(x) = \lim_{n \rightarrow \infty} \|x + x_n\|$. Types give information on the asymptotic behavior of a sequence acting on the whole space. This contrasts with the notion of a spreading model which involves only the asymptotic behavior of the sequence (x_n) itself. In this paper we characterize the three properties considered in the theorem above in terms solely of the asymptotic behavior of the sequences themselves. The price that must necessarily be paid is that we have to consider this behavior under all equivalent norms on X .

Let $(X, \|\cdot\|)$ be a Banach space over \mathbb{R} . If $x \in X$ we define the *symmetrized type* norm $\|\cdot\|_x : X \rightarrow [0, \infty)$ by

$$\|y\|_x = \left\| \|x\|y\| + y \right\| + \left\| \|x\|y\| - y \right\| \text{ for } y \in X .$$

Lemma 2.1. *For all $x \in X$, $\|\cdot\|_x$ is an equivalent norm on X satisfying $2\|y\| \leq \|y\|_x \leq 2(1 + \|x\|)\|y\|$ for all $y \in X$.*

Proof. The only property not evident is that $\|\cdot\|_x$ satisfies the triangle inequality. It is easy to check that for fixed $u, v \in X$ the function $r \mapsto \|ru + v\| + \|ru - v\|$ is symmetric and convex on \mathbb{R} and thus increasing on $[0, \infty)$. Hence for $y_1, y_2 \in X$

$$\begin{aligned} \|y_1 + y_2\|_x &= \left\| \|x\|y_1 + y_2\| + y_1 + y_2 \right\| + \left\| \|x\|y_1 + y_2\| - y_1 - y_2 \right\| \\ &\leq \left\| x(\|y_1\| + \|y_2\|) + y_1 + y_2 \right\| + \left\| x(\|y_1\| + \|y_2\|) - y_1 - y_2 \right\| \\ &\leq \left\| x\|y_1\| + y_1 \right\| + \left\| x\|y_2\| + y_2 \right\| + \left\| x\|y_1\| - y_1 \right\| + \left\| x\|y_2\| - y_2 \right\| \\ &= \|y_1\|_x + \|y_2\|_x . \quad \square \end{aligned}$$

Let X be a separable Banach space. It is well known that X admits an equivalent *strictly convex* norm $\|\cdot\|$, i.e., $\|x\| = \|y\| = 1$ and $\|x+y\| = 2$ implies that $x = y$. Fix a countable dense subset C in X which is closed under rational linear combinations. Choose $(p_c)_{c \in C} \subseteq (0, \infty)$ so that $\sum_{c \in C} p_c(1 + \|c\|) < \infty$ for some (and thus for any) equivalent norm on X . If $\|\cdot\|$ is an equivalent norm on X , define $\|\cdot\| : X \rightarrow [0, \infty)$

by $\|x\| = \sum_{c \in C} p_c \|x\|_c$. By Lemma 2.1, $\|\cdot\|$ is an equivalent norm on X . Since $0 \in C$ and since the sum of a strictly convex norm and any other equivalent norm is also strictly convex, $\|\cdot\|$ is strictly convex.

Remark. We have assumed that X is a real Banach space. Similar results in the complex case can be obtained using

$$\|y\|_x = \int_0^{2\pi} \left\| \|y\|_x + e^{-i\theta} y \right\| d\theta .$$

Our goal is to show that $\|\cdot\|$ satisfies the main theorem if $\|\cdot\|$ is strictly convex.

Lemma 2.2. *Let $\|\cdot\| = \sum_{c \in C} p_c \|\cdot\|_c$ and let $(x_n) \subseteq X$ be $\|\cdot\|$ -normalized.*

a) *If*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 2 \lim_{m \rightarrow \infty} \|x_m\| ,$$

then there exists a subsequence (x'_n) of (x_n) satisfying for all $y \in X$ and $\beta_1, \beta_2 \geq 0$ that

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + \beta_1 x'_m + \beta_2 x'_n\| = \lim_{m \rightarrow \infty} \|y + (\beta_1 + \beta_2) x'_m\| .$$

b) *If*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m \pm x_n\| = 2 \lim_{m \rightarrow \infty} \|x_m\| ,$$

then there exists a subsequence (x'_n) of (x_n) satisfying for all $y \in X$ and $\beta_1, \beta_2 \in \mathbb{R}$ with $|\beta_1| + |\beta_2| \neq 0$ that

$$\begin{aligned} & \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + \beta_1 x'_m + \beta_2 x'_n\| \\ &= \lim_{m \rightarrow \infty} \left(\left\| y \frac{|\beta_1|}{|\beta_1| + |\beta_2|} + \beta_1 x'_m \right\| + \left\| y \frac{|\beta_2|}{|\beta_1| + |\beta_2|} + \beta_2 x'_m \right\| \right) . \end{aligned}$$

Proof. a) We may choose $(x'_n) \subseteq (x_n)$ so that for all $c \in C$, $y \in C$ and $\beta_1, \beta_2 \in [0, \infty) \cap \mathbb{Q}$ the limits

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + \beta_1 x'_m + \beta_2 x'_n\|_c$$

exist. Indeed this is easily done for fixed parameters and then one applies a diagonal argument.

Our hypothesis is that

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \sum_{c \in C} p_c \|x'_m + x'_n\|_c = 2 \lim_{m \rightarrow \infty} \sum_{c \in C} p_c \|x'_m\|_c .$$

Since $\lim_{m \rightarrow \infty} \|x'_m\|_c$ exists, $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x'_m + x'_n\|_c$ exists, $\|x'_m + x'_n\|_c \leq \|x'_m\|_c + \|x'_n\|_c$ and $p_c \|\cdot\|_c \leq 2p_c(1 + \|c\|) \|\cdot\|$ for all $c \in C$, it follows that, since $\sum_{c \in C} p_c(1 + \|c\|) < \infty$, for all $c \in C$,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x'_m + x'_n\|_c = 2 \lim_{m \rightarrow \infty} \|x'_m\|_c .$$

In particular taking $c = 0$ we obtain that

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x'_m + x'_n\| = 2 .$$

It follows that since $\|x'_n\| = 1$,

$$(1) \quad \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|\beta_1 x'_m + \beta_2 x'_n\| = \beta_1 + \beta_2$$

for all $\beta_1, \beta_2 \in [0, \infty)$. Similarly we have for all $c \in C$ and $\beta_1, \beta_2 \in [0, \infty)$ that

$$(2) \quad \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|\beta_1 x'_m + \beta_2 x'_n\|_c = (\beta_1 + \beta_2) \lim_{m \rightarrow \infty} \|x'_m\|_c .$$

Let $y \in C$ and $\beta_1, \beta_2 \in [0, \infty) \cap \mathbb{Q}$ with $\beta_1 + \beta_2 > 0$. Setting $c = \frac{y}{\beta_1 + \beta_2}$ in (2) we obtain using (1) that

$$(3) \quad \begin{aligned} & \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} (\|y + \beta_1 x'_m + \beta_2 x'_n\| + \|y - \beta_1 x'_m - \beta_2 x'_n\|) \\ &= \lim_{m \rightarrow \infty} (\beta_1 + \beta_2) \left(\left\| \frac{y}{\beta_1 + \beta_2} + x'_m \right\| + \left\| \frac{y}{\beta_1 + \beta_2} - x'_m \right\| \right) \\ &= \lim_m (\|y + (\beta_1 + \beta_2)x'_m\| + \|y - (\beta_1 + \beta_2)x'_m\|) . \end{aligned}$$

The triangle inequality yields

$$\begin{aligned} & \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y \pm (\beta_1 x'_m + \beta_2 x'_n)\| \\ & \leq \lim_{m \rightarrow \infty} \left\| y \frac{\beta_1}{\beta_1 + \beta_2} \pm \beta_1 x'_m \right\| + \lim_{n \rightarrow \infty} \left\| y \frac{\beta_2}{\beta_1 + \beta_2} \pm \beta_2 x'_n \right\| \\ & = \lim_{m \rightarrow \infty} \beta_1 \left\| \frac{y}{\beta_1 + \beta_2} \pm x'_m \right\| + \beta_2 \left\| \frac{y}{\beta_1 + \beta_2} \pm x'_m \right\| \\ & = \lim_{m \rightarrow \infty} \|y \pm (\beta_1 + \beta_2)x'_m\| . \end{aligned}$$

Thus from (3) we have

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + \beta_1 x'_m + \beta_2 x'_n\| = \lim_{m \rightarrow \infty} \|y + (\beta_1 + \beta_2)x'_m\| .$$

This proves a) for $y \in C$ and $\beta_1, \beta_2 \in \mathbb{Q} \cap [0, \infty)$, and hence by a density argument we obtain a) in general.

b) We may assume a) holds for (x_n) . The only remaining case we need consider is where $\beta_1 > 0$ and $\beta_2 < 0$. Actually we shall consider “ $\beta_1 x'_m - \beta_2 x'_n$ ” when $\beta_1, \beta_2 > 0$. Arguing as above using $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m - x_n\| = 2 \lim_{m \rightarrow \infty} \|x_m\|$, we may assume that (x_n) satisfies

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|\beta_1 x_m - \beta_2 x_n\|_c = (\beta_1 + \beta_2) \lim_{m \rightarrow \infty} \|x_m\|_c$$

for all $c \in C$. Letting $y \in C$ and $c = \frac{y}{\beta_1 + \beta_2}$ we obtain

$$(4) \quad \begin{aligned} & \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} (\|y + \beta_1 x_m - \beta_2 x_n\| + \|y - \beta_1 x_m + \beta_2 x_n\|) \\ &= (\beta_1 + \beta_2) \lim_{m \rightarrow \infty} \left(\left\| \frac{y}{\beta_1 + \beta_2} + x_m \right\| + \left\| \frac{y}{\beta_1 + \beta_2} - x_m \right\| \right) \\ &= \lim_{m \rightarrow \infty} (\|y + (\beta_1 + \beta_2)x_m\| + \|y - (\beta_1 + \beta_2)x_m\|) . \end{aligned}$$

Again by the triangle inequality we have

$$(5) \quad \begin{aligned} & \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y \pm (\beta_1 x_m - \beta_2 x_n)\| \\ & \leq \lim_{m \rightarrow \infty} \left[\left\| y \frac{\beta_1}{\beta_1 + \beta_2} \pm \beta_1 x_m \right\| + \left\| y \frac{\beta_2}{\beta_1 + \beta_2} \mp \beta_2 x_m \right\| \right] . \end{aligned}$$

Since

$$(6) \quad \lim_{m \rightarrow \infty} \left[\left\| y \frac{\beta_1}{\beta_1 + \beta_2} + \beta_1 x_m \right\| + \left\| y \frac{\beta_2}{\beta_1 + \beta_2} - \beta_2 x_m \right\| \right. \\ \left. + \left\| y \frac{\beta_1}{\beta_1 + \beta_2} - \beta_1 x_m \right\| + \left\| y \frac{\beta_2}{\beta_1 + \beta_2} + \beta_2 x_m \right\| \right] \\ = \lim_{m \rightarrow \infty} (\|y + (\beta_1 + \beta_2)x_m\| + \|y - (\beta_1 + \beta_2)x_m\|)$$

it follows from (4) and (5) that we have

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + \beta_1 x_m - \beta_2 x_n\| \\ = \lim_{m \rightarrow \infty} \left(\left\| y \frac{\beta_1}{\beta_1 + \beta_2} + \beta_1 x_m \right\| + \left\| y \frac{\beta_2}{\beta_1 + \beta_2} - \beta_2 x_m \right\| \right)$$

which completes the proof of b). \square

Remark. In the language of types ([R1], [M]) a) may be restated as

$$\text{if } \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 2 \lim_{m \rightarrow \infty} \|x_m\|, \quad \text{then} \\ (x_m) \text{ generates an } \ell_1^+ \text{ type on } (X, \|\cdot\|) \text{ (equivalently a double dual type).}$$

The first part of our next lemma is not new (see e.g., [M], [R1]) but we include the proof for the sake of completeness. The second part is a slight twist of Maurey's result that a symmetric ℓ_1^+ -type yields ℓ_1 . In addition the second part of the next lemma establishes that b) of the main theorem holds for $\|\cdot\|$.

Lemma 2.3. *Let $(x_n) \subseteq X$ be $\|\cdot\|$ -normalized and let $\varepsilon_i \subseteq (0, 1)$ decrease to 0.*

a) *If*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x + \beta_1 x_m + \beta_2 x_n\| = \lim_{m \rightarrow \infty} \|x + (\beta_1 + \beta_2)x_m\|$$

for all $x \in X$ and $\beta_1, \beta_2 > 0$, then there exists a subsequence (x_{n_i}) of (x_i) satisfying for all $1 \leq i_0 \leq k$ and $(\alpha_{i_0}, \alpha_{i_0+1}, \dots, \alpha_k) \subseteq [0, \infty)$ that

$$\left\| \sum_{i=i_0}^k \alpha_i x_{n_i} \right\| \geq (1 - \varepsilon_{i_0}) \sum_{i=i_0}^k \alpha_i.$$

In particular (x_n) has no weakly null subsequence.

b) *If*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x + \beta_1 x_m + \beta_2 x_n\| \\ = \lim_{m \rightarrow \infty} \left(\left\| y \frac{|\beta_1|}{|\beta_1| + |\beta_2|} + \beta_1 x_m \right\| + \left\| y \frac{|\beta_2|}{|\beta_1| + |\beta_2|} + \beta_2 x_m \right\| \right)$$

for all $x \in X$ and $\beta_1, \beta_2 \in \mathbb{R}$ with $|\beta_1| + |\beta_2| \neq 0$, then there is a subsequence (x_{n_i}) so that for all $1 \leq i_0 \leq k$ and $(\alpha_i)_{i_0}^k \subseteq \mathbb{R}$,

$$\left\| \sum_{i=i_0}^k \alpha_i x_{n_i} \right\| \geq (1 - \varepsilon_{i_0}) \sum_{i=i_0}^k |\alpha_i|.$$

In particular (x_{n_i}) is equivalent to the unit vector basis of ℓ_1 .

Proof. a) Given $\delta_i \downarrow 0$ we can choose $(x_{n_i}) \subseteq (x_i)$ satisfying the following. For all $m < \ell$ and $y \in \text{Ball}\langle x_{n_i} \rangle_{i=1}^{m-1}$, we have

$$\|y + \beta_1 x_{n_m} + \beta_2 x_{n_\ell}\| \geq (1 - \delta_m) \|y + (\beta_1 + \beta_2) x_{n_m}\|$$

if $\beta_1, \beta_2 \in [0, 1]$. Thus if $(\beta_i)_{i_0}^k \subseteq [0, 1]$, $\sum_{i_0}^k \beta_i = 1$, then

$$\begin{aligned} \left\| \sum_{i=i_0}^k \beta_i x_{n_i} \right\| &= \left\| \sum_{i=i_0}^{k-2} \beta_i x_{n_i} + \beta_{k-1} x_{n_{k-1}} + \beta_2 x_{n_k} \right\| \\ &\geq (1 - \delta_{k-1}) \left\| \sum_{i=i_0}^{k-2} \beta_i x_{n_i} + (\beta_{k-1} + \beta_k) x_{n_{k-1}} \right\| \\ &\geq (1 - \delta_{k-1})(1 - \delta_{k-2}) \left\| \sum_{i=i_0}^{k-3} \beta_i x_{n_i} + (\beta_{k-2} + \beta_{k-1} + \beta_k) x_{n_{k-2}} \right\| \\ &\geq \cdots \geq \prod_{i=i_0}^{k-1} (1 - \delta_i) \left\| \sum_{i=i_0}^k \beta_i x_{n_{i_0}} \right\| = \prod_{i=i_0}^{k-1} (1 - \delta_i). \end{aligned}$$

a) follows if we choose the δ_i 's to satisfy $\prod_{i=i_0}^{\infty} (1 - \delta_i) \geq 1 - \varepsilon_{i_0}$ for all i_0 . The ‘‘in particular’’ assertion is immediate from Mazur’s theorem.

b) The argument here is similar but slightly more complicated than a) in as much as the condition in b) is not as nice as the one in a). Let $\delta_i \downarrow 0$ satisfy $\prod_{i=i_0}^{\infty} (1 - \delta_i) > 1 - \varepsilon_{i_0}$ for all i_0 and using the assumption choose $(x_{n_i}) \subseteq (x_i)$ to satisfy for all $m < \ell$ and $y \in \text{Ball}\langle x_{n_i} \rangle_{i=1}^{m-1}$,

$$(1) \quad \begin{aligned} &\|y + \beta_1 x_{n_m} + \beta_2 x_{n_\ell}\| \\ &> (1 - \delta_m) \left[\left\| \frac{|\beta_1|}{|\beta_1| + |\beta_2|} y + \beta_1 x_{n_m} \right\| + \left\| \frac{|\beta_2|}{|\beta_1| + |\beta_2|} y + \beta_2 x_{n_m} \right\| \right] \end{aligned}$$

if $\beta_1, \beta_2 \in [-1, 1]$ with $|\beta_1| + |\beta_2| \neq 0$.

We now show by induction on k that

$$\left\| \sum_{i=i_0}^k \beta_i x_{n_i} \right\| \geq \prod_{i=i_0}^{k-1} (1 - \delta_i)$$

if $i_0 \leq k$ and $\sum_{i=i_0}^k |\beta_i| = 1$. The claim is trivial for $k = 1$ (taking $\prod_{i=i_0}^0 (1 - \delta_i) \equiv 1$). Assume validity of the claim for k and let $\sum_{i=i_0}^{k+1} |\beta_i| = 1$. For simplicity of the exposition assume $\beta_i \neq 0$ for $i_0 \leq i \leq k+1$ (the general case follows by a density

argument). Thus letting $y = \sum_{i=i_0}^{k-1} \beta_i x_{n_i}$ in (1),

$$\begin{aligned} \left\| \sum_{i=i_0}^{k+1} \beta_i x_{n_i} \right\| &\geq (1 - \delta_k) \left[\left\| \frac{|\beta_k|}{|\beta_k| + |\beta_{k+1}|} \sum_{i=i_0}^{k-1} \beta_i x_{n_i} + \beta_k x_{n_k} \right\| \right. \\ &\quad \left. + \left\| \frac{|\beta_{k+1}|}{|\beta_k| + |\beta_{k+1}|} \sum_{i=i_0}^{k-1} \beta_i x_{n_i} + \beta_{k+1} x_{n_k} \right\| \right] \\ &\geq (1 - \delta_k) \prod_{i=i_0}^{k-1} (1 - \delta_i) \left[\frac{|\beta_k|}{|\beta_k| + |\beta_{k+1}|} \sum_{i=i_0}^{k-1} |\beta_i| + |\beta_k| \right. \\ &\quad \left. + \frac{|\beta_{k+1}|}{|\beta_k| + |\beta_{k+1}|} \sum_{i=i_0}^{k-1} |\beta_i| + |\beta_{k+1}| \right] \\ &= \prod_{i=i_0}^k (1 - \delta_i). \quad \square \end{aligned}$$

Lemma 2.4. Let $\|\cdot\| = \sum_{c \in C} p_c \|\cdot\|_c$, where $\|\cdot\|$ is an equivalent strictly convex norm on X . Let $(x_n) \subseteq X$ be a relatively weakly compact sequence. If

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 2 \lim_{m \rightarrow \infty} \|x_m\|,$$

then (x_n) is norm convergent.

Proof. Since $\|\cdot\|$ is a strictly convex norm we need only show that (x_n) has a convergent subsequence. Indeed if then (x_n) were not convergent it would have two subsequences converging to $x \neq y$ respectively. But our hypothesis yields $\|x + y\| = 2 \lim \|x_m\| = \|x\| + \|y\|$ which is impossible.

By passing to a subsequence of (x_n) we may assume that $x_n = x + y_n$ where (y_n) is weakly null and $\lim_{n \rightarrow \infty} \|y_n\|$ exists. If (y_n) were not norm null, we may also assume $\|y_n\| = 1$ for all n . From Lemma 2.2, passing to a further subsequence, we may assume that for all $y \in X$,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + x_m + x_n\| = \lim_{m \rightarrow \infty} \|y + 2x_m\|.$$

For $z \in X$, letting $y = z - 2x$ we obtain

$$(1) \quad \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|z + y_m + y_n\| = \lim_{m \rightarrow \infty} \|z + 2y_m\|.$$

Since in particular $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y_m + y_n\| = 2$ it follows from (1) and the definition of $\|\cdot\|$ that

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y_m + y_n\| = 2 \lim_{m \rightarrow \infty} \|y_m\|.$$

By Lemma 2.2 a) and Lemma 2.3 a) we conclude that (y_n) is not weakly null which is a contradiction. \square

Summarizing our progress thus far we have shown that b) of the main theorem is satisfied for $\|\cdot\| = \sum_{c \in C} p_c \|\cdot\|_c$ and in addition a) holds if $\|\cdot\|$ is furthermore a strictly convex norm on X .

Lemma 2.5. Let $\|\cdot\| = \sum_{c \in C} \|\cdot\|_c$. If $(x_n) \subseteq X$ is weakly null and satisfies

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = \lim_{m \rightarrow \infty} \|x_m\| > 0,$$

then (x_n) admits a subsequence which is equivalent to the unit vector basis of c_0 .

Proof. Let (x_n) satisfy the hypothesis of the lemma for $\|\cdot\| = \sum_{c \in C} p_c \|\cdot\|_c$. We may assume (x_n) is basic, $\|x_n\| = 1$ for all n , and that for all $y \in X$ and $\beta_1, \beta_2 \in \mathbb{R}$ the following limits exist:

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + \beta_1 x_m + \beta_2 x_n\|.$$

Since (x_n) is weakly null for all $y \in X$,

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\|_y \geq \lim_{m \rightarrow \infty} \|x_m\|_y.$$

As in the proof of Lemma 2.2 since

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \sum_{c \in C} p_c \|x_m + x_n\|_c = \lim_{m \rightarrow \infty} \sum_{c \in C} p_c \|x_m\|_c$$

we obtain for all $y \in C$ and hence in X that

$$(1) \quad \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\|_y = \lim_{m \rightarrow \infty} \|x_m\|_y.$$

In particular, $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 1$. Thus by (1) for all $y \in X$,

$$(2) \quad \begin{aligned} & \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} (\|y + x_m + x_n\| + \|-y + x_m + x_n\|) \\ &= \lim_{m \rightarrow \infty} (\|y + x_m\| + \|-y + x_m\|). \end{aligned}$$

Since

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|\pm y + x_m + x_n\| \geq \lim_{m \rightarrow \infty} \|\pm y + x_m\|$$

we have from (2) that for all $y \in X$,

$$(3) \quad \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|y + x_m + x_n\| = \lim_{m \rightarrow \infty} \|y + x_m\|.$$

Choose $\varepsilon_i \downarrow 0$ with $\prod_1^\infty (1 + \varepsilon_i) < 2$ and choose, using (3), a subsequence (x_{n_i}) of (x_n) so that for any integer $k \geq 0$, $k < i < j$, and $F \subseteq \{1, \dots, k\}$ then

$$\left\| \sum_{s \in F} x_{n_s} + x_{n_i} + x_{n_j} \right\| \leq (1 + \varepsilon_i) \left\| \sum_{s \in F} x_{n_s} + x_{n_i} \right\|.$$

It follows by iterating this inequality that for all finite $F \subseteq \mathbb{N}$, $\|\sum_{s \in F} x_{n_s}\| \leq \prod_1^\infty (1 + \varepsilon_i) < 2$. This implies that (x_{n_i}) is equivalent to the unit vector basis of c_0 . \square

Remark. The proof yields that for any $\varepsilon > 0$ by judiciously choosing the p_c 's and the original strictly convex norm one can choose the norm $\|\cdot\|$ satisfying the conclusion of the main theorem to satisfy for all $x \in X$,

$$\|x\| \leq \|x\| \leq (1 + \varepsilon)\|x\|.$$

We give one final corollary of Lemma 2.5. Recall that the summing basis (s_n) for c_0 is defined for all n by $s_n = \sum_{i=1}^n e_i$.

Corollary 2.6. *Let $\|\cdot\| = \sum_{c \in C} p_c \|\cdot\|_c$ and let $(x_n) \subseteq X$ satisfy*

$$\begin{aligned} & \lim_{n_1 \rightarrow \infty} \lim_{n_2 \rightarrow \infty} \lim_{n_3 \rightarrow \infty} \lim_{n_4 \rightarrow \infty} \|x_{n_1} - x_{n_2} + x_{n_3} - x_{n_4}\| \\ &= \lim_{n_1 \rightarrow \infty} \lim_{n_2 \rightarrow \infty} \|x_{n_1} - x_{n_2}\| > 0. \end{aligned}$$

- a) *If (x_n) is weak Cauchy but not weakly convergent, then some subsequence of (x_n) is equivalent to the summing basis.*

b) If (x_n) is weakly null, then some subsequence of (x_n) is equivalent to the unit vector basis of c_0 .

Proof. Lemma 2.5 yields the following. There exists $C < \infty$ so that for all subsequences of (x_n) there exists a further subsequence (y_n) so that for all finite $F \subseteq \mathbb{N}$,

$$\left\| \sum_{n \in F} (y_{2n} - y_{2n-1}) \right\| \leq C.$$

Let

$$\mathcal{A} = \left\{ (n_i) \in [\mathbb{N}] : \text{for all finite } F \subseteq \mathbb{N}, \left\| \sum_{i \in F} (x_{n_{2i}} - x_{n_{2i-1}}) \right\| \leq C \right\}.$$

Here $[\mathbb{N}]$ denotes the set of all subsequences of \mathbb{N} . \mathcal{A} is a Ramsey set (see e.g., [O]) and thus by our remark above there exists $M \in [\mathbb{N}]$ so that $[M] \subseteq \mathcal{A}$. Thus by passing to a subsequence we may assume that if $n_1 < \dots < n_{2k}$, then

$$\left\| \sum_{i=1}^k (x_{n_{2i}} - x_{n_{2i-1}}) \right\| \leq C.$$

a) By passing to a subsequence of x_n we may assume that (x_n) is basic and moreover $(x_1, x_2 - x_1, x_3 - x_2, \dots)$ is seminormalized basic (see e.g., [Be, Theorem 8] or [R2]). Calling this sequence (y_n) we have that $\left\| \sum_{n \in F} y_n \right\| \leq 2C + \|x_{n_1}\|$ for all finite F , and so (y_n) is equivalent to the unit vector basis of c_0 . Hence (x_n) is equivalent to the summing basis: $x_n = \sum_{i=1}^n y_i$.

b) By Elton's theorem (see [O]) we have that either a subsequence of (x_n) is equivalent to the unit vector basis of c_0 or some subsequence (y_n) of (x_n) satisfies

$$\lim_{k \rightarrow \infty} \left\| \sum_{i=1}^k (-1)^i y_{n_i} \right\| = \infty \text{ for all } n_1 < n_2 < \dots.$$

From our above remarks we have that a subsequence is the unit vector basis of c_0 . \square

Remark. 1. We do not know if X can be given a norm $\|\cdot\|$ satisfying:

$$\text{if } \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m \pm x_n\| = \lim_{m \rightarrow \infty} \|x_m\| > 0,$$

then some subsequence of (x_n) is equivalent to the unit vector basis of c_0 .

We can show that this is the case for $\|\cdot\|$ provided in addition one has

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m \pm x_n\| = \lim_{m \rightarrow \infty} \|x_m\|.$$

However the hypothesis of Corollary 2.6 a) does require the assertion that (x_n) not be weakly convergent. Indeed if X contains c_0 , then there exists a normalized sequence $(y_n) \subseteq X$ which is asymptotically 1-equivalent to the unit vector basis of c_0 and hence

$$1 = \lim_{n_1 \rightarrow \infty} \lim_{n_2 \rightarrow \infty} \|y_{n_1} - y_{n_2}\| = \lim_{n_1 \rightarrow \infty} \lim_{n_2 \rightarrow \infty} \lim_{n_3 \rightarrow \infty} \lim_{n_4 \rightarrow \infty} \|y_{n_1} - y_{n_2} + y_{n_3} - y_{n_4}\|.$$

Thus $x_n = y + y_n$ satisfies the same condition for any $y \neq 0$ but (x_n) admits no basic subsequence.

2. As we have noted parts b) and c) of the main theorem hold for any equivalent norm $\|\cdot\|$ on X where $\|\cdot\| = \sum_{c \in C} p_c \|\cdot\|_c$. From the proof of Lemma 2.4 it follows that whenever $(x_n) \subseteq X$ is relatively weakly compact and satisfies

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = \lim_{m \rightarrow \infty} \|x_m\|,$$

then (x_n) is relatively norm compact.

§3. COROLLARIES

We now give some corollaries. Part a) of Corollary 3.1 yields a positive answer to Milman's problem mentioned above.

Corollary 3.1. *Let X be a separable Banach space. X is reflexive (if and) only if there exists an equivalent norm $\|\cdot\|$ on X satisfying the following for any bounded $(x_n) \subseteq X$:*

a) *If $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 2 \lim_n \|x_n\|$, then (x_n) is norm convergent.*

Furthermore the norm $\|\cdot\|$ in a) satisfies

b) *if (x_n) is weakly null but not norm null, then*

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| > \lim_{m \rightarrow \infty} \|x_m\|$$

provided both limits exist.

Proof. The main theorem (a), c) yields such a norm if X is reflexive. Conversely if a) holds let $x^* \in X^*$ with $\|x^*\| = 1$. Choose $(x_n) \subseteq X$, $\|x_n\| = 1$ with $\lim_{n \rightarrow \infty} x^*(x_n) = 1$. It follows that $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \|x_m + x_n\| = 2$ and so by a), (x_n) converges to some x with $\|x\| = 1$ and $x^*(x) = 1$. Thus x^* achieves its norm. By James' theorem [J1] X must be reflexive. \square

Corollary 3.2. *Let X be a separable Banach space. Then there exists an equivalent norm $\|\cdot\|$ on X such that if Y is a subspace of X , then Y is reflexive iff a) (and b)) of Corollary 3.1 hold for all bounded $(x_n) \subseteq Y$.*

From b) and c) of the main theorem we obtain

Corollary 3.3. *Let X be a separable Banach space. The following are equivalent.*

- 1) *X contains an isomorph of ℓ_1 (respectively, c_0).*
- 2) *For all equivalent norms $\|\cdot\|$ on X there exists a normalized sequence in X having spreading model (e_n) which is 1-equivalent to the unit vector basis of ℓ_1 (respectively, c_0).*
- 3) *For all equivalent norms $\|\cdot\|$ on X there exists a normalized (and respectively, weakly null) sequence in X having spreading model (e_n) satisfying $\|e_1 \pm e_2\| = 2$ (respectively $\|e_1 + e_2\| = 1$).*

In addition to the main theorem the proof requires James' proof that ℓ_1 and c_0 are not distortable ([J2] or [LT, p.97]). Indeed 1) \Rightarrow 2) or 3) is well known from James' result. Our discovery is the reverse implications.

Our work also yields the following corollaries.

Corollary 3.4. *Let X be a separable Banach space. The following are equivalent.*

- 1) *X is not reflexive.*

- 2) For all equivalent norms $\|\cdot\|$ on X there exists a $\|\cdot\|$ -normalized basic sequence (x_i) having spreading model $((e_i), \|\cdot\|)$ satisfying for all $(a_i) \subseteq [0, \infty)$,

$$\left\| \sum a_i e_i \right\| = \sum a_i .$$

- 3) For all equivalent norms $\|\cdot\|$ on X there exists a $\|\cdot\|$ -normalized basic sequence (x_i) having spreading model $((e_i), \|\cdot\|)$ satisfying

$$\|e_1 + e_2\| = 2 .$$

Corollary 3.5. *Let X be a separable Banach space. The following are equivalent.*

- a) X is reflexive.
- b) There exists an equivalent norm $\|\cdot\|$ on X such that if $((e_i), \|\cdot\|)$ is a spreading model of any $\|\cdot\|$ -normalized basic sequence in X , then $1 < \|e_1 + e_2\| < 2$.

REFERENCES

- [BL] B. Beauzamy and J.-T. Lapresté, *Modèles étalés des espaces de Banach*, Travaux en Cours, Herman, Paris, 1984. MR **86h**:46024
- [Be] S. Bellenot, *Somewhat quasireflexive Banach spaces*, Arkiv för matematik **22** (1984), 175–183. MR **86b**:46014
- [BP] C. Bessaga and A. Pełczyński, *On bases and unconditional convergence of series in Banach spaces*, Studia Math. **17** (1958), 151–164. MR **22**:5872
- [B] A. Brunel, *Espaces associés à une suite bornée dans un espace de Banach*, Séminaire Maurey-Schwartz, exposés 15, 16, 18, Ecole Polytechnique, 1973/4. MR **53**:8866
- [BS] A. Brunel and L. Sucheston, *B-convex Banach spaces*, Math. Systems Th. **7** (1974), 294–299. MR **55**:11004
- [CS] P.G. Casazza and T.J. Shura, *Tsirelson's Space*, Lectures Notes in Math., vol. 1363, Springer-Verlag, Berlin and New York, 1989. MR **90b**:46030
- [D1] M.M. Day, *Normed linear spaces*, Springer-Verlag, New York, 1973. MR **49**:9588
- [D2] M.M. Day, *Reflexive spaces not isomorphic to uniformly convex Banach spaces*, Bull. Amer. Math. Soc. **47** (1941), 313–317. MR **2**:221b
- [DGZ] R. Deville, G. Godefroy and V. Zizler, *Smoothness and Renormings in Banach Spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics **64** (1993). MR **94d**:46012
- [FG] K. Fan and I. Glicksberg, *Fully convex normed linear spaces*, Proc. Nat. Acad. Sci. USA **41** (1955), 947–953. MR **17**:386h
- [FJ] T. Figiel and W.B. Johnson, *A uniformly convex Banach space which contains no ℓ_p* , Comp. Math. **29** (1974), 179–190. MR **50**:8011
- [G1] W.T. Gowers, *A new dichotomy for Banach spaces*, Geom. Funct. Anal. **6** (1996), 1083–1093. CMP 97:04
- [G2] W.T. Gowers, *A space not containing c_0 , ℓ_1 or a reflexive subspace*, Trans. Amer. Math. Soc. **344** (1994), 407–420. MR **94j**:46024
- [GM] W.T. Gowers and B. Maurey, *The unconditional basic sequence problem*, J. Amer. Math. Soc. **6** (1993), 851–874. MR **94k**:46021
- [HM] R. Haydon and B. Maurey, *On Banach spaces with strongly separable types*, J. London Math. Soc. (2) **33** (1986), 484–498. MR **87g**:46026
- [J1] R.C. James, *Reflexivity and the sup of linear functionals*, Israel J. Math. **13** (1972), 289–300. MR **49**:3506
- [J2] R.C. James, *Uniformly nonsquare Banach spaces*, Ann. Math. **80** (1964), 542–550. MR **30**:4139
- [J3] R.C. James, *Bases and reflexivity of Banach spaces*, Annals of Math. **52** (1950), 518–527. MR **12**:616b
- [KM] J.L. Krivine and B. Maurey, *Espaces de Banach stables*, Israel J. Math. **39** (1981), 273–295. MR **83a**:46030

- [LT] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces I*, Springer-Verlag, New York, 1977. MR **58**:17766
- [M] B. Maurey, *Types and ℓ_1 -subspaces*, Longhorn Notes, Texas Functional Analysis Seminar 1982-83, The University of Texas at Austin, 123–137.
- [MMT] B. Maurey, V.D. Milman and N. Tomczak-Jaegermann, *Asymptotic infinite-dimensional theory of Banach spaces*, Operator Theory: Advances and Applications **77** (1995), 149–175. MR **97g**:46015
- [Mil] D.P. Milman, *On some criteria for the regularity of spaces of the type (B)*, Dokl. Akad. Nauk SSSR **20** (1938), 243–246, (in Russian).
- [Mi] V.D. Milman, *Geometric theory of Banach spaces II, geometry of the unit sphere*, Russian Math. Survey **26** (1971), 79–163, (trans. from Russian). MR **54**:8240
- [MT] V.D. Milman and N. Tomczak-Jaegermann, *Asymptotic ℓ_p spaces and bounded distortion*, (Bor-Luh Lin and W.B. Johnson, eds.), Contemporary Math. **144** Amer. Math. Soc. (1993), 173–195. MR **94m**:46014
- [O] E. Odell, *Applications of Ramsey theorems to Banach space theory*, Notes in Banach spaces (H.E. Lacey, ed.), U.T. Press, Austin, pp. 379–404. MR **83g**:46018
- [OS] E. Odell and Th. Schlumprecht, *A problem on spreading models*, to appear in J. Funct. Anal.
- [P] B.J. Pettis, *A proof that every uniformly convex space is reflexive*, Duke Math. J. **5** (1939), 249–253.
- [R1] H. Rosenthal, *Double dual types and the Maurey characterization of Banach spaces containing ℓ^1* , Longhorn Notes, Texas Functional Analysis Seminar 1983-84, The University of Texas at Austin, 1–37. CMP 18:10
- [R2] H. Rosenthal, *A characterization of Banach spaces containing c_0* , J. Amer. Math. Soc. **7** (1994), 707–747. MR **94i**:46032
- [R3] H. Rosenthal, *A characterization of Banach spaces containing ℓ^1* , Proc. Nat. Acad. Sci. USA **71** (1974), 2411–2413. MR **50**:10773
- [T] B.S. Tsirelson, *Not every Banach space contains ℓ_p or c_0* , Funct. Anal. Appl. **8** (1974), 138–141.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TEXAS 78712-1082

E-mail address: `odell@math.utexas.edu`

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843-3368

E-mail address: `schlump@math.tamu.edu`