

ON THE EQUATION $\operatorname{div} Y = f$ AND APPLICATION TO CONTROL OF PHASES

JEAN BOURGAIN AND HAÏM BREZIS

1. INTRODUCTION

The purpose of this paper is to present new results concerning the equation

$$(1.1) \quad \operatorname{div} Y = f \quad \text{on } \mathbb{T}^d,$$

i.e., we work on \mathbb{R}^d with 2π -periodic functions in all variables. In what follows we will always assume that $d \geq 2$ and that

$$(1.2) \quad \int_Q f = 0$$

where $Q = (0, 2\pi)^d$. The notations $L^p, W^{1,p}$, etc. refer to $L^p(\mathbb{T}^d), W^{1,p}(\mathbb{T}^d)$, etc. or to 2π -periodic functions in $L^p_{loc}(\mathbb{R}^d), W^{1,p}_{loc}(\mathbb{R}^d)$, etc. We denote by $L^p_{\#}$ the space of functions in L^p satisfying (1.2).

Clearly, (1.1) is an underdetermined problem which admits many solutions. A standard way of tackling (1.1) is to look for a vector field Y satisfying the *additional* condition

$$\operatorname{curl} Y = 0,$$

i.e., one looks for a *special* Y of the form

$$Y = \operatorname{grad} u.$$

Equation (1.1) then becomes

$$(1.3) \quad \Delta u = f$$

and the standard L^p -regularity theory yields a solution $u \in W^{2,p}$ when $f \in L^p_{\#}, 1 < p < \infty$. Consequently (1.1) has a solution $Y \in W^{1,p}$ for every $f \in L^p_{\#}, 1 < p < \infty$. More precisely, the operator $\operatorname{div} : W^{1,p} \rightarrow L^p_{\#}$ admits a right inverse which is a bounded linear operator $K : L^p_{\#} \rightarrow W^{1,p}$. Strictly speaking, we should write $Y \in (W^{1,p})^d (= d\text{-fold copy of } W^{1,p}), \operatorname{div} : (W^{1,p})^d \rightarrow L^p$, etc. But we will often omit the superscript d to alleviate notation.

Three *limiting* cases are of interest:

Received by the editors January 14, 2002 and, in revised form, October 2, 2002.

2000 *Mathematics Subject Classification.* Primary 35C99, 35F05, 35F15, 42B05, 46E35.

Key words and phrases. Divergence equations, gradient equations, critical Sobolev norms.

The first author was partially supported by NSF Grant DMS-9801013.

The second author was partially sponsored by a European Grant ERB FMRX CT98 0201. He is also a member of the Institut Universitaire de France.

The authors thank C. Fefferman, P. Lax, P. Mironescu, L. Nirenberg, T. Rivière, M. Vogelius and D. Ye for useful comments.

Case 1: $\mathbf{p} = \mathbf{1}$. It is well known that when $f \in L^1$ equation (1.3) does not necessarily admit a solution $u \in W^{2,1}$. However, one might still hope to have some solution Y of (1.1) in $W^{1,1}$ or at least in BV . This is not true: for some f 's in L^1 , equation (1.1) has no solution in BV and not even in $L^{d/(d-1)}$; see Section 2.1.

Case 2: $\mathbf{p} = \infty$. It is well known that when $f \in L^\infty$ equation (1.3) does not necessarily admit a solution $u \in W^{2,\infty}$. However, one might hope to find a solution Y of (1.1) in $W^{1,\infty}$. This is not true: McMullen [13] has shown that for some f 's in L^∞ (even continuous f) equation (1.1) has no solution in $W^{1,\infty}$. This is proved using a duality argument and a “non-estimate” of Ornstein [16]; see Section 2.2.

Case 3: $\mathbf{p} = \mathbf{d}$. This is the heart of our work. For every $f \in L^d_\#$, equation (1.3) admits a solution $u \in W^{2,d}$ and thus equation (1.1) admits a solution $Y = \text{grad } u \in W^{1,d}$. Since $W^{1,d}$ is *not* contained in L^∞ (this is a limiting case for the Sobolev imbedding), we *cannot* assert that this Y belongs to L^∞ . In fact, we give in Section 3 (Remark 7) an explicit $f \in L^d$ such that the corresponding $Y = \text{grad } u$ does *not* belong to L^∞ . However one might still hope that given any $f \in L^d_\#$ there is *some* $Y \in L^\infty$ solving (1.1). This is indeed true:

Proposition 1. *Given any $f \in L^d_\#$ there exists some $Y \in L^\infty$ solving (1.1) (in the sense of distributions) with*

$$(1.4) \quad \|Y\|_{L^\infty} \leq C(d)\|f\|_{L^d}.$$

Remark 1. A more precise statement established in the course of the proof says that there exists $Y \in C^0$ satisfying (1.1) and (1.4).

The proof of Proposition 1 is quite elementary; see Section 3. It relies on the Sobolev-Nirenberg imbedding $W^{1,1} \subset L^{d/(d-1)}$ (and even $BV \subset L^{d/(d-1)}$) combined with duality, i.e., Hahn-Banach. As a consequence, the argument is *not constructive*, and Y is not obtained as above via a bounded linear operator acting on f . In fact, surprisingly, the operator div has no bounded right inverse in this setting:

Proposition 2. *There exists no bounded linear operator $K: L^d_\# \rightarrow L^\infty$ such that $\text{div } Kf = f \quad \forall f \in L^d_\#$ (in the sense of distributions).*

Remark 2. Another way of formulating Proposition 2 is to say that the subspace $\{Y \in L^\infty; \text{div } Y = 0\}$ admits no complement in the space $\{Y \in L^\infty; \text{div } Y \in L^d\}$ equipped with its natural norm. Alternatively, the closed subspace $\{\text{grad } u; u \in W^{1,1}\}$ has no complement in L^1 ; see Section 3.

To summarize: for every $f \in L^d_\#$, equation (1.1) admits

- a) a solution $Y_1 \in W^{1,d}$,
- b) a solution $Y_2 \in L^\infty$.

A natural question is whether there exists a solution Y of (1.1) in $L^\infty \cap W^{1,d}$. This is indeed one of our main results.

Theorem 1. *For every $f \in L^d_\#$ there exists a solution $Y \in L^\infty \cap W^{1,d}$ of (1.1) satisfying*

$$(1.5) \quad \|Y\|_{L^\infty} + \|Y\|_{W^{1,d}} \leq C(d)\|f\|_{L^d}.$$

Despite the simplicity of this statement the argument is rather involved and a simpler proof would be desirable.

We will present two techniques to tackle Theorem 1.

First proof of Theorem 1 when $d = 2$ (see Section 4). It relies on Hahn-Banach (via duality) and thus it is *not* constructive. But it is rather elementary; the main ingredient is the new estimate (1.6) which is established by L^2 -Fourier methods.

Lemma 1. *On \mathbb{T}^2 we have*

$$(1.6) \quad \|u - fu\|_{L^2} \leq C \|\operatorname{grad} u\|_{L^1 + H^{-1}}, \quad \forall u \in L^2,$$

for some absolute constant C .

The main difficulty, in proving (1.6), stems from the fact that if we decompose

$$\operatorname{grad} u = h_1 + h_2$$

with $h_1 \in L^1$ and $h_2 \in H^{-1}$, then h_1 and h_2 need *not* be gradients themselves; it is only their sum which is a gradient.

The analogue of Lemma 1 for $d > 2$ is the estimate on \mathbb{T}^d ,

$$(1.7) \quad \|u - fu\|_{L^{d/(d-1)}} \leq C(d) \|\operatorname{grad} u\|_{L^1 + W^{-1, d/(d-1)}}.$$

We have no direct proof of (1.7). But it can be deduced by duality from the statement of Theorem 1 (and thus from the second proof presented in Section 7).

Second proof of Theorem 1, valid for all $d \geq 2$ (see Sections 5 and 6). We exhibit via a *constructive* (nonlinear) argument some explicit $Y \in W^{1,d} \cap L^\infty$ satisfying (1.1) and (1.5). The argument for $d = 2$ is simpler and we start with this case for expository reasons.

One should observe a certain analogy with the Fefferman-Stein [10] decomposition of BMO-functions and Uchiyama's [21] constructive proof. Indeed, returning to equation (1.1) and defining F by $|\xi|\hat{F}(\xi) = \hat{f}(\xi)$, we obtain that $F \in W^{1,d} \subset BMO$ and (1.1) becomes

$$(1.8) \quad F = \sum_{j=1}^d R_j Y_j$$

with $R_j = j^{th}$ Riesz transform ($\widehat{R_j \psi}(\xi) = \hat{\psi}(\xi) \frac{\xi_j}{|\xi|}$), $Y = (Y_1, \dots, Y_d)$.

The statement of Theorem 1 is that (1.8) has a solution $Y \in L^\infty \cap W^{1,d}$. Recall that according to Fefferman-Stein [10] any $F \in BMO$ has a decomposition of the form

$$(1.9) \quad F = Y_0 + \sum_{j=1}^d R_j Y_j \quad \text{with } Y_0, Y_1, \dots, Y_d \in L^\infty.$$

The proof of this decomposition is again by duality and nonconstructive. The later constructive approach from Uchiyama [21] gives a different proof of (1.9). If we assume moreover that $F \in W^{1,d}$, Uchiyama's argument gives that (1.9) has a solution $Y_0, Y_1, \dots, Y_d \in L^\infty \cap W^{1,d}$. The new result in this paper shows that, in fact, for $F \in W^{1,d}$, the Y_0 -component is unnecessary and (1.8) holds for some $Y_1, \dots, Y_d \in L^\infty \cap W^{1,d}$.

It should be mentioned that to achieve our decomposition we do use significantly different methods from Uchiyama. This raises the question what are the function

spaces $X, W^{1,d} \subset X \subset BMO$, such that every $F \in X$ has a decomposition

$$(1.10) \quad F = \sum_{j=1}^d R_j Y_j$$

where $Y_j \in L^\infty$ or (assuming the Riesz transforms bounded on X) the stronger property $Y_j \in L^\infty \cap X$.

Remark 3. Using Theorem 1 we will prove (in Sections 4 and 6) that a slightly stronger conclusion holds:

Theorem 1'. *For every $f \in L^d_{\#}$ there exists a solution $Y \in C^0 \cap W^{1,d}$ of (1.1) satisfying (1.5).*

The original motivation for studying (1.1) comes from the following question about lifting discussed in Bourgain-Brezis-Mironescu [3], [4], [5]. Consider the equation

$$g = e^{i\varphi} \quad \text{on } \mathbb{T}^d$$

where φ is a smooth real-valued function.

Question. Assuming g is controlled in $H^{1/2}$, what kind of estimate can we deduce for φ ?

Here is a first easy consequence of Theorem 1.

Corollary 1. *We have*

$$(1.11) \quad \|\varphi - f\varphi\|_{L^{d/(d-1)}} \leq C(d)(1 + \|g\|_{H^{1/2}})\|g\|_{H^{1/2}}.$$

Proof. Write

$$\text{grad } g = ie^{i\varphi} \text{ grad } \varphi$$

and thus

$$(1.12) \quad \text{grad } \varphi = -i\bar{g}(\text{grad } g).$$

Multiplying by Y gives

$$(1.13) \quad \int_Q \varphi \text{ div } Y = \int_Q i\bar{g}Y \cdot \text{grad } g.$$

Given $f \in L^d$ we obtain from Theorem 1 some Y satisfying (1.1) (with f replaced by $f - f\varphi$) and (1.5). Thus we have

$$(1.14) \quad \left| \int (\varphi - f\varphi)f \right| \leq \|g\|_{H^{1/2}}(\|\bar{g}Y\|_{H^{1/2}}).$$

But

$$(1.15) \quad \|\bar{g}Y\|_{H^{1/2}} \leq \|g\|_{H^{1/2}}\|Y\|_{L^\infty} + \|g\|_{L^\infty}\|Y\|_{H^{1/2}}$$

$$\text{(by (1.5))} \leq C(\|g\|_{H^{1/2}}\|f\|_{L^d} + \|f\|_{L^d})$$

where we have used the obvious fact that $\|Y\|_{H^{1/2}} \leq C\|Y\|_{W^{1,d}}$. Combining (1.14) and (1.15) yields (1.11). □

Remark 4. Estimate (1.11) cannot be improved, replacing the norm $\|\cdot\|_{L^{d/(d-1)}}$ by $\|\cdot\|_{L^p, p > d/(d-1)}$. This may be seen by choosing $g = e^{i\varphi}$ with $\varphi(x) = (|x|^2 + \varepsilon^2)^{-\alpha/2}$ with $\alpha < d-1, \alpha$ close to $(d-1)$ and ε close to 0 (the same example has already been used in Bourgain-Brezis-Mironescu [3], Lemma 5). There is however a better estimate than (1.11), namely

Theorem 4. *Let φ be a smooth real-valued function on \mathbb{T}^d and set $g = e^{i\varphi}$, then*

$$\|\varphi\|_{H^{1/2}+W^{1,1}} \leq C(d)(1 + \|g\|_{H^{1/2}})\|g\|_{H^{1/2}}.$$

Theorem 4 has been announced in Bourgain-Brezis-Mironescu [4] (Theorem 3) and is proved in Section 8. Our proof of Theorem 4 is a direct estimate based on paraproducts. In view of the preceding argument one may wonder whether Theorem 4 can be proved by solving a divergence equation. After duality the required statement would be

$$(1.16) \quad \|u - fu\|_{H^{1/2}+W^{1,1}} \leq C\|\operatorname{grad} u\|_{H^{-1/2}+L^1}$$

but we do not know whether (1.16) holds.

We now turn to the question of coupling equation (1.1) with the Dirichlet condition

$$(1.17) \quad Y = 0 \quad \text{on } \partial Q.$$

This question was addressed (in various forms) by a few authors; see e.g. Arnold–Scott–Vogelius [2], Duvaut–Lions [9] (Theorem 3.2), X. Wang [22], Temam [20] (Proposition 1.2(ii) and Lemma 2.4) and the references therein to Magenes–Stampacchia [12] and Nečas [14]. Our aim is to establish the analogue of Theorem 1' under the Dirichlet condition. We start with the following known fact (see e.g. Arnold–Scott–Vogelius [2] for $d = 2$).

Theorem 2. *Given $f \in L^p_{\#}(Q), 1 < p < \infty$, there exists some $Y \in W^{1,p}_0(Q)$ satisfying (1.1) with*

$$(1.18) \quad \|Y\|_{W^{1,p}} \leq C(p)\|f\|_{L^p}.$$

Moreover Y can be chosen, depending linearly on f .

The operator and the estimate do not depend on p assuming we stay away from the end points.

For the convenience of the reader we include a new proof; our technique is extremely elementary and can be adapted to establish, for the limiting case $p = d$,

Theorem 3. *Given $f \in L^d_{\#}(Q)$ there exists some $Y \in C^0(\bar{Q}) \cap W^{1,d}_0(Q)$ satisfying (1.1) with*

$$\|Y\|_{L^\infty} + \|Y\|_{W^{1,d}} \leq C\|f\|_{L^d}.$$

Theorem 3 is stronger than Theorem 1'. However it will be deduced from Theorem 1'. There are variants of Theorems 2 and 3 when Q is replaced by a Lipschitz domain in \mathbb{R}^d (see Section 7.2).

The plan of the paper is the following:

1. Introduction.
2. The cases $f \in L^p$ with $p = 1$ and $p = \infty$.
3. Proofs of Propositions 1 and 2 and related questions.
4. Proof of Theorem 1 when $d = 2$ via duality.
5. Proof of Theorem 1 when $d = 2$ (explicit construction).
6. Proof of Theorem 1 when $d > 2$ (explicit construction).
7. The equation $\operatorname{div} Y = f$ with Dirichlet condition. Proof of Theorems 2 and 3.
8. Estimation of the phase in $H^{1/2} + W^{1,1}$. Proof of Theorem 4.

2. THE CASES $f \in L^p$ WITH $p = 1$ AND $p = \infty$

We consider here equation (1.1) with $f \in L^p_{\#}$ and ask whether there exists a solution $Y \in W^{1,p}$ of (1.1) when $p = 1$ and $p = \infty$. As we have already mentioned in the Introduction the answer is negative. Here is the proof.

2.1. The case $p = 1$. Assume by contradiction that for every $f \in L^1_{\#}$ there is some $Y \in W^{1,1}$ satisfying (1.1). It follows that the linear operator

$$Tu = \operatorname{div} u \text{ from } E = W^{1,1} \text{ into } F = L^1_{\#}$$

is bounded and surjective. By the open mapping principle there is a constant C such that for every $f \in F$ there exists a solution $Y \in E$ of (1.1) satisfying

$$\|Y\|_{W^{1,1}} \leq C\|f\|_{L^1}.$$

We now use a duality argument which occurs frequently in the rest of the paper. We will deduce that $W^{1,d} \subset L^\infty$ with continuous injection, and since this is false, we infer that for some f 's in F there is no $Y \in W^{1,1}$ satisfying (1.1).

Let $u \in W^{1,d}$ and set

$$(2.1) \quad \operatorname{grad} u = h \in L^d.$$

Given any $f \in L^1$, let $Y \in W^{1,1}$ be such that

$$\operatorname{div} Y = f - \int_Q f$$

and

$$\|Y\|_{W^{1,1}} \leq C\|f - \int_Q f\|_{L^1}.$$

Taking the scalar product of (2.1) with Y and integrating yields

$$\int_Q (u - \int_Q u) f = - \int_Q hY.$$

Consequently

$$(2.2) \quad \left| \int_Q (u - \int_Q u) f \right| \leq \|h\|_{L^d} \|Y\|_{L^{d/(d-1)}}.$$

By the Sobolev-Nirenberg imbedding we have $W^{1,1} \subset L^{d/(d-1)}$ and thus

$$(2.3) \quad \|Y\|_{L^{d/(d-1)}} \leq C\|Y\|_{W^{1,1}} \leq C\|f\|_{L^1}.$$

Combining (2.2) and (2.3) we deduce that $(u - \int_Q u) \in L^\infty$ with

$$\|u - \int_Q u\|_{L^\infty} \leq C\|\operatorname{grad} u\|_{L^d}.$$

Impossible.

Remark 5. The same argument shows that equation (1.1) with $f \in L^1_{\#}$ need not have a solution Y in the sense of distributions with $Y \in L^{d/(d-1)}$. (Note, however, that the solution Y given via (1.3) belongs to L^p , $\forall p < d/(d-1)$, and even to weak- $L^{d/(d-1)}$). It suffices to follow the above argument with $E = W^{1,1}$ replaced by

$$\tilde{E} = \{Y \in L^{d/(d-1)}; \operatorname{div} Y \in L^1\}$$

equipped with its natural norm.

2.2. The case $p = \infty$. This case has been settled negatively by McMullen [13] (the interest in this kind of problem grew out of the study of the equation $\det(\nabla\varphi) = f$ with φ bi-Lipschitz and also from a question of Gromov [11] on separated nets; see Dacorogna-Moser [18], Ye [24], Rivière-Ye [17],[18], Burago-Kleiner [7]).

For the convenience of the reader we sketch a proof when $d = 2$, which is essentially similar to the one of McMullen [13]. We argue by contradiction as above. Then, for every $f \in L^\infty$ there is a $Y \in W^{1,\infty}$ satisfying

$$\operatorname{div} Y = f - ff$$

and

$$\|Y\|_{W^{1,\infty}} \leq C\|f\|_{L^\infty}.$$

Let ψ be a smooth function on \mathbb{T}^2 and set $g = \psi_{x_1x_2}$. Write

$$\int g_{x_1}Y_1 + g_{x_2}Y_2 = - \int gf = - \int \psi_{x_1x_1}Y_{1x_2} + \psi_{x_2x_2}Y_{2x_1}.$$

Consequently

$$\left| \int gf \right| \leq C(\|\psi_{x_1x_1}\|_{L^1} + \|\psi_{x_2x_2}\|_{L^1})\|f\|_{L^\infty}$$

and thus

$$\|g\|_{L^1} = \|\psi_{x_1x_2}\|_{L^1} \leq C(\|\psi_{x_1x_1}\|_{L^1} + \|\psi_{x_2x_2}\|_{L^1}).$$

This contradicts a celebrated “non-inequality” of Ornstein [16] and completes the proof.

Remark 6. The same argument shows that equation (1.1) with $f \in C^0$ and $\int f = 0$ need not have a solution $Y \in W^{1,\infty}$.

3. PROOFS OF PROPOSITIONS 1 AND 2 AND RELATED QUESTIONS

Proof of Proposition 1. Recall the Sobolev-Nirenberg imbedding $W^{1,1} \subset L^{d/(d-1)}$ and, more generally, $BV \subset L^{d/(d-1)}$ with

$$(3.1) \quad \|u - fu\|_{L^{d/(d-1)}} \leq C(d)\|\operatorname{grad} u\|_{\mathcal{M}} \quad \forall u \in BV,$$

where \mathcal{M} denotes the space of measures. Set

$$E = C^0, \quad F = L^d_{\#}$$

and consider the unbounded linear operator $A = D(A) \subset E \rightarrow F$, defined by

$$D(A) = \{Y \in E; \operatorname{div} Y \in L^d\}, \quad AY = \operatorname{div} Y,$$

so that A is densely defined and has closed graph. Clearly we have

$$E^* = \mathcal{M}, \quad F^* = L^{d/(d-1)}_{\#},$$

$$D(A^*) = F^* \cap BV, \quad A^*u = \operatorname{grad} u.$$

By (3.1) we have

$$\|u\|_{F^*} \leq C(d)\|A^*u\|_{E^*} \quad \forall u \in D(A^*).$$

It follows from the closed-range theorem (see e.g. Brezis [6], Section II.7) that A is surjective. More precisely, we claim that for any $f \in F$ there is some $Y \in E$ satisfying (1.1) and

$$\|Y\|_{L^\infty} \leq 2C(d)\|f\|_{L^d},$$

where $C(d)$ is the constant in (3.1). □

Indeed, let $f \in F$ with $\|f\|_{L^d} = 1$ and consider the two convex sets

$$B = \{Y \in E; \|Y\|_E < 2C(d)\}$$

and

$$L = \{Y \in E; \operatorname{div} Y = f\}.$$

We have to prove that $B \cap L \neq \emptyset$. Suppose not, and $B \cap L = \emptyset$. Then, by Hahn-Banach there exists $\mu \in E^*$, $\mu \neq 0$, and $\alpha \in \mathbb{R}$ such that

$$(3.2) \quad \langle \mu, Y \rangle \leq \alpha \quad \forall Y \in B$$

and

$$(3.3) \quad \langle \mu, Y \rangle \geq \alpha \quad \forall Y \in L.$$

From (3.2) we have $\|\mu\| \leq \alpha/2C(d)$ and from (3.3) we deduce, in particular, that $\langle \mu, Z \rangle = 0 \quad \forall Z \in N(A)$. It follows that $\mu \in N(A)^\perp = R(A^*)$. Hence there exists some $u \in F^* \cap BV$ such that $\operatorname{grad} u = \mu$. Applying (3.1) we see that

$$(3.4) \quad \|u\|_{L^{d/(d-1)}} \leq C(d)\|\mu\| \leq \alpha/2.$$

On the other hand, by (3.3), $\forall Y \in L$,

$$\alpha \leq \langle \mu, Y \rangle = \langle \operatorname{grad} u, Y \rangle = - \int u \operatorname{div} Y = - \int u f \leq \|u\|_{L^{d/(d-1)}} \leq \alpha/2.$$

This is impossible since $\alpha > 0$ (because $\mu \neq 0$).

Remark 7. The special solution of (1.1) given by $Y = \operatorname{grad} u$, where u is the solution of (1.3), belongs to $W^{1,d}$ when $f \in L^d$; however, in general, it does not belong to L^∞ . Here is an example due to L. Nirenberg. Using (x_1, x_2, \dots, x_d) as coordinates in \mathbb{R}^d consider the function

$$u = x_1 |\log r|^{\alpha} \zeta$$

where ζ is a smooth cut-off function with support near 0 and $0 < \alpha < (d-1)/d$. Note that $Y = \operatorname{grad} u$ does not belong to L^∞ while

$$|\Delta u| \leq \frac{C}{r} |\log r|^{\alpha-1},$$

so that $\Delta u \in L^d$.

We now turn to the proof of Proposition 2, i.e., the non-existence of a bounded right inverse $K : L^d_{\#} \rightarrow L^\infty$ for the operator div . We present two proofs. The first is the simplest: after a standard averaging trick we obtain a bounded multiplier $L^d \rightarrow L^\infty$ and we reach a contradiction by a direct summability consideration. The second proof is related to Remark 2: the existence of K would yield a factorization of the identity map $I : W^{1,1} \rightarrow L^{d/(d-1)}$ through the Banach space L^1 ; however no such factorization exists by a general argument from the geometry of Banach spaces.

First proof of Proposition 2. Assume $K : L^d_{\#} \rightarrow L^\infty$ is a bounded operator satisfying $\operatorname{div} K = I$ on $L^d_{\#}$. Then the averaged operator

$$\tilde{K} = \int_{\mathbb{T}^d} \tau_{-x} K \tau_x dx,$$

where $\tau_x f(y) = f(y+x)$, still satisfies

$$(3.5) \quad \operatorname{div} \tilde{K} = I \quad \text{on } L^d.$$

On the other hand, \tilde{K} is clearly a multiplier

$$\tilde{K}(e^{in \cdot x}) = (\lambda_1(n), \lambda_2(n), \dots, \lambda_d(n))e^{in \cdot x}$$

which is bounded from L^d into L^∞ and hence from L^1 into $L^{d'}$ where $d' = d/(d-1)$. By (3.5) we have

$$\sum_{j=1}^d n_j \lambda_j(n) = 1 \quad \forall n \in \mathbb{Z}^d$$

so that

$$(3.6) \quad |\lambda(n)|^2 = \sum_{j=1}^d |\lambda_j(n)|^2 \geq 1/|n|^2 \quad \forall n.$$

Consider the multiplier

$$M(e^{in \cdot x}) = \frac{1}{|n|^{\frac{d}{2}-1}} e^{in \cdot x}, \quad n \neq 0.$$

Then M is bounded from $L^{d'}$ into L^2 . Hence $M\tilde{K}$ is a bounded multiplier from L^1 into L^2 . Thus

$$\sum_{\substack{n \in \mathbb{Z}^d \\ n \neq 0}} \frac{|\lambda_j(n)|^2}{|n|^{d-2}} < \infty, \quad \forall j.$$

Summing over $j = 1, 2, \dots, d$, and using (3.6) we deduce

$$\sum_{\substack{n \in \mathbb{Z}^d \\ n \neq 0}} \frac{1}{|n|^d} < \infty.$$

A contradiction. □


Second proof of Proposition 2. Assuming the existence of $K : L_{\#}^d \rightarrow L^\infty$ we obtain a factorization of the identity map $I : W^{1,1} \rightarrow L^{d'}$ as

$$I = K^* \circ \operatorname{grad}$$

which, in particular, gives a factorization of I through the Banach space L^1 . We claim that there is no such factorization, as a consequence of Grothendieck's theorem on absolutely summing operators. Both the result and the method are well known and we briefly recall them (see Wojtaszczyk [23] for details). First take $d = 2$. Then $I : W^{1,1} \rightarrow L^2$ and we consider the operator $I \circ D$ where $D : L^2 \rightarrow W^{1,1}$ is defined by

$$D(e^{in \cdot x}) = \frac{1}{\sqrt{1 + |n|^2}} e^{in \cdot x}.$$

Thus D is clearly bounded as an operator into H^1 , hence into $W^{1,1}$. Since I is assumed to factor through L^1 , so does $I \circ D$:



□

Next, recall Grothendieck’s theorem that any bounded operator $B : L^1 \rightarrow L^2$ is 1-summing, i.e.,

$$\pi_1(B) \equiv \sup \left\{ \sum \|Bx_i\|; (x_i) \subset L^1 \text{ and } \max_{x^* \in L^\infty, \|x^*\| \leq 1} \sum |\langle x_i, x^* \rangle| \leq 1 \right\} \leq K_G \|B\|,$$

where K_G is Grothendieck’s constant.

From the usual ideal properties, we obtain

$$\begin{aligned}
 \left(\sum_{n \in \mathbb{Z}^2} \frac{1}{1 + |n|^2} \right)^{1/2} &= \|I \circ D\|_{HS} = \pi_2(I \circ D) \leq \pi_1(I \circ D) \\
 &= \pi_1(B \circ A) \leq \|A\| \pi_1(B) \leq K_G \|A\| \|B\| < \infty,
 \end{aligned}$$

which in an obvious contradiction.

For $d > 2$, we have $I : W^{1,1} \rightarrow L^d$ and we consider the multiplication operator $M : L^d \rightarrow L^2$ given by $M(e^{in \cdot x}) = (1 + |n|)^{1 - \frac{d}{2}} e^{in \cdot x}$. Hence, considering now $M \circ I \circ D : L^2 \rightarrow L^2$ factoring through L^1 , we obtain a contradiction again:

$$\left(\sum \frac{1}{(1 + |n|)^{d-2}(1 + |n|^2)} \right)^{1/2} = \|M \circ I \circ D\|_{HS} = \pi_2(M \circ I \circ D) \leq \pi_1(M \circ I \circ D) < \infty.$$

Proof of Remark 2. Consider the Banach space

$$E = \{Y \in L^\infty; \operatorname{div} Y \in L^d\}$$

equipped with its natural norm $\|Y\|_{L^\infty} + \|\operatorname{div} Y\|_{L^d}$. Then

$$N = \{Y \in L^\infty; \operatorname{div} Y = 0\}$$

is a closed subspace of E which admits no complement in E . Indeed, set

$$F = L^d_{\#}$$

and consider the bounded linear operator $T : E \rightarrow F$ defined by $TY = \operatorname{div} Y$. By Proposition 1, T is surjective. If $N = N(T)$ admits a complement in E , then T has a bounded right inverse, i.e., an operator $S : F \rightarrow E$ such that

$$\operatorname{div} (Sf) = f \quad \forall f \in F$$

(see e.g. Brezis [6], Théorème II.10). But this is impossible by Proposition 2.

Similarly, the subspace

$$R = \{\operatorname{grad} u; u \in W^{1,1}\}$$

of L^1 is closed and admits no complement in L^1 . Indeed, consider the spaces $E = \{u \in W^{1,1}; \int u = 0\}$, $F = L^1$ and the operator $T = \operatorname{grad}$, a bounded linear injective operator from E into F . If $R = R(T)$ admits a complement in F , then

T has a bounded left inverse $S : F \rightarrow E$ (see e.g. Brezis [6], Théorème II.11). In particular, $S : F \rightarrow L^d_{\#}$ satisfies

$$S(\operatorname{grad} u) = u, \quad \forall u \in W^{1,1} \text{ with } \int u = 0.$$

Then $S^* : L^d_{\#} \rightarrow L^\infty$ satisfies

$$\operatorname{div} (S^* f) = f, \quad \forall f \in L^d_{\#},$$

and this is again impossible by Proposition 2. □

4. PROOF OF THEOREM 1 WHEN $d = 2$ VIA DUALITY

We now return to the periodic setting and we will prove the slightly stronger form of Theorem 1,

Theorem 1' (for $d = 2$). *For every $f \in L^2_{\#}$ there exists a solution $Y \in C^0 \cap H^1$ of (1.1) with*

$$(4.1) \quad \|Y\|_{L^\infty} + \|Y\|_{H^1} \leq C \|f\|_{L^2}$$

for some absolute constant C .

Theorem 1' is proved by duality from

Lemma 2. *On \mathbb{T}^2 we have*

$$(4.2) \quad \|u - \int u\|_{L^2} \leq C \|\operatorname{grad} u\|_{L^1 + H^{-1}}, \forall u \in L^2$$

where C is an absolute constant.

Assuming the lemma we turn to the

Proof of Theorem 1'. First observe that

$$L^1 + H^{-1} \subset \mathcal{M} + H^{-1}$$

and that

$$(4.3) \quad \|\cdots\|_{L^1 + H^{-1}} = \|\cdots\|_{\mathcal{M} + H^{-1}} \text{ on } L^1 + H^{-1}$$

(this may be easily seen using regularization by convolution).

Let $E = C^0 \cap H^1, F = L^2_{\#}$ and consider the bounded operator $T : E \rightarrow F$ defined by $TY = \operatorname{div} Y$. Clearly, $T^* : F^* = F \rightarrow E^* = \mathcal{M} + H^{-1}$ is given by $T^*u = \operatorname{grad} u$. By Lemma 2 we have

$$\|u\|_{F^*} \leq C \|T^*u\|_{E^*} \quad \forall u \in F^*,$$

and therefore T is surjective from E onto F . Estimate (4.1) follows from the open mapping principle or one could argue directly using (4.2) and Hahn-Banach as in the proof of Proposition 1. □

Proof of Lemma 2. Assume

$$(4.4) \quad u \in L^2_{\#},$$

$$(4.5) \quad \partial_x u = F_1 + h_1, \quad \partial_y u = F_2 + h_2$$

and

$$(4.6) \quad \|F_1\|_{L^1} + \|F_2\|_{L^1} + \|h_1\|_{H^{-1}} + \|h_2\|_{H^{-1}} \leq 1.$$

We have to prove that

$$(4.7) \quad \|u\|_{L^2} \leq C.$$

□

The main ingredient is

Lemma 3. *Under assumptions (4.4)–(4.6) we have*

$$(4.8) \quad \sum_{n_1, n_2 \in \mathbb{Z}} \frac{n_1^2 n_2^2}{(n_1^2 + n_2^2)^2} |\hat{u}(n_1, n_2)|^2 \leq C(\|u\|_{L^2} + 1).$$

Assuming Lemma 3 we may now complete the proof of Lemma 2. Define

$$(4.9) \quad u'(x', y') = u(x' + y', x' - y') = \sum_{n_1, n_2} \hat{u}(n_1, n_2) e^{i[(n_1 + n_2)x' + (n_1 - n_2)y']}$$

so that

$$(4.10) \quad \widehat{u}'(n_1 + n_2, n_1 - n_2) = \hat{u}(n_1, n_2)$$

and

$$\begin{aligned} \partial_{x'} u'(x', y') &= \partial_x u(x' + y', x' - y') + \partial_y u(x' + y', x' - y') \\ &= (F_1 + F_2)(x' + y', x' - y') + (h_1 + h_2)(x' + y', x' - y') \\ &\in L^1 + H^{-1} \end{aligned}$$

and similarly for $\partial_{y'} u'$.

From (4.8) and (4.10) we obtain

$$(4.11) \quad \begin{aligned} \sum_{n_1, n_2} \frac{(n_1 + n_2)^2 (n_1 - n_2)^2}{4(n_1^2 + n_2^2)^2} |\hat{u}(n_1, n_2)|^2 &= \sum_{n'_1, n'_2} \frac{(n'_1)^2 (n'_2)^2}{((n'_1)^2 + (n'_2)^2)^2} |\widehat{u}'(n'_1, n'_2)|^2 \\ &\leq C(\|u'\|_{L^2} + 1) = C(\|u\|_{L^2} + 1). \end{aligned}$$

Addition of (4.8) and (4.11) implies that

$$\|u\|_{L^2}^2 = \sum_{n_1, n_2} |\hat{u}(n_1, n_2)|^2 \leq C(\|u\|_{L^2} + 1)$$

and the desired estimate (4.7) follows.

We now turn to the

Proof of Lemma 3. We have

$$\begin{aligned} \sum_{n \neq 0} \frac{n_1^2 n_2^2}{(n_1^2 + n_2^2)^2} |\hat{u}(n)|^2 &= \frac{1}{i} n \sum \frac{n_1 n_2^2}{(n_1^2 + n_2^2)^2} \widehat{\partial_x u}(n) \hat{u}(-n) \\ &\stackrel{\text{by (4.5)}}{=} \frac{1}{i} \sum \frac{n_1 n_2^2}{(n_1^2 + n_2^2)^2} \hat{F}_1(n) \hat{u}(-n) + \frac{1}{i} \sum \frac{n_1 n_2^2}{(n_1^2 + n_2^2)^2} \hat{h}_1(n) \hat{u}(-n) \\ &= (4.12) + (4.13). \end{aligned}$$

Estimate

$$(4.14) \quad |(4.13)| \leq \sum_{n_1, n_2} \frac{|\hat{h}_1(n)|}{\sqrt{n_1^2 + n_2^2}} |\hat{u}(-n)| \leq \|h_1\|_{H^{-1}} \|u\|_{L^2}.$$

Write

$$\begin{aligned}
 (4.12) &= \sum \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \widehat{F}_1(n) \widehat{\partial_y u}(-n) \\
 &= \sum \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \widehat{F}_1(n) \widehat{F}_2(-n) + \sum \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \widehat{F}_1(n) \widehat{h}_2(-n) \\
 &= (4.15) + (4.16).
 \end{aligned}$$

Estimate

$$\begin{aligned}
 (4.16) &\leq \sum \frac{|n_1| |n_2|}{(n_1^2 + n_2^2)^2} (|\widehat{\partial_x u}(n)| + |\widehat{h}_1(n)|) |\widehat{h}_2(-n)| \\
 (4.17) &\leq \sum \frac{n_1^2 |n_2|}{(n_1^2 + n_2^2)^2} |\widehat{u}(n)| |\widehat{h}_2(-n)| + \sum \frac{|\widehat{h}_1(n)|}{\sqrt{n_1^2 + n_2^2}} \frac{|\widehat{h}_2(-n)|}{\sqrt{n_1^2 + n_2^2}} \\
 &\leq \|f\|_{L^2} \|h_2\|_{H^{-1}} + \|h_1\|_{H^{-1}} \|h_2\|_{H^{-1}}.
 \end{aligned}$$

□

Estimation of (4.15). This is the key point. Since $\|F_1\|_{L^1} \leq 1, \|F_2\|_{L^1} \leq 1$, it suffices (by convexity) to replace $\widehat{F}_i(n)$ by

$$(4.18) \quad \widehat{F}_1(n) = e^{in \cdot a}, \quad \widehat{F}_2(n) = e^{in \cdot b}$$

for some $a, b \in \mathbb{T}^2$ (this amounts to replacing F_1, F_2 by the Dirac measures δ_a, δ_b , respectively).

Thus we obtain

$$\begin{aligned}
 \sum_{n_1, n_2 \in \mathbb{Z}} \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \widehat{F}_1(n) \widehat{F}_2(-n) &= \sum \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} e^{i[n_1(a_1 - b_1) + n_2(a_2 - b_2)]} \\
 (4.19) \quad &= - \sum \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \sin n_1(a_1 - b_1) \sin n_2(a_2 - b_2)
 \end{aligned}$$

by parity considerations.

Claim. For all $\theta_1, \theta_2 \in \mathbb{T}$

$$(4.20) \quad \left| \sum_{n_1, n_2} \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \sin n_1 \theta_1 \sin n_2 \theta_2 \right| \leq C.$$

From the claim, we conclude that $|(4.15)|, |(4.19)| \leq C$ and, recalling also (4.14), (4.17), inequality (4.8) follows.

Proof of the Claim. Splitting \mathbb{Z} in dyadic intervals, we obtain

$$(4.21) \quad \sum_{k_1, k_2 \geq 0} \left| \sum_{n_1 \sim 2^{k_1}, n_2 \sim 2^{k_2}} \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \sin n_1 \theta_1 \sin n_2 \theta_2 \right|.$$

Recall the inequality

$$(4.22) \quad \left| \sum_{n \in I} \sin n \theta \right| \lesssim 4^k |\theta| \wedge \frac{1}{|\theta|}$$

if $\theta \in \mathbb{T}$ and $I \subset [2^{k-1}, 2^k]$ is an interval (where \wedge denotes min).

From (4.22), assuming $k_1 \geq k_2$, we have

$$(4.23) \quad \left| \sum_{n_1 \sim 2^{k_1}, n_2 \sim 2^{k_2}} \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \sin n_1 \theta_1 \sin n_2 \theta_2 \right| \leq \left(4^{k_1} |\theta_1| \wedge \frac{1}{|\theta_1|} \right) \left(4^{k_2} |\theta_2| \wedge \frac{1}{|\theta_2|} \right) \left\| \left\{ \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \right\} \right\|_{\ell^\infty(n_1 \sim 2^{k_1}) \hat{\otimes} \ell^\infty(n_2 \sim 2^{k_2})}$$

where $\ell^\infty(I) \hat{\otimes} \ell^\infty(J)$ denotes the usual projective tensor product. Thus the last factor in (4.23) may be bounded by

$$(4.24) \quad \left\| \partial_{n_1 n_2}^2 \frac{n_1 n_2}{(n_1^2 + n_2^2)^2} \right\|_{\ell^1(n_1 \sim 2^{k_1}, n_2 \sim 2^{k_2})} \leq C \left\| \frac{1}{(n_1^2 + n_2^2)^2} \right\|_{\ell^1(n_1 \sim 2^{k_1}, n_2 \sim 2^{k_2})} \leq C \frac{2^{k_2}}{8^{k_1}}.$$

Substitution of (4.23), (4.24) in (4.21) gives the bound

$$(4.20), (4.21) \leq C \sum_{k_1 \geq k_2 \geq 0} 4^{k_2 - k_1} \left(2^{k_1} |\theta_1| \wedge \frac{1}{2^{k_1} |\theta_1|} \right) \left(2^{k_2} |\theta_2| \wedge \frac{1}{2^{k_2} |\theta_2|} \right) \lesssim C \prod_{i=1}^2 \left[\sum_{k \in \mathbb{Z}_+} \left(2^k |\theta_i| \wedge \frac{1}{2^k |\theta_i|} \right) \right] \leq C.$$

This completes the proof of the Claim and of Theorem 1' for $d = 2$. □

5. PROOF OF THEOREM 1 WHEN $d = 2$ (EXPLICIT CONSTRUCTION)

Our aim is to construct $Y \in L^\infty \cap H^1$ such that

$$(5.1) \quad \operatorname{div} Y = f \in L^2_{\#}(\mathbb{T}^2).$$


Write

$$\mathbb{Z}^2 = \bigcup_{j \geq 0} (\Lambda_j^1 \cup \Lambda_j^2)$$

where

$$\Lambda_j^1 = [2^{j-1} < |n_1| \leq 2^j; |n_2| \leq 2^j]$$

$$\Lambda_j^2 = [2^j < |n_2| \leq 2^{j+1}; |n_1| \leq 2^j].$$



Let

$$\Lambda^\alpha = \bigcup_j \Lambda_j^\alpha \quad (\alpha = 1, 2).$$

Decompose

$$f = f^1 + f^2 \text{ where } f^\alpha = P_{\Lambda^\alpha} f \equiv \sum_{n \in \Lambda^\alpha} \hat{f}(n) e^{in \cdot x}.$$

Claim. Let $\delta > 0$ be small enough and $\|f\|_2 \leq \delta$. Then there are Y_1, Y_2 such that

$$(5.2) \quad \|Y_\alpha\|_{L^\infty \cap H^1} \leq 1$$

and

$$(5.3) \quad \|\partial_\alpha Y_\alpha - f^\alpha\|_2 \leq \delta^{4/3} \quad (\alpha = 1, 2).$$

Thus if $\|f\|_2 = \delta$, then

$$\|f - \partial_1 Y_1 - \partial_2 Y_2\|_2 \leq \delta^{1/3} \|f\|_2$$

and iteration of this gives (5.1).

The construction of Y_1, Y_2 is explicit but *nonlinear* (see Proposition 2).

Take $\alpha = 1$ and denote f^1 by f, Λ_j^1 by Λ_j .

Define


$$\begin{aligned} f_j &= P_{\Lambda_j} f, \\ c_j &= \|f_j\|_2, \\ F_j &= D_{x_1}^{-1} f_j \equiv \sum \frac{1}{n_1} \hat{f}_j(n) e^{in \cdot x}. \end{aligned}$$

Hence

$$(5.4) \quad \begin{aligned} \left(\sum c_j^2 \right)^{1/2} &= \|f\|_2, \\ \|F_j\|_\infty &\leq \sum_{n \in \Lambda_j} \frac{1}{|n_1|} |\hat{f}(n)| \lesssim 2^{-j} |\Lambda_j|^{1/2} \|f_j\|_2 \lesssim c_j. \end{aligned}$$

Fix $\varepsilon > 0$ a small constant and partition

$$\Lambda_j = \bigcup_{r < \frac{1}{\varepsilon} + 1} \Lambda_{j,r}$$



in stripes $\Lambda_{j,r}$ such that

$$(5.5) \quad |\operatorname{Proj}_{n_1} \Lambda_{j,r}| \sim \varepsilon 2^j.$$

Define first

$$(5.6) \quad \tilde{F}_j(x) = \sum_r \left| \sum_{n \in \Lambda_{j,r}} \frac{1}{n_1} \hat{f}_j(n) e^{in \cdot x} \right|.$$

Thus

$$(5.7) \quad |F_j(x)| \leq \tilde{F}_j(x) \lesssim c_j.$$

From Cauchy-Schwarz

$$(5.8) \quad \|\tilde{F}_j\|_2 \leq \varepsilon^{-1/2} \|F_j\|_2 \lesssim \varepsilon^{-1/2} 2^{-j} c_j.$$

Observe that if $\text{Proj}_{n_1} \Lambda_{j,r} = [a_r, b_r]$, $b_r - a_r \sim \varepsilon 2^j$, then

$$|\partial_1 \tilde{F}_j| \leq \sum_r \left| \sum_{n \in \Lambda_{j,r}} \frac{n_1 - a_r}{n_1} \hat{f}_j(n) e^{in \cdot x} \right|$$

where

$$\left| \frac{n_1 - a_r}{n_1} \right| < \varepsilon.$$

Therefore

$$(5.9) \quad \|\partial_1 \tilde{F}_j\|_2 \lesssim \sum_r \varepsilon \|P_{\Lambda_{j,r}} f\|_2 \lesssim \varepsilon^{1/2} \|P_{\Lambda_j} f\|_2 = \varepsilon^{1/2} c_j$$

(this is the purpose of the construction of \tilde{F}_j).

We also need to make an appropriate localization of the Fourier transform of \tilde{F}_j . Denote

$$K_N(y) = \sum_{|n| < N} \frac{N - |n|}{N} e^{iny},$$

the usual Féjer kernel on \mathbb{T} . It is easy to see that if

$$P(y) = \sum_{|n| < N} \hat{P}(n) e^{iny}$$

is a trigonometric polynomial, then

$$(5.10) \quad |P| \leq 3(|P| * K_N).$$

Using this fact in the variables x_1, x_2 , we see that

$$(5.11) \quad |F_j| \leq \tilde{F}_j \leq G_j$$

denoting

$$(5.12) \quad G_j = 9\tilde{F}_j * (K_{N_1} \otimes K_{N_2})$$

where each $\Delta_{j,r}$ is an $N_1 \times N_2$ rectangle, $N_1 \sim \varepsilon 2^j$, $N_2 \sim 2^j$.

Thus, by construction

$$(5.13) \quad \text{supp } \hat{G}_j \subset [-N_1, N_1] \times [-N_2, N_2] \subset \{|n| \leq 2^j\}$$

and inequalities (5.7), (5.8), (5.9) remain preserved.

Therefore,

$$(5.14) \quad \|G_j\|_\infty \leq 9\|\tilde{F}_j\|_\infty \lesssim c_j \quad (0 < \delta < 1),$$

$$(5.15) \quad \|G_j\|_2 \lesssim \varepsilon^{-1/2} 2^{-j} c_j,$$

$$(5.16) \quad \|\partial_1 G_j\|_2 \lesssim \varepsilon^{1/2} c_j,$$

$$(5.17) \quad \|\nabla G_j\|_2 \lesssim \varepsilon^{-1/2} c_j.$$

Assume that $\{f_j \mid j \leq K\}$ is a finite sequence (which is no restriction).

Define

$$\begin{aligned}
 Y_1 &= F_K + F_{K-1}(1 - G_K) \\
 &\quad + F_{K-2}(1 - G_{K-1})(1 - G_K) + \cdots \\
 (5.18) \quad &= \sum_{j \leq K} F_j \prod_{k > j} (1 - G_k).
 \end{aligned}$$

Thus from (5.11)

$$\begin{aligned}
 |Y_1| &\leq |F_K| + (1 - |F_K|)|F_{K-1}| \\
 &\quad + (1 - |F_K|)(1 - |F_{K-1}|)|F_{K-2}| + \cdots \leq 1.
 \end{aligned}$$

One may also rewrite (5.18) as

$$(5.19) \quad Y_1 = \sum F_j - \sum G_j H_j$$

with

$$\begin{aligned}
 H_j &= F_{j-1} + F_{j-2}(1 - G_{j-1}) \\
 &\quad + F_{j-3}(1 - G_{j-2})(1 - G_{j-1}) + \cdots \\
 (5.20) \quad &= \sum_{k < j} F_k \prod_{k < k' < j} (1 - G_{k'}).
 \end{aligned}$$

Clearly

$$|H_j| < 1.$$

By construction

$$(5.21) \quad \partial_1 Y_1 = \sum f_j - \sum \partial_1(G_j H_j).$$

Next, we estimate the second term in (5.21) that will appear as an error term.

Observe that since $\operatorname{supp} \hat{F}_j \subset [|n| \sim 2^j]$ and (5.13), also

$$(5.22) \quad \operatorname{supp} \hat{H}_j \subset [|n| \lesssim 2^j].$$

Denote P_k Fourier projection operators on $[|n| \sim 2^k]$ such that $Id = \sum_{k \geq 0} P_k$.

From the preceding, we may thus ensure that

$$(5.23) \quad G_j H_j = \sum_{k \leq j} P_k(G_j H_j).$$

Estimate then

$$(5.24) \quad \left\| \sum_j \partial_1(G_j H_j) \right\|_2 \leq \sum_{s \geq 0} \left(\sum_j \|\partial_1 P_{j-s}(G_j H_j)\|_2^2 \right)^{1/2}$$

(since for fixed s , the P_{j-s} have disjoint ranges).

Returning to the parameter $0 < \varepsilon < 1$ introduced earlier, write

$$(5.25) \quad \varepsilon = 2^{-s_*} \quad (s_* > 0)$$

and estimate (5.24) in the ranges

$$(5.26) \quad s > s_*$$

$$(5.27) \quad 0 \leq s \leq s_*.$$

Contribution of (5.26). Since $|H_j| \leq 1$ and (5.15),

$$(5.28) \quad \begin{aligned} \|\partial_1 P_{j-s}(G_j H_j)\|_2 &\lesssim 2^{j-s} \|G_j H_j\|_2 \\ &\leq 2^{j-s} \|G_j\|_2 \leq \varepsilon^{-1/2} 2^{-s} c_j. \end{aligned}$$

Substitution in (5.24) gives the contribution

$$(5.29) \quad \sum_{s \geq s_*} 2^{-s} \varepsilon^{-1/2} \left(\sum c_j^2 \right)^{1/2} < 2^{-s_*} \varepsilon^{-1/2} \|f\|_2 < \varepsilon^{1/2} \|f\|_2.$$

Contribution of (5.27). Estimate now

$$(5.30) \quad \begin{aligned} \|\partial_1 P_{j-s}(G_j H_j)\|_2 &\leq \|\partial_1(G_j H_j)\|_2 \leq \|\partial_1 G_j\|_2 + \|G_j \partial_1 H_j\|_2 \\ &\leq \varepsilon^{1/2} c_j + \|G_j \partial_1 H_j\|_2 \end{aligned}$$

using (5.16).

Recalling definition (5.20) of H_j , one easily verifies that

$$(5.31) \quad |\nabla H_j| \leq \sum_{k < j} (|\nabla F_k| + |\nabla G_k|).$$

Hence

$$(5.32) \quad \|\nabla H_j\|_\infty \leq \sum_{k < j} 2^k c_k$$

and from (5.15)

$$(5.33) \quad \|G_j \partial_1 H_j\|_2 \leq \varepsilon^{-1/2} c_j \left(\sum_{k < j} 2^{-(j-k)} c_k \right).$$

Substitution of (5.30), (5.33) in (5.24) gives the following bound on the contribution of (5.27):

$$(5.34) \quad \begin{aligned} &s_* \varepsilon^{1/2} \left(\sum c_j^2 \right)^{1/2} + s_* \varepsilon^{-1/2} \left[\sum_j c_j^2 \left(\sum_{k < j} 2^{-(j-k)} c_k \right)^2 \right]^{1/2} \\ &\leq \left(\log \frac{1}{\varepsilon} \right) \varepsilon^{1/2} \|f\|_2 + \left(\log \frac{1}{\varepsilon} \right) \varepsilon^{-1/2} \|f\|_2^2. \end{aligned}$$

Consequently, from (5.21), (5.29), (5.34),

$$(5.35) \quad \|f - \partial_1 Y_1\|_2 = \left\| \sum_j \partial_1(G_j H_j) \right\|_2 \leq \log \frac{1}{\varepsilon} (\varepsilon^{1/2} \|f\|_2 + \varepsilon^{-1/2} \|f\|_2^2).$$

Under the assumption $\|f\|_2 \leq \delta$, letting $\varepsilon = \delta$ in (5.35), we obtain thus

$$(5.36) \quad \|f - \partial_1 Y_1\|_2 \leq \delta^{\frac{3}{2}-} \leq \delta^{\frac{4}{3}}$$

which is (5.3).

It remains to estimate $\|Y_1\|_{H^1} = \|\nabla Y_1\|_2$.

By (5.19)

$$(5.37) \quad \|\nabla Y_1\|_2 \leq \left\| \sum_j \nabla F_j \right\|_2 + \left\| \sum \nabla(G_j H_j) \right\|_2.$$

From the definition of F_j and since $\operatorname{supp} \hat{F}_j \subset \Lambda_j^1$, it follows that

$$(5.38) \quad \left\| \sum_j \nabla F_j \right\|_2 \sim \left(\sum_j \|f_j\|_2^2 \right)^{1/2} = \|f\|_2.$$

Estimate the second term in (5.37) as in (5.24),

$$(5.39) \quad \left\| \sum_j \nabla(G_j H_j) \right\|_2 \leq \sum_{s \geq 0} \left(\sum_j \|\nabla P_{j-s}(G_j H_j)\|_2^2 \right)^{1/2}$$

and

$$(5.40) \quad \|\nabla P_{j-s}(G_j H_j)\|_2 \lesssim 2^{j-s} \|G_j H_j\|_2 \leq \varepsilon^{-1/2} 2^{-s} c_j.$$

Thus

$$(5.41) \quad (5.39) \leq \varepsilon^{-1/2} \sum_{s \geq 0} 2^{-s} \left(\sum_j c_j^2 \right)^{1/2} \leq \varepsilon^{-1/2} \|f\|_2$$

and

$$(5.42) \quad \|\nabla Y_1\|_2 \leq \delta^{-1/2} \|f\|_2 \leq \delta^{1/2}.$$

Since $\|Y_1\|_\infty \lesssim 1$, this establishes (5.2).

This proves the Claim and completes the proof of Theorem 1 for $d = 2$.

6. PROOF OF THEOREM 1 WHEN $d > 2$ (EXPLICIT CONSTRUCTION)

Let $f \in L_{\#}^d(\mathbb{T}^d)$. Our aim is to construct a solution Y of $\operatorname{div} Y = f$ satisfying

$$(6.1) \quad \|Y\|_\infty \leq C \|f\|_d,$$

$$(6.2) \quad \|\nabla Y\|_d \leq C \|f\|_d.$$

We do this by standard modification of the previous L^2 -argument with the Littlewood-Paley square function theory as main additional ingredient. Consider again a partition

$$\mathbb{Z}^d = \bigcup_{j \geq 0} (\Lambda_j^1 \cup \dots \cup \Lambda_j^d)$$

of disjoint d -rectangles Λ_j^α of side length $\sim 2^j$.

We formulate the analogue of the Claim with Y_α satisfying bounds (6.1), (6.2). Letting $\alpha = 1$, $f = f^1$, define again

$$(6.3) \quad F_j = D_{x_1}^{-1} f_j$$

satisfying

$$(6.4) \quad \|F_j\|_\infty \lesssim (2^{j/d})^d \|F_j\|_d = 2^j \|D_{x_1}^{-1} f_j\|_d \sim \|f_j\|_d \equiv c_j.$$

Define \tilde{F}_j and G_j as in (5.6), (5.12). Thus (5.11), (5.13) hold. Also

$$\begin{aligned}
\|G_j\|_\infty &\lesssim \|\tilde{F}_j\|_\infty \leq \varepsilon^{-1/d'} \left(\sum_{r < \frac{1}{\varepsilon}} \left\| \sum_{n \in \Lambda_{j,r}} \frac{1}{n_1} \hat{f}_j(n) e^{inx} \right\|_\infty^d \right)^{1/d} \\
&\leq \varepsilon^{-1/d'} \left(\sum_{r < \frac{1}{\varepsilon}} \left(2^{j \frac{d-1}{d}} (\varepsilon 2^j)^{\frac{1}{d}} \left\| \sum_{n \in \Lambda_{j,r}} \frac{1}{n_1} \hat{f}_j(n) e^{inx} \right\|_d^d \right)^{1/d} \right)^{1/d} \\
&\lesssim \varepsilon^{-1/d'+1/d} \left(\sum_{r < \frac{1}{\varepsilon}} \left\| \sum_{n \in \Lambda_{j,r}} \hat{f}_j(n) e^{inx} \right\|_d^d \right)^{\frac{1}{d}} \\
(6.5) \quad &\lesssim \varepsilon^{\frac{2}{d}-1} \|f_j\|_d = \varepsilon^{\frac{2}{d}-1} c_j \leq \varepsilon^{\frac{2}{d}-1} \delta.
\end{aligned}$$

(We assume that δ is small enough compared with ε to ensure, in particular, that $\varepsilon^{\frac{2}{d}-1} \delta \ll 1$.)

Repeat the construction from Section 5. In place of estimate (5.24) we now have

$$(6.6) \quad \left\| \sum_j \partial_1(G_j H_j) \right\|_d \leq \sum_{s \geq 0} \left\| \sum_j |\partial_1 P_{j-s}(G_j H_j)|^2 \right\|_d^{1/2}$$

and distinguish between the cases (5.26), (5.27).

Contribution of (5.26). Estimate

$$\begin{aligned}
&\left\| \left(\sum_j |\nabla P_{j-s}(G_j H_j)|^2 \right)^{1/2} \right\|_d \\
&\lesssim \left\| \left(\sum_j 4^{j-s} |P_{j-s}(G_j H_j)|^2 \right)^{1/2} \right\|_d \\
&\lesssim 2^{-s} \left\| \left(\sum_j 4^j |G_j H_j|^2 \right)^{1/2} \right\|_d \\
(6.7) \quad &\lesssim 2^{-s} \left\| \left(\sum_j 4^j (\tilde{F}_j * K_j)^2 \right)^{1/2} \right\|_d
\end{aligned}$$

where K_j is a product of Féjer kernels

$$K_{N_1} \otimes K_{N_2} \otimes \cdots \otimes K_{N_d}, \quad N_1 \sim \varepsilon 2^j, \quad \text{and } N_2, \dots, N_d \sim 2^j.$$

Again from standard square function inequalities

$$(6.8) \quad (6.7) \lesssim 2^{-s} \left\| \left(\sum_j 4^j (\tilde{F}_j)^2 \right)^{1/2} \right\|_d.$$

Recalling the definition of \tilde{F}_j , estimate

$$(6.9) \quad (\tilde{F}_j)^2 \leq \varepsilon^{-1} \sum_{r \leq \varepsilon^{-1}} \left| \sum_{n \in \Lambda_{j,r}^1} \frac{1}{n_1} \hat{f}(n) e^{inx} \right|^2.$$

Substituting in (6.8), this gives

$$\begin{aligned}
 (6.10) \quad & \varepsilon^{-1/2} 2^{-s} \left\| \left(\sum_j \sum_{r < \varepsilon^{-1}} \left| \sum_{n \in \Lambda_{j,r}^1} \frac{2^j}{n_1} \hat{f}(n) e^{inx} \right|^2 \right)^{1/2} \right\|_d \\
 & \lesssim \varepsilon^{-1/2} 2^{-s} \left\| \left(\sum_j \sum_{r < \varepsilon^{-1}} \left| \sum_{n \in \Lambda_{j,r}^1} \hat{f}(n) e^{inx} \right|^2 \right)^{1/2} \right\|_d.
 \end{aligned}$$

We use here the fact that $|n_1| \sim |n| \sim 2^j$ for $n \in \Lambda_j^1$.

Recall also the definition of $\Lambda_{j,r}$ obtained by partitioning the n_1 -variable in intervals of size $\varepsilon 2^j$.

At this stage, we use the following (1-variable) inequality due to Rubio de Francia [19], which generalizes the Littlewood-Paley inequality to arbitrary intervals.

Proposition 3. *Let $\{I_\alpha\}$ be disjoint intervals in \mathbb{Z} and*

$$P_I f = \sum_{n \in I} \hat{f}(n) e^{inx}$$

the corresponding Fourier projection.

Then, for $2 \leq d < \infty$, there is the (one-sided) inequality

$$(6.11) \quad \left\| \left(\sum |P_{I_\alpha} f|^2 \right)^{1/2} \right\|_d \leq C \|f\|_d.$$

Since $\{\operatorname{Proj}_{n_1} \Lambda_{j,r}^1\}$ are disjoint intervals in \mathbb{Z} , application of (6.11) in the x_1 -variable implies that

$$(6.12) \quad (6.6) \lesssim \varepsilon^{-1/2} 2^{-s} \|f\|_d.$$

Summation of (6.12) for $s \geq s_*$ gives then

$$(6.13) \quad (5.26)\text{-contribution} \leq \varepsilon^{1/2} \|f\|_d.$$

Remark 8. We used the general Proposition 3 for convenience; the present case could in fact be treated by more elementary means.

Contribution of (5.27). Estimate

$$\begin{aligned}
 & \left\| \left(\sum_j |\partial_1 P_{j-s}(G_j H_j)|^2 \right)^{1/2} \right\|_d \lesssim \left\| \left(\sum_j |\partial_1(G_j H_j)|^2 \right)^{1/2} \right\|_d \\
 & \leq \left\| \left(\sum_j |\partial_1 G_j|^2 \right)^{1/2} \right\|_d + \left\| \left(\sum_j |G_j(\partial_1 H_j)|^2 \right)^{1/2} \right\|_d = (6.14) + (6.15).
 \end{aligned}$$

Estimate (6.14) by

$$(6.16) \quad \left\| \left(\sum_j |\partial_1 \tilde{F}_j|^2 \right)^{1/2} \right\|_d.$$

We have that

$$\begin{aligned} |\partial_1 \tilde{F}_j| &\leq \sum_{r < \varepsilon^{-1}} \left| \sum_{n \in \Lambda_{j,r}^1} \frac{n_1 - a_{j,r}}{n_1} \hat{f}(n) e^{inx} \right| \\ &\leq \varepsilon^{-1/2} \left(\sum_{r < \varepsilon^{-1}} \left| \sum_{n \in \Lambda_{j,r}^1} \frac{n_1 - a_{j,r}}{n_1} \hat{f}(n) e^{inx} \right|^2 \right)^{1/2} \end{aligned}$$

where $\text{Proj}_{n_1} \Lambda_{j,r}^1 = [a_{j,r}, b_{j,r}]$, $b_{j,r} - a_{j,r} \sim \varepsilon 2^j$. Thus $|\frac{n_1 - a_{j,r}}{n_1}| \leq \varepsilon$.

We get therefore

$$\begin{aligned} (6.16) &\leq \varepsilon^{-1/2} \cdot \varepsilon \left\| \left(\sum_j \sum_{r < \varepsilon^{-1}} \left| \sum_{n \in \Lambda_{j,r}^1} \hat{f}(n) e^{inx} \right|^2 \right)^{1/2} \right\|_d \\ (6.17) &\lesssim \varepsilon^{1/2} \|f\|_d. \end{aligned}$$

To estimate (6.15), use again inequality (5.31), together with (6.4), (6.5). Thus

$$(6.18) \quad \|\nabla H_j\|_\infty \leq \varepsilon^{\frac{2}{d}-1} \sum_{k < j} 2^k c_k < \varepsilon^{\frac{2}{d}-1} 2^j \|f\|_d.$$

Hence

$$\begin{aligned} (6.15) &\leq \varepsilon^{\frac{2}{d}-1} \|f\|_d \left\| \left(\sum_j 4^j G_j^2 \right)^{1/2} \right\|_d \\ &\leq \varepsilon^{\frac{2}{d}-1} \|f\|_d \left\| \left(\sum_j (2^j \tilde{F}_j)^2 \right)^{1/2} \right\|_d \\ (6.19) &\leq \varepsilon^{\frac{2}{d}-\frac{3}{2}} \|f\|_d^2 \end{aligned}$$

applying again the (6.8)-bound using Proposition 3.

Thus the (5.27)-contribution is

$$(6.20) \quad \leq \varepsilon^{1/2} \log \frac{1}{\varepsilon} \|f\|_d + \varepsilon^{\frac{2}{d}-\frac{3}{2}} \log \frac{1}{\varepsilon} \|f\|_d^2.$$

Collecting estimates (6.13), (6.20), it follows that

$$\begin{aligned} \|f - \partial_1 Y\|_d &= \left\| \sum_j \partial_1 (G_j H_j) \right\|_d \\ (6.21) &\leq \varepsilon^{1/2} \log \frac{1}{\varepsilon} \|f\|_d + \varepsilon^{\frac{2}{d}-\frac{3}{2}} \log \frac{1}{\varepsilon} \|f\|_d^2 \end{aligned}$$

which is the analogue of (5.35). Assuming $\|f\|_d = \delta$, take $\varepsilon = \delta^{1/2}$ to obtain

$$(6.22) \quad \|f - \partial_1 Y\|_d \leq \delta^{1/5} \|f\|_d.$$

It remains to estimate

$$\|\nabla Y\|_d \leq \left\| \sum \nabla F_j \right\|_d + \left\| \sum \nabla (G_j H_j) \right\|_d = (6.23) + (6.24).$$

We have

$$(6.23) \sim \left\| \left(\sum |\nabla F_j|^2 \right)^{1/2} \right\|_d \sim \left\| \left(\sum |f_j|^2 \right)^{1/2} \right\|_d \lesssim \|f\|_d.$$

Estimate (6.24) as

$$(6.25) \quad \left\| \sum_{s \geq 0} \left(\sum_j |\nabla P_{j-s}(G_j H_j)|^2 \right)^{1/2} \right\|_d \lesssim \varepsilon^{-1/2} \|f\|_d$$

using (6.7)–(6.12).

This completes the argument.

We conclude this section with a

Proof of Theorem 1' when $d > 2$. The argument is somewhat bizarre: one uses duality twice! First, from Theorem 1 we easily deduce the estimate on \mathbb{T}^d

$$(6.26) \quad \|u - fu\|_{L^{d/(d-1)}} \leq C(d) \|\operatorname{grad} u\|_{L^1 + W^{-1, d/(d-1)}}, \forall u \in L^{d/(d-1)}.$$

Next, we argue as in the beginning of Section 4. Observe that

$$L^1 + W^{-1, d/(d-1)} \subset \mathcal{M} + H^{-1}$$

and that

$$(6.27) \quad \|\cdots\|_{L^1 + W^{-1, d/(d-1)}} = \|\cdots\|_{\mathcal{M} + W^{-1, d/(d-1)}} \text{ on } L^1 + W^{-1, d/(d-1)}$$

(this may be easily seen using regularization by convolution).

Let $E = C^0 \cap W^{1, d}$, $F = L^d_{\#}$ and consider the bounded operator $T : E \rightarrow F$ defined by $TY = \operatorname{div} Y$. Clearly $T^* : F^* \rightarrow E^* = \mathcal{M} + W^{-d, d/(d-1)}$ is given by $T^*u = \operatorname{grad} u$. By (6.26) and (6.27) we obtain

$$\|u\|_{F^*} \leq C \|T^*u\|_{E^*} \quad \forall u \in F^*$$

and therefore T is surjective from E onto F . Applying the open mapping principle (or use Hahn-Banach as in the proof of Proposition 1), we see that for every $f \in F$ there is some $Y \in E$ satisfying $TY = f$ and $\|Y\|_E \leq C \|f\|_F$. \square

Remark 9. Alternatively, one may approximate $f \in L^d_{\#}(\mathbb{T}^d)$ by trigonometric polynomials. If f is a trigonometric polynomial, we may clearly obtain Y as a trigonometric polynomial (after convolution). A standard limit procedure permits then to complete the argument.

7. THE EQUATION $\operatorname{div} Y = f$ WITH DIRICHLET CONDITION. PROOF OF THEOREMS 2 AND 3

So far we have studied problem (1.1) coupled with a periodic condition. We consider here problem (1.1) coupled with a Dirichlet condition. Usually one associates with (1.1) the “partial” Dirichlet condition

$$(7.1) \quad Y \cdot n = 0 \quad \text{on } \partial Q$$

(n is normal to ∂Q). It is quite standard that for every $f \in L^p_{\#}$, $1 < p < \infty$, there is some $Y \in W^{1, p}$ satisfying (1.1), (7.1) and

$$\|Y\|_{W^{1, p}} \leq C \|f\|_{L^p}.$$

Indeed, one may look for a *special* Y of the form $Y = \operatorname{grad} u$ and one is led to the Neumann problem

$$(7.2) \quad \begin{cases} \Delta u = f & \text{in } Q, \\ \frac{\partial u}{\partial n} = 0 & \text{on } \partial Q, \end{cases}$$

which admits a solution $u \in W^{2,p}$ such that

$$\|u\|_{W^{2,p}} \leq C\|f\|_{L^p}.$$

It is also possible to couple problem (1.1) with the *full* Dirichlet condition

$$(7.3) \quad Y = 0 \quad \text{on } \partial Q.$$

For simplicity we investigate first the case where the domain is a cube and then the case of a Lipschitz bounded domain.

7.1. The case of a cube. Let $Q = (0, 1)^d$. Here is the first result:

Theorem 2. *Given $f \in L^p_{\#}(Q)$, $1 < p < \infty$, there exists some $Y \in W_0^{1,p}(Q)$ solving (1.1) with*

$$\|Y\|_{W^{1,p}} \leq C(p, d)\|f\|_{L^p},$$

where we use the standard notation

$$W_0^{1,p}(Q) = \{Y \in W^{1,p}(Q); Y = 0 \text{ on } \partial Q\}.$$

Moreover Y can be chosen, depending linearly on f .

We will make use of the following lemma (which is a special case of Theorem 2).

Lemma 4. *Given $f \in W_0^{1,p}(Q)$, $1 < p < \infty$, with $\int f = 0$, there exists $Y \in W_0^{1,p}(Q)$, such that*

$$\operatorname{div} Y = f$$

and

$$(7.4) \quad \|Y\|_{W^{1,p}(Q)} \leq C(d)\|f\|_{W^{1,p}(Q)}.$$

Moreover Y can be chosen, depending linearly on f .

Proof. Following a known construction (see Adams [1], p. 58 and Nirenberg [15]), we construct Y by induction on the dimension d . The assertion is obvious for $d = 1$. Assume that it holds in dimension $(d - 1)$. Let $f \in W_0^{1,p}(Q_d)$, where $Q_d = (0, 1)^d$, with $\int_{Q_d} f = 0$.

Set

$$g(x') = \int_0^1 f(x', t) dt, \quad \text{where } x' = (x_1, \dots, x_{d-1}) \in Q_{d-1}.$$

Clearly, $g \in W_0^{1,p}(Q_{d-1})$ with

$$\|g\|_{W^{1,p}(Q_{d-1})} \leq C\|f\|_{W^{1,p}(Q_d)}$$

and also $\int_{Q_{d-1}} g = 0$. By the induction assumption there is some $Z \in W_0^{1,p}(Q_{d-1})$ such that

$$(7.5) \quad \operatorname{div}_{x'} Z = g \quad \text{on } Q_{d-1}$$

and

$$\|Z\|_{W^{1,p}(Q_{d-1})} \leq C\|g\|_{W^{1,p}(Q_{d-1})} \leq C\|f\|_{W^{1,p}(Q_d)}.$$

Fix a function $\zeta \in C_0^\infty(0, 1)$ such that

$$(7.6) \quad \int_0^1 \zeta(t) dt = 1.$$

For $x = (x', x_d) \in Q_d$ set

$$h(x) = \int_0^{x_d} (f(x', t) - \zeta(t)g(x')) dt.$$

It is easy to see (using (7.6)) that $h \in W_0^{1,p}(Q_d)$ and

$$\|h\|_{W^{1,p}(Q_d)} \leq C\|f\|_{W^{1,p}(Q_d)}.$$

Moreover

$$\frac{\partial h}{\partial x_d}(x) = f(x) - \zeta(x_d)g(x').$$

Combining this with (7.5) yields

$$f(x) = \operatorname{div}_{x'}(\zeta(x_d)Z(x')) + \frac{\partial h}{\partial x_d}$$

i.e., the conclusion holds with

$$Y(x) = (\zeta(x_d)Z(x'), h(x)).$$

□

Proof of Theorem 2. For simplicity we assume that $d = 2$; the argument is similar for $d > 2$.

Let

$$Q = \{(x, y) \in \mathbb{R}^2; \quad 0 < x < 1, 0 < y < 1\}.$$

Given $f \in L_{\#}^p(Q)$, $1 < p < \infty$, we will construct a solution $Y \in W_0^{1,p}(Q)$ of (1.1); moreover

$$(7.7) \quad \|Y\|_{W^{1,p}} \leq C_p\|f\|_{L^p}$$

and Y depends linearly on f . This is done in three steps. □

Step 1. Construct a solution $Y \in W^{1,p}(Q)$ of (1.1) satisfying (7.7) and

$$(7.8) \quad Y = 0 \quad \text{on the edge } \{(x, 0); 0 < x < 1\}.$$

Proof. Set

$$\tilde{Q} = \{(x, y); 0 < x < 1, -2 < y < 1\}$$

and

$$(7.9) \quad \tilde{f} = \begin{cases} f & \text{in } Q, \\ 0 & \text{in } \tilde{Q} \setminus Q. \end{cases}$$

Let $Z \in W^{1,p}(\tilde{Q})$ be the solution of

$$(7.10) \quad \operatorname{div} Z = \tilde{f} \quad \text{in } \tilde{Q}$$

obtained via (7.2) (or via periodic conditions on \tilde{Q}).

The heart of the matter is the following construction. Write $Z = (Z_1, Z_2)$ and define $Y = (Y_1, Y_2)$ in Q , where

$$(7.11) \quad \begin{aligned} Y_1(x, y) &= Z_1(x, y) + 3Z_1(x, -y) - 4Z_1(x, -2y), \\ Y_2(x, y) &= Z_2(x, y) - 3Z_2(x, -y) + 2Z_2(x, -2y). \end{aligned}$$

(This type of “reflection” is reminiscent of standard extension techniques in $W^{m,p}$, $m \geq 2$; see e.g. Adams [1]).

It is easy to see using (7.9), (7.10) and (7.11) that

$$\operatorname{div} Y = f \quad \text{in } Q$$

while (7.8) is clear from the definition of Y .

It is important (for the next step) to observe that if we had started with the additional information

$$Z = 0 \quad \text{on the edge } \{(0, y); -2 < y < 1\} \text{ of } \tilde{Q},$$

then we could infer that Y also vanishes on the edge $\{(0, y); 0 < y < 1\}$ of Q . \square

Step 2. Construct a solution $Y \in W^{1,p}(Q)$ of (1.1) satisfying (7.7) and (7.12)

$$Y = 0 \text{ on the 2 adjacent edges } \{(x, 0); 0 < x < 1\} \text{ and } \{(0, y); 0 < y < 1\}.$$

Proof. Set

$$\hat{Q} = \{(x, y); -2 < x < 1, 0 < y < 1\}$$

and

$$\hat{f} = \begin{cases} f & \text{in } Q, \\ 0 & \text{in } \hat{Q} \setminus Q. \end{cases}$$

From Step 1 applied to \hat{f} in \hat{Q} we obtain a solution \hat{Z} of

$$\operatorname{div} \hat{Z} = \hat{f} \quad \text{in } \hat{Q}$$

such that

$$\hat{Z} = 0 \quad \text{on the edge } \{(x, 0); -2 < x < 1\} \text{ of } \hat{Q}.$$

Starting with \hat{Z} (instead of Z) we repeat the construction of Step 1 changing the roles of x and y . We thus obtain a $Y \in W^{1,p}(Q)$ satisfying (1.1) in Q , (7.7) and (7.12). \square

Step 3. Proof of Theorem 2 completed.

Consider a smooth partition of unity $(\theta_i), i = 1, 2, 3, 4$, subordinate to the covering of Q consisting of the 4 discs of radius 1 centered at the 4 vertices. Let $Y_i \in W^{1,p}(Q)$ be the solution constructed in Step 2 relative to each vertex.

Set

$$Z = \sum_{i=1}^4 \theta_i Y_i.$$

It is easy to see from this construction that $\theta_i Y_i \in W_0^{1,p}(Q), \forall i$ and thus $Z \in W_0^{1,p}(Q)$. Moreover

$$\operatorname{div} Z = f + \sum_i \nabla \theta_i \cdot Y_i$$

and $\sum_i \nabla \theta_i \cdot Y_i \in W_0^{1,p}(Q)$. By Lemma 4 we may construct $X \in W_0^{1,p}(Q)$ satisfying

$$\operatorname{div} X = \sum_i \nabla \theta_i \cdot Y_i$$

and $Y = Z - X$ has all the desired properties in Theorem 2.

Next we have a variant of Theorem 1' for the full Dirichlet condition.

Theorem 3. Given $f \in L^d_{\#}(Q)$ there exists some $Y \in C^0(\bar{Q}) \cap W_0^{1,d}(Q)$ satisfying (1.1) with

$$\|Y\|_{L^\infty} + \|Y\|_{W^{1,d}} \leq C \|f\|_{L^d}.$$

Remark 10. Clearly, Theorem 3 implies Theorem 1' since the function Y extended by periodicity belongs to $C^0(\mathbb{T}^d) \cap W^{1,d}(\mathbb{T}^d)$ and satisfies (1.1) on \mathbb{T}^d . However its proof relies heavily on Theorem 1'.

Proof of Theorem 3. Follow the same strategy as in the proof of Theorem 2. The only difference is that in Step 1 use Theorem 1' to obtain Z (instead of taking the special Z in the form of a gradient). Of course the dependence of Y on f is not linear anymore.

In Step 3 rely on the following variant of Lemma 4 (with an identical proof). \square

Lemma 4'. *Given $f \in C^0(\bar{Q}) \cap W_0^{1,p}(Q), 1 < p < \infty$, with $\int f = 0$, there exists $Y \in C^0(\bar{Q}) \cap W_0^{1,p}(Q)$ such that*

$$\operatorname{div} Y = f$$

and

$$\|Y\|_{L^\infty} + \|Y\|_{W^{1,p}} \leq C(\|f\|_{L^\infty} + \|f\|_{W^{1,p}}).$$

7.2. The case of Lipschitz domains. Let Ω be a Lipschitz, connected, bounded domain in \mathbb{R}^d . Recall that Ω is Lipschitz if there is a $\delta > 0$ such that for every point $p \in \partial\Omega$, $\partial\Omega \cap B_\delta(p)$ is the graph of a Lipschitz function (in an appropriate coordinate system varying with p).

We have the following variants of Theorems 2 and 3.

Theorem 2'. *Given any $f \in L_{\#}^p(\Omega), 1 < p < \infty$, there exists some $Y \in W_0^{1,p}(\Omega)$ solving (1.1) with*

$$(7.13) \quad \|Y\|_{W^{1,p}} \leq C(p, \Omega) \|f\|_{L^p}.$$

Moreover Y can be chosen, depending linearly on f .

Theorem 3'. *For every $f \in L_{\#}^d(\Omega)$ there exists some $Y \in C^0(\bar{\Omega}) \cap W_0^{1,d}(\Omega)$ solving (1.1) with*

$$(7.14) \quad \|Y\|_{L^\infty} + \|Y\|_{W^{1,d}} \leq C(p, \Omega) \|f\|_{L^d}.$$

The heart of the argument (for both theorems) is the following.

Lemma 5. *There is a bounded operator $S : L^p(\Omega) \rightarrow W_0^{1,p}(\Omega)$ such that*

$$f - \operatorname{div} Sf \in W_0^{1,p} \quad \forall f \in L^p$$

and

$$(7.15) \quad \|f - \operatorname{div} Sf\|_{W^{1,p}} \leq C \|f\|_{L^p}.$$

The variant needed for the proof of Theorem 3' is

Lemma 5'. *There is a nonlinear map $S : L^d(\Omega) \rightarrow C^0(\bar{\Omega}) \cap W_0^{1,d}(\Omega)$ such that*

$$(7.16) \quad \|Sf\|_{L^\infty} + \|Sf\|_{W^{1,d}} \leq C \|f\|_{L^d}$$

and

$$(7.17) \quad \|f - \operatorname{div} Sf\|_{W^{1,d}} \leq C \|f\|_{L^d}.$$

The proof of Lemma 5 relies on the following construction. Let Q' be a cube of side δ in \mathbb{R}^{d-1} and set

$$U = \{(x', y) \in Q' \times \mathbb{R}; \psi(x') < y < \psi(x') + \delta\}$$

where $\psi \in \operatorname{Lip}(Q')$.

Lemma 6. *Assume*

$$(7.18) \quad \|\nabla\psi\|_{L^\infty(Q')} \leq \varepsilon_0(d) \text{ sufficiently small (depending only on } d).$$

Then, given any } g \in L^p(U) \text{ there is some } Z \in W^{1,p}(U) \text{ satisfying}

$$(7.19) \quad \operatorname{div} Z = g \quad \text{in } U,$$

$$(7.20) \quad Z = 0 \text{ on } \{y = \psi(x'); x' \in Q'\} \text{ and on the lateral boundary of } U, \\ \text{with}$$

$$\|Z\|_{W^{1,p}(U)} \leq C(p, d)\|g\|_{L^p(U)}.$$

Moreover } Z \text{ can be chosen to depend linearly on } g.

Proof. For $x' \in Q'$ and $0 < y < \delta$ set

$$\tilde{g}(x', y) = g(x', y + \psi(x')).$$

Note that

$$\|\tilde{g}\|_{L^p(Q)} = \|g\|_{L^p(U)}$$

where $Q = Q' \times (0, \delta)$.

By Theorem 2 there exists $\tilde{Z} \in W^{1,p}(Q)$ such that

$$\begin{cases} \operatorname{div} \tilde{Z} = \tilde{g} & \text{in } Q, \\ \tilde{Z} = 0 & \text{on } \{(x', 0); x' \in Q'\} \cup (\partial Q' \times (0, \delta)) \end{cases}$$

with

$$(7.21) \quad \|\tilde{Z}\|_{W^{1,p}(Q)} \leq C(d)\|\tilde{g}\|_{L^p(Q)}.$$

Note that here $\int \tilde{g} = 0$ is not required since we may consider in $\hat{Q} = Q' \times (0, 2\delta)$ the function

$$\hat{g}(x', y) = \begin{cases} \tilde{g}(x', y) & \text{for } x' \in Q' \text{ and } 0 < y < \delta, \\ -\tilde{g}(x', y - \delta) & \text{for } x' \in Q' \text{ and } \delta < y < 2\delta, \end{cases}$$

and then solve (using Theorem 2)

$$\begin{aligned} \operatorname{div} \hat{Z} &= \hat{g} & \text{in } \hat{Q}, \\ \hat{Z} &= 0 & \text{on } \partial\hat{Q}, \end{aligned}$$

with

$$\|\hat{Z}\|_{W^{1,p}(\hat{Q})} \leq C(d)\|\hat{g}\|_{L^p(\hat{Q})}.$$

The restriction \tilde{Z} of \hat{Z} to $Q' \times (0, \delta)$ satisfies the desired properties.

Also, it is clear by scaling that the constant in (7.21) is independent of δ .

Returning to $(x', y) \in U$, set

$$Z(x', y) = \tilde{Z}(x', y - \psi(x'));$$

it is easy to see, using (7.18) and (7.21), that

$$\|\operatorname{div} Z - g\|_{L^p(U)} \leq C(d)\varepsilon_0\|g\|_{L^p(U)}$$

and

$$\|Z\|_{W^{1,p}(U)} \leq C(d)(1 + \varepsilon_0)\|g\|_{L^p(U)}.$$

Choosing ε_0 such that $C(d)\varepsilon_0 < 1$ and iterating this construction yields the lemma. \square

The variant necessary for Theorem 3' is

Lemma 6'. *Assume (7.18). Then given $g \in L^d(U)$ there is some $Z \in C^0(\bar{U}) \cap W^{1,p}(U)$ satisfying (7.19), (7.20) and*

$$\|Z\|_{L^\infty(U)} + \|Z\|_{W^{1,d}(U)} \leq C(d)\|g\|_{L^d(U)}.$$

Next, we remove the smallness condition (7.18) on the Lipschitz constant of ψ .

Lemma 7. *With the same notation as in Lemma 6, assume only that $\psi \in \operatorname{Lip}(Q')$. Then, given any $g \in L^p(U)$, there is some $Z \in W^{1,p}(U)$ satisfying (7.19), (7.20) and*

$$\|Z\|_{W^{1,p}(U)} \leq C(p, d, \|\nabla\psi\|_{L^\infty(Q')})\|g\|_{L^p(U)}.$$

Moreover Z can be chosen to depend linearly on g .

Proof. Consider the dilation $x' \mapsto \tilde{x}' = Nx'$ (only in x' , not in the full x -variable). Set $\tilde{Q}' = NQ'$ and define on \tilde{Q}' the function

$$\tilde{\psi}(\tilde{x}') = \psi(\tilde{x}'/N).$$

Fix an integer N sufficiently large so that

$$\|\nabla\tilde{\psi}\|_{L^\infty(\tilde{Q}')} = \frac{1}{N}\|\nabla\psi\|_{L^\infty(Q')} \leq \varepsilon_0(d)$$

where $\varepsilon_0(d)$ comes from (7.18).

Set

$$\tilde{g}(\tilde{x}', y) = g\left(\frac{\tilde{x}'}{N}, y\right).$$

Divide the cube \tilde{Q}' (of side $N\delta$) into N^{d-1} cubes of side δ and apply, in each of them, Lemma 6 to $\tilde{\psi}$ and \tilde{g} . By gluing the corresponding solutions (this is possible because all these solutions vanish on the lateral boundaries of their domains), we obtain some $\tilde{Z}(\tilde{x}', y) \in W^{1,p}(\tilde{U})$ satisfying

$$\begin{cases} \operatorname{div}_{\tilde{x}', y} \tilde{Z} = \tilde{g} & \text{in } \tilde{U} = \{(\tilde{x}', y) \in \tilde{Q}' \times \mathbb{R}; \tilde{\psi}(\tilde{x}') < y < \tilde{\psi}(\tilde{x}') + \delta\}, \\ \tilde{Z} = 0 & \text{on } \{y = \tilde{\psi}(\tilde{x}'); \tilde{x}' \in \tilde{Q}'\}, \end{cases}$$

and the corresponding $W^{1,p}$ -estimate for \tilde{Z} .

We now return to the variables $(x', y) \in U$. Write the components of \tilde{Z} as

$$\tilde{Z} = (\tilde{Z}', \tilde{Z}_d)$$

and set

$$Z(x', y) = \left(\frac{1}{N} \tilde{Z}'(Nx', y), \tilde{Z}_d(Nx', y) \right).$$

It is easy to check that Z satisfies all the required properties. \square

The variant necessary for Theorem 3' is

Lemma 7'. *With the same notation as in Lemma 6, assume only that $\psi \in \operatorname{Lip}(Q')$. Then, given any $g \in L^d(U)$, there is some $Z \in C^0(\bar{U}) \cap W^{1,p}(U)$ satisfying (7.19), (7.20) and*

$$\|Z\|_{L^\infty(U)} + \|Z\|_{W^{1,p}(U)} \leq C(d, \|\nabla\psi\|_{L^\infty(Q')})\|g\|_{L^p(U)}.$$

We now return to the

Proof of Lemma 5. Consider a finite covering of $\partial\Omega$ by a collection of cubes $Q_i, i = 1, \dots, k$, of side δ such that in each $Q_i, \partial\Omega \cap Q_i$ admits a Lipschitz parametrization ψ_i . To this covering we associate functions $\theta_0, \theta_1, \dots, \theta_k$ such that

$$\theta_0 + \sum_{i=1}^k \theta_i = 1 \quad \text{on } \Omega,$$

$$\theta_0 \in C_0^\infty(\Omega) \text{ and } \theta_i \in C_0^\infty(Q_i) \text{ for } i = 1, \dots, k.$$

Given $g \in L^p(\Omega)$ solve, using Lemma 7, for $i = 1, 2, \dots, k$,

$$\begin{cases} \operatorname{div} Z_i = g & \text{in } U_i, \\ Z_i = 0 & \text{on } \partial\Omega \cap Q_i. \end{cases}$$

Next solve

$$\operatorname{div} Z_0 = g \quad \text{in } \Omega,$$

for example $Z_0 = \operatorname{grad}(\Delta)^{-1}$ where Δ^{-1} is used with zero Dirichlet condition on $\partial\Omega$.

Note that

$$Z = \sum_{i=0}^k \theta_i Z_i \in W_0^{1,p}$$

and

$$\operatorname{div} Z = g + \sum_{i=0}^k \nabla\theta_i \cdot Z_i.$$

All the conclusions of Lemma 5 hold with

$$Sg = Z.$$

□

Proof of Lemma 5'. We make the same construction as above, using Lemma 7' in place of Lemma 7 and Theorem 2 to solve $\operatorname{div} Z_0 = g$ in any large cube containing Ω . □

Theorem 2' is an immediate consequence of Lemma 5 and the following general functional analysis argument applied with $E = W_0^{1,p}, F = L_{\#}^p$ and $T = \operatorname{div}$. (Note that $T^* = \operatorname{grad}$ is injective on $F^* = L_{\#}^q$, since Ω is connected.)

Lemma 8. *Let E, F be two Banach spaces and let T be a bounded operator from E into F . Assume*

$$(7.22) \quad N(T^*) = \{0\}.$$

$$(7.23) \quad \begin{cases} \text{There is a bounded operator } S \text{ from } F \text{ into } E \text{ and} \\ \text{a compact operator } K \text{ from } F \text{ into itself such that} \\ T \circ S = I + K. \end{cases}$$

Then T admits a right inverse.

Proof. First we note that T is onto. Indeed, in view of (7.22) it suffices to show that T (or equivalently T^*) has closed range. This is an obvious consequence of the inequality

$$\|f\| \leq C\|T^*f\| + \|K^*f\| \quad \forall f \in F^*$$

(which follows from (7.23)).

Next, let X be a complementing subspace for $N(I + K)$ in F and set $Y = R(I + K)$. Since $u = (I + K)|_X$ is an isomorphism onto Y , its inverse $u^{-1} : Y \rightarrow X \subset F$ satisfies

$$(7.24) \quad (I + K) \circ u^{-1} = I \text{ on } Y.$$

Let Q be a projector from F onto Y ; since $R(I - Q)$ is finite dimensional, we may choose a base (e_α) of $R(I - Q)$ and write

$$(7.25) \quad f = Qf + \sum_{\alpha} \langle e_{\alpha}^*, f \rangle e_{\alpha} \quad \forall f \in F,$$

for some e_{α}^* 's in F^* .

Since we showed that T is onto, one has, for each α , some $\bar{e}_{\alpha} \in E$ satisfying

$$(7.26) \quad T\bar{e}_{\alpha} = e_{\alpha} \quad \forall \alpha.$$

Consider the operator $S_1 : F \rightarrow E$ defined for every $f \in F$, by

$$S_1 f = S \circ u^{-1} \circ Qf + \sum_{\alpha} \langle e_{\alpha}^*, f \rangle \bar{e}_{\alpha}.$$

Using (7.24), (7.25) and (7.26) we see that

$$\begin{aligned} T \circ S_1 f &= (I + K) \circ u^{-1} \circ Qf + \sum_{\alpha} \langle e_{\alpha}^*, f \rangle e_{\alpha} \\ &= Qf + \sum_{\alpha} \langle e_{\alpha}^*, f \rangle e_{\alpha} = f \end{aligned}$$

for every $f \in F$. Thus S_1 is a right inverse for T . \square

Proof of Theorem 3'. Given $f \in L^d$ write, using Lemma 5',

$$f = \operatorname{div} Y_1 + R$$

with $Y_1 \in C^0(\bar{\Omega}) \cap W_0^{1,d}(\Omega)$ and $R \in W_0^{1,d}(\Omega)$ (and the corresponding estimates).

If $\int f = 0$, then $\int R = 0$ and we may apply Theorem 2' in any L^p (since $W^{1,d} \subset L^p$, $\forall p < \infty$). In particular, if we choose $p > d$, we obtain $Y_2 \in W_0^{1,p}(\Omega)$ such that

$$R = \operatorname{div} Y_2.$$

By the Sobolev imbedding, $Y_2 \in C^0(\bar{\Omega})$ and $Y = Y_1 + Y_2$ satisfies all the required properties. \square

8. ESTIMATION OF THE PHASE IN $H^{1/2} + W^{1,1}$. PROOF OF THEOREM 4

We return in this last section to the question discussed in the Introduction concerning the control of the phase φ in terms of $\|e^{i\varphi}\|_{H^{1/2}}$.

Let φ be a smooth real-valued function on \mathbb{T}^d and set $g = e^{i\varphi}$. The main result is the estimate

$$(8.1) \quad \|\varphi\|_{H^{1/2} + W^{1,1}} \leq C(d)(1 + \|g\|_{H^{1/2}})\|g\|_{H^{1/2}}.$$


Write g as a Fourier series

$$g = \sum_{\xi \in \mathbb{Z}^d} \hat{g}(\xi) e^{ix\xi}.$$

The $H^{1/2}$ -component in the decomposition of φ will be obtained as a paraproduct of g and \bar{g} ,

$$(8.2) \quad P = \sum_k \left[\sum_{\xi_2} \lambda_k(|\xi_2|) \overline{\hat{g}(\xi_2)} e^{-ix\xi_2} \right] \left[\sum_{2^k \leq |\xi_1| < 2^{k+1}} \hat{g}(\xi_1) e^{ix\xi_1} \right],$$

where for each k we let $0 \leq \lambda_k \leq 1$ be a smooth function on \mathbb{R}_+ :



We claim that

$$(8.3) \quad \|P\|_{H^{1/2}} \leq C \|g\|_\infty \|g\|_{H^{1/2}}$$

and

$$(8.4) \quad \|\varphi - \frac{1}{i} P\|_{W^{1,1}} \leq C \|g\|_{H^{1/2}}^2.$$

Proof of (8.3). This is totally obvious from the construction

$$(8.5) \quad \begin{aligned} \|P\|_{H^{1/2}}^2 &\sim \sum_k 2^k \left\| \left[\sum_{\xi_2} \lambda_k(|\xi_2|) \overline{\hat{g}(\xi_2)} e^{-ix\xi_2} \right] \left[\sum_{2^k \leq |\xi_1| < 2^{k+1}} \hat{g}(\xi_1) e^{ix\xi_1} \right] \right\|_2^2 \\ &\leq \sum_k 2^k \left\| \sum_{\xi} \lambda_k(|\xi|) \overline{\hat{g}(\xi)} e^{-ix\xi} \right\|_\infty^2 \left[\sum_{|\xi| \sim 2^k} |\hat{g}(\xi)|^2 \right] \\ &\leq C \|g\|_\infty^2 \|g\|_{H^{1/2}}^2. \end{aligned}$$

□

Proof of (8.4). We estimate for instance

$$(8.6) \quad \|\partial_1 \varphi - \frac{1}{i} \partial_1 P\|_{L^1}.$$

Thus, letting $\xi = (\xi^1, \dots, \xi^d) \in \mathbb{Z}^d$,

$$(8.7) \quad \partial_1 \varphi = \frac{1}{i} \bar{g} \partial_1 g = \sum_{\xi_1, \xi_2 \in \mathbb{Z}^d} \xi_1^1 \hat{g}(\xi_1) \overline{\hat{g}(\xi_2)} e^{ix \cdot (\xi_1 - \xi_2)}$$

and by (8.2)

$$(8.8) \quad \frac{1}{i} \partial_1 P = \sum_k \sum_{2^k \leq |\xi_1| < 2^{k+1}, \xi_2} (\xi_1^1 - \xi_2^1) \lambda_k(|\xi_2|) \hat{g}(\xi_1) \overline{\hat{g}(\xi_2)} e^{ix \cdot (\xi_1 - \xi_2)},$$

$$(8.9) \quad \partial_1 \varphi - \frac{1}{i} \partial_1 P = \sum_k \sum_{2^k \leq |\xi_1| < 2^{k+1}, \xi_2} m_k(\xi_1, \xi_2) \hat{g}(\xi_1) \overline{\hat{g}(\xi_2)} e^{ix \cdot (\xi_1 - \xi_2)},$$

where by definition of λ_k

$$(8.10) \quad m_k(\xi_1, \xi_2) = \xi_1^1 - \lambda_k(|\xi_2|)(\xi_1^1 - \xi_2^1) = \begin{cases} \xi_2^1 & \text{if } |\xi_2| \leq 2^{k-2}, \\ \xi_1^1 & \text{if } |\xi_2| \geq 2^{k-1}. \end{cases}$$

Estimate

$$(8.11) \quad \left\| \partial_1 \varphi - \frac{1}{i} \partial_1 P \right\|_1 \leq \sum_{k_1, k_2} \left\| \sum_{|\xi_1| \sim 2^{k_1}, |\xi_2| \sim 2^{k_2}} m_{k_1}(\xi_1, \xi_2) \hat{g}(\xi_1) \overline{\hat{g}(\xi_2)} e^{ix \cdot (\xi_1 - \xi_2)} \right\|_1.$$

Distinguish the contributions of

$$\sum_{k_1 \sim k_2} + \sum_{k_1 < k_2 - 4} + \sum_{k_1 > k_2 + 4} = (8.12) + (8.13) + (8.14).$$

Clearly $2^{-k} m_k(\xi_1, \xi_2)$ restricted to $[|\xi_1| \sim 2^k] \times [|\xi_2| \sim 2^k]$ is a smooth multiplier satisfying the usual derivative bounds. Therefore

$$(8.15) \quad (8.12) \leq C \sum_k 2^k \left\| \sum_{|\xi_1| \sim 2^k} \hat{g}(\xi_1) e^{ix \xi_1} \right\|_2 \left\| \sum_{|\xi_2| \sim 2^k} \hat{g}(\xi_2) e^{ix \xi_2} \right\|_2 \sim \|g\|_{H^{1/2}}^2.$$

If $k_1 < k_2 - 4$, then $|\xi_2| > 2^{k_1}$ and $m_{k_1}(\xi_1, \xi_2) = \xi_1^1$ by (8.10). Therefore

$$(8.13) = \sum_{k_1 < k_2 - 4} \left\| \sum_{|\xi_1| \sim 2^{k_1}, |\xi_2| \sim 2^{k_2}} \xi_1^1 \hat{g}(\xi_1) \overline{\hat{g}(\xi_2)} e^{ix \cdot (\xi_1 - \xi_2)} \right\|_1$$

$$\leq \sum_{k_1 < k_2 - 4} 2^{k_1} \left\| \sum_{|\xi_1| \sim 2^{k_1}} \hat{g}(\xi_1) e^{ix \xi_1} \right\|_2 \cdot \left\| \sum_{|\xi_2| \sim 2^{k_2}} \hat{g}(\xi_2) e^{ix \xi_2} \right\|_2$$

$$(8.16) \quad \leq \sum_{k_1 < k_2} 2^{k_1} \left(\sum_{|\xi_1| < 2^{k_1}} |\hat{g}(\xi_1)|^2 \right)^{1/2} \left(\sum_{|\xi_2| \sim 2^{k_2}} |\hat{g}(\xi_2)|^2 \right)^{1/2} \leq C \|g\|_{H^{1/2}}^2.$$

If $k_1 > k_2 + 4$, then $|\xi_2| < 2^{k_1 - 2}$ and $m_{k_1}(\xi_1, \xi_2) = \xi_2^1$ and the bound on (8.14) is similar. \square

REFERENCES

[1] R.A. Adams, "Sobolev spaces", Acad. Press, 1975. MR **56**:9247
 [2] D.N. Arnold, L.R. Scott and M. Vogelius, *Regular inversion of the divergence operator with Dirichlet boundary condition on a polygon*, Ann. Sc. Norm. Pisa, Serie IV, **15** (1988), 169–192. MR **91i**:35043
 [3] J. Bourgain, H. Brezis and P. Mironescu, *Lifting in Sobolev spaces*, J. d'Analyse **80** (2000), 37-86. MR **2001h**:46044
 [4] ———, *On the structure of the Sobolev space $H^{1/2}$ with values into the circle*, C. R. Acad. Sc. Paris **331** (2000), 119-124. MR **2001m**:46068
 [5] ———, in preparation.
 [6] H. Brezis, "Analyse fonctionnelle, théorie et applications", Masson, 1983. MR **2001m**:46068
 [7] D. Burago and B. Kleiner, *Separated nets in Euclidean space and Jacobians of bi-Lipschitz maps*, Geom. Funct. Anal. **8** (1998), 273-282. MR **99d**:26018
 [8] B. Dacorogna and J. Moser, *On a partial differential equation involving the Jacobian determinant*, Ann. Inst. H. Poincaré, Anal. Nonlinéaire, **7** (1990), 1-26. MR **91i**:58148
 [9] G. Duvaut and J.L. Lions, "Les inéquations en mécanique et en physique", Dunod, 1972; English translation, Springer, 1976. MR **57**:4778
 [10] C. Fefferman and E. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), 137-193. MR **56**:6263
 [11] M. Gromov, *Asymptotic invariants of infinite groups*, in "Geometric Group Theory", Vol.2 (G. A. Niblo, M. A. Roller, eds.), Cambridge Univ. Press, 1993. MR **95m**:20041
 [12] E. Magenes and G. Stampacchia, *I problemi al contorno per le equazioni differenziali di tipo ellittico*, Ann. Sc. Norm. Pisa **12** (1958), 247–357. MR **23**:A1140
 [13] C. T. McMullen, *Lipschitz maps and nets in Euclidean space*, Geom. Funct. Anal. **8** (1998), 304-314. MR **99e**:58017

- [14] J. Nečas, *Sur les normes équivalentes dans $W_p^{(k)}(\Omega)$ et sur la coercitivité des formes formellement positives*, in “Equations aux dérivées partielles”, Presses de l’Université de Montreal, 1966.
- [15] L. Nirenberg, “Topics in nonlinear functional analysis”, New York Univ. Lecture Notes, 1973–74. MR **58**:7672
- [16] D. Ornstein, *A non-inequality for differential operators in the L_1 norm*, Arch. Rat. Mech. Anal. **11** (1962), 40-49. MR **26**:6821
- [17] T. Rivière and D. Ye, *Une résolution de l’équation à forme volume prescrite*, C. R. Acad. Sc. Paris **319** (1994), 25-28. MR **95f**:35055
- [18] ———, *Resolutions of the prescribed volume form equations*, Nonlinear Differential Equations Appl. **3** (1996), 323-369. MR **97g**:35045
- [19] J.L. Rubio de Francia, *A Littlewood-Paley theorem for arbitrary intervals*, Rev. Mat. Iberoamericana **1** (1985), 1–14. MR **87j**:42057
- [20] R. Temam, “Navier-Stokes equations”, North-Holland, revised edition, 1979. MR **82b**:35133
- [21] A. Uchiyama, *A constructive proof of the Fefferman-Stein decomposition of $BMO(\mathbb{R}^n)$* , Acta Math. **148** (1982), 215-241. MR **84h**:42037
- [22] X. Wang, *A remark on the characterization of the gradient of a distribution*, Applic. Anal. **51** (1993), 35-40. MR **95k**:46064
- [23] R. Wojtaszczyk, “Banach spaces for analysts”, Cambridge Univ. Press, 1991. MR **93d**:46001
- [24] D. Ye, *Prescribing the Jacobian determinant in Sobolev spaces*, Ann. Inst. H. Poincaré, Anal. Nonlinéaire **11** (1994), 275-296. MR **95g**:35058

INSTITUTE FOR ADVANCED STUDY, PRINCETON, NEW JERSEY 08540

E-mail address: bourgain@math.ias.edu

ANALYSE NUMÉRIQUE, UNIVERSITÉ P. ET M. CURIE, B.C. 187, 4 PL. JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: brezis@ccr.jussieu.fr

Current address: Department of Mathematics, Rutgers University, Hill Center, Busch Campus, 110 Frelinghuysen Rd., Piscataway, New Jersey 08854

E-mail address: brezis@math.rutgers.edu