

ON NEUMANN EIGENFUNCTIONS IN LIP DOMAINS

RAMI ATAR AND KRZYSZTOF BURDZY

1. INTRODUCTION AND MAIN RESULT

A planar set D will be called a *lip domain* if it is Lipschitz, open, bounded, connected, and given by

$$(1) \quad D = \{(x_1, x_2) : f_1(x_1) < x_2 < f_2(x_1)\},$$

where f_1, f_2 are Lipschitz functions with constant 1. The assumption that D is a Lipschitz domain puts an extra constraint on the functions f_k ; we discuss this issue in greater detail later in this section.

Let μ_2 denote the second eigenvalue for the Laplacian in D with Neumann boundary conditions. Here is our main result.

Theorem 1. (i) *The second eigenvalue μ_2 is simple in all lip domains except squares.*

(ii) (“Hot spots conjecture”) *For every lip domain, every Neumann eigenfunction corresponding to μ_2 attains its maximum and minimum at boundary points only.*

The “hot spots” conjecture was proposed by J. Rauch at a conference in 1974 and was never published in print so it is somewhat vague; roughly speaking, it says that in Euclidean domains, the second Neumann eigenfunction attains its maximum at the boundary (see [3] for a discussion of different versions of the conjecture). It was shown by Kawohl ([16]) that the conjecture holds in cylinders. Bañuelos and Burdzy ([3]) proved the conjecture for a large class of lip domains and some convex domains. However, their techniques were limited to polygonal domains. In the proof of Theorem 1 (ii), we will show how one can drop this assumption using the results of Burdzy and Chen ([8]). Bañuelos and Burdzy ([3]) were able to prove the “hot spots” conjecture for only one of the eigenfunctions corresponding to μ_2 . This naturally lead to the problem of characterization of those lip and convex domains where μ_2 is simple. They gave a partial answer in [3]—it was shown that the eigenvalue is simple in a convex domain if the ratio of length to width of the domain is larger than 3.07. The critical ratio is only 1.53 if the convex domain is assumed in addition to have a line of symmetry.

Received by the editors December 17, 2001.

2000 *Mathematics Subject Classification.* Primary 35J05; Secondary 60H30.

Key words and phrases. Neumann eigenfunctions, reflected Brownian motion, couplings, hot spots problem.

Research partially supported by the fund for the promotion of research at the Technion.

The second author gratefully acknowledges the hospitality and financial support of Technion (Israel) and Institut Mittag-Leffler (Sweden). This research was partially supported by NSF Grant DMS-0071486 and ISF Grant 12/98.

The “hot spots” conjecture does not hold in arbitrary planar domains ([10], [5]) but it is still an open problem as to whether it holds in convex domains. See [2], [14] and [20] for some recent results on this problem.

Our main result solves two problems posed by Jerison and Nadirashvili ([14]).

Corollary 1. *The following conjectures are true.*

(i) ([14], *Conj. 8.2*) *If D is a convex polygon with two axes of symmetry and a right angle, then the only case in which the eigenvalue is multiple is when D is a square.*

(ii) ([14], *Conj. 8.5*) *Let D be a convex, centrally symmetric domain contained in a circular sector of acute angle and vertex P . If $P \in \partial D$, then the maximum or the minimum of the eigenfunction corresponding to μ_2 is achieved at P , and μ_2 is simple.*

Our proof of Theorem 1 uses a probabilistic technique called “coupling.” Two large sections of the article are devoted to the construction (Section 2) and analysis (Section 3) of “mirror couplings” of reflected Brownian motions. Mirror couplings (without reflection) were used by Kendall ([17]) to obtain analytic estimates. The same couplings for reflected diffusions were applied in [22], [3] and [9]. Unfortunately, none of the last three quoted papers paid sufficient attention to the construction of mirror couplings. The paper by Wang [22] does not offer any proof of existence of mirror couplings for reflected diffusions. The papers by Bañuelos and Burdzy [3] and Burdzy and Kendall [9] give an explicit construction for the mirror coupling of Brownian motions reflected on a straight line. This construction breaks down in polygonal domains when two Brownian particles hit two different line segments on the boundary at the same time (proving that this event has a positive probability is nontrivial; we omit the proof as it is not needed here). Although some readers may find our construction of mirror couplings in piecewise C^2 domains exceedingly detailed, we believe that the history of the problem justifies the extra care we give to it.

The most challenging part of our paper is a probabilistic version of the “parabolic boundary Harnack principle,” i.e., Section 3. We will show that if two mirror coupled Brownian motions are conditioned on not meeting by a certain time, then they are likely to be far apart at that time. This intuitively obvious statement seems to have no easy proof. The core of our argument is taken from [4]. The same approach has already been used in [1] and [9] but each application of the same basic idea needs new estimates and some minor changes. The present application of the argument from [4] needs completely new estimates for a few fundamental quantities—this is what makes that part of the paper tedious. An additional complication stems from the fact that our construction of mirror couplings of reflected Brownian motions is limited to piecewise C^2 domains. We have to consider a sequence of mirror couplings in a sequence of piecewise smooth domains approximating the lip domain. Known results on the existence of strong solutions to the Skorohod equation indicate that the construction of a mirror coupling with the strong Markov property in an arbitrary lip domain may be a very difficult task—we do not try to undertake it in this article.

The proof of the main theorem, using ideas of Atar [2], is given in Section 4. The core argument is the following. Let (e_1, e_2) be the usual orthonormal basis for \mathbb{R}^2 . It was observed in [3] (see also Section 4 in this article) that in every lip domain, there exists an “increasing” Neumann eigenfunction corresponding to

μ_2 , i.e., an eigenfunction whose directional derivatives in directions $e_1 + e_2$ and $e_1 - e_2$ are nonnegative on D . We prove that if μ_2 is not simple, then there exists a corresponding eigenfunction which is “increasing” in a large part of the domain but not at all points. The key estimate, Lemma 6, shows that if this condition is satisfied by the gradient in the bulk of the domain, then it has to be satisfied everywhere in the domain. In other words, Lemma 6 provides a link between the local and global estimates for the direction of the gradient of an eigenfunction—this is where the results from Sections 2 and 3 on mirror couplings are used in a crucial way.

We end this introduction with a few remarks on lip domains. This class of domains appeared for the first time in [3] and [9]. The name “lip domains” was coined in [7].

It follows easily from the definition of a lip domain D that there exist leftmost and rightmost extreme points $x^*, y^* \in \overline{D}$, i.e., for every $x = (x_1, x_2) \in \overline{D} \setminus \{x^*, y^*\}$, we have $x_1^* < x_1 < y_1^*$. Recall that the definition of a lip domain D involves a requirement that D be a Lipschitz domain. This means that every $x \in \partial D$ has a neighborhood U such that $\partial D \cap U$ is the graph of a Lipschitz function in some orthonormal coordinate system. Hence, $f_2(x_1) - f_1(x_1)$ cannot go to 0 very fast as $x_1 \downarrow x_1^*$ or $x_1 \uparrow y_1^*$. Without this condition, one could construct domains satisfying (1) and containing infinite sequences of “rooms and passages” close to x^* and y^* . The results of [13] indicate that such domains might not have discrete spectrum. The exclusion of these domains from our consideration does not seem to be an essential limitation because lip domains with shapes significantly different from the square are not likely to have multiple second eigenvalues.

We will write $B_r(x) \subset \mathbb{R}^2$ to denote an open ball of radius r around x and we will use c to denote positive constants whose values are unimportant and may change from line to line.

2. CONSTRUCTION OF MIRROR COUPLINGS

Before we go into technical proofs, we briefly and informally describe the goal of this section. Suppose for a moment that $D \in \mathbb{R}^d$, $d \geq 2$, is a smooth domain. We are looking for a pair of reflected Brownian motions (X_t, Y_t) satisfying the system of stochastic differential equations

$$dX = dW + dL, \quad dY = dZ + dM, \quad dZ = dW - 2m \frac{m \cdot dW}{|Y - X|}, \quad m = \frac{Y - X}{|Y - X|}$$

for times less than $\zeta = \inf\{s : X_s = Y_s\}$. Here W is a d -dimensional Brownian motion, and Z is another Brownian motion for which the increments are mirror images of those of W , the mirror being the $(d - 1)$ -dimensional hyperplane with respect to which X and Y are symmetric. The processes X and Y are reflecting Brownian motions with boundary terms L and M , respectively. We will prove pathwise uniqueness and strong existence for this system of equations. As a corollary, it will be proved that (X_t, Y_t) is strong Markov. The final proposition in this section will show that mirror couplings in lip domains have an “order preserving” property which is crucially used in this paper.

An SDE with reflecting boundary conditions. The starting point for our analysis is some results from [19] regarding unique solvability of an SDE with reflection. The

results in that paper apply to a class of domains satisfying complicated assumptions—we refer the reader to [19] for details. It will suffice to say that the results of [19] apply to all $D \subset \mathbb{R}^d$, $d \geq 3$, which are C^2 -smooth and all planar piecewise C^2 -smooth domains with a finite number of convex corners (more precisely, domains for which the boundary consists of finitely many smooth parts and for which the uniform exterior sphere condition and the uniform interior cone condition are satisfied; see [19]). All results in this section hold true for all domains which satisfy the conditions listed in [19], so they are not limited to the cases we have just listed.

Fix a bounded open set $D \subset \mathbb{R}^d$ for some $d \geq 2$ and assume that its boundary is C^2 -smooth if $d \geq 3$ or piecewise smooth with a finite number of convex corners if $d = 2$. For $x \in \partial D$, let $\nu(x)$ be the inward unit normal vector, if it exists. If x is a corner, we let $\nu(x)$ be an arbitrary unit vector pointing inside the domain at x (this is just for the sake of completeness of the definition—the reflected Brownian motion does not visit corners with probability 1 so the definition of $\nu(x)$ for such points is irrelevant).

Let $(\Omega, \mathcal{F}, (F_t), P)$ be a complete filtered probability space, on which an (F_t) -Brownian motion (W_t) is given. It was shown in [19] that for any $x \in \overline{D}$ there exists a unique continuous (F_t) -semimartingale (X_t) with values in \overline{D} for all $t \geq 0$ a.s., satisfying

$$(2) \quad X_t = x + W_t + L_t, \quad |L|_t = \int_0^t 1_{X_s \in \partial D} d|L|_s, \quad L_t = \int_0^t \nu(X_s) d|L|_s,$$

where L_t is a continuous process with values in \mathbb{R}^d , with variation $|L|_t$ which is bounded on each finite interval. In fact, there is a map $\Gamma : C([0, \infty) : \mathbb{R}^d) \rightarrow C([0, \infty) : \mathbb{R}^d)$ (the “Skorohod map”) such that $X = \Gamma(x + W)$ a.s. Moreover, for every $T > 0$, there exists a map Γ_T from $C([0, T] : \mathbb{R}^d)$ into itself, such that

$$(3) \quad X|_{[0, T]} = \Gamma(x + W)|_{[0, T]} = \Gamma_T(x + W|_{[0, T]}),$$

and Γ_T is Hölder continuous of order $1/2$ on compact subsets of $C([0, T] : \mathbb{R}^d)$. This implies that the function $x \mapsto \Gamma_T(x + W|_{[0, T]})$ is continuous. With an abuse of notation, we will refer below to Γ_T as Γ . We will call the process X a reflected Brownian motion in D (with normal reflection), driven by W , starting from x .

Denote

$$\|Z\|_t = [E \sup_{0 \leq s \leq t} |Z_s|^4]^{1/4}.$$

Let \mathbf{Z} be the space of continuous adapted processes Z for which $\|Z\|_t < \infty$ for all $t > 0$. We shall need the following version of Lemma 3.1 of [19].

Lemma 1. *Let σ be an (F_t) -stopping time. Then for each $T > 0$ there exists C such that for all $t \leq T$ and all $Z, Z' \in \mathbf{Z}$,*

$$\|\Gamma(Z)_{\cdot \wedge \sigma} - \Gamma(Z')_{\cdot \wedge \sigma}\|_t^4 \leq C \int_0^t \|Z_{\cdot \wedge \sigma} - Z'_{\cdot \wedge \sigma}\|_s^4 ds.$$

Proof. The only difference between our lemma and Lemma 3.1 of [19] is that the processes are stopped at a stopping time σ in our version of the inequality. The original proof given in [19] applies, with minor adjustments needed to take into account the presence of the stopping time σ . \square

Equations for the mirror coupling. We will formulate a system of stochastic differential equations for a pair of “mirror coupled” reflecting Brownian motions, i.e., satisfying the following condition. On any time interval $[s, t]$, on the event A that both processes do not hit the boundary during $[s, t]$, there is a $(d - 1)$ -dimensional hyperplane (the “mirror”), depending on $\omega \in A$ and s, t , with respect to which the processes are symmetric at every time in $[s, t]$.

Let $(\Omega, F, (F_t), P, W)$ be given, where W is an (F_t) -Brownian motion. We are looking for processes Z, X and Y with the following properties. The processes X and Y are reflecting Brownian motions in D , starting from x and y , and driven by W and Z , respectively. Here Z is another Brownian motion on the same probability space. We will work with deterministic initial conditions, but we note that replacing them with F_0 -measurable initial conditions would add no difficulty. We want to have

$$(4) \quad X = \Gamma(x + W),$$

as in (2). We also need,

$$(5) \quad Y = \Gamma(y + Z).$$

Let m_t be a unit vector perpendicular to the mirror with respect to which X_t and Y_t are symmetric, namely $m_t = (Y_t - X_t)/|Y_t - X_t|$. Here and later, all vectors will be column vectors. For m in the unit sphere of \mathbb{R}^d , let $H(m)$ denote the $d \times d$ matrix

$$(6) \quad H(m) = I - 2mm',$$

where m' denotes the transpose of m . Then $H(m)v = v - 2(m \cdot v)m$ is the mirror image of v about the hyperplane through the origin, perpendicular to m . We would like Z to depend on W and on the mirror, in such a way that, before X and Y first meet, Z 's increments are mirror images of W 's:

$$Z_t = \int_0^t H(m_s)dW_s.$$

For $x \in \mathbb{R}^d$, let $G(x) = H(x/|x|)$ if $x \neq 0$, and let $G(0) = 0$ (the value of $G(0)$ is in fact irrelevant). Equip the space of $d \times d$ matrices with the norm $\|A\|^2 = \sum_{i,j} A_{i,j}^2$. Then it is easy to check that for $|m| = |\tilde{m}| = 1$,

$$(7) \quad \|H(m) - H(\tilde{m})\| \leq c|m - \tilde{m}|.$$

Consider the equation

$$(8) \quad Z_t = \int_0^{t \wedge \zeta} G(\Gamma(y + Z)_s - X_s)dW_s + 1_{t \geq \zeta}(W_t - W_\zeta), \quad \zeta = \inf\{s : \Gamma(y + Z)_s = X_s\}.$$

Definition 1. (i) We say that *pathwise uniqueness holds for (8)* if whenever $(\Omega, F, (F_t), P)$ is a filtered probability space, W is an (F_t) -Brownian motion, $X = \Gamma(x + W)$, processes Z and Z' are (F_t) -adapted and have continuous sample paths, and both Z and Z' satisfy (8) for $t \geq 0$ a.s., then $Z(t) = Z'(t)$ for all $t \geq 0$, a.s.

(ii) A *strong solution* to (8) on a given probability space (Ω, F, P) relative to the Brownian motion W is a process Z with continuous sample paths, adapted to the (augmented) filtration (F_t^W) , generated by W , and with $X = \Gamma(x + W)$, which satisfies (8) for all $t \geq 0$, a.s.

Theorem 2. *Pathwise uniqueness holds for (8). Let W be a Brownian motion on a complete probability space (Ω, \mathcal{F}, P) . Then there exists a strong solution of (8) relative to W .*

We will refer to the pair (X, Y) as a *mirror coupling* of reflecting Brownian motions.

Proof of Theorem 2. Pathwise uniqueness. We will first consider the problem on a finite time interval $[0, T]$. Let a Brownian motion W on $(\Omega, \mathcal{F}, (F_t), P)$ be given, and let $X = \Gamma(x+W)$. Assume Z and \tilde{Z} are two (F_t) -adapted processes with continuous sample paths, satisfying (8) for all $t \in [0, T]$ a.s. Denote $Y = \Gamma(y+Z)$, $\tilde{Y} = \Gamma(y+\tilde{Z})$, $V = Y - X$, $\tilde{V} = \tilde{Y} - X$. Let $\tau_n = \inf\{t : |V_t| \leq 1/n\}$, $\tilde{\tau}_n = \inf\{t : |\tilde{V}_t| \leq 1/n\}$, $S_n = \tau_n \wedge \tilde{\tau}_n$. Then $S_n \uparrow \zeta \wedge \tilde{\zeta}$. To prove pathwise uniqueness, it is enough to show that $P(Z_t = \tilde{Z}_t, 0 \leq t < \zeta \wedge \tilde{\zeta}) = 1$. On $[0, S_n]$, one has

$$(9) \quad \left| \frac{V}{|V|} - \frac{\tilde{V}}{|\tilde{V}|} \right| \leq c \frac{|V - \tilde{V}|}{|V| \wedge |\tilde{V}|} \leq cn|V - \tilde{V}| = cn|Y - \tilde{Y}|.$$

We will use the following version of the Burkholder-Davis-Gundy inequality ([15] p. 163),

$$(10) \quad E \left| \int_0^T A_t dW_t \right|^{2m} \leq c_m T^{m-1} E \int_0^T |A_t|^{2m} dt$$

for $m \geq 1$ and A_t adapted. Since Z is adapted, it follows from (3) that so is the integrand in (8), and therefore Z is a martingale. Doob's inequality for the martingale $Z - \tilde{Z}$, along with Lemma 1, (7), (9) and (10), yields for any $t \in (0, T]$,

$$\begin{aligned} E \sup_{0 \leq s \leq t} |Z_{s \wedge S_n} - \tilde{Z}_{s \wedge S_n}|^4 &\leq cE|Z_{t \wedge S_n} - \tilde{Z}_{t \wedge S_n}|^4 \\ &\leq cTE \int_0^{t \wedge S_n} \|G(Y_s - X_s) - G(\tilde{Y}_s - X_s)\|^4 ds \\ &= cTE \int_0^{t \wedge S_n} \left\| H\left(\frac{V_s}{|V_s|}\right) - H\left(\frac{\tilde{V}_s}{|\tilde{V}_s|}\right) \right\|^4 ds \\ &\leq cTE \int_0^{t \wedge S_n} \left| \frac{V_s}{|V_s|} - \frac{\tilde{V}_s}{|\tilde{V}_s|} \right|^4 ds \\ &\leq cTn^4 E \int_0^{t \wedge S_n} |Y_s - \tilde{Y}_s|^4 ds \\ &\leq cTn^4 \int_0^t E \sup_{0 \leq u \leq s} |Y_{u \wedge S_n} - \tilde{Y}_{u \wedge S_n}|^4 ds \\ &\leq cT^2 n^4 \int_0^t E \sup_{0 \leq u \leq s} |Z_{u \wedge S_n} - \tilde{Z}_{u \wedge S_n}|^4 ds. \end{aligned}$$

Gronwall's inequality now shows that $E \sup_{0 \leq s \leq T} |Z_{s \wedge S_n} - \tilde{Z}_{s \wedge S_n}|^4 = 0$. Hence $P(Z_s = \tilde{Z}_s, 0 \leq s \leq S_n) = 1$, for all n , and therefore $P(Z_s = \tilde{Z}_s, 0 \leq s < \zeta \wedge \tilde{\zeta}) = 1$. This shows that Z and \tilde{Z} are indistinguishable on $[0, T]$. Since T is arbitrary, the same is true on $[0, \infty)$. This completes the proof of pathwise uniqueness.

Existence of strong solutions. By (7), the restriction of the matrix-valued mapping $x \mapsto G(x)$ to $\mathbb{R}^d \setminus B_\varepsilon(0)$ is globally Lipschitz with respect to $\|\cdot\|$. Therefore, for every $n \in \mathbb{N}$, one can find a globally Lipschitz function G^n on \mathbb{R}^d that agrees with G outside $B_{1/n}(0)$. Let $\lambda_n < \infty$ denote the Lipschitz constant of G^n . We are given a probability space (Ω, F, P) with a Brownian motion W and the filtration (F_t^W) generated by W . Denote $X = \Gamma(x + W)$. We will first show that for any n there exists a continuous adapted process Z^n such that

$$(11) \quad Z_t^n = \int_0^t G^n(\Gamma(y + Z^n)_s - X_s) dW_s.$$

But before that, let us note that pathwise uniqueness holds for (11). The proof is similar to that of pathwise uniqueness for (8) but simpler, so it is omitted.

In this paragraph we drop the dependence on n in the notation of the processes involved. Recall the space \mathbf{Z} defined before the statement of Lemma 1. Let F map $Z \in \mathbf{Z}$ to

$$F(Z)_t = \int_0^t G^n(\Gamma(y + Z)_s - X_s) dW_s.$$

Suppose $Z, Z' \in \mathbf{Z}$. By Lemma 1, Doob's inequality for the martingale $F(Z) - F(Z')$ and (10), we have for $t \in (0, T]$,

$$(12) \quad \begin{aligned} E \sup_{s \leq t} |F(Z)_s - F(Z')_s|^4 &\leq cE|F(Z)_t - F(Z')_t|^4 \\ &\leq ctE \int_0^t \|G^n(\Gamma(y + Z)_s - X_s) - G^n(\Gamma(y + Z')_s - X_s)\|^4 ds \\ &\leq c\lambda_n^4 tE \int_0^t |\Gamma(y + Z)_s - \Gamma(y + Z')_s|^4 ds \\ &\leq c\lambda_n^4 T^2 \int_0^t E \sup_{0 \leq u \leq s} |Z_u - Z'_u|^4 ds. \end{aligned}$$

Let $Z^{(0)} = \mathbf{0}$, and for $k \in \mathbb{N}$ let $Z^{(k)} = F(Z^{(k-1)})$. Since G^n is bounded, there is c such that $\|Z^{(1)} - Z^{(0)}\|_T \leq c$. Iterating (12), it easily follows that

$$\|Z^{(k+1)} - Z^{(k)}\|_T^4 \leq c \frac{c_1^k}{k!},$$

where c_1 is a constant that may depend on n and T . Hence by Chebyshev's inequality,

$$P(\sup_{s \leq T} |Z_s^{(k+1)} - Z_s^{(k)}| \geq 2^{-k}) \leq c(16c_1)^k / k!,$$

and the Borel-Cantelli lemma shows that with probability one,

$$\sup_{0 \leq s \leq T} |Z_s^{(k+1)} - Z_s^{(k)}| \leq 2^{-k}$$

for all sufficiently large k . Hence the paths of $Z^{(k)}$ converge in the uniform topology, a.s., say, to Z . Again using (12), we have $\|F(Z^{(k)}) - F(Z)\|_T^4 \leq c\|Z^{(k)} - Z\|_T^4$, and as a result $F(Z^{(k)})$ converge a.s. to $F(Z)$. We see that Z is a fixed point of F . Since T is arbitrary, we conclude that there exists a continuous, (F_t^W) -adapted process satisfying (11) for all $t \in [0, \infty)$ a.s.

We reintroduce the dependence on n in our notation. Let Z^n denote the process in (11) and set $Y^n = \Gamma(y + Z^n)$. Let $\tau_n = \inf\{s : |Y_s^n - X_s| \leq 1/n\}$. Since we have

uniqueness for (11), it is clear that on $[0, \tau_n]$, the processes Z^n and Z^{n+1} agree a.s. Setting $\zeta = \lim_n \tau_n$ and defining

$$Z_t = 1_{t < \zeta} \lim_{n \rightarrow \infty} Z^n(t) + 1_{t \geq \zeta} (W_t - W_\zeta),$$

we obtain a continuous and adapted process Z . We see that (8) is satisfied for all $t \in [0, \infty)$ a.s. Therefore there exists a strong solution relative to W . \square

Strong Markov property. Let $(\Omega, \mathcal{F}, (F_t), P, W)$ be a complete filtered probability space and a Brownian motion. Let $U = (X, Y)$ denote the unique solution to (4), (5) and (8). For $u = (x, y) \in (\overline{D})^2$, let P^u denote the measure induced by U on $(C([0, \infty) : \mathbb{R}^{2d}), \mathcal{B}(C([0, \infty) : \mathbb{R}^{2d})))$, assuming $U_0 = u$. Let E^u denote the expectation with respect to P^u . The transition function for the process U is defined as

$$P_t(u, f) = E^u(f(U_t)),$$

for all $u \in (\overline{D})^2$ and bounded Borel measurable functions f . Given the results on existence and uniqueness, and the properties of the Skorohod map quoted before, we have the following.

Corollary 2. *The process U is strong Markov, i.e., for any a.s. finite (F_t) -stopping time T and bounded Borel function f , one has P -a.s., for all $s \geq 0$,*

$$E[f(U_{T+s})|F_T] = P_s(U_T, f).$$

Proof. We take here the approach of [21], Theorem V.32. For each finite stopping time T , note that $\widetilde{W}_s = W_{T+s} - W_T$, $s \geq 0$, is a Brownian motion (cf., e.g., [21], Theorem I.32), and consider the unique solution \widetilde{Z} to the following equation, with $u = (x, y) \in (\overline{D})^2$,

$$(13) \quad \widetilde{Z}_t = \int_0^{t \wedge \widetilde{\zeta}} G(\Gamma(y + \widetilde{Z})_s - \Gamma(x + \widetilde{W})_s) d\widetilde{W}_s + 1_{t \geq \widetilde{\zeta}} (\widetilde{W}_t - \widetilde{W}_{\widetilde{\zeta}}),$$

where $\widetilde{\zeta} = \inf\{s : \Gamma(y + \widetilde{Z})_s = \Gamma(x + \widetilde{W})_s\}$. To take into account the dependence on the stopping time and the initial condition, we denote \widetilde{Z}_t above by $Z(u, T, t)$, and we also let

$$U(u, T, t) = (\Gamma(x + W_{\cdot+T})_t, \Gamma(y + Z(u, T, \cdot))_t).$$

Denote the Borel sigma-field of subsets of $(\overline{D})^2$ [resp., \mathbb{R}_+] by \mathcal{U} [resp., \mathcal{R}_+]. We will show that, for a given stopping time, the mapping $(u, t, \omega) \mapsto Z(u, T, t)$ has a $\mathcal{U} \otimes \mathcal{R}_+ \otimes \mathcal{F}$ jointly measurable version, and hence so does $U(u, T, t)$. We will use these versions in the rest of the proof. Recall the processes $Z^{(k)}$ and Z^n of the proof of Theorem 2. Recall also that for each n , $Z^{(k)}$ converge uniformly on compacts in probability to Z^n . Since each $Z^{(k)}$ is jointly measurable, this implies that Z^n is also (cf. [21]). Since Z is the pointwise limit of Z^n as $n \rightarrow \infty$, this shows that Z is jointly measurable. The same argument applies to $Z(u, T, t)$ for any fixed stopping time T . A similar measurability property holds for $U(u, T, t)$ by the uniform continuity of Γ on compact sets.

Fix a finite stopping time T and set $F^* = \sigma\{W_{T+u} - W_T : u \geq 0\}$. Then F^* is independent of F_T under P , since W is an (F_t) -Brownian motion. Note that $Z(u, T, t)$ is measurable with respect to F^* . By the uniqueness results for the Skorohod map (the uniqueness holds in the deterministic sense), $\Gamma(R)_{T+s} = \Gamma(\Gamma(R)_T - R_T + R_{T+\cdot})_s$, $s \geq 0$, whenever R is an (F_t) -Brownian motion. Fix a u

for the moment and denote $Z_t = Z(u, 0, t)$. Then we have for $t \geq 0$, on the event $\{\zeta > T + t\}$,

$$\begin{aligned} Z_{T+t} - Z_T &= \int_T^{T+t} G(\Gamma(y + Z)_s - \Gamma(x + W)_s) dW_s \\ (14) \quad &= \int_0^t G(\Gamma(\Gamma(y + Z)_T + (Z_{T+} - Z_T))_s - \Gamma(\Gamma(x + W)_T + \widetilde{W}))_s) d\widetilde{W}_s. \end{aligned}$$

Let $\bar{\zeta} = \inf\{s \geq 0 : \Gamma(y + Z_{T+})_s = \Gamma(x + W_{T+})_s\}$, and note that $\bar{\zeta} = 0$ on $\zeta \leq T$. On the event $\{\zeta \leq T + t\} = \{\bar{\zeta} \leq t\}$, we have,

$$\begin{aligned} Z_{T+t} - Z_T &= Z_{\bar{\zeta}} + W_{T+t} - W_{\bar{\zeta}} \\ (15) \quad &= Z_{\bar{\zeta}} + \widetilde{W}_t - \widetilde{W}_{\bar{\zeta}}. \end{aligned}$$

Equations (14) and (15) show that $Z_{T+t} - Z_T$ is a solution to (13) with the initial condition $U(u, 0, T)$. By uniqueness of solutions we therefore have

$$Z_{T+t} - Z_T \equiv Z(u, 0, T + t) - Z(u, 0, T) = Z(U(u, 0, T), T, t).$$

It follows that P -a.s., for all $t \geq 0$,

$$U(u, 0, T + t) = U(U(u, 0, T), T, t).$$

We will use the following standard fact. If $\Phi(h, \cdot)$ is independent of the σ -field \mathcal{H} for every h and H is \mathcal{H} -measurable, then $E(\Phi(H, \cdot) | \mathcal{H}) = \phi(H)$ a.s., where $\phi(h) = E(\Phi(h, \cdot))$. Hence, for any bounded Borel f ,

$$\begin{aligned} E\{f(U(u, 0, T + t)) | F_T\} &= E\{f(U(U(u, 0, T), T, t)) | F_T\} \\ &= g(U(u, 0, T)), \end{aligned}$$

where $g(r) = Ef(U(r, T, t))$. However, by pathwise uniqueness of solutions to (13), under P , $Z(u, T, \cdot)$ and $Z(u, 0, \cdot)$ are equal in law, and therefore so are $U(u, T, \cdot)$ and $U(u, 0, \cdot)$. Consequently, $g(r) = Ef(U(r, 0, t)) = P_t(r, f)$. This shows that

$$E\{f(U(u, 0, T + t)) | F_T\} = P_t(U(u, 0, T), f).$$

□

Remark. Fix some u , consider the probability space (Ω, F, P^u) , where P^u is the measure induced by U when started at u , and let (\widehat{F}_t) be the filtration generated by U . Then by Corollary 2, for any a.s. finite (\widehat{F}_t) -stopping time T , the random variable $E[f(U_{T+s}) | F_T] = P_s(U_T, f)$ is measurable with respect to $\sigma(U_T)$. It follows that it is measurable with respect to \widehat{F}_T , and therefore P^u -a.s.,

$$E^u[f(U_{T+s}) | \widehat{F}_T] = E[f(U_{T+s}) | F_T] = P_s(U_T, f).$$

Next we will show that the “mirror coupling” (X, Y) has in fact the mirror property.

Proposition 1. *Let (X, Y) be a mirror coupling of reflecting Brownian motions, let A be an event in F , and let σ, τ be two a.s. finite F -measurable times with $\sigma \leq \tau$ a.s. on A , such that $X_s, Y_s \notin \partial D$ and $X_s \neq Y_s$ for all $s \in [\sigma, \tau]$ a.s. on A . Then a.s. on A ,*

$$(16) \quad (X_s + Y_s) \cdot (Y_\sigma - X_\sigma) = (X_\sigma + Y_\sigma) \cdot (Y_\sigma - X_\sigma), \quad \text{for all } s \in [\sigma, \tau].$$

Proof. Recall that in our notation, $dX = dW + dL$, $dY = dZ + dM$, L and M are the boundary terms for X and Y , and $dZ = H(m)dW = dW - 2m \cdot m \cdot dW$. Here $m_s = V_s/|V_s|$ on $\{s < \zeta\}$, where $V_s = Y_s - X_s$. Applying Itô's formula, one finds that

$$(17) \quad dm_s = |V_s|^{-1}(dM_s - dL_s) - |V_s|^{-1}m_s m_s \cdot (dM_s - dL_s).$$

This shows that the mirror does not move within $[\sigma, \tau]$ on A . Hence for $s \in [\sigma, \tau]$,

$$Z_s = Z_\sigma + G(Y_\sigma - X_\sigma)(W_s - W_\sigma).$$

Since for $s \in [\sigma, \tau]$,

$$X_s = X_\sigma + W_s - W_\sigma, \quad Y_s = Y_\sigma + Z_s - Z_\sigma,$$

(16) follows. □

Let (e_1, e_2) be the usual orthonormal basis for \mathbb{R}^2 and let

$$e'_1 = (e_1 - e_2)/\sqrt{2}, \quad e'_2 = (e_1 + e_2)/\sqrt{2}.$$

For $x, y \in \mathbb{R}^2$, let $x \leq y$ mean

$$x \cdot e'_i \leq y \cdot e'_i, \quad i = 1, 2.$$

The following “order preserving” property of mirror couplings in lip domains is one of the crucial ingredients of the main argument.

Proposition 2. *Suppose $D \subset \mathbb{R}^2$ is a piecewise C^2 -smooth lip domain for which the defining functions f_1, f_2 (cf. equation (1)) are both Lipschitz with constants strictly less than 1. Let (X, Y) be a mirror coupling of reflecting Brownian motions in D , starting from $(x, y) \in (\overline{D})^2$. If $x \leq y$, then $X_t \leq Y_t$ for all $t \geq 0$, a.s.*

Proof. Recall the definition of m_s from the proof of Proposition 1. Let a_s be the unit vector perpendicular to m_s such that, in complex number notation, $ia_s = m_s$. Then the identity (17) is equivalent to

$$dm_s = |V_s|^{-1}a_s a_s \cdot (dM_s - dL_s).$$

Let $\tau = \inf\{t : X_t \not\leq Y_t\}$ and $\Omega_1 = \{\tau < \infty\}$. We will show that Ω_1 has probability 0. Suppose $\tau < \infty$. Since a normally reflecting Brownian motion does not hit a fixed point on the boundary, we can assume that X_τ and Y_τ are not at any of the vertices of ∂D . Let $\partial_+ D$ [$\partial_- D$] denote the intersection of the boundary ∂D with the graph of f_2 [respectively, f_1]. Because the Lipschitz constants of f_1 and f_2 are assumed to be less than 1, it cannot happen that both X_τ and Y_τ are on the same side $\partial_+ D$ or $\partial_- D$ of the boundary. Assume without loss of generality that $Y_\tau \in \partial_+ D$. Then $X_\tau \in D \cup \partial_- D$, $m_\tau = e'_2$ and $a_\tau = e'_1$. It follows from the definition of τ that for every $\varepsilon > 0$ there exists $t \in (\tau, \tau + \varepsilon)$ with $m_t \cdot e'_1 < 0$. Since a_t is continuous, we can find small $\varepsilon_0 > 0$ such that for $s \in (\tau, \tau + \varepsilon_0]$, (i) $a_s \cdot e'_1 \geq 0$; (ii) $a_s \cdot \nu(Y_s) \geq 0$ if $Y_s \in \partial D$; and (iii) $a_s \cdot \nu(X_s) \leq 0$ if $X_s \in \partial D$. In (ii) and (iii) we have used the fact that $Y_\tau \in \partial_+ D$ and $X_\tau \notin \partial_+ D$ and the assumption that the Lipschitz constants of f_k 's are less than 1. For all $t \in (\tau, \tau + \varepsilon_0]$,

$$\begin{aligned} m_t \cdot e'_1 &= m_\tau \cdot e'_1 + \int_\tau^t |V_s|^{-1} a_s \cdot e'_1 a_s \cdot (dM_s - dL_s) \\ &= \int_\tau^t |V_s|^{-1} a_s \cdot e'_1 a_s \cdot \nu(Y_s) d|M|_s - \int_\tau^t |V_s|^{-1} a_s \cdot e'_1 a_s \cdot \nu(X_s) d|L|_s. \end{aligned}$$

The first integral in the last line is nonnegative and the second one is nonpositive. This contradicts the fact that $m_t \cdot e'_1 < 0$ for some $t \in (\tau, \tau + \varepsilon_0)$ and so it shows that Ω_1 has probability zero. \square

3. CONDITIONED MIRROR COUPLINGS

Throughout this section, we fix a lip domain $D \subset \mathbb{R}^2$ and a sequence of lip domains D^n increasing to D , such that the defining functions f_1^n, f_2^n (cf. equation (1)) are C^2 -smooth and Lipschitz with the Lipschitz constants strictly less than 1. Recall that a lip domain is a Lipschitz domain, i.e., its boundary can be represented as the graph of a Lipschitz function with constant λ in some neighborhood of every boundary point (the point here is that the extreme left and right points have to be included). We will assume without loss of generality that there exist constants $\lambda < \infty$ and $r_0 > 0$, not depending on n , such that for every n and $x \in \partial D^n$, the set $\partial D^n \cap B_{r_0}(x)$ can be represented as the graph of a (single) function with Lipschitz constant λ . Moreover, the same applies to any $x \in \partial D$ and $\partial D \cap B_{r_0}(x)$. For each D^n , Corollary 2 and Propositions 1 and 2 apply.

Let $D_\varepsilon = \{x \in D : \text{dist}(x, \partial D) \geq \varepsilon\}$ and $D_\varepsilon^n = \{x \in D^n : \text{dist}(x, \partial D^n) \geq \varepsilon\}$. The hitting time of a set A by a process W will be denoted $T^W(A)$, or simply $T(A)$, if no confusion may arise.

Lemma 2. *Let $Q_{x,y}^n$ denote any probability measure under which (X, Y) is a strong Markov process, and X [respectively, Y] is a reflected Brownian motion in D^n starting from x [y]. Then there are constants $\delta, \sigma > 0$, such that for any $\rho > 0, n, x, y \in \overline{D^n}, |x - y| \geq \rho$,*

$$Q_{x,y}^n(T_0 \leq \sigma \rho^2) \geq 1/2,$$

where

$$T_0 = \inf\{t > 0 : X_t, Y_t \in D_{\delta\rho}^n, X_t \in B_{\rho/16}(x), Y_t \in B_{\rho/16}(y)\}.$$

Proof. The lemma will be proved in two steps.

Step 1. In this step, we will show that there exist $\gamma, \varepsilon_0 > 0$, such that for any $\varepsilon \in (0, \varepsilon_0), n$, and $x \in \overline{D^n}$,

$$(18) \quad Q_{x,y}^n(T^X(D_\varepsilon \cap B_{\gamma\varepsilon}(x)) < \varepsilon^2) > 9/10.$$

Recall the Lipschitz constant λ defined before the lemma and suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz with constant λ , and $f(0) = 0$. Let $A = \{x \in \mathbb{R}^2 : x_2 > f(x_1)\}$,

$$\begin{aligned} \Delta(x, r, h) &= \{y \in A : |y_1 - x_1| < r, y_2 < f(y_1) + h\}, \\ \partial_u \Delta(x, r, h) &= \{y \in \partial \Delta(x, r, h) : y_2 = f(y_1) + h\}, \\ \partial_s \Delta(x, r, h) &= \{y \in \partial \Delta(x, r, h) : |y_1 - x_1| = r\}. \end{aligned}$$

Let P_x^n be the distribution of the reflected Brownian motion in A starting from x . The function

$$g(y) = P_y^n(T^X(\partial_u \Delta((0, 0), 1, 1)) \leq T^X(\partial_s \Delta((0, 0), 1, 1)))$$

is harmonic in $\Delta((0, 0), 1, 1)$. By a Harnack inequality proved for such functions in [6], $g(y) \geq cg((0, 1/2))$ for some c depending only on λ and all $y \in \Delta((0, 0), 1/2, 1)$. Hence, for some $p > 0$ depending only on λ , the reflected Brownian motion starting from any point of $\Delta((0, 0), 1/2, 1)$ exits $\Delta((0, 0), 1, 1)$ through $\partial_u \Delta((0, 0), 1, 1)$ with probability greater than p . By the strong Markov property applied at the hitting time of $\partial \Delta((0, 0), 1, 1) \setminus \partial A$, the process starting from any point in $\Delta((0, 0), 1/2, 1)$

exits $\Delta((0, 0), 2, 1)$ through $\partial_u \Delta((0, 0), 2, 1)$ with probability greater than $1 - (1 - p)^2$. By induction, the process starting from a point of $\Delta((0, 0), 1/2, 1)$ exits $\Delta((0, 0), \gamma, 1)$ through $\partial_u \Delta((0, 0), \gamma, 1)$ with probability greater than $1 - (1 - p)^\gamma$, for integer $\gamma > 1$. Fix a γ so large that $1 - (1 - p)^\gamma > 1 - 1/20$.

By Theorem 2.3 of [6], $P_x^n(X_1 \in dy) \leq c_1 \exp(c_2|x - y|^2)dy$, where the constants c_1 and c_2 depend only on λ . Choose $h > 0$ so small that $P_x^n(X_1 \in \Delta((0, 0), \gamma, h)) < 1/20$ for $x \in \Delta((0, 0), 1/2, h)$. We see that the probability that the process starting from $\Delta((0, 0), 1/2, h)$ is still in $\Delta((0, 0), \gamma, h)$ at time 1 or exited it through $\partial_s \Delta((0, 0), \gamma, h)$ by that time is bounded by $1/20 + 1/20 = 1/10$. Now (18) follows by scaling.

Step 2. Consider small $\bar{\delta} > 0$, large $\bar{t} < \infty$, and finite stopping times

$$T'_1 = T^X(D_{\bar{\delta}\rho}^n \cap B_{\gamma\bar{\delta}\rho}(x)) \wedge \bar{t}, \quad T''_1 = T^Y(D_{\bar{\delta}\rho}^n \cap B_{\gamma\bar{\delta}\rho}(y)) \wedge \bar{t}, \quad T_1 = T'_1 \wedge T''_1.$$

Using Step 1 with $\varepsilon = \bar{\delta}\rho$, one has for large \bar{t} ,

$$(19) \quad Q_{x,y}^n(T'_1 < (\bar{\delta}\rho)^2) > 9/10,$$

and, similarly, $Q_{x,y}^n(T''_1 < (\bar{\delta}\rho)^2) > 9/10$. Let

$$C_{\bar{\delta},\eta} = \{X_t \in B_{\bar{\delta}\rho/2}(X_{T_1}), t \in [T_1, T_1 + \eta^2\bar{\delta}^2\rho^2/4]\}.$$

If $T'_1 \leq T''_1$, then $B_{\bar{\delta}\rho/2}(X_{T_1}) \subset D^n$ and by choosing $\eta \in (0, 1)$ small, one can have

$$Q_{x,y}^n(C_{\bar{\delta},\eta} | \mathcal{F}_{T_1}, T'_1 \leq T''_1) \geq 9/10.$$

Let $T_2 = \inf\{t > T_1 : Y_t \in D_{\eta\bar{\delta}\rho/2} \cap B_{\gamma\eta\bar{\delta}\rho/2}(Y_{T_1})\}$. Using Step 1 with $\varepsilon = \eta\bar{\delta}\rho/2$ and the strong Markov property,

$$Q_{x,y}^n(T_2 < T_1 + (\eta\bar{\delta}\rho)^2/4 | \mathcal{F}_{T_1}, T'_1 \leq T''_1) > 9/10.$$

Hence,

$$Q_{x,y}^n(C_{\bar{\delta},\eta}, T_2 < T_1 + (\eta\bar{\delta}\rho)^2/4 | \mathcal{F}_{T_1}, T'_1 \leq T''_1) > 8/10.$$

A similar formula holds in the case $T''_1 \leq T'_1$. Combining this with (19), we obtain

$$Q_{x,y}^n(T_0 \leq (\bar{\delta}\rho)^2 + (\eta\bar{\delta}\rho)^2/4) \geq (9/10)(8/10) > 1/2.$$

Now make $\bar{\delta} > 0$ smaller, if necessary, so that $(\gamma\bar{\delta}\rho + \bar{\delta}\rho/2) \vee (\gamma\bar{\delta}\rho + \gamma\eta\bar{\delta}\rho/2) < \rho/16$, and thus the lemma holds with $\delta = \eta\bar{\delta}/2$. \square

In the rest of this section, we denote by $P_{x,y}^n$ a measure under which (X, Y) is a mirror coupling of reflected Brownian motions in D^n . Let τ be the coupling time for X and Y . Following [9], we let $\mathcal{D}^n = \{(x, y) \in \overline{D^n} \times \overline{D^n} : x \neq y\}$, $\mathcal{D}^n(\varepsilon) = \{(x, y) \in \overline{D^n} \times \overline{D^n} : |x - y| \geq \varepsilon\}$, and $\widehat{\mathcal{D}}^n(\varepsilon) = \mathcal{D}^n \setminus \mathcal{D}^n(\varepsilon)$.

Lemma 3. (i) *There are constants $\alpha, c, a > 0$ such that for all $\varepsilon \in (0, a)$ and all $(x, y) \in \mathcal{D}^n(\varepsilon)$,*

$$P_{x,y}^n(T(\mathcal{D}^n(a)) < \tau) \geq c\varepsilon^\alpha.$$

(ii) *There is a constant $c < \infty$ such that for all $\varepsilon > 0$ and $(x, y) \in \widehat{\mathcal{D}}^n(\varepsilon)$,*

$$E_{x,y}^n T((\widehat{\mathcal{D}}^n(\varepsilon))^c) \leq c\varepsilon^2.$$

(iii) *For any $u > 0$ we can find $\varepsilon_0, c > 0$ such that $\mathcal{D}^1(2\varepsilon_0)$ has a nonempty interior and*

$$P_{x,y}^n(\tau > t, (X_t, Y_t) \in \mathcal{D}^n(\varepsilon_0)) \geq c,$$

for all n , $(x, y) \in \mathcal{D}^n(2\varepsilon_0)$, and $t \in [u/4, u]$.

Proof. To keep the notation simple, the superscript n will be dropped from $D^n, D_\varepsilon^n, P_{x,y}^n, \mathcal{D}^n, \mathcal{D}^n(\varepsilon)$ and $\widehat{\mathcal{D}}^n(\varepsilon)$ throughout this proof. All constants in all estimates will be tacitly assumed to be independent of n .

Let us outline the strategy of the proof. The main step in the proof of (i) will be to show that starting at any $x, y \in \overline{D}, x \leq y, |x - y| = \rho > 0$, the probability of hitting $\mathcal{D}(2\rho)$ before exiting \mathcal{D} is bounded below by some $p > 0$, independent of ρ and n . The main estimate in the proof of (ii) will be that the probability of exiting \mathcal{D} before time $c\rho^2$ is bounded below by some $p > 0$, where c and p are independent of ρ and n . These two estimates will be obtained by identifying events H, K and a time S , with the properties

$$(20) \quad H \subset \{S < \tau, |X_S - Y_S| \geq 2\rho\},$$

while

$$(21) \quad P_{x,y}(H) \geq p,$$

and

$$(22) \quad K \subset \{\tau \leq c\rho^2\}, \quad P_{x,y}(K) \geq p.$$

Recall the notation x^* and y^* for the unique points in \overline{D} for which $x^* \leq x \leq y^*$, all $x \in D$. See the beginning of this section for the definition of the Lipschitz constant λ characterizing D . Let ∂^+D [respectively, ∂^-D] denote the upper [lower] part of the boundary ∂D . Let $a > 0$ be so small that (I) whenever $B_{32a}(x)$ intersects both ∂^+D and ∂^-D , one has $|x - x^*| \wedge |x - y^*| \leq 64\lambda a$; (II) there is a coordinate system in which $D \cap B_{128\lambda a}(x^*)$ is equal to the set of points above the graph of a Lipschitz function with constant λ ; and (III) a similar statement holds for y^* . We will consider $x, y \in \overline{D}$ with $|x - y| = \rho$ and $\rho \in (0, a)$. There are three cases: (a) $B_{32\rho}(x)$ intersects either ∂^+D or ∂^-D , but not both, (b) $B_{32\rho}(x)$ intersects both parts of the boundary, (c) $B_{32\rho}(x)$ is contained in D . In case (c), the existence of H, K, S for which (20), (21) and (22) are valid follows from a simple scaling argument for the planar Brownian motion, and we therefore omit the details.

(a) From now on, we will refer to X and Y as “Brownian particles” or simply “particles.” In case (a) there are two possibilities according to whether the particles X and Y are close to the upper or lower boundary. We consider only the lower boundary, since the other case can be treated similarly.

Recall that (e_1, e_2) is the usual coordinate system in \mathbb{R}^2 . Within case (a) we consider three subcases: (a1) the angle θ , which $y - x$ forms with e_1 , is within $[-\pi/4, -\pi/16]$; (a2) $\theta \in [\pi/16, \pi/4]$; and (a3) $\theta \in (-\pi/16, \pi/16)$.

(a1) Assume x and y are as in case (a1). Let $\sigma, \delta > 0$ and let T_0 be as in Lemma 2, and let H_0 denote the event $\{T_0 \leq \sigma\rho^2\}$. By definition, on $H_0, X_{T_0}, Y_{T_0} \in D_{\delta\rho}$, and $X_{T_0} \in B_{\rho/16}(x), Y_{T_0} \in B_{\rho/16}(y)$. A simple calculation, using the fact that $\theta \in [-\pi/4, -\pi/16]$, shows that $X_{T_0} \cdot e_2 \geq Y_{T_0} \cdot e_2$. Define

$$A^X(x, s, v, \gamma, T) = \{|X_t - x - vt| < \gamma, t \in [s, T]\}.$$

If H_0 holds, then $Y_{T_0} - X_{T_0} \neq 0$ and we can define orthogonal vectors (ξ_1, ξ_2) as $\xi_1 = (Y_{T_0} - X_{T_0})/|Y_{T_0} - X_{T_0}|$ and ξ_2 is the unit vector orthogonal to ξ_1 for which $\xi_2 \cdot e_2 > 0$. We will define an event $H_1 \subset H_0$ which represents, intuitively speaking, a motion of X along a tube that starts from X_{T_0} and is parallel to the mirror. Namely,

$$H_1 = H_0 \cap A^X(X_{T_0}, T_0, \rho^{-1}\xi_2, h\rho, T_1),$$

where $T_1 = T_0 + 8\rho^2$, and $h = (\delta/2) \wedge (1/16)$. Note that on H_1 , neither X nor Y hits the boundary ∂D within the time interval $[T_0, T_1]$ —this follows easily from the fact that $B_{h\rho}(X_{T_0})$ and $B_{h\rho}(Y_{T_0})$ are contained in D and that, locally, D is the set of points above the graph of a Lipschitz function with constant 1. Also note that the fact that $X_{T_0} \cdot e_2 \geq Y_{T_0} \cdot e_2$ on H_0 implies that $\xi_2 \cdot e_1 \geq 0$, which shows that, on H_1 ,

$$(23) \quad X_{T_1} \cdot e_1 \geq X_{T_0} \cdot e_1 - h\rho.$$

On H_1 , the distance between the particles at time T_1 is within $[(1 - 1/8 - 2h)\rho, (1 + 1/8 + 2h)\rho]$. To make the distance larger, we consider an event H , on which X moves within a tube starting from X_{T_1} in the direction of $-e_1$. Specifically,

$$H = H_1 \cap A^X(X_{T_1}, T_1, -\rho^{-1}e_1, h\rho, T_2),$$

where $T_2 = T_1 + 5\rho^2$. Let $\Lambda \subset \mathbb{R}^2$ consist of two line segments—the first one starts from X_{T_0} and goes 8ρ units in the ξ_2 direction; the second one starts at the endpoint of the first one and goes 5ρ units in the $-e_1$ direction. If H_0 holds, then the distance of Λ from ∂D is greater than $h\rho$, because the boundary is locally Lipschitz with constant 1. Hence, if H occurs, then X does not hit the boundary within the time interval $[T_0, T_2]$.

Recall that the “mirror” is the line of symmetry between X_t and Y_t . We will now show that assuming H , the distance between the particles exceeds 2ρ at time T_2 or earlier than that. First, if $Y \in D$ on $[T_1, T_2]$, then within the time interval $[T_0, T_2]$ the mirror is fixed. Note that the angle of the mirror relative to e_1 is within $[\pi/4, \pi/2]$. Therefore X_{T_2} , which is inside $B_{h\rho}(X_{T_1} - 5\rho e_1)$, is at a distance of at least $5\rho/\sqrt{2} - h\rho > \rho$ from the mirror. Hence the two particles are at a distance greater than 2ρ from each other. If Y hits the boundary during $[T_1, T_2]$, let

$$U = \sup\{t \in [T_1, T_2] : Y_t \in \partial D\}.$$

There are two cases: (I) If $U \in [T_1, T_1 + 3\rho^2]$, then the mirror does not move within $[U, T_2]$. Recall that the angle that the mirror forms with the horizontal axis is always in the range from $\pi/4$ to $3\pi/4$. It follows that on the interval $[U, T_2]$, the distance from X_{T_2} to the mirror must grow to at least $2\rho/\sqrt{2} - 2h\rho > \rho$. Hence at time T_2 the particles are more than 2ρ units apart. (II) If $U \in (T_1 + 3\rho^2, T_2]$, then X_U is at a distance greater than $3\rho/\sqrt{2} - h\rho > 2\rho$ from the boundary, while Y_U is on the boundary, and again we find a time prior to T_2 when the particles are more than 2ρ apart. We conclude that there is an S satisfying (20). For a standard planar Brownian motion W , the probability of $\{|W_t - \rho^{-1}rt| \leq \gamma\rho, t \in [0, \rho^2T]\}$ is strictly positive by the “support theorem” and independent of ρ , by scaling. For H to happen, two events of this form have to happen, with X in place of W . This, the strong Markov property, and the fact that H_0 has probability greater than $1/2$ imply that (21) holds, with some $p > 0$ that is independent of ρ .

Next we discuss the event K which we define as

$$K = H_1 \cap \{\tau \leq T_1 + \rho^2\}.$$

It follows immediately from the definitions of H_0 and H_1 that on K one has $\tau \leq c\rho^2$. If H_1 occurs, then the convex hull of $B_{h\rho}(X_{T_1}) \cup B_{h\rho}(Y_{T_1})$ is contained in D . This, Brownian scaling and the strong Markov property easily imply that the probability that $\tau \leq T_1 + \rho^2$ conditional on \mathcal{F}_{T_1} is bounded below on H_1 by a positive constant independent of ρ . Since the probability of H_1 is itself bounded away from zero,

(22) follows. This completes the discussion of case (a1). Case (a2) can be treated in a similar fashion, by interchanging the roles of X and Y .

(a3) We let H and K be defined as in case (a1). A simple calculation shows that in this case, on H , the angle which ξ_1 forms with e_1 is within $[-\pi/8, \pi/8]$. Moreover, neither of the particles hits the boundary before time $T_1 + \rho^2$, and at this time the distance between the particles is greater than 2ρ . Hence (20) holds true. The validity of (21) and (22) can be proved as in case (a1).

(b) In case (b), both points x and y must be in one of the discs $B_{128a}(x^*)$ or $B_{128a}(y^*)$. We will assume the latter, without loss of generality. Let $(\tilde{e}_1, \tilde{e}_2)$ be a coordinate system in which D is locally the set of points above the graph of a Lipschitz function with constant λ . Let H_0 be defined as in case (a1). Let

$$H = H_0 \cap A^X(X_{T_1}, T_1, \rho^{-1}\tilde{e}_2, h\rho, T_2),$$

where $T_1 = T_0$, $T_2 = T_1 + 5\lambda\rho^2$. Then the analysis based on the last visit of Y to ∂D within $[T_1, T_2]$, carried out for case (a1), applies in the current case in an analogous way and shows that there is an S for which (20) holds. Again, (21) is a simple consequence of scaling.

Finally, we describe the event K for case (b). Let $\overline{y^*}$ denote the line segment emanating from y^* defined as $\{y^* + r\tilde{e}_2 : r \in [0, 129a]\}$. Then it is easy to see that if $x_1 \in \overline{y^*}$, then for any $x \in B_{\rho/16}(x_1)$, and $y \in D_{\delta\rho}$ with $y \geq x$, the convex hull of $B_{h\rho}(x) \cup B_{h\rho}(y)$ is contained in D . Let

$$K_0 = H_0 \cap A^X(X_{T_0}, T_0, \rho^{-1}r\tilde{e}_1, h\rho, \tilde{T}_1),$$

where $r = 1$ if $(X_{T_0} - y^*) \cdot \tilde{e}_1 < 0$, and $r = -1$ otherwise, and \tilde{T}_1 is the first hitting time of $\overline{y^*}$ by X after time T_0 . It is easy to check that

$$H_0 \cap A^X(X_{T_0}, T_0, \rho^{-1}r\tilde{e}_1, h\rho, T_0 + 3\rho^2) \subset K_0.$$

This and scaling show that the probability of K_0 is bounded below. Next, let \tilde{T}_2 be defined analogously to T_0 , namely,

$$\tilde{T}_2 = \inf\{t > \tilde{T}_1 : X_t, Y_t \in D_{\delta\rho}, X_t \in B_{\rho/16}(X_{\tilde{T}_1}), Y_t \in B_{\rho/16}(Y_{\tilde{T}_1})\}.$$

Lemma 2 and the strong Markov property imply that conditional on $\mathcal{F}_{\tilde{T}_1}$, one has $\tilde{T}_2 \leq \tilde{T}_1 + \sigma\rho^2$ on $\{\tilde{T}_1 < \tau\}$ with probability at least $1/2$. It follows that the event $K_1 = K_0 \cap \{\tilde{T}_2 \leq \tilde{T}_1 + \sigma\rho^2\}$ has probability bounded below by a positive constant independent of ρ . According to an earlier remark, under K_1 , one has that the convex hull of $B_{h\rho}(X_{\tilde{T}_2}) \cup B_{h\rho}(Y_{\tilde{T}_2})$ is contained in ρ . We conclude that

$$K = K_1 \cap \{\tau \leq \tilde{T}_2 + \rho^2\}$$

has probability bounded below by a constant independent of ρ . This completes the proof of (22).

Let us show how (20) and (21) imply part (i) of the lemma. Let $(x, y) \in \mathcal{D}(\varepsilon)$, and let $\rho = |x - y| \geq \varepsilon$. Then

$$P_{x,y}(T(\mathcal{D}(2\rho)) < \tau) \geq p,$$

and by the strong Markov property,

$$P_{x,y}(T(\mathcal{D}(2^k\rho)) < \tau) \geq p^k.$$

Taking $k = \lceil \log(a/\varepsilon) \rceil + 1$, we obtain

$$P_{x,y}(T(\mathcal{D}(a)) < \tau) \geq p(\varepsilon/a)^{-\log p},$$

which completes the proof of (i).

Consider part (ii) next. By (22), we have $P_{x,y}(\tau \geq c\rho^2) \leq 1 - p$. By the Markov property,

$$P_{x,y}(\tau \geq kc\rho^2) \leq (1 - p)^k,$$

and therefore for any x, y with $|x - y| = \rho \leq \varepsilon$,

$$E_{x,y}T(\widehat{\mathcal{D}}_\varepsilon^c) \leq E_{x,y}\tau \leq \sum_{k \geq 1} kc\rho^2(1 - p)^k = c_1\rho^2 \leq c_1\varepsilon^2.$$

Finally, we will prove part (iii) of the lemma. Suppose $t \in [u/4, u]$ and recall δ and σ from Lemma 2. Let $h = (\delta/2) \wedge (1/16)$ and suppose that $\varepsilon_0 > 0$ is so small that $\sigma(2\varepsilon_0)^2 < u/4$. Assume that $|x - y| = \rho \geq 2\varepsilon_0$. Then Lemma 2 and the strong Markov property applied at T_0 imply that

$$\begin{aligned} P_{x,y}(\tau > t, (X_t, Y_t) \in \mathcal{D}^n(\varepsilon_0)) &\geq P_{x,y}(\tau > u, (X_s, Y_s) \in \mathcal{D}^n(\varepsilon_0), s \in [u/4, u]) \\ &\geq P_{x,y}(T_0 \leq \sigma(2\varepsilon_0)^2, X_s \in B_{h\varepsilon_0}(X_{T_0}), Y_s \in B_{h\varepsilon_0}(Y_{T_0}), s \in [T_0, T_0 + u]) \\ &\geq (1/2)P_x(W_s \in B_{h\varepsilon_0}(x), s \in [0, u]) \\ &\geq c. \end{aligned}$$

Here W denotes a planar Brownian motion starting from x . The third inequality follows from the fact that X and Y do not visit the boundary within $[T_0, T_0 + u]$, if the event on the third line holds, because of the definition of T_0 given in Lemma 2. □

The following lemma is almost the same as Lemma 5.1 in [4]. We reproduce it here because it is one of the most important elements of our argument. The present form of the result is very close to Lemma 3.10 in [9]. The intuitive meaning of the lemma is that if we condition X and Y on not coupling before time $u > 0$, then the processes are likely to move apart for a considerable distance at time u . Hence, Lemma 4 below is a version of the parabolic boundary Harnack principle for the process (X, Y) .

Lemma 4. *For every $u > 0$ there exist constants $c_1, c_2 > 0$ such that for all n large and for all $x, y \in D^n$ with $x \geq y$ and $x \neq y$,*

$$P_{x,y}^n(|X_u - Y_u| > c_1|\tau > u) > c_2.$$

Proof. The proof of this lemma uses methods considerably different from those in the rest of the paper so it will be convenient to introduce some notation specific only to this proof. First of all recall the following convention from the proof of Lemma 3—the superscript n will be dropped from all pieces of notation. This should not cause any confusion because the constants in Lemma 3 do not depend on n .

We will write $Z = (X, Y)$ since the separate components of Z will play no role in this proof. The state space of Z is $\overline{D} \times \overline{D}$. Recall \mathcal{D} , $\mathcal{D}(\varepsilon)$ and $\widehat{\mathcal{D}}(\varepsilon)$ defined before Lemma 3. The distribution of Z starting from z and the corresponding expectation will be denoted P^z and E^z . Conditioning by a harmonic function h will be reflected in the notation by writing P_h^z and E_h^z . See [12] for the discussion of conditioned Brownian motion and [18] for conditioning of general Markov processes.

We will denote the space-time counterpart of Z by V . More precisely, if Z has law P^z , then the law of the space-time process $\{V_t = (Z_t, s - t), t \geq 0\}$ will be denoted $P^{z,s}$. The distribution of the space-time process conditioned by a parabolic

function g will be denoted $P_g^{z,s}$. By abuse of notation, $T(A)$ will denote the first hitting time of A for V as well as for Z .

Fix some small $\varepsilon_0 > 0$ such that Lemma 3 applies with ε_0 , and let $M = \mathcal{D}(\varepsilon_0)$ and $\mathcal{D}_1 = \widehat{\mathcal{D}}(2\varepsilon_0)$. Let $h(z) = P^z(T(\mathcal{D}(2\varepsilon_0)) < \tau)$ and $U_k = \{z \in \mathcal{D}_1 : h(z) \in [2^{k-1}, 2^k]\}$ for integer k .

By Lemma 3 (i), $U_k \subset \widehat{\mathcal{D}}(c_1 2^{k/\alpha})$, for some $c_1, \alpha > 0$. Then Lemma 3 (ii) shows that $\sup_{z \in U_k} E^z(T(U_k^c)) \leq c_2 2^{2k/\alpha}$. It follows that

$$\sum_{k=0}^{\infty} \sup_{z \in U_{-k}} E^z(T(U_{-k}^c)) < \infty.$$

An argument of Chung [11] (see also [4]) shows that for suitable c_3 ,

$$c_3 \sum_{k=0}^{\infty} \sup_{z \in U_{-k}} E^z(T(U_{-k}^c))$$

is an upper bound for $E_h^z(T(\mathcal{D}_1^c))$. It follows that for a suitable $c_4 > 0$ and every $z \in \mathcal{D}$,

$$(24) \quad P_h^z(T(\mathcal{D}_1^c) < u/4) > c_4.$$

Recall the discussion of space-time processes at the beginning of the proof. The function

$$(z, t) \mapsto g(z, t) = P^z(\tau > t)$$

is parabolic in $\mathcal{D} \times [0, \infty)$ with boundary values 1 on $\mathcal{D} \times \{0\}$ and 0 otherwise. Let g_1 be a parabolic function in $\mathcal{D} \times [0, \infty)$ which has the same boundary values as g except that $g_1(z, 0)$ is changed from 1 to δ for $z \in \mathcal{D} \setminus M$, where $\delta \in (0, 1)$ will be chosen later.

We will use the following notation convention in the next displayed formula. Suppose that $V_t = (Z_t, s - t)$ has the distribution $P^{x,s}$. Then $T_*(\mathcal{D}_1^c) = s - \inf\{t > 0 : V_t \in \mathcal{D}_1^c \times \mathbb{R}\}$. Note that, typically, $T_*(\mathcal{D}_1^c) < s$.

By Lemma 3 (iii), we have $g_1(z, s) > c_5$ for all $z \in \mathcal{D}(2\varepsilon_0)$ and $s \in [u/4, u]$, where c_5 does not depend on δ . We obviously have $h(y) \leq 1$ for all y . Let $h(x, s) = h(x)$. For $x \in \mathcal{D}_1$ and $s \geq u/2$ we have, by (24),

$$\begin{aligned} g_1(x, s) &\geq \int_{\substack{t \in [u/4, s] \\ y \in \partial \mathcal{D}(2\varepsilon_0)}} g_1(y, t) P^{x,s}(T_*(\mathcal{D}_1^c) \in dt, Z(T_*(\mathcal{D}_1^c)) \in dy) \\ &= \int_{\substack{t \in [u/4, s] \\ y \in \partial \mathcal{D}(2\varepsilon_0)}} \frac{h(x, s)}{h(y, t)} \frac{h(y, t)}{h(x, s)} g_1(y, t) P^{x,s}(T_*(\mathcal{D}_1^c) \in dt, Z(T_*(\mathcal{D}_1^c)) \in dy) \\ &= \int_{\substack{t \in [u/4, s] \\ y \in \partial \mathcal{D}(2\varepsilon_0)}} \frac{h(x, s)}{h(y, t)} g_1(y, t) P_h^{x,s}(T_*(\mathcal{D}_1^c) \in dt, Z(T_*(\mathcal{D}_1^c)) \in dy) \\ &\geq \int_{\substack{t \in [u/4, s] \\ y \in \partial \mathcal{D}(2\varepsilon_0)}} h(x, s) c_5 P_h^{x,s}(T_*(\mathcal{D}_1^c) \in dt, Z(T_*(\mathcal{D}_1^c)) \in dy) \\ &= h(x, s) c_5 P_h^{x,s}(T_*(\mathcal{D}_1^c) \in [u/4, s]) \\ &\geq h(x, s) c_5 c_4 = c_6 h(x, s) = c_6 h(x). \end{aligned}$$

Let $U_k^* = \{z \in \mathcal{D}_1 : h(z) \leq 2^k\}$ for integer k . We argue as in the case of U_k 's. By Lemma 3 (i), $U_k^* \subset \widehat{\mathcal{D}}(c_1 2^{k/\alpha})$, for some $c_1, \alpha > 0$. Lemma 3 (ii) shows that $\sup_{z \in U_k^*} E^z(T((U_k^*)^c)) \leq c_2 2^{2k/\alpha}$. It follows that

$$\sum_{k=0}^{\infty} \sup_{z \in U_{-k}^*} E^z(T((U_{-k}^*)^c)) < \infty.$$

Let

$$W_k = \{(z, s) : g_1(z, s) \in [2^k, 2^{k+1}], s \in [u/2, u]\},$$

$$W = \bigcup_{k=-\infty}^{k_1} W_k,$$

where $k_1 < 0$ will be chosen later. If $2^{-m} < c_6$, then $W_k \subset U_{k+m}^* \times [u/2, u]$. Let such m be fixed. Using the estimate of Chung [11], we obtain for small k_1 and all $z \in \mathcal{D}$,

$$\begin{aligned} E_{g_1}^{z,u}(T(W^c)) &\leq c_7 \sum_{k=-\infty}^{k_1} \sup_{(y,s) \in W_k} E^{y,s} T(W_k^c) \\ &\leq c_7 \sum_{k=-\infty}^{k_1} \sup_{y \in U_{k+m}^*} E^y T((U_{k+m}^*)^c) \\ &\leq c_2 c_7 \sum_{k=-\infty}^{k_1} 2^{2(k+m)/\alpha} =: R(k_1) < \infty. \end{aligned}$$

Choose k_1 so small that $R(k_1) < u/8$, and note that k_1 is independent of n and δ . Then for any $z \in \mathcal{D}$,

$$(25) \quad E_{g_1}^{z,u} T(W^c) < u/8.$$

Let

$$Q = \{(x, s) : g_1(x, s) \geq 2^{k_1}, s \in [u/2, u]\}.$$

Since the g_1 -process cannot exit $\mathcal{D} \times [0, \infty)$ through $\partial\mathcal{D} \times [0, \infty)$, (25) implies

$$(26) \quad P_{g_1}^{z,u}(T(Q) > u/4) < 1/2.$$

Now let $\delta = 2^{k_1-1}$. Since $0 \leq g_1 \leq 1$, the process $g_1(V_t)$ is a martingale under $P^{z,s}$, and $g_1(z, s) \geq 2^{k_1}$ for $(z, s) \in Q$, we see that there is at least a $2^{k_1-1}/2$ chance that V under $P^{z,s}$ will hit $M \times \{0\}$ before hitting any other part of $\partial(\mathcal{D} \times [0, \infty))$. Thus we have for $(z, s) \in Q$,

$$\begin{aligned} P_{g_1}^{z,s}[V_s \in M \times \{0\}] &= \int_M (g_1(y, 0)/g_1(z, s)) P^{z,s}[V_s \in dy, T(\partial(\mathcal{D} \times [0, \infty))) = s] \\ &\geq \int_M P^{z,s}[V_s \in dy, T(\partial(\mathcal{D} \times [0, \infty))) = s] \\ &\geq 2^{k_1-1}/2. \end{aligned}$$

This and (26) yield, by the strong Markov property, for all $z \in \mathcal{D}$,

$$P_{g_1}^{z,u}[V_u \in M \times \{0\}] \geq c_8 > 0.$$

The ratio of g and g_1 is bounded away from 0 and ∞ on the accessible boundary of $\mathcal{D} \times [0, \infty)$, so

$$P_g^{z,u} [V_u \in M \times \{0\}] \geq c_9 > 0$$

for all $z \in \mathcal{D}$. This is equivalent to the statement in the lemma. □

4. NEUMANN EIGENFUNCTIONS

We will consider lip domains $D \subset \mathbb{R}^2$ in this section.

Recall the coordinate systems (e_1, e_2) and (e'_1, e'_2) , and the partial order “ \leq ” on \mathbb{R}^2 , defined before Proposition 2. For a function $u \in C^1(D)$ write

$$\partial' u = \min \left\{ \frac{\partial u}{\partial e'_1}, \frac{\partial u}{\partial e'_2} \right\}.$$

Let

$$S = \{u \in C^1(D) : \partial' u(x) \geq 0, x \in D\},$$

$$\tilde{S} = \{u \in C^1(D) : \partial' u(x) > 0, x \in D\}.$$

Lemma 5. *Let D be a lip domain other than a rectangle. Consider a Neumann eigenfunction ψ in D corresponding to the second eigenvalue. If $\psi \in S$, then $\psi \in \tilde{S}$.*

Proof. We will first prove the following claim.

(A) There exists a nonempty ball $B \subset D$ and a constant $a > 0$ such that $\partial' \psi \geq a$ on B .

Assume that (A) does not hold. Since $\psi \in S$ and ψ is nonconstant, there is $x \in D$ where $\partial\psi(x)/\partial e'_i > 0$ for either $i = 1, 2$. Assume without loss of generality that $\partial\psi(x)/\partial e'_1 > 0$. The inequality holds in a neighborhood of x , by continuity of $\nabla\psi$. We have assumed that (A) fails, so there exists an open set where $\partial\psi/\partial e'_2 = 0$. Since ψ , as an eigenfunction, is real-analytic in D , $\partial\psi/\partial e'_2 = 0$ holds everywhere in D . Hence $\psi(x) = g(x_1)$, where x_1 refers to the first coordinate of $x = (x_1, x_2)$ in the coordinate system (e'_1, e'_2) . The only function of the form $\psi(x) = g(x_1)$ which satisfies the equation $\Delta\psi = -\mu_2\psi$ is $\psi(x) = \cos x_1$, upon appropriate translation and scaling of the coordinate system. As a generalized solution to the eigenfunction equation, ψ satisfies for any $f \in C^2(D)$,

$$\mu_2 \int_D f\psi dx + \int_D \psi\Delta f dx + \int_{\partial D} \psi \frac{\partial f}{\partial \nu} \sigma(dx) = 0,$$

where ν is the inward unit normal vector field, and σ is the surface measure on ∂D . The divergence theorem implies that

$$\mu_2 \int_D f\psi dx = \int_D \nabla f \cdot \nabla \psi dx.$$

We will use a test function of the form $f(x) = \theta(x_2)$. For any such function f , the right-hand side of the last formula is zero, and so we have

$$(27) \quad \int_D \cos x_1 \theta(x_2) dx = 0.$$

The eigenfunction ψ must vanish at some points of D . It follows from the periodicity of $\psi(x) = \cos x_1$ that we can assume without loss of generality that ψ vanishes on $\{x \in D : x_1 = \pi/2\}$. By Courant’s Nodal Line Theorem, ψ cannot vanish anywhere else in D . Hence, $D \subset \{x : -\pi/2 < x_1 < 3\pi/2\}$. Recall from Section 1 the left and right vertices of D , called x^* and y^* . Let α_1 [resp., α_2] denote the line through x^*

[resp., y^*], parallel to the e'_1 -axis. The fact that D is a lip domain which is not a rectangle implies that the projection onto the e'_1 -axis of $\alpha_1 \cap \partial D$ is not equal to that of $\alpha_2 \cap \partial D$. Hence, for either $i = 1$ or 2 , $\alpha_i \cap \partial D$ is not symmetric about the line through the point $(\pi/2, 0)$ parallel to e'_2 . Assume without loss of generality that there is no symmetry for $i = 1$. Then either $\int_{\alpha \cap D} \psi(x_1, x_2) dx_1 > 0$ for all lines α which intersect D and are sufficiently close to α_1 or the integral is strictly negative for all such α . Now let θ be such that $f(x) = \theta(x_2)$ is zero off an ε -neighborhood of α_1 , it is equal to 1 in a neighborhood of α_1 , and it is nonnegative in D . Then, if ε is small enough, $\int_D f\psi dx \neq 0$. This contradicts (27), and therefore (A) holds.

We will need the following result from [8] (Theorem 2.4) on “synchronous couplings.” Suppose $x, y \in D$. Then there is a probability space $(\Omega', \mathcal{F}', P'_{x,y})$ and a process (W, X, Y) such that X [respectively, Y] is a reflected Brownian motion in \overline{D} starting from x [y] and W is a Brownian motion on the filtration generated by (W, X, Y) , such that (X, Y) admits the following Skorohod representation:

$$X_t = x + W_t + \int_0^t \nu(X_s) d|L|_s, \quad Y_t = y + W_t + \int_0^t \nu(Y_s) d|M|_s,$$

where $|L|$ and $|M|$ are boundary local times for X and Y , respectively, and where ν is the inward normal vector field on ∂D . Theorem 2.4 of [8] shows that the synchronous coupling (X, Y) in D may be obtained as a limit of synchronous couplings in a sequence of polygonal lip domains D_k approximating D . For polygonal lip domains, the following “order preserving property” has been proved in [3], and so it holds for the coupling (X, Y) in D ; if $x \leq y$, then $X_t \leq Y_t$ for all $t \geq 0$, $P'_{x,y}$ -a.s. We have for any $t > 0$,

$$e^{-\mu_2 t} \psi(x) = E'_{x,y} \psi(X_t), \quad e^{-\mu_2 t} \psi(y) = E'_{x,y} \psi(Y_t).$$

Let $B = B_\rho(x_0)$ be a ball satisfying claim (A) proved at the beginning of the proof, and let $\tilde{B} = B_{\rho/2}(x_0)$. Let $x \in D$, $y = x + re'_1$ and $T = \inf\{t : \text{dist}(X_t, \partial D) < \text{dist}(x, \partial D)/2\}$. Note that $\psi(Y_1) - \psi(X_1) \geq 0$ with probability 1, because $X_1 \leq Y_1$ and $\psi \in S$. We have $Y_1 - X_1 = Y_0 - X_0$ if $T > 1$. Hence for all $r \in (0, \rho/2)$ such that $y \in D$,

$$\begin{aligned} r^{-1}(\psi(y) - \psi(x)) &\geq r^{-1} e^{\mu_2} E'_{x,y} \left[(\psi(Y_1) - \psi(X_1)) 1_{\{X_1 \in \tilde{B}, T > 1\}} \right] \\ &\geq (e^{\mu_2} a/2) P'_{x,y}(X_1 \in \tilde{B}, T > 1), \end{aligned}$$

where $a > 0$ is the constant in (A). Since the right-hand side is positive and independent of r , it follows that $\partial\psi(x)/\partial e'_1 > 0$. A similar argument applies to $\partial\psi/\partial e'_2$, and we conclude that $\partial'\psi > 0$ on D . □

Recall that $D_\varepsilon = \{x \in D : \text{dist}(x, \partial D) \geq \varepsilon\}$.

Lemma 6. *Suppose D is a lip domain other than a rectangle. For any $\varepsilon_1 > 0$ such that the interior of D_{ε_1} is nonempty, and any $\delta, b > 0$, there exists $\varepsilon_2 > 0$ with the following property. If ψ is a Neumann eigenfunction corresponding to μ_2 with*

$$\begin{cases} \partial'\psi \geq \delta & \text{on } D_{\varepsilon_1}, \\ \partial'\psi \geq 0 & \text{on } D_{\varepsilon_2}, \\ |\psi| \leq b & \text{on } D, \end{cases}$$

then

$$\partial' \psi \geq 0 \quad \text{on } D.$$

Proof. Let E_x be the expectation corresponding to the distribution of reflected Brownian motion in D , starting from x . Then for any $t \geq 0$, and $x, y \in \overline{D}$,

$$e^{-\mu_2 t}(\psi(x) - \psi(y)) = E_x \psi(X_t) - E_y \psi(Y_t).$$

Hence, it will suffice to show that for all $x, y \in D$, $x \geq y$,

$$E_x \psi(X_2) - E_y \psi(Y_2) \geq 0.$$

Recall the domains D^n from Section 3. We will denote by $P_{x,y}^n$ a measure under which (X, Y) is a pair of reflected Brownian motions in D^n , starting from (x, y) , which is mirror coupled on the time interval $[0, 1)$ and synchronously coupled on $[1, 2]$ (see the proof of Lemma 5 for the definition of a synchronous coupling).

If $x \in D$, then the distributions of X under $P_{x,y}^n$ converge weakly to the distribution of the reflected Brownian motion in D in the uniform topology on $C([0, 2], \mathbb{R}^2)$, as $n \rightarrow \infty$, by Theorem 2 of [7]. Since a similar remark applies to Y , it is enough to show that there exists $\varepsilon_2 > 0$ such that for all large n ,

$$E_{x,y}^n(\psi(X_2) - \psi(Y_2)) \geq 0.$$

Denote the coupling time by τ . Since $X_2 = Y_2$ on $\{\tau \leq 1\}$, it suffices to show that

$$(28) \quad E_{x,y}^n[\psi(X_2) - \psi(Y_2) \mid \tau > 1] \geq 0.$$

By the uniform continuity of transition probabilities in D^n 's proved in Theorem 2.1 of [8],

$$(29) \quad P_{x,y}^n(X_1 \in C) \leq c|C|,$$

for any Borel subset C of \overline{D} , where $|C|$ denotes the Lebesgue measure of C and c does not depend on x, C and n . Let c_1 and c_2 be the constants from the statement of Lemma 4. We can assume that $\varepsilon_2 < \varepsilon_1$. By Proposition 2, $X_2 \geq Y_2$ a.s. Recall that $\partial' \psi \geq 0$ on D_{ε_2} . This implies that $\psi(X_2) - \psi(Y_2) \geq 0$ on $\{X_2, Y_2 \in D_{\varepsilon_2}\}$. By (29),

$$\begin{aligned} E_{x,y}^n(\psi(X_2) - \psi(Y_2) \mid \tau > 1) &\geq E_{x,y}^n[(\psi(X_2) - \psi(Y_2))1_{\{X_2, Y_2 \in D_{\varepsilon_2}\}} \mid \tau > 1] \\ &\quad - 2b[P_{x,y}(X_2 \notin D_{\varepsilon_2} \mid \tau > 1) + P_{x,y}(Y_2 \notin D_{\varepsilon_2} \mid \tau > 1)] \\ &\geq E_{x,y}^n[(\psi(X_2) - \psi(Y_2))1_{\{X_2, Y_2 \in D_{\varepsilon_1}, |X_2 - Y_2| > c_1/2\}} \mid \tau > 1] - c\varepsilon_2. \end{aligned}$$

Since $\partial' \psi \geq \delta$ on D_{ε_1} , we have $\psi(X_2) - \psi(Y_2) \geq \delta c_1/2$ on $\{X_2, Y_2 \in D_{\varepsilon_1}, |X_2 - Y_2| > c_1/2\}$, so

$$E_{x,y}^n(\psi(X_2) - \psi(Y_2) \mid \tau > 1) \geq cP_{x,y}^n(X_2, Y_2 \in D_{\varepsilon_1}, |X_2 - Y_2| > c_1/2 \mid \tau > 1) - c\varepsilon_2.$$

It is clear that Lemma 4 remains valid if we replace c_1 in that lemma with any smaller constant. Assume that $c_1 > 0$ is so small that there exists $z \in D$ such that $B_{2c_1}(z) \subset D_{\varepsilon_1}$. An easy argument based on Lemma 2 and the strong Markov property shows that if $|X_1 - Y_1| > c_1$, then with some probability $p > 0$, independent of n , we have $X_2, Y_2 \in D_{\varepsilon_1}$ and $|X_2 - Y_2| > c_1/2$. Hence,

$$\begin{aligned} E_{x,y}^n(\psi(X_2) - \psi(Y_2) \mid \tau > 1) &\geq cP_{x,y}^n[|X_1 - Y_1| > c_1 \mid \tau > 1] - c\varepsilon_2 \\ &\geq cc_2 - c\varepsilon_2, \end{aligned}$$

where the last inequality follows from Lemma 4. Note that c_2 and the other constants in the last formula do not depend on x and y . Taking $\varepsilon_2 > 0$ small, (28) follows, and Lemma 6 is proved. \square

Proof of Theorem 1. One can easily verify that both parts of the theorem hold in rectangles so we will assume that D is not a rectangle.

(i) The argument is similar to that in the proof of Theorem 4.1 in [2]. Arguing by contradiction, assume that μ_2 is multiple.

Recall from the proof of Lemma 5 that there exists a synchronous coupling of reflecting Brownian motions in D with the property that $X_t \leq Y_t$ for all $t \geq 0$, if $X_0 \leq Y_0$. This and the argument given in the proof of Theorem 3.3 of [3] show that at least one of the eigenfunctions corresponding to μ_2 belongs to S . Fix one of these eigenfunctions and call it ϕ .

We have assumed that μ_2 is multiple so there exists an eigenfunction ϕ^\perp which is orthogonal to ϕ . Since ϕ^\perp and $-\phi^\perp$ cannot both be in S , we can assume without loss of generality that $\phi^\perp \notin S$. Let

$$\phi^a = (1 - a)\phi + a\phi^\perp, \quad a \in [0, 1],$$

and

$$a^* = \inf\{a \in [0, 1] : \phi^a \notin S\}.$$

We claim that $a^* < 1$. Since $\partial' \phi^a \geq 0$ on D for $a < a^*$ and $\partial' \phi^a \rightarrow \partial' \phi^{a^*}$ pointwise in D when $a \uparrow a^*$, we have that $\phi^{a^*} \in S$. Hence a^* cannot be equal to 1. Therefore there is a sequence a_k with

$$(30) \quad a_k \downarrow a^*, \quad a_k \in (a^*, 1), \quad \phi^{a_k} \notin S.$$

For $a \in (a^*, 1)$ let

$$\varepsilon(a) = \sup\{\text{dist}(x, \partial D) : \partial' \phi^a(x) < 0\}.$$

Note that as $a \downarrow a^*$, $\partial'(\phi^a) \rightarrow \partial'(\phi^{a^*})$ uniformly on compact subsets of D . Since $\phi^{a^*} \in S$, Lemma 5 implies that $\phi^{a^*} \in \tilde{S}$. Therefore for any compact set C , we have $\partial'(\phi^a) > 0$ on C for all a sufficiently close to a^* . Hence

$$(31) \quad \lim_{a \downarrow a^*} \varepsilon(a) = 0.$$

Let $\varepsilon_1 > 0$ be as in Lemma 6. Using again the facts that the convergence $\partial' \phi^a \rightarrow \partial' \phi^{a^*}$ is uniform on compacts and that $\phi^{a^*} \in \tilde{S}$, we have that there are constants $\delta > 0$ and $a_1 \in (a^*, 1)$ such that $\partial' \phi^a \geq \delta$ on D_{ε_1} for all $a \in (a^*, a_1)$. Since by definition, $\partial' \phi^a \geq 0$ on $D_{\varepsilon(a)}$ for all $a \in (a^*, a_1)$, Lemma 6 and (31) show that $\phi^a \in S$ for all $a \in (a^*, a_2)$, where $a_2 \in (a^*, a_1)$. This gives a contradiction to (30). As a result, the eigenvalue is simple, and Theorem 1 (i) follows.

(ii) We have shown in the first part of the proof that there is only one eigenfunction corresponding to μ_2 and that it belongs to S . This immediately implies part (ii) of the theorem. \square

ACKNOWLEDGMENTS

We would like to express our gratitude to the following colleagues for very useful advice: R. Bañuelos, R. Bass, Z. Chen, S. Jacka, D. Jerison, C. Kenig, J.-B. Lasserre, T. Pakes, P. Pollett, and R. Tweedie.

REFERENCES

- [1] O. Adelman, K. Burdzy and R. Pemantle (1998). Sets avoided by Brownian motion. *Ann. Probab.* 26, 429–464. MR **99i**:60152
- [2] R. Atar (2001). Invariant wedges for a two-point reflecting Brownian motion and the “hot spots” problem. *Elect. J. of Probab.* 6, paper 18, 1–19. MR **2002i**:60146
- [3] R. Bañuelos and K. Burdzy (1999). On the “hot spots” conjecture of J. Rauch. *J. Funct. Anal.* 164, 1–33. MR **2000m**:35085
- [4] R. Bass and K. Burdzy (1992). Lifetimes of conditioned diffusions. *Probab. Theory Related Fields* 91, 405–443. MR **93e**:60155
- [5] R. Bass and K. Burdzy (2000). Fiber Brownian motion and the ‘hot spots’ problem. *Duke Math. J.* 105, 25–58. MR **2001g**:60190
- [6] R. F. Bass and E. P. Hsu (1991). Some potential theory for reflecting Brownian motion in Hölder and Lipschitz domains. *Ann. Probab.* 19, 486–508. MR **92i**:60142
- [7] K. Burdzy and Z.-Q. Chen (1998). Weak convergence of reflecting Brownian motions. *Elect. Comm. in Probab.* 3, 29–33. MR **99d**:60091
- [8] K. Burdzy and Z.-Q. Chen (2002). Coalescence of synchronous couplings. *Probab. Theory Related Fields* **123**, 553–578. MR **2003e**:60184
- [9] K. Burdzy and W. S. Kendall (2000). Efficient Markovian couplings: examples and counterexamples. *Ann. Appl. Probab.* 10, 362–409. MR **2002b**:60129
- [10] K. Burdzy and W. Werner (1999). A counterexample to the “hot spots” conjecture. *Ann. Math.* 149, 309–317. MR **2000b**:35044
- [11] K. L. Chung (1984). The lifetime of conditioned Brownian motion in the plane. *Ann. Inst. Henri Poincaré* 20, 349–351. MR **86d**:60088
- [12] J. L. Doob. *Classical Potential Theory and Its Probabilistic Counterpart*. Springer-Verlag, New York, 1984. MR **85k**:31001
- [13] R. Hempel, L. A. Seco and B. Simon (1991). The essential spectrum of Neumann Laplacians on some bounded singular domains. *J. Func. Anal.* 102, 448–483. MR **93h**:35144
- [14] D. Jerison and N. Nadirashvili (2000). The “hot spots” conjecture for domains with two axes of symmetry. *J. Amer. Math. Soc.* 13, 741–772. MR **2001f**:35110
- [15] I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus*, second edition, Springer, New-York, 1991. MR **92h**:60127
- [16] B. Kawohl. *Rearrangements and Convexity of Level Sets in PDE*. Lecture Notes in Mathematics 1150, Springer, Berlin, 1985. MR **87a**:35001
- [17] W. S. Kendall (1989). Coupled Brownian motions and partial domain monotonicity for the Neumann heat kernel. *J. Funct. Anal.* 86, 226–236. MR **90m**:58203
- [18] M. Sharpe. *General Theory of Markov Processes*. Academic Press, New York, 1989. MR **89m**:60169
- [19] P. L. Lions and A. S. Sznitman (1984). Stochastic differential equations with reflecting boundary conditions. *Comm. Pure appl. Math.* 37, 511–537. MR **85m**:60105
- [20] M. Pascu (2002). Scaling coupling of reflecting Brownian motions and the hot spots problem. *Trans. Amer. Math. Soc.* **354**, 4681–4702. MR **2003i**:60141
- [21] P. Protter. *Stochastic Integration and Differential Equations. A New Approach*. Springer-Verlag, Berlin (1990). MR **91i**:60148
- [22] F.-Y. Wang (1994). Application of coupling methods to the Neumann eigenvalue problem. *Probab. Theory Related Fields* 98, 299–306. MR **94k**:58153

DEPARTMENT OF ELECTRICAL ENGINEERING, TECHNION–ISRAEL INSTITUTE OF TECHNOLOGY,
HAIFA 32000, ISRAEL

E-mail address: atar@ee.technion.ac.il

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON 98195-
4350

E-mail address: burdzy@math.washington.edu