

ERRATUM TO “REAL BOUNDS, ERGODICITY
AND NEGATIVE SCHWARZIAN FOR MULTIMODAL MAPS”

SEBASTIAN VAN STRIEN AND EDSON VARGAS

In Part 1 of Theorem C of the paper *Real bounds, ergodicity and negative Schwarzian for multimodal maps*, see [1], the assumption that V is nice was, by mistake, omitted. We would like to thank Weixiao Shen for pointing this out. The correct version of Theorem C(1) is as follows:

Theorem C(1) (Improved Macroscopic Koebe Principle). *Assume that $f: M \rightarrow M$ is contained in $\mathcal{A}^{1+\text{Zygmund}}$. Then for each $\xi > 0$, there exists $\xi' > 0$ such that if I is a nice interval, V is nice and ξ -well-inside I and $x \in I$, $f^k(x) \in V$ (with $k \geq 1$ not necessarily minimal), then the pullback of V along $\{x, \dots, f^k(x)\}$ is ξ' -well-inside the return domain to I containing x .*

Here, as before, we define an open interval K to be nice if no iterate of ∂K enters K . This implies that if K_1 and K_2 are pullbacks of K , then they are either disjoint or nested.

In Lemma 9 (page 762) it was implicitly assumed that V is disjoint from J_n . It is for this reason that the proof of Theorem C(1) does not work unless we assume V is nice (or something similar). The proof of Theorem C(1) as stated above is essentially the same as before, using Lemma 6' below instead of Lemma 6; then in Lemma 9 (page 762) we do not need to require that k_{n+1} is a jump time provided we assume that V is nice. Making the additional assumption that V is nice, Proposition 1 (and its proof) and the rest of the paper go through unchanged.

Lemma 6'. *For each $\rho > 0$ sufficiently small, there exists $\delta_3 > 0$ such that if I is a ρ -scaled neighbourhood of a nice interval $V \subset I$, then J is a δ_3 -scaled neighbourhood of any component A of $\phi_{|J}^{-k}(V)$ (where $k \geq 1$ is arbitrary).*

Proof. Let V_i , $i = 0, \dots, k$ be the component of $\phi_{|J}^{-(k-i)}(V)$ containing $\phi^i(A)$. Of course, we may assume that k is large and that V_0, \dots, V_k are disjoint.

Claim. There exists $\alpha > 0$ such that if $0 \leq j < k$ and V_{j+1} is contained in a neighbourhood of V_j of size $(1 + \alpha)|V_j|$, then V_j is α -well-inside I . Similarly, if $0 \leq j < k - 1$ and V_j lies between V_{j+1} and V_{j+2} , then V_j is α -well-inside I .

Proof of Claim. If V_{j+1} is contained in a neighbourhood of V_j of size $(1 + \alpha)|V_j|$ and α is small enough, then ϕ' is close to zero on a definite neighbourhood of V_j . So V_j is contained in the basin of an attracting fixed point with multiplier close to zero. Since V_k is nice and δ_3 -well-inside I , we easily get that V_j is δ'_3 -well-inside

Received by the editors February 18, 2006.

2000 *Mathematics Subject Classification.* Primary 37Exx, 37Fxx.

Key words and phrases. Dynamical systems, interval dynamics, holomorphic dynamics.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

I , proving the first part of the claim. The second part of the claim follows in the same way by first applying part 1 to V_j, V_{j+1} and then applying it to V_j, V_{j+2} while considering ϕ^2 instead of ϕ , completing the proof of the claim.

If both sides of J are small, then $|\phi'|$ is bounded on J . There are three possibilities.

(a) V_1 lies between V_2 and V_3 , in which case, by the second part of the Claim, V_1 is well-inside I .

(b) V_2 lies between V_1 and V_3 ; in this case since $|V_1|/|V_2|$ is not small and by the first part of the Claim, V_2 is well-inside I .

(c) V_3 lies between V_1 and V_2 ; then, because $|V_1|/|V_2|$ and $|V_2|/|V_3|$ are not small, V_3 is well-inside I . In all cases, we get that V_0 is well-inside J .

So assume one of the sides, say the right side, of J is not small. Let $1 \leq j \leq k$ be the largest integer so that V_1, \dots, V_j are all not α -well-inside I . By taking $\alpha > 0$ small, we may assume $j \leq k - 2$. Since V_{j+1} is α -well-inside I , we may assume that $j \geq 1$ and that there exists $\alpha' > 0$ so that V_j, V_{j-1} are α' -well-inside $J \subset I$. By the claim, for each $i = 1, \dots, j$, V_i has an α -small and an α -big side, and V_{i+1} is contained in the α -big side. Since the right side of J is not small, V_1, \dots, V_{j+1} lie therefore ordered from left to right. If for each $i = 1, \dots, j-1$, V_{i-1} is contained in a β -scaled neighbourhood of V_i , then V_1 is in a $(\beta + \beta^2 + \dots + \beta^j)$ -scaled neighbourhood of V_{j-1} . So taking $\beta \in (0, 1)$ so small that $\beta/(1 - \beta) < \alpha'/2$, then, because V_{j-1} is α' -well-inside I , the left component of $I \setminus V_1$ has at least size $\frac{\alpha'}{2}|V_{j-1}| > \frac{\alpha'}{2}|V_1|$, i.e., V_1 is well-inside I , and V_0 is well-inside J . Hence we may assume there exists $i \in \{1, \dots, j-1\}$ so that V_{i-1} is not contained in a β -scaled neighbourhood of V_i . This and the first part of the Claim imply that V_i is well-inside the convex hull $H_i := [V_{i-1}, V_{i+1}]$ of V_{i-1} and V_{i+1} . Because the intervals V_1, \dots, V_{i+2} lie ordered, it follows that the pullback of H_i along V_1, \dots, V_i has intersection multiplicity at most 4 and therefore that V_1 is well-inside I . This again gives that V_0 is well-inside J . (This method of proof can also be used to provide a slightly shorter proof of Lemma 5.) \square

REFERENCE

- [1] S. van Strien, E. Vargas, *Real bounds, ergodicity and negative Schwarzian for multimodal maps*, Journal of the American Mathematical Society, **17**, 2004, 749–782. MR2083467 (2005i:37043)

DEPARTMENT OF MATHEMATICS, WARWICK UNIVERSITY, COVENTRY CV4 7AL, ENGLAND
E-mail address: strien@maths.warwick.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SÃO PAULO, SÃO PAULO, BRAZIL
E-mail address: vargas@ime.usp.br