

$\mathcal{B}(\ell^p)$ IS NEVER AMENABLE

VOLKER RUNDE

INTRODUCTION

The theory of amenable Banach algebras begins with B. E. Johnson’s memoir [Joh 1]. The choice of terminology comes from [Joh 1, Theorem 2.5]: a locally compact group is amenable (in the usual sense) if and only if its L^1 -algebra satisfies a certain cohomological triviality condition, which is then used to define the class of amenable Banach algebras.

The notion of Banach algebraic amenability has turned out to be extremely fruitful ever since the publication of [Joh 1]. The class of amenable Banach algebras is small enough to allow for strong theorems of a general nature, yet large enough to encompass a diverse collection of examples, such as the nuclear C^* -algebras (see [Run 2, Chapter 6] for an account), Banach algebras of compact operators on certain well-behaved Banach spaces ([G–J–W]), and even radical Banach algebras ([Run 1] and [Rea 1]).

The Memoir [Joh 1] concludes with a list of suggestions for further research. One of them ([Joh 1, 10.4]) asks:

Is $\mathcal{B}(E)$ —the Banach algebra of all bounded linear operators on a Banach space E —ever amenable for any infinite-dimensional E ?

From a philosophical point of view, the answer to this question ought to be a clear “no”. As for groups, amenability for Banach algebras can be viewed as a weak finiteness condition: amenable Banach algebras tend to be “small”—whatever that may mean precisely—and $\mathcal{B}(E)$ simply feels too “large” to be amenable. It seems, however, as if Johnson’s question has recently—somewhat surprisingly—found a positive answer: in [A–H], S. A. Argyros and R. G. Haydon construct an infinite-dimensional Banach space E with few bounded linear operators, i.e., $\mathcal{B}(E) = \mathcal{K}(E) + \mathbb{C} \text{id}_E$ (with $\mathcal{K}(E)$ denoting the compact operators on E). As H. G. Dales pointed out to the author, E has property (A) introduced in [G–J–W], so that $\mathcal{K}(E)$ is an amenable Banach algebra for this space, as is, consequently, $\mathcal{B}(E)$.

Still, infinite-dimensional Banach spaces E with $\mathcal{B}(E)$ amenable ought to be the exception rather than the rule. Indeed, it follows from work by S. Wassermann ([Was]) and the equivalence of amenability and nuclearity for C^* -algebras that $\mathcal{B}(\ell^2)$ cannot be amenable. With ℓ^2 being the “best behaved” of all ℓ^p -spaces, one is led to expect that $\mathcal{B}(\ell^p)$ fails to be amenable for all $p \in [1, \infty]$. However, until recently

Received by the editors July 4, 2009 and, in revised form, December 5, 2009, December 7, 2009, and December 8, 2009.

2010 *Mathematics Subject Classification*. Primary 47L10; Secondary 46B07, 46B45, 46H20.

Key words and phrases. Amenability, Kazhdan’s property (T), \mathcal{L}^p -spaces.

The author’s research was supported by NSERC.

©2010 American Mathematical Society
Reverts to public domain 28 years from publication

it was not known for any $p \in [1, \infty]$ other than 2 whether or not $\mathcal{B}(\ell^p)$ is amenable. The first to establish the non-amenability of $\mathcal{B}(\ell^p)$ for any $p \in [1, \infty] \setminus \{2\}$ was C. J. Read in [Rea 2], where he showed that $\mathcal{B}(\ell^1)$ is not amenable. Subsequently, Read's proof was simplified by G. Pisier ([Pis]). Eventually, N. Ozawa simplified Pisier's argument even further and gave a proof that simultaneously establishes the non-amenability of $\mathcal{B}(\ell^p)$ for $p = 1, 2, \infty$ ([Oza]).

In [D–R], M. Daws and the author investigated the question of whether $\mathcal{B}(\ell^p)$ is amenable for $p \in (1, \infty) \setminus \{2\}$. Among the results obtained in [D–R] are the following: If $\mathcal{B}(\ell^p)$ is amenable, then so are $\ell^\infty(\mathcal{B}(\ell^p))$ and $\ell^\infty(\mathcal{K}(\ell^p))$. The amenability of $\ell^\infty(\mathcal{K}(\ell^p))$, in turn, forces $\ell^\infty(\mathcal{K}(E))$ to be amenable for every infinite-dimensional \mathcal{L}^p -space in the sense of [L–P]; in particular, if $\mathcal{B}(\ell^p)$ is amenable, then so is $\ell^\infty(\mathcal{K}(\ell^2 \oplus \ell^p))$.

This last implication is the starting point of this paper. Through a modification of Ozawa's approach from [Oza], we show that $\ell^\infty(\mathcal{K}(\ell^2 \oplus E))$ is not amenable for certain Banach spaces E , including $E = \ell^p$ for all $p \in (1, \infty)$. As a consequence, $\mathcal{B}(\ell^p)$ cannot be amenable for such p (and neither can $\mathcal{B}(L^p[0, 1])$). Together with the results from [Rea 2] and [Oza], this proves that $\mathcal{B}(\ell^p)$ is not amenable for any $p \in [1, \infty]$.

1. AMENABLE BANACH ALGEBRAS

The original definition of an amenable Banach algebra from [Joh 1] is given in terms of first-order Hochschild cohomology. An equivalent, but more intrinsic characterization—through approximate and virtual diagonals—was given soon thereafter in [Joh 2]. For the work done in this paper, however, yet another equivalent characterization of amenability due to A. Ya. Helemskiĭ turns out to be best suited ([Hel, Theorem VII.2.20]).

We denote the algebraic tensor product by \otimes and use the symbol $\hat{\otimes}$ for the projective tensor product of Banach spaces. If \mathfrak{A} and \mathfrak{B} are Banach algebras, then so is $\mathfrak{A} \hat{\otimes} \mathfrak{B}$ in a canonical fashion. For any Banach algebra \mathfrak{A} , we use \mathfrak{A}^{op} for its *opposite algebra*, i.e., the underlying Banach space is the same as for \mathfrak{A} , but multiplication has been reversed. Multiplication in \mathfrak{A} induces a bounded linear map $\Delta: \mathfrak{A} \hat{\otimes} \mathfrak{A} \rightarrow \mathfrak{A}$; it is immediate that $\ker \Delta$ is a left ideal in $\mathfrak{A} \hat{\otimes} \mathfrak{A}^{\text{op}}$.

Definition 1.1. A Banach algebra \mathfrak{A} is said to be amenable if

- (a) \mathfrak{A} has a bounded approximate identity and
- (b) the left ideal $\ker \Delta$ of $\mathfrak{A} \hat{\otimes} \mathfrak{A}^{\text{op}}$ has a bounded right approximate identity.

Definition 1.1 makes the proof of the following lemma, which we will require later on, particularly easy:

Lemma 1.2. *Let \mathfrak{A} be an amenable Banach algebra, and let $e \in \mathfrak{A}$ be an idempotent. Then, for any $\epsilon > 0$ and any finite subset F of $e\mathfrak{A}e$, there are $a_1, b_1, \dots, a_r, b_r \in \mathfrak{A}$ such that*

$$(1) \quad \sum_{k=1}^r a_k b_k = e$$

and

$$(2) \quad \left\| \sum_{k=1}^r x a_k \otimes b_k - a_k \otimes b_k x \right\|_{\mathfrak{A} \hat{\otimes} \mathfrak{A}} < \epsilon \quad (x \in F).$$

Proof. Let \bullet denote the product in $\mathfrak{A} \hat{\otimes} \mathfrak{A}^{\text{op}}$. Since $F \subset e\mathfrak{A}e$, we have that $\{x \otimes e - e \otimes x : x \in F\} \subset \ker \Delta$. By Definition 1.1(b), there is $\mathbf{r} \in \ker \Delta$ such that

$$(3) \quad \|(x \otimes e - e \otimes x) \bullet \mathbf{r} - (x \otimes e - e \otimes x)\| < \epsilon.$$

Set $\mathbf{d} := e \otimes e - (e \otimes e) \bullet \mathbf{r}$. Without loss of generality, we can suppose that $\mathbf{d} \in \mathfrak{A} \otimes \mathfrak{A}$, i.e., there are $a_1, b_1, \dots, a_r, b_r \in \mathfrak{A}$ with $\mathbf{d} = \sum_{j=1}^r a_j \otimes b_j$. From the definition of \mathbf{d} , it is then immediate that (1) holds while (3) translates into (2). \square

Remark. Since $\ker \Delta$ has bounded right approximate identity, there is $C \geq 0$, depending only on \mathfrak{A} , but not on F or ϵ , such that $a_1, b_1, \dots, a_r, b_r \in \mathfrak{A}$ satisfying (1) and (2) can be chosen such that $\|\sum_{k=1}^r a_k \otimes b_k\|_{\mathfrak{A} \hat{\otimes} \mathfrak{A}} \leq C$. We shall make no use of this, however.

2. OZAWA'S PROOF REVISITED

In [Oza], Ozawa presents a proof that simultaneously establishes the non-amenability of the Banach algebras $\mathcal{B}(\ell^p)$ for $p = 1, 2, \infty$ and $\ell^\infty\text{-}\bigoplus_{n=1}^\infty \mathcal{B}(\ell_N^p)$ for all $p \in [1, \infty]$.

In this section, we recast the final step of his proof as a lemma, which does not make any reference to particular Banach algebras.

A pivotal rôle in Ozawa's argument is played by the fact that the group $\text{SL}(3, \mathbb{Z})$ has *Kazhdan's property (T)* ([B-dlH-V, Theorem 4.2.5]) and thus, in particular, is finitely generated ([B-dlH-V, Theorem 1.3.1]) by g_1, \dots, g_m , say. For the definition (and more) on Kazhdan's property (T), we refer to [B-dlH-V].

What we require is the following consequence of $\text{SL}(3, \mathbb{Z})$ having (T) ([Oza, Theorem 3.1]; compare also [B-dlH-V, Proposition 1.1.9]):

Proposition 2.1. *For any g_1, \dots, g_m generating $\text{SL}(3, \mathbb{Z})$, there is a constant $\kappa > 0$ such that, for any unitary representation π of $\text{SL}(3, \mathbb{Z})$ on a Hilbert space \mathfrak{H} and any $\xi \in \mathfrak{H}$, there is a π -invariant vector $\eta \in \mathfrak{H}$, i.e., satisfying $\pi(g)\eta = \eta$ for all $g \in \text{SL}(3, \mathbb{Z})$, such that*

$$\|\xi - \eta\| \leq \kappa \max_{j=1, \dots, m} \|\pi(g_j)\xi - \xi\|.$$

We briefly recall the setup laid out in [Oza, Section 3], which we will require both for the lemma at the end of this section and in the proof of Theorem 3.2 below.

Let \mathbb{P} denote the set of all prime numbers, and let, for each $p \in \mathbb{P}$, the projective plane over the finite field $\mathbb{Z}/p\mathbb{Z}$ be denoted by Λ_p . Obviously, $\text{SL}(3, \mathbb{Z})$ acts on Λ_p through matrix multiplication, which, in turn, induces a unitary representation π_p of $\text{SL}(3, \mathbb{Z})$ on $\ell^2(\Lambda_p)$. The action of $\text{SL}(3, \mathbb{Z})$ on Λ_p is 2-transitive, i.e., the product action of $\text{SL}(3, \mathbb{Z})$ on $\Lambda_p \times \Lambda_p$ has exactly two orbits: the diagonal and its complement. Consequently, whenever $\boldsymbol{\eta} \in \ell^2(\Lambda_p \times \Lambda_p)$ is $\pi_p \otimes \pi_p$ -invariant, there are $\alpha, \beta \in \mathbb{C}$ such that

$$(4) \quad \boldsymbol{\eta} = \alpha \boldsymbol{\zeta}_1 + \beta \boldsymbol{\zeta}_2$$

with

$$(5) \quad \boldsymbol{\zeta}_1 = |\Lambda_p|^{-\frac{1}{2}} \sum_{\lambda \in \Lambda_p} e_\lambda \otimes e_\lambda \quad \text{and} \quad \boldsymbol{\zeta}_2 = |\Lambda_p|^{-1} \sum_{\lambda, \mu \in \Lambda_p} e_\lambda \otimes e_\mu.$$

(Here, e_λ for $\lambda \in \Lambda_p$ is the point mass at λ , as usual.) Finally, choose a subset S_p of Λ_p such that $|S_p| = \frac{|\Lambda_p|-1}{2}$, and define an invertible isometry $v_p \in \mathcal{B}(\ell^2(\Lambda_p))$

through

$$v_p e_\lambda = \begin{cases} e_\lambda, & \lambda \in S_p, \\ -e_\lambda, & \lambda \notin S_p. \end{cases}$$

With generators g_1, \dots, g_m of $SL(3, \mathbb{Z})$ fixed, we shall write $\pi_p(g_{m+1})$ instead of v_p for notational convenience.

The following statement is proven ([Oza, pp. 561–562]), albeit not explicitly stated in [Oza, Section 5]. For the reader’s convenience, we include a proof.

Ozawa’s Lemma. *It is impossible to find, for each $\epsilon > 0$, a number $r \in \mathbb{N}$ with the following property: for each $p \in \mathbb{P}$, there are $\xi_{1,p}, \eta_{1,p}, \dots, \xi_{r,p}, \eta_{r,p} \in \ell^2(\Lambda_p)$ such that $\sum_{k=1}^r \xi_{k,p} \otimes \eta_{k,p} \neq 0$ and*

$$(6) \quad \left\| \sum_{k=1}^r \xi_{j,p} \otimes \eta_{k,p} - (\pi_p(g_j) \otimes \pi_p(g_j))(\xi_{k,p} \otimes \eta_{k,p}) \right\|_{\ell^2(\Lambda_p) \hat{\otimes} \ell^2(\Lambda_p)} \leq \epsilon \left\| \sum_{k=1}^r \xi_{k,p} \otimes \eta_{k,p} \right\|_{\ell^2(\Lambda_p) \hat{\otimes} \ell^2(\Lambda_p)} \quad (j = 1, \dots, m + 1).$$

Proof. Assume towards a contradiction that, for each $\epsilon > 0$, there is $r \in \mathbb{N}$ and, for each $p \in \mathbb{P}$, there are $\xi_{1,p}, \eta_{1,p}, \dots, \xi_{r,p}, \eta_{r,p} \in \ell^2(\Lambda_p)$ with $\sum_{k=1}^r \xi_{k,p} \otimes \eta_{k,p} \neq 0$ such that (6) holds.

Suppose without loss of generality that $\sum_{k=1}^r \xi_{k,p} \otimes \eta_{k,p}$ has norm one in $\ell^2(\Lambda_p) \hat{\otimes} \ell^2(\Lambda_p)$, and let $\Phi : \ell^2 \hat{\otimes} \ell^2 \rightarrow \ell^2 \tilde{\otimes}_2 \ell^2$ be the non-commutative Mazur map of [Oza, Section 4] (with $\tilde{\otimes}_2$ denoting the Hilbert space tensor product). Then $\xi := \Phi(\sum_{k=1}^r \xi_{k,p} \otimes \eta_{k,p}) \in \ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)$ is a unit vector satisfying

$$\|\xi - (\pi_p(g_j) \otimes \pi_p(g_j))\xi\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} \leq \omega_\Phi(\epsilon) \quad (j = 1, \dots, m + 1),$$

where ω_Φ is the *modulus of continuity* of Φ ([Oza, p. 562]). With $\kappa > 0$ as in Proposition 2.1, there is a $\pi_p \otimes \pi_p$ -invariant vector $\eta \in \ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)$ such that

$$\|\xi - \eta\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} \leq \kappa \omega_\Phi(\epsilon)$$

which must be of the form (4) with ζ_1 and ζ_2 as in (5). From the definition of $\pi_p(g_{m+1})$, it follows that

$$\begin{aligned} |\beta| &\leq |\beta| (2 - 2|\Lambda_p|^{-2})^{\frac{1}{2}} \\ &= |\beta| \|\zeta_2 - (\pi_p(g_{m+1}) \otimes \pi_p(g_{m+1}))\zeta_2\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} \\ &= \|\eta - (\pi_p(g_{m+1}) \otimes \pi_p(g_{m+1}))\eta\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} \\ &\leq 2\kappa \omega_\Phi(\epsilon) + \|\xi - (\pi_p(g_{m+1}) \otimes \pi_p(g_{m+1}))\xi\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} \\ &\leq (2\kappa + 1) \omega_\Phi(\epsilon) \end{aligned}$$

and thus

$$(7) \quad \|\xi - \alpha \zeta_1\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} \leq \|\xi - \eta\|_{\ell^2(\Lambda_p) \tilde{\otimes}_2 \ell^2(\Lambda_2)} + |\beta| \leq (3\kappa + 1) \omega_\Phi(\epsilon);$$

in particular,

$$(8) \quad |\alpha| \geq 1 - (3\kappa + 1) \omega_\Phi(\epsilon)$$

holds.

On the other hand, we may view ξ as a Hilbert–Schmidt operator on $\ell^2(\Lambda_p)$ of rank at most r , so that $|\text{Tr } \xi| \leq r^{\frac{1}{2}}$, where Tr is the canonical trace on $\mathcal{B}(\ell^2(\Lambda_p))$. The Cauchy–Schwarz inequality then yields

$$|\Lambda_p|^{\frac{1}{2}} \|\xi - \alpha\zeta_1\|_{\ell^2(\Lambda_p) \otimes_2 \ell^2(\Lambda_2)} \geq |\text{Tr}(\alpha\zeta_1 - \xi)| \geq |\alpha| |\Lambda_p|^{\frac{1}{2}} - r^{\frac{1}{2}}.$$

Dividing by $|\Lambda_p|^{\frac{1}{2}}$ and taking (7) and (8) into account, we obtain

$$(3\kappa + 1) \omega_\Phi(\epsilon) \geq \|\xi - \alpha\zeta_1\|_{\ell^2(\Lambda_p) \otimes_2 \ell^2(\Lambda_2)} \geq 1 - (3\kappa + 1) \omega_\Phi(\epsilon) - \left(\frac{r}{|\Lambda_p|}\right)^{\frac{1}{2}}$$

and thus

$$2(3\kappa + 1) \omega_\Phi(\epsilon) \geq 1 - \left(\frac{r}{|\Lambda_p|}\right)^{\frac{1}{2}}.$$

Since Φ is uniformly continuous by [Oza, Theorem 4.1], $\lim_{\epsilon \rightarrow 0} \omega_\Phi(\epsilon) = 0$ holds. Choosing $\epsilon > 0$ so small that $(3\kappa + 1) \omega_\Phi(\epsilon) \leq \frac{1}{4}$, we get

$$\frac{1}{2} \geq 1 - \left(\frac{r}{|\Lambda_p|}\right)^{\frac{1}{2}}$$

for all $p \in \mathbb{P}$, which is impossible. □

Remark. It is crucial that $r \in \mathbb{N}$ depend only on $\epsilon > 0$, but not on any particular p .

3. THE NON-AMENABILITY OF $\ell^\infty(\mathcal{K}(\ell^2 \oplus E))$

It is possible to adapt the argument from [Oza] to prove the non-amenability of $\ell^\infty(\mathcal{K}(\ell^2))$. In this section, we shall go one step further: we shall see that, for certain Banach spaces E , the Banach algebra $\ell^\infty(\mathcal{K}(\ell^2 \oplus E))$ is not amenable.

Recall (from [L–T], for instance) that a sequence $(x_n)_{n=1}^\infty$ in a Banach space E is called a *basis* for E if, for each $x \in E$, there is a unique sequence $(\lambda_n)_{n=1}^\infty$ of scalars such that

$$(9) \quad x = \sum_{n=1}^\infty \lambda_n x_n$$

(with convergence in the norm topology of E). For instance, the canonical basis $(e_n)_{n=1}^\infty$ of ℓ^p is a basis in this sense if $p \in [1, \infty)$ (but not if $p = \infty$). For each $x \in E$ and $n \in \mathbb{N}$, set $\langle x, x_n^* \rangle := \lambda_n$, where λ_n is the coefficient of x_n in the expansion (9) of x . Then $(x_n^*)_{n=1}^\infty$ is a sequence in E^* .

Let E be a Banach space, and let $\mathcal{F}(E)$ denote the bounded finite rank operators on it. Identifying $\mathcal{F}(E)$ with the algebraic tensor product $E \otimes E^*$, we define the *trace* on $\mathcal{F}(E)$ via

$$\text{Tr}: \mathcal{F}(E) \rightarrow \mathbb{C}, \quad x \otimes x^* \mapsto \langle x, x^* \rangle.$$

We collect the following (mostly well-known) facts for later use:

Lemma 3.1. *Let E be a Banach space. Then the following hold:*

- (i) $\text{Tr } ST = \text{Tr } TS \quad (S \in \mathcal{F}(E), T \in \mathcal{B}(E));$
- (ii) *if F is another Banach space and $T = \begin{bmatrix} T_{1,1} & T_{1,2} \\ T_{2,1} & T_{2,2} \end{bmatrix} \in \mathcal{F}(E \oplus F)$, then*

$$\text{Tr } T = \text{Tr } T_{1,1} + \text{Tr } T_{2,2};$$

(iii) if E has a basis $(x_n)_{n=1}^\infty$, then

$$\text{Tr } T = \sum_{n=1}^\infty \langle Tx_n, x_n^* \rangle \quad (T \in \mathcal{F}(E)).$$

Proof. Parts (i) and (ii) are immediate from the definition (and are well known).

For (iii), note that, by linearity, it suffices to check the claim for $T = x \otimes x^*$ with $x \in E$ and $x^* \in E^*$. Since $(x_n)_{n=1}^\infty$ is a basis for E , there is a unique sequence $(\lambda_n)_{n=1}^\infty$ in \mathbb{C} such that (9) holds. It follows that

$$\text{Tr } T = \left\langle \sum_{n=1}^\infty \lambda_n x_n, x^* \right\rangle = \sum_{n=1}^\infty \lambda_n \langle x_n, x^* \rangle = \sum_{n=1}^\infty \langle x, x_n^* \rangle \langle x_n, x^* \rangle = \sum_{n=1}^\infty \langle Tx_n, x_n^* \rangle$$

as claimed. □

Theorem 3.2. *Let E be a Banach space with a basis $(x_n)_{n=1}^\infty$ such that there is $C > 0$ with*

$$(10) \quad \sum_{n=1}^\infty \|Sx_n\| \|Tx_n^*\| \leq CN \|S\| \|T\| \quad (N \in \mathbb{N}, S \in \mathcal{B}(E, \ell_N^2), T \in \mathcal{B}(E^*, \ell_N^2)).$$

Then $\ell^\infty(\mathcal{K}(\ell^2 \oplus E))$ is not amenable.

Proof. We identify $\ell^\infty(\mathcal{K}(\ell^2 \oplus E))$ with $\ell^\infty(\mathbb{P}, \mathcal{K}(\ell^2 \oplus E))$, which we will denote simply by \mathfrak{A} . Each summand of \mathfrak{A} has a canonical block matrix structure

$$(11) \quad \mathcal{K}(\ell^2 \oplus E) = \begin{bmatrix} \mathcal{K}(\ell^2) & \mathcal{K}(E, \ell^2) \\ \mathcal{K}(\ell^2, E) & \mathcal{K}(E) \end{bmatrix}.$$

For each $p \in \mathbb{P}$, we embed $\mathcal{B}(\ell^2(\Lambda_p))$ into $\mathcal{K}(\ell^2 \oplus E)$ as upper left corner of the block matrix (11). This allows us to consider $\ell^\infty\text{-}\bigoplus_{p \in \mathbb{P}} \mathcal{B}(\ell^2(\Lambda_p))$ as a closed subalgebra of \mathfrak{A} . In particular, we can identify

$$F := \{(\pi_p(g_j))_{p \in \mathbb{P}} : j = 1, \dots, m + 1\}$$

with a finite subset of \mathfrak{A} . Furthermore, we let \mathfrak{A} act (as block diagonal matrices) on the space

$$\ell^2(\mathbb{P}, \ell^2 \oplus E) \cong \ell^2(\mathbb{P}, \ell^2) \oplus \ell^2(\mathbb{P}, E).$$

We shall identify $\ell^2(\mathbb{P}, \ell^2)$ and $\ell^2(\mathbb{P}, E)$ with completions of the algebraic tensor products $\ell^2(\mathbb{P}) \otimes \ell^2$ and $\ell^2(\mathbb{P}) \otimes E$.

For any $p \in \mathbb{P}$, let $P_p \in \mathcal{B}(\ell^2)$ be the canonical projection onto the first $|\Lambda_p|$ coordinates of the p^{th} ℓ^2 -summand of $\ell^2(\mathbb{P}, \ell^2) \oplus \ell^2(\mathbb{P}, E)$. Let $a = (a_p)_{p \in \mathbb{P}}$ and $b = (b_p)_{p \in \mathbb{P}}$ be elements of \mathfrak{A} . By [Oza, Lemma 2.1(i)], we have

$$(12) \quad \sum_{q \in \mathbb{P}} \sum_{n=1}^\infty \|P_p a(e_q \otimes e_n)\| \|P_p^* b^*(e_q^* \otimes e_n^*)\| = \sum_{n=1}^\infty \|P_p a_p e_n\| \|P_p^* b_p^* e_n^*\| \leq |\Lambda_p| \|a_p\| \|b_p\| \leq |\Lambda_p| \|a\| \|b\|,$$

and from (10), we conclude that

$$(13) \quad \sum_{q \in \mathbb{P}} \sum_{n=1}^\infty \|P_p a(e_q \otimes x_n)\| \|P_p^* b^*(e_q^* \otimes x_n^*)\| = \sum_{n=1}^\infty \|P_p a_p x_n\| \|P_p^* b_p^* x_n^*\| \leq C |\Lambda_p| \|a_p\| \|b_p\| \leq C |\Lambda_p| \|a\| \|b\|.$$

Set $e = (P_p)_{p \in \mathbb{P}}$. Then e is an idempotent in \mathfrak{A} with

$$e\mathfrak{A}e = \ell^\infty - \bigoplus_{p \in \mathbb{P}} \mathcal{B}(\ell^2(\Lambda_p)).$$

Assume now towards a contradiction that $\ell^\infty(\mathbb{P}, \mathcal{K}(\ell^2 \oplus E))$ is amenable, and let $\epsilon > 0$ be arbitrary. By Lemma 1.2 there are thus $a_1, b_1, \dots, a_r, b_r \in \mathfrak{A}$ such that (1) holds and

$$(14) \quad \left\| \sum_{k=1}^r x a_k \otimes b_k - a_k \otimes b_k x \right\| < \frac{\epsilon}{(C+1)(m+1)} \quad (x \in F).$$

For $p, q \in \mathbb{P}$ and $n \in \mathbb{N}$, define

$$T_p(q, n) := \sum_{k=1}^r P_p a_k (e_q \otimes e_n) \otimes P_p^* b_k^* (e_q^* \otimes e_n^*) + P_p a_k (e_q \otimes x_n) \otimes P_p^* b_k^* (e_q^* \otimes x_n^*) \in \ell^2(\Lambda_p) \hat{\otimes} \ell^2(\Lambda_p).$$

For $j = 1, \dots, m+1$ and fixed $p \in \mathbb{P}$, we then have by (12) and (13)

$$\begin{aligned} & \sum_{q \in \mathbb{P}} \sum_{n=1}^{\infty} \|T_p(q, n) - (\pi_p(g_j) \otimes \pi_p(g_j))T_p(q, n)\| \\ & \leq (1+C)|\Lambda_p| \left\| \sum_{k=1}^r (\pi_p(g_j))_{p \in \mathbb{P}} a_k \otimes b_k - a_k \otimes b_k (\pi_p(g_j))_{p \in \mathbb{P}} \right\| \leq \frac{\epsilon}{m+1} |\Lambda_p| \end{aligned}$$

and thus

$$(15) \quad \sum_{q \in \mathbb{P}} \sum_{n=1}^{\infty} \sum_{j=1}^{m+1} \|T_p(q, n) - (\pi_p(g_j) \otimes \pi_p(g_j))T_p(q, n)\| \leq \epsilon |\Lambda_p|.$$

For $k = 1, \dots, r$, write $a_k = (a_{k,q})_{q \in \mathbb{P}}$ and $b_k = (b_{k,q})_{q \in \mathbb{P}}$, and note that

$$(16) \quad \begin{aligned} \sum_{q \in \mathbb{P}} \sum_{n=1}^{\infty} \|T_p(q, n)\| & \geq \sum_{n=1}^{\infty} \left| \sum_{k=1}^r \langle P_p a_{k,p} e_n, P_p^* b_{k,p}^* e_n^* \rangle + \langle P_p a_{k,p} x_n, P_p^* b_{k,p}^* x_n^* \rangle \right| \\ & \geq \left| \sum_{n=1}^{\infty} \sum_{k=1}^r \langle P_p a_{k,p} e_n, P_p^* b_{k,p}^* e_n^* \rangle + \sum_{n=1}^{\infty} \sum_{k=1}^r \langle P_p a_{k,p} x_n, P_p^* b_{k,p}^* x_n^* \rangle \right| \\ & = \left| \text{Tr} \sum_{k=1}^r b_{k,p} P_p a_{k,p} \right|, \quad \text{by Lemma 3.1(ii) and (iii),} \\ & = \left| \text{Tr} \sum_{k=1}^r P_p a_{k,p} b_{k,p} \right|, \quad \text{by Lemma 3.1(i),} \\ & = |\text{Tr } P_p|, \quad \text{because } \sum_{k=1}^r a_{k,p} b_{k,p} = P_p, \\ & = |\Lambda_p|. \end{aligned}$$

Of course, the inequalities (15) and (16) remain intact if we only take the sum over those $q \in \mathbb{P}$ and $n \in \mathbb{N}$ such that $T_p(q, n) \neq 0$. Both inequalities together then

imply that there are $q \in \mathbb{P}$ and $n \in \mathbb{N}$ with $T_p(q, n) \neq 0$ such that

$$\sum_{j=1}^{m+1} \|T_p(q, n) - (\pi_p(g_j) \otimes \pi_p(g_j))T_p(q, n)\| \leq \epsilon \|T_p(q, n)\|$$

and thus

$$(17) \quad \|T_p(q, n) - (\pi_p(g_j) \otimes \pi_p(g_j))T_p(q, n)\| \leq \epsilon \|T_p(q, n)\| \quad (j = 1, \dots, m + 1).$$

From the definition of $T_p(q, n)$ it is clear that there are $\xi_{1,p}, \eta_{1,p}, \dots, \xi_{2r,p}, \eta_{2r,p} \in \ell^2(\Lambda_p)$ with $T_p(q, n) = \sum_{k=1}^{2r} \xi_{k,p} \otimes \eta_{k,p}$. In view of (17), we thus have $\xi_{1,p}, \eta_{1,p}, \dots, \xi_{2r,p}, \eta_{2r,p} \in \ell^2(\Lambda_p)$ with $\sum_{k=1}^{2r} \xi_{k,p} \otimes \eta_{k,p} \neq 0$ such that

$$\begin{aligned} & \left\| \sum_{k=1}^{2r} \xi_{j,p} \otimes \eta_{k,p} - (\pi_p(g_j) \otimes \pi_p(g_j))(\xi_{k,p} \otimes \eta_{k,p}) \right\|_{\ell^2(\Lambda_p) \hat{\otimes} \ell^2(\Lambda_p)} \\ & \leq \epsilon \left\| \sum_{k=1}^{2r} \xi_{k,p} \otimes \eta_{k,p} \right\|_{\ell^2(\Lambda_p) \hat{\otimes} \ell^2(\Lambda_p)} \quad (j = 1, \dots, m + 1). \end{aligned}$$

Since $p \in \mathbb{P}$ was arbitrary, this contradicts Ozawa’s lemma. □

4. THE NON-AMENABILITY OF $\mathcal{B}(\ell^p(E))$

With Theorem 3.2 proven, we shall now see that the condition imposed on the basis of E in that theorem is automatically satisfied by the canonical basis $(e_n)_{n=1}^\infty$ of ℓ^p with $p \in (1, \infty)$. (As is customary, we set $p' := \frac{p}{p-1}$, so that $(\ell^p)^* = \ell^{p'}$.)

Let $p \in [1, \infty)$, and let E and F be Banach spaces. Recall (from [D–J–T], for instance) that a linear operator $T: E \rightarrow F$ is called *p-summing* if the amplification $\text{id}_{\ell^p} \otimes T: \ell^p \otimes E \rightarrow \ell^p \otimes F$ extends to a bounded linear operator from $\ell^p \check{\otimes} E$ ($\check{\otimes}$ stands for the injective Banach space tensor product) to $\ell^p(F)$. In this case, the operator norm of $\text{id}_{\ell^p} \otimes T: \ell^p \check{\otimes} E \rightarrow \ell^p(F)$ is called the *p-summing norm* of T and is denoted by $\pi_p(T)$.

Lemma 4.1. *Let $p \in (1, \infty)$. Then there is $C > 0$ such that*

$$\sum_{n=1}^\infty \|S e_n\|_{\ell_N^2} \|T e_n^*\|_{\ell_N^2} \leq C N \|S\| \|T\| \quad (N \in \mathbb{N}, S \in \mathcal{B}(\ell^p, \ell_N^2), T \in \mathcal{B}(\ell^{p'}, \ell_N^2)).$$

Proof. We can, for each $N \in \mathbb{N}$, algebraically identify $\mathcal{B}(\ell^p, \ell_N^2)$ and $\mathcal{B}(\ell^{p'}, \ell_N^2)$ with the algebraic tensor products $\ell^{p'} \otimes \ell_N^2$ and $\ell^p \otimes \ell_N^2$, respectively. On the other hand, those tensor products can also be identified with the spaces $\ell^{p'}(\ell_N^2)$ and $\ell^p(\ell_N^2)$, respectively. Hölder’s inequality yields immediately that

$$\begin{aligned} & \sum_{n=1}^\infty \|S e_n\|_{\ell_N^2} \|T e_n^*\|_{\ell_N^2} \leq \|S\|_{\ell^{p'}(\ell_N^2)} \|T\|_{\ell^p(\ell_N^2)} \\ & \quad (N \in \mathbb{N}, S \in \mathcal{B}(\ell^{p'}, \ell_N^2), T \in \mathcal{B}(\ell^p, \ell_N^2)). \end{aligned}$$

With $\check{\otimes}$ denoting the injective tensor product, we have the isometric identifications

$$\mathcal{B}(\ell^p, \ell_N^2) \cong \ell^{p'} \check{\otimes} \ell_N^2 \quad \text{and} \quad \mathcal{B}(\ell^{p'}, \ell_N^2) \cong \ell^p \check{\otimes} \ell_N^2,$$

so that

$$\|S\|_{\ell^{p'}(\ell_N^2)} \leq \pi_{p'}(\text{id}_{\ell_N^2})\|S\| \quad \text{and} \quad \|T\|_{\ell^p(\ell_N^2)} \leq \pi_p(\text{id}_{\ell_N^2})\|T\|$$

$$(N \in \mathbb{N}, S \in \mathcal{B}(\ell^p, \ell_N^2), T \in \mathcal{B}(\ell^{p'}, \ell_N^2)).$$

By [Gor, Theorem 5] (compare also [T-J, Theorem 10.3])

$$\pi_{p'}(\text{id}_{\ell_N^2}) \sim N^{\frac{1}{2}} \sim \pi_p(\text{id}_{\ell_N^2})$$

holds; in particular, there are $C_{p'}, C_p > 0$ with

$$\pi_{p'}(\text{id}_{\ell_N^2}) \leq C_{p'}\sqrt{N} \quad \text{and} \quad \pi_p(\text{id}_{\ell_N^2}) \leq C_p\sqrt{N} \quad (N \in \mathbb{N}).$$

Letting $C := C_{p'}C_p$ yields the desired constant. □

By Lemma 4.1, ℓ^p therefore satisfies the hypothesis of Theorem 3.2 for any $p \in (1, \infty)$, so that $\ell^\infty(\mathcal{K}(\ell^2 \oplus \ell^p))$ is not amenable, but more is true.

Recall the definition of an \mathcal{L}^p -space from [L-P]:

Definition 4.2. Let $p \in [1, \infty]$.

- (a) A Banach space E is called an \mathcal{L}_λ^p -space with $\lambda \geq 1$ if, for any finite-dimensional subspace X of E , there are $N \in \mathbb{N}$ and an N -dimensional subspace Y of E containing X such that $d(Y, \ell_N^p) \leq \lambda$, where d is the Banach–Mazur distance.
- (b) We call E an \mathcal{L}^p -space if it is an \mathcal{L}_λ^p -space for some $\lambda \geq 1$.

All L^p -spaces are \mathcal{L}^p -spaces, but the converse fails: For $p \in (1, \infty) \setminus \{2\}$, the space $\ell^2 \oplus \ell^p$ is an \mathcal{L}^p -space but not an L^p -space ([L-P, Example 8.2]). By [D-R, Theorem 4.3], all infinite-dimensional \mathcal{L}^p -spaces E are equivalent when it comes to the question of whether $\ell^\infty(\mathcal{K}(E))$ is amenable: either $\ell^\infty(\mathcal{K}(E))$ is amenable for all infinite-dimensional \mathcal{L}^p -spaces E or it is non-amenable for all such spaces.

In view of Theorem 3.2 and Lemma 4.1, we thus have

Proposition 4.3. Let $p \in (1, \infty)$, and let E be an infinite-dimensional \mathcal{L}^p -space. Then $\ell^\infty(\mathcal{K}(E))$ is not amenable.

We can finally deduce the main result of this paper:

Theorem 4.4. Let $p \in (1, \infty)$, and let E be an \mathcal{L}^p -space. Then $\mathcal{B}(\ell^p(E))$ is not amenable.

Proof. Assume towards a contradiction that $\mathcal{B}(\ell^p(E))$ is amenable. By [D-R, Theorem 2.1], this forces $\ell^\infty(\mathcal{B}(\ell^p(E)))$ to be amenable as well. Since E is an \mathcal{L}^p -space, so is $\ell^p(E)$; in particular, $\ell^p(E)$ is reflexive and has the approximation property, so that $\mathcal{K}(\ell^p(E))$ has a bounded approximate identity. Consequently, $\ell^\infty(\mathcal{K}(\ell^p(E)))$ is an ideal in $\ell^\infty(\mathcal{B}(\ell^p(E)))$ with a bounded approximate identity and thus is an amenable Banach algebra in its own right ([Run 2, Proposition 2.3.3]), which is impossible by Proposition 4.3. □

Corollary 4.5. Let $p \in (1, \infty)$, and let E be an \mathcal{L}^p -space such that $E \cong \ell^p(E)$. Then $\mathcal{B}(E)$ is not amenable.

For any $p \in (1, \infty)$, the spaces $E = \ell^p$ and $E = L^p[0, 1]$ satisfy the hypothesis of Corollary 4.5, so that $\mathcal{B}(\ell^p)$ and $\mathcal{B}(L^p[0, 1])$ are not amenable. Since any infinite-dimensional, separable L^p -space is isomorphic to either ℓ^p or $L^p[0, 1]$ ([Woj, p. 83]),

we even obtain

Corollary 4.6. *Let $p \in (1, \infty)$, and let E be an infinite-dimensional, separable L^p -space. Then $\mathcal{B}(E)$ is not amenable.*

Remark. If one is only interested in the non-amenability of $\mathcal{B}(\ell^p)$ for $p \in (1, \infty)$, a more direct route is possible: According to [L–T, p. 73], the isomorphism

$$(18) \quad \ell^p \cong \ell^p \text{-}\bigoplus_{n=1}^{\infty} \ell_n^2$$

holds. Using, for $q \in \mathbb{P}$, the projection from ℓ^p onto $\ell^2(\Lambda_q) = \ell_{|\Lambda_q|}^2$ according to (18), then invoking Lemma 4.1, and finally following more or less [Oza] will also yield a proof of the non-amenability of $\mathcal{B}(\ell^p)$. However, unlike the argument presented here, we do not see how this way of reasoning will yield the non-amenability of $\mathcal{B}(E)$ for every \mathcal{L}^p -space with $E \cong \ell^p(E)$, such as $L^p[0, 1]$.

ACKNOWLEDGMENTS

I am grateful to Gilles Pisier and Seytek Tabaldyev for pointing out to me that the original requirement of Theorem 3.2 that E have a shrinking basis was stronger than necessary. I owe special thanks to Seytek Tabaldyev also for discovering an error in an earlier version of the proof of that theorem.

REFERENCES

- [A–H] S. A. ARGYROS and R. G. HAYDON, A hereditarily indecomposable \mathcal{L}_∞ -space that solves the scalar-plus-compact problem. [arXiv:0903.3921](https://arxiv.org/abs/0903.3921).
- [B–dlH–V] B. BEKKA, P. DE LA HARPE, and A. VALETTE, *Kazhdan’s Property (T)*. New Mathematical Monographs **11**, Cambridge University Press, Cambridge, 2008. MR2415834 (2009i:22001)
- [D–R] M. DAWS and V. RUNDE, Can $\mathcal{B}(\ell^p)$ ever be amenable? *Studia Math.* **188** (2008), 151–174. Erratum. *Studia Math.* **195** (2009), 297–298. MR2431000; MR2559179
- [D–J–T] J. DIESTEL, H. JARCHOW, and A. TONGE, *Absolutely Summing Operators*. Cambridge Studies in Advanced Mathematics **43**, Cambridge University Press, Cambridge–New York–Melbourne, 1995. MR1342297 (96i:46001)
- [Gor] Y. GORDON, On p -absolutely summing constants of Banach spaces. *Israel J. Math.* **7** (1969), 151–163. MR0250028 (40:3269)
- [G–J–W] N. GRØNBÆK, B. E. JOHNSON, and G. A. WILLIS, Amenability of Banach algebras of compact operators. *Israel J. Math.* **87** (1994), 289–324. MR1286832 (95k:46078)
- [Hel] A. YA. HELEMSKIĬ, *The Homology of Banach and Topological Algebras* (translated from the Russian). Mathematics and Its Applications (Soviet Series) **41**, Kluwer Academic Publishers Group, Dordrecht, 1989. MR1093462 (92d:46178)
- [Joh 1] B. E. JOHNSON, Cohomology in Banach algebras. *Mem. Amer. Math. Soc.* **127** (1972). MR0374934 (51:11130)
- [Joh 2] B. E. JOHNSON, Approximate diagonals and cohomology of certain annihilator Banach algebras. *Amer. J. Math.* **94** (1972), 685–698. MR0317050 (47:5598)
- [L–P] J. LINDENSTRAUSS and A. PELCZYŃSKI, Absolutely summing operators in \mathcal{L}_p -spaces and their applications. *Studia Math.* **29** (1968), 275–326. MR0231188 (37:6743)
- [L–T] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach Spaces*, I. Ergebnisse der Mathematik und ihrer Grenzgebiete **92**, Springer Verlag, Berlin–Heidelberg–New York, 1977. MR0500056 (58:17766)
- [Oza] N. OZAWA, A note on non-amenability of $\mathcal{B}(\ell_p)$ for $p = 1, 2$. *Internat. J. Math.* **15** (2004), 557–565. MR2078880 (2005g:46135)
- [Pis] G. PISIER, On Read’s proof that $B(l_1)$ is not amenable. In: *Geometric Aspects of Functional Analysis*, pp. 269–275. Lecture Notes in Mathematics **1850**, Springer Verlag, Berlin–Heidelberg–New York, 2004. MR2087163 (2005g:46136)

- [Rea 1] C. J. READ, Commutative, radical amenable Banach algebras. *Studia Math.* **140** (2000), 199–212. MR1784151 (2001f:46078)
- [Rea 2] C. J. READ, Relative amenability and the non-amenability of $B(l^1)$. *J. Austral. Math. Soc.* **80** (2006), 317–333. MR2236040 (2007h:46062)
- [Run 1] V. RUNDE, The structure of contractible and amenable Banach algebras. In: E. ALBRECHT and M. MATHIEU (eds.), *Banach Algebras '97 (Blaubeuren)*, pp. 415–430, de Gruyter, Berlin, 1998. MR1656619 (2000d:46062)
- [Run 2] V. RUNDE, *Lectures on Amenability*. Lecture Notes in Mathematics **1774**, Springer Verlag, Berlin–Heidelberg–New York, 2002. MR1874893 (2003h:46001)
- [T-J] N. TOMCZAK-JAEGERMANN, *Banach–Mazur Distances and Finite-Dimensional Operator Ideals*. Pitman Monographs and Surveys in Pure and Applied Mathematics **38**, Longman Scientific & Technical, Essex, 1988. MR993774 (90k:46039)
- [Was] S. WASSERMANN, On tensor products of certain group C^* -algebras. *J. Functional Anal.* **23** (1976), 239–254. MR0425628 (54:13582)
- [Woj] P. WOJTASZCZYK, *Banach Spaces for Analysts*. Cambridge Studies in Advanced Mathematics **25**, Cambridge University Press, Cambridge, 1991. MR1144277 (93d:46001)

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA T6G 2G1

E-mail address: vrunde@ualberta.ca

URL: <http://www.math.ualberta.ca/~runde/>