

NEW BOUNDS ON CAP SETS

MICHAEL BATEMAN AND NETS HAWK KATZ

1. INTRODUCTION

A set $A \subset F_3^N$ is called a cap set if it contains no lines. In this paper, we will be concerned with proving the following theorem:

Theorem 1.1. *There exist an $\epsilon > 0$ and $C < \infty$ such that if $A \subseteq F_3^N$ is a cap set, then*

$$\frac{|A|}{3^N} \leq \frac{C}{N^{1+\epsilon}}.$$

The problem of the maximal size of cap sets is a characteristic 3 model for the problem of finding arithmetic progressions of length 3 in rather dense sets of integers. Meshulam [M95], through a direct use of ideas of Roth, was able to show that there is a constant C so that any cap set A has density at most $\frac{C}{N}$. Our result may be viewed as a very modest improvement over Meshulam's result.

Sanders [S11] recently showed that any subset of the integers whose density in $\{1, \dots, M\}$ is at least $\frac{C(\log \log M)^5}{\log M}$ must contain arithmetic progressions of length 3. This may be thought of as bringing the results known for arithmetic progressions almost to the level of Meshulam's result. This has spurred further interest in improving Meshulam's result in hopes that it might suggest a way of improving the results on arithmetic progressions.

A rather concrete, though perhaps still out of reach, goal in this direction is a conjecture of Erdős and Turan:

Conjecture 1.2. *Suppose $A \subseteq \mathbb{Z}$ is such that*

$$\sum_{n \in A} \frac{1}{n} = \infty.$$

Then A contains an arithmetic progression of every length.

It is clear that the present paper is directly relevant only for finding 3-term progressions. However it is also easy to see, based purely on density considerations, that proving an estimate of the type in Theorem 1.1 in the integer setting would yield the 3-term case of the conjecture stated above. In fact, Polymath 6 [PM6] has recently been started with the goal of adapting the ideas of this paper to the integer setting. See [PM] for more information about so-called "polymath" projects in general.

Received by the editors April 2, 2011 and, in revised form, October 28, 2011.

2010 *Mathematics Subject Classification.* Primary 11T71; Secondary 05D40.

The first author is supported by an NSF postdoctoral fellowship, DMS-0902490.

The second author is partially supported by NSF grant DMS-1001607.

©2011 American Mathematical Society
Reverts to public domain 28 years from publication

While the research in this paper was well underway, Gowers [G] wrote a post on his blog suggesting that one could attack the problem of bounding cap sets by studying the additive structure of their large spectrum. This had been our approach as well and we wrote a reply [K1] sketching our rather strong results regarding that structure. In the course of a few days, we realized that we actually could convert our structural theory into an estimate on cap sets. We recorded this [K2] in a second reply to Gowers's blog. The current paper should be viewed as an elaboration of these two posts.

We describe our plan for proving Theorem 1.1. To prove this theorem we will prove a theorem about sets without unusually dense subspaces, a notion we make precise below.

Definition 1.3. Suppose $A \subseteq F_3^N$ has density ρ ; i.e., $|A| = \rho 3^n$. We say that A has *no strong increments* if for every subspace $V \subseteq F_3^N$ with $d = \text{codim} V \leq \frac{N}{2}$, we have

$$\frac{|A \cap V|}{|V|} \leq \rho + \frac{20d\rho}{N}.$$

Theorem 1.4. *There exist an $\epsilon > 0$ and $C < \infty$ such that if $A \subseteq F_3^N$ is a cap set with no strong increments, then*

$$\frac{|A|}{3^N} \leq \frac{C}{N^{1+\epsilon}}.$$

The major ingredients needed to prove this theorem are Proposition 3.3, Lemma 5.4, and Theorem 7.1. We combine them with a Fourier analytic argument in Section 8.

Proof. We deduce Theorem 1.1 from Theorem 1.4 using induction. Suppose that for every $n \leq N - 1$ we have shown that if $B \subseteq F_3^n$ is a cap set, then

$$\frac{|B|}{3^n} \leq \frac{C}{n^{1+\epsilon}}.$$

We aim for a contradiction: assume there exists a cap set $A \subseteq F_3^N$ such that $\frac{|A|}{3^N} > \frac{C}{N^{1+\epsilon}}$. By Theorem 1.4, this implies that A has a strong increment. Since A has a strong increment, there exists an affine subspace $V \subseteq F_3^N$ with codimension $\leq \frac{N}{2}$ such that

$$\begin{aligned} \frac{|A \cap V|}{|V|} &\geq \rho + \frac{20d\rho}{N} \\ &= \rho \left(1 + \frac{20d}{N}\right) \\ &> \frac{C}{(N-d)^{1+\epsilon}} \end{aligned}$$

since the derivative of $\frac{C}{x^{1+\epsilon}}$ is uniformly bounded by $16CN^{-2-\epsilon} = 16\rho$ whenever $0 < \epsilon < 1$. But we know that $A \cap V$ (in fact any subset of A) is a cap set. This contradicts the induction hypothesis, yielding Theorem 1.1. \square

1.1. **Proof sketch.** We sketch our plan for proving Theorem 1.4.

We will study the large spectrum Δ of a cap set A with no strong increments. The reader should think of Δ , for a cap set of the size $\frac{3^N}{N}$ given by Meshulam's estimate, as consisting of the positions at which the absolute value of the Fourier transform of A is around $\frac{1}{N^2}$. The set Δ should have cardinality approximately N^3 (established in Section 3) and have about N^7 additive quadruples (established in Section 4). The estimate on the number of quadruples is essentially the same as a result in the integer setting obtained independently by Shkredov [Shk08]. Recall that an additive quadruple is a quadruple (x_1, x_2, x_3, x_4) of elements of Δ with the property that

$$x_1 + x_2 = x_3 + x_4.$$

Similarly an additive octuple is an octuple $(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8)$ of elements of Δ with

$$x_1 + x_2 + x_3 + x_4 = x_5 + x_6 + x_7 + x_8.$$

It is easy to see *a priori* that a set with many additive quadruples will have many additive octuples. In our case, a set with size N^3 , having N^7 quadruples must have at least N^{15} octuples. (But it may have more octuples.) The number of octuples it has should be taken as an indication of its structure. If there are many octuples, it means that the sum set $\Delta + \Delta$ looks like it has more additive structure than the set Δ . We then say that the set Δ is additively smoothing. (It becomes smoother under addition.) We show, however, that this cannot be the case for Δ , the large spectrum. We use the probabilistic method to do this, finding too much of the spectrum contained in a small subspace in the additively smoothing case. We establish this in Section 5. (This is somewhat reminiscent of the paper of Croot and Sisask [CS11], where random selections are used to uncover structure.)

Thus our set Δ is entirely additively nonsmoothing. This means it is already as smooth as it will become under a small number of additions. This makes its additive structure particularly easy to uncover as it is already present without adding the set to itself. This kind of idea was first exploited in a paper of the second author with Koester [KK10], and we use techniques quite similar to those found in that paper. We end up showing that the set Δ should be thought of as looking like the sum of a very structured set K of size N (that is to say that K is almost additively closed) with a very random set Λ of size N^2 . Section 6 contains the proof of a structural theorem for sets with substantial additive energy (i.e., many additive quadruples) but no additive smoothing.

We find that this structure of Δ is inconsistent with A 's being a cap set with no strong increments. The reason is that we can use Freiman's theorem to place K inside a subspace H with relatively low dimension. We can essentially mod out by H , examining the "fibers", the intersections of A with translates of H^\perp . We find that the structure of Δ makes the behavior of the fibers unrealistic. This argument is suggested by a paper of Sanders [S10] and is carried out in Section 8.

One final remark about the value of ϵ obtained in this paper: it is necessarily rather small (at least with our current argument). We discuss the reasons for this in a brief final section as well as giving some conjectures that if true would greatly improve the efficiency. We have not attempted to optimize ϵ (or even keep track of the exact dependence on ϵ) throughout the paper.

2. PRELIMINARIES

In this section, we record a few general results of which we will make frequent use throughout the paper. We begin with our favorite form of the Cauchy-Schwarz inequality.

Lemma 2.1. *Let S and T be finite sets and ρ a map,*

$$\rho : S \longrightarrow T.$$

Let P be the set of pairs

$$P = \{(s_1, s_2) : \rho(s_1) = \rho(s_2)\}.$$

Then

$$|P| \geq \frac{|S|^2}{|T|}.$$

Proof. Note that we can express $|P|$ by

$$|P| = \sum_{t \in T} |\rho^{-1}(t)|^2.$$

Applying the Cauchy-Schwarz inequality we obtain

$$|P| \geq \frac{1}{|T|} \left(\sum_{t \in T} |\rho^{-1}(t)| \right)^2 = \frac{|S|^2}{|T|}.$$

□

Here we introduce another variant of Cauchy-Schwarz:

Lemma 2.2. *Let (X, m) be a measure space with total measure M . Let A_1, \dots, A_k be measurable subsets of X and $0 < \rho < 1$ be a number (the density), so that $m(A_j) \gtrsim \rho M$ for each j . Suppose $k \gtrsim \frac{1}{\rho^2}$. Then*

$$\sum_{j=1}^k \sum_{l \neq j} m(A_j \cap A_l) \gtrsim k^2 \rho^2 M.$$

Proof. Note that

$$\sum_{j=1}^k m(A_j) = \rho k M \ll k^2 \rho^2 M,$$

unless $\rho \sim 1$. Thus we may estimate the full sum

$$\sum_{j=1}^k \sum_{l=1}^k m(A_j \cap A_l).$$

Define $c(x)$ to be the measurable function giving, for each x , the number of sets A_j which contain x . Thus we would like to estimate

$$\int c(x)^2 \geq \frac{1}{M} \left(\int c(x) \right)^2 = k^2 \rho^2 M.$$

□

In what follows μ shall be a small exponent. We will frequently use expressions such as $N^{O(\mu)}$. The exponent will be bounded by $C\mu$ for C a universal constant which varies from line to line in the paper. We illustrate this by the following version of the large families principle (this is the principle which says that most children belong to large families) which will be used extremely often in this paper.

Lemma 2.3 (Large Families Principle). *Let $M_1, \dots, M_K > 0$ be real numbers and let $R > 0$ be a real number. Suppose that $M_j \leq RN^\mu$ for each j and suppose that*

$$\sum_{j=1}^K M_j \geq RKN^{-\mu}.$$

Then there exists a subset J of $\{1, \dots, K\}$ with $|J| \gtrsim N^{-O(\mu)}K$ so that for each $j \in J$, we have $M_j \gtrsim N^{-O(\mu)}R$.

Proof. Let

$$J = \{j : M_j \gtrsim N^{-10\mu}R\}.$$

Suppose that $|J| \lesssim N^{-10\mu}K$. Then by the upper bound on M_j , we have that

$$\sum_{j \in J} M_j \lesssim N^{-9\mu}RK,$$

while

$$\sum_{j \notin J} M_k \lesssim N^{-10\mu}RK.$$

Combining these two estimates gives us a contradiction. □

We take a moment to state the asymmetric Balog-Szemerédi-Gowers theorem, which we will have occasion to use. First, we give a slightly nonstandard definition.

Definition 2.4. A set $B \subset F_3^N$ will be said to be μ -additively closed if

$$|B + B| \lesssim N^{O(\mu)}|B|.$$

Lemma 2.5. *Let $B, C \subset F_3^N$ so that there are at least $N^\eta|B||C|^2$ additive quadruples of the form*

$$b_1 + c_1 = b_2 + c_2$$

with $b_1, b_2 \in B$ and $c_1, c_2 \in C$. Let $L = \frac{|B|}{|C|}$ and assume $L \lesssim N^{10}$. Then there exists μ depending only on η , with $\mu \rightarrow 0$ as $\eta \rightarrow 0$, and there exist a μ -additively closed set $K \subset F_3^N$ and a set $X \subset F_3^N$ with

$$|X| \lesssim N^{O(\mu)} \frac{|B|}{|C|},$$

so that

$$|B \cap (X + K)| \gtrsim N^{-O(\mu)}|B|,$$

and an element $x \in F_3^N$ so that

$$|C \cap (x + K)| \gtrsim N^{-O(\mu)}|C|.$$

In particular, the last inequality implies that

$$|K| \gtrsim N^{-O(\mu)}|C|.$$

An only slightly stronger form of this lemma appears in the book of Tao and Vu as Lemma 2.35. See [TV06].

Finally, we record the form of Freiman's theorem which we shall use.

Theorem 2.6. *A μ -additively closed set is contained in a subspace of dimension $N^{O(\mu)}$.*

Various improvements of the finite characteristic Freiman's theorem have appeared, such as the result by Sanders [S08], but these only affect the constant in our formulation. Even Ruzsa's original version [R99] suffices. We also mention a remarkable recent result in the integer setting due to Schoen [Sch11].

3. REVIEW OF MESHULAM'S ARGUMENT, FOURIER ANALYSIS IN F_3^N , AND SPARSITY OF THE SPECTRUM

For the remainder of this paper, A will be a subset of F_3^N with $|A| = \rho 3^N \gg \frac{3^N}{N^{1+\epsilon}}$ with $\epsilon > 0$ to be determined later. Moreover, A shall be a cap set, meaning that it contains no lines. A line in F_3^N is characterized by being a set with exactly three distinct elements a, b, c satisfying $a + b + c = 0$.

In this section we will establish some basic facts needed for our proof and that are enough to obtain Meshulam's bound of $\sim \frac{1}{N}$ on the density of cap sets. Further, we shall prove a statement of the form "The spectrum Δ does not have too much intersection with any small subspace."

We will assume that there are no strong increments for A in the sense of Definition 1.3. Precisely, we assume there is no hyperplane H so that $A \cap H$ has density $\geq \rho + \frac{20\rho}{N}$ in H and no subspace H of codimension $d < \frac{N}{2}$ so that $A \cap H$ has density $\geq \rho + \frac{20\rho d}{N}$. We recall that a contradiction of this assumption will mean that every large cap set A has strong increments. This will contradict the existence of large cap sets.

We define the character $e : F_3 \rightarrow \mathbf{C}$ by $e(0) = 1$, $e(1) = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$, and $e(2) = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$. We will study the Fourier transform of the set A , namely

$$\hat{A}(x) = \frac{1}{3^N} \sum_{a \in A} e(a \cdot x).$$

As a consequence of the assumption that A is a large cap set without strong increments, we shall see that there is a significant set Δ of x for which $|\hat{A}(x)|$ is fairly large (the set Δ will be called the spectrum of A) and we shall see that Δ does not concentrate too much in any fairly low dimensional subspace of F_3^N . Our first nontrivial fact about \hat{A} is that $\widehat{A - \rho}$ has large L^3 norm, and moreover that this large L^3 norm is accounted for by the set of x where $|\hat{A}(x)|$ is large. Precisely:

Definition 3.1. Define

$$\Delta = \{x \neq 0 : |\hat{A}(x)| \gtrsim \rho^2\}.$$

We shall refer to the set Δ as the *spectrum* of A and it shall be our central object of study for the remainder of the paper.

Note that with this definition, Δ is symmetric, that is,

$$\Delta = -\Delta.$$

It is worth noting that the following proposition is the only place in the paper where we use the assumption that A is a cap set specifically; the other parts of the paper use only the assumption that A has no strong increments.

Proposition 3.2. *If A is a cap set, then*

$$\sum_{x \neq 0} |\hat{A}(x)|^3 \gtrsim \rho^3.$$

Moreover,

$$\sum_{x \in \Delta} |\hat{A}(x)|^3 \gtrsim \rho^3,$$

and

$$N^3 \lesssim |\Delta| \lesssim N^{3+3\epsilon}.$$

Note that this proposition is already enough to obtain Meshulam’s estimate: in particular, it guarantees that the set Δ defined in the statement of the proposition is nonempty. This means that there is at least one x such that $|\hat{A}(x)| \gtrsim \rho^2$. This guarantees the existence of a hyperplane P such that the density of $A \cap P$ inside P is at least $\rho + c\rho^2$. Taking ρ large enough compared to $\frac{1}{N}$ already contradicts the no-strong-increment hypothesis, yielding Meshulam’s estimate.

In what follows, we prove this proposition. We consider

$$\sum_{x \in F_3^N} \hat{A}(x)^3 = \frac{1}{3^{3N}} \sum_{x \in F_3^N} \sum_{a \in A} \sum_{b \in A} \sum_{c \in A} e((a + b + c) \cdot x).$$

Summing first in x , we see that this expression yields 3^{-2N} multiplied by the number of solutions of the equation $a + b + c = 0$ with a, b, c taken from A . Since we have assumed that A is a cap set, the only solutions occur when $a = b = c$. Thus

$$\sum_{x \in F_3^N} \hat{A}(x)^3 = 3^{-2N} |A| = 3^{-N} \rho.$$

However, we observe that $\hat{A}(0) = \rho$. Thus, given the size of A , we see that ρ^3 dominates $3^{-N} \rho$ and we conclude that

$$(3.1) \quad \sum_{x \neq 0} |\hat{A}(x)|^3 \gtrsim \rho^3.$$

However, following the proof of Plancherel’s inequality, we see that

$$\sum_{x \neq 0} |\hat{A}(x)|^2 \leq \sum_{x \in F_3^N} |\hat{A}(x)|^2 = \frac{1}{3^{2N}} \sum_{x \in F_3^N} \sum_{a \in A} \sum_{b \in B} e((a - b) \cdot x).$$

Summing first in x , we conclude that

$$(3.2) \quad \sum_{x \neq 0} |\hat{A}(x)|^2 \leq 3^{-N} |A| \leq \rho.$$

By the assumption that A has no strong codimension 1 increment, we conclude that

$$(3.3) \quad |\hat{A}(x)| \lesssim \frac{|A|}{N^{3N}} = \frac{\rho}{N}.$$

Recall that

$$\Delta = \{x \neq 0 : |\hat{A}(x)| \gtrsim \rho^2\}.$$

By selecting the implicit constant in the definition of Δ correctly, we see by combining inequalities (3.1) and (3.2) that

$$(3.4) \quad \sum_{x \in \Delta} |\hat{A}(x)|^3 \gtrsim \rho^3.$$

Combining inequalities (3.3) and (3.4), we see that

$$|\Delta| \gtrsim N^3.$$

Combining the definition of Δ with Proposition 3.2 we get

$$|\Delta| \lesssim N^{3+3\epsilon}.$$

Now we prove a statement of the form “The spectrum Δ does not have too much intersection with any small subspace.” Precisely:

Proposition 3.3. *Let A be a set without strong increments. Let Δ be the spectrum, as in Definition 3.1. Then for any subspace W of F_3^N having dimension $d \leq \frac{N}{2}$, we have*

$$|\Delta \cap W| \lesssim dN^{1+2\epsilon}.$$

Moreover for such a subspace W , we have the estimate

$$\sum_{w \neq 0 \in W} |\hat{A}(w)|^2 \lesssim \rho^2 \frac{d}{N}.$$

Here we see what the assumption of no higher codimension strong increments implies about the spectrum Δ . Letting H be a subspace with codimension $d < \frac{N}{2}$, we let V be a dimension d subspace which is transverse to H (i.e., $V + H = F_3^N$) and we let W be the annihilator space of H . Then for any $w \neq 0 \in W$, we see that

$$\hat{A}(w) = \frac{1}{3^N} \sum_{v \in V} (|A \cap (H + v)| - 3^{-d}|A|)e(v \cdot w).$$

Then we have

$$(3.5) \quad \sum_{w \neq 0 \in W} |\hat{A}(w)|^2 = 3^{d-2N} \sum_{v \in V} (|A \cap (H + v)| - 3^{-d}|A|)^2.$$

By getting an upper bound on the right-hand side of (3.5), we can obtain an upper bound on $|\Delta \cap W|$, which is our goal.

To estimate the right-hand side, we subdivide $V = V^+ \cup V^-$, where

$$V^+ = \{v : |A \cap (H + v)| - 3^{-d}|A| \geq 0\}$$

and

$$V^- = \{v : |A \cap (H + v)| - 3^{-d}|A| < 0\}.$$

We observe that since

$$\sum_{v \in V} (|A \cap (H + v)| - 3^{-d}|A|) = 0,$$

we have

$$(3.6) \quad \sum_{v \in V^+} (|A \cap (H + v)| - 3^{-d}|A|) = \sum_{v \in V^-} (|A \cap (H + v)| - 3^{-d}|A|).$$

We observe that the hypothesis that A has no strong increments implies that for $v \in V^+$, we have the estimate

$$||A \cap (H + v)| - 3^{-d}|A|| \lesssim 3^{-d}|A| \frac{d}{N}.$$

Thus simply using that $|V| = 3^d$, we get the estimates

$$(3.7) \quad \sum_{v \in V^+} ||A \cap (H + v)| - 3^{-d}|A|| \lesssim |A| \frac{d}{N}$$

and

$$(3.8) \quad \sum_{v \in V^+} ||A \cap (H + v)| - 3^{-d}|A||^2 \lesssim 3^{-d}|A|^2 \frac{d^2}{N^2}.$$

Now for $v \in V^-$, we have the trivial estimate

$$||A \cap (H + v)| - 3^{-d}|A|| \leq 3^{-d}|A|.$$

In light of (3.6) and (3.7) this yields

$$(3.9) \quad \sum_{v \in V^-} ||A \cap (H + v)| - 3^{-d}|A||^2 \lesssim 3^{-d}|A|^2 \frac{d}{N}.$$

Combining (3.8) and (3.9) gives the estimate

$$\sum_{v \in V} ||A \cap (H + v)| - 3^{-d}|A||^2 \lesssim 3^{-d}|A|^2 \frac{d}{N}.$$

Thus (3.5) gives

$$\sum_{w \neq 0 \in W} |\hat{A}(w)|^2 \lesssim \rho^2 \frac{d}{N}.$$

However, we recall that if $w \in \Delta$, we have that

$$|\hat{A}(w)| \gtrsim \rho^2 = \frac{1}{N^{2+2\epsilon}}.$$

Thus we get the desired estimate:

$$|\Delta \cap W| \lesssim dN^{1+2\epsilon}.$$

4. ADDITIVE STRUCTURE IN THE SPECTRUM OF LARGE CAP SETS

In this section we establish that the spectrum has some nontrivial additive structure. Specifically, we prove that it has $N^{7-O(\epsilon)}$ additive quadruples. We mention that similar results in the integer setting were obtained independently by Shkredov; see [Shk08].

Corollary 4.1. *Let A be a large cap set. Let Δ be the spectrum of A and let Δ' be any symmetric subset of Δ with*

$$|\Delta'| \geq \frac{1}{2}|\Delta|.$$

Let $E_4(\Delta')$ be the number of additive quadruplets $x_1 + x_2 = x_3 + x_4$ with $x_1, x_2, x_3, x_4 \in \Delta'$. Then

$$E(\Delta') \gtrsim \frac{|\Delta|^4}{N^{5+5\epsilon}}.$$

The argument for a major subset Δ' of Δ is no different, so for convenience of notation we assume in fact $\Delta' = \Delta$.

We retain the notation of the previous section, considering Δ to be the spectrum of a large cap set A . In particular, we have $|A| \gg \frac{3^N}{N^{1+\epsilon}}$, we have

$$N^3 \lesssim |\Delta| \lesssim N^{3+3\epsilon},$$

we have for every $x \in \Delta$, that $|\widehat{A}(x)| \gtrsim \frac{|A|}{N^{1+\epsilon}}$, and we have that Δ is symmetric, namely $\Delta = -\Delta$.

From the lower bound on $|\widehat{A}(x)|$, we have for each x , an affine hyperplane H_x , annihilated by x so that

$$|A \cap H_x| - \frac{1}{3}|A| \gtrsim 3^N \frac{\rho}{N^{1+\epsilon}} = \frac{|A|}{N^{1+\epsilon}}.$$

Summing over Δ , we obtain

$$\sum_{x \in \Delta} (|A \cap H_x| - \frac{|A|}{3}) \gtrsim \frac{|A||\Delta|}{N^{1+\epsilon}}.$$

We wish to rewrite this as a double sum by introducing 1_{H_x} , the indicator function of H_x :

$$\sum_{x \in \Delta} \sum_{y \in A} (1_{H_x}(y) - \frac{1}{3}) \gtrsim \frac{|A||\Delta|}{N^{1+\epsilon}}.$$

We interchange the order of the sum:

$$\sum_{y \in A} (\sum_{x \in \Delta} (1_{H_x}(y) - \frac{1}{3})) \gtrsim \frac{|A||\Delta|}{N^{1+\epsilon}}.$$

Now we apply Hölder's inequality:

$$|A|^{\frac{3}{4}} (\sum_{y \in A} |\sum_{x \in \Delta} (1_{H_x}(y) - \frac{1}{3})|^4)^{\frac{1}{4}} \gtrsim \frac{|A||\Delta|}{N^{1+\epsilon}}.$$

Taking everything to the fourth power and simplifying, we get

$$\sum_{y \in A} |\sum_{x \in \Delta} (1_{H_x}(y) - \frac{1}{3})|^4 \gtrsim \frac{|A||\Delta|^4}{N^{4+4\epsilon}}.$$

Crudely expanding the sum, we get the apparently weaker inequality

$$\sum_{y \in F_3^N} |\sum_{x \in \Delta} (1_{H_x}(y) - \frac{1}{3})|^4 \gtrsim \frac{|A||\Delta|^4}{N^{4+4\epsilon}}.$$

We can rewrite this as

$$(4.1) \quad \sum_{y \in F_3^N} \sum_{x_1, x_2, x_3, x_4 \in F_3^N} \prod_{\alpha=1}^4 (1_{H_{x_\alpha}}(y) - \frac{1}{3}) \gtrsim \frac{|A||\Delta|^4}{N^{4+4\epsilon}}.$$

We claim that (4.1) says that the spectrum Δ has substantial additive structure. This will be demonstrated by the following proposition.

Proposition 4.2. *Let x_1, x_2, x_3 , and x_4 be nonzero elements of F_3^N . Then the expression*

$$\sum_{y \in F_3^N} \prod_{\alpha=1}^4 (1_{H_{x_\alpha}}(y) - \frac{1}{3}) \lesssim 3^N.$$

Moreover it vanishes unless an equality of the form

$$\pm x_1 \pm x_2 \pm x_3 \pm x_4 = 0$$

holds.

Proof. We introduce the Fourier transforms of the balanced function of the hyperplanes H_{x_α} setting

$$f_\alpha(z) = \frac{1}{3^N} \sum_{x \in F_3^N} (1_{H_{x_\alpha}}(y) - \frac{1}{3})e(y \cdot z).$$

Then we use the standard Fourier identity

$$\sum_{y \in F_3^N} \prod_{\alpha=1}^4 (1_{H_{x_\alpha}}(y) - \frac{1}{3}) = 3^N \sum_{z_1+z_2+z_3+z_4=0} f_1(z_1)f_2(z_2)f_3(z_3)f_4(z_4).$$

We observe that $f_j(z_j)$ vanishes unless $z_j = \pm x_j$.

The upper bound on the sum just follows from the triangle inequality. □

To finish the proof of Proposition 4.1, we apply (4.1), the fact that $|A| \gg \frac{3^N}{N^{1+\epsilon}}$, Proposition 4.2 and the fact that the spectrum Δ is symmetric.

5. RANDOM SELECTION ARGUMENT FOR ADDITIVELY SMOOTHING SPECTRUM

In this section we study the additive properties of random subsets of the spectrum. We will show that they typically have very poor additive structure. This will allow us to conclude that, although the spectrum has many 4-tuples, it cannot have too many 8-tuples. The significance of this will only be made clear in Section 6.

We defined $E_4(\Delta)$ to be the number of additive quadruplets in Δ .

Definition 5.1. We define $E_{2m}(\Delta)$ to be the number of additive $2m$ -tuples

$$x_1 + x_2 + \dots + x_m = x_{m+1} + x_{m+2} + \dots + x_{2m},$$

such that $x_1, x_2, \dots, x_{2m} \in \Delta$.

We let $\widehat{\Delta}(x)$ be the Fourier transform:

$$\widehat{\Delta}(y) = \frac{1}{3^N} \sum_{x \in \Delta} e(y \cdot x).$$

Then

$$E_{2m}(\Delta) = 3^{(2m-1)N} \sum_{y \in F_3^N} |\widehat{\Delta}(y)|^{2m}.$$

We always have $E_2(\Delta) = |\Delta|$. When we have nontrivial amounts of additive structure in the sense that say $E_{2k}(\Delta) \gg |\Delta|^k$, we can lift this up to counts of higher-tuples using Hölder’s inequality. (We use the inequality to bound the $2k$ -norm by the 2-norm and the $2m$ -norm.) We can view this process as a poor man’s Plunnecke theorem. We record this result for high E_4 and high E_8 .

Lemma 5.2. *Let $m > 2$. Then*

$$\frac{E_4(\Delta)^{m-1}}{|\Delta|^{m-2}} \leq E_{2m}(\Delta).$$

Suppose $m > 3$. Then

$$\frac{E_8(\Delta)^{\frac{m-1}{3}}}{|\Delta|^{\frac{m-4}{3}}} \leq E_{2m}(\Delta).$$

5.1. Discussion of additive smoothing. We are now ready to introduce the notion of additive smoothing. We keep in mind two examples of kinds of sets having additive structure. One kind of set consists of a subspace plus a random set. The other consists of a random subset of a subspace. We think of the first kind of set as not being additively smoothing because as you add it to itself, its expansion rate stays essentially constant. This is the kind of example for which Lemma 5.2 is close to sharp. But the second kind of set, when added to itself, will quickly fill out the subspace, and its rate of additive expansion will shrink dramatically. The lack of additive structure smooths out under addition. This is the kind of example for which Lemma 5.2 is far from sharp.

We will momentarily define Δ to be additively smoothing if $E_8(\Delta)$ is substantially larger than expected from Lemma 5.2 and our lower bound for $E_4(\Delta)$ obtained in Section 4. (Nonetheless the gain in the exponent need only be $O(\epsilon)$.) We will define additive smoothing so that if Δ is additively smoothing, then for some not very large m , we may expect to find additive m -tuplets of Δ in a randomly chosen set S of d elements.

Before we formally define the property of additive smoothing, we illustrate how the calculation works in the case $\epsilon = 0$. In that case $d \sim N$ so that an element of Δ (which has size N^3) is chosen with probability N^{-2} . We have a lower bound of N^7 on $E_4(\Delta)$. Suppose that we can improve on the lower bound of N^{15} for $E_8(\Delta)$, which we get from the first part of Lemma 5.2, and in fact

$$E_8(\Delta) > N^{15+\delta}$$

for some $\delta > 0$. Then from the second part of Lemma 5.2, we obtain the estimate

$$N^{(4+\frac{\delta}{3})m+7-\frac{\delta}{3}} \leq E_{2m}(\Delta).$$

Thus there is some m which depends only on δ so that

$$E_{2m}(\Delta) \gg N^{4m}.$$

Thus the expected number of $2m$ -tuplets in S is $\gg 1$. We will formally define additive smoothing to achieve the same effect when ϵ is different from 0.

5.2. Nonsmoothing of the spectrum. In this subsection we make rigorous the arguments of the last subsection.

Definition 5.3. We define Δ to be additively smoothing if there is some $\sigma > 30\epsilon$ so that $E_8(\Delta) \gg N^{15+\sigma}$.

We are now in a position to state the main result of this section.

Lemma 5.4. *If Δ is the spectrum of a large cap set without strong increments, then Δ is not additively smoothing.*

We begin with a few comments about our proof strategy in this section. If S is a “random” subset of Δ , then we expect

$$E_{2m}(S) = \left(\frac{|S|}{|\Delta|} \right)^{2m} E_{2m}(\Delta).$$

Thus we can show that $E_{2m}(\Delta)$ is small by showing that E_R is small for a typical (somewhat large) subset R of Δ .

Now we fix a particular number d and consider random subsets of Δ of size d . We will take

$$(5.1) \quad d \sim N^{1-\epsilon}$$

with the explicit constant to be determined later. Our first goal is to prove that we expect this subset to span a space of dimension d . More precisely:

Definition 5.5. Let S be a set of d vectors $x_1, \dots, x_d \in F_3^N$. We say that the set S has nullity k if the dimension of the span of S is $d - k$.

We will consider uniform random selections of sets of d elements from Δ . We can view these selections as d -fold repetitions of uniform selection without replacement. We will prove

Lemma 5.6. *A random selection S of size d from the spectrum Δ has nullity at least k with probability $\lesssim 2^{-k}$.*

Proof. Once we have completed our first m choices, our selections x_1, \dots, x_m span a vector space W_m with dimension no more than m . Thus $|\Delta \cap W_m| \lesssim mN^{1+2\epsilon}$ by Proposition 3.3. We choose the constant in (5.1) so that the probability that the $(m + 1)$ st element of S lies in W_m is bounded by $\frac{1}{d}$ for all $m \leq d - 1$. Note that since $m \leq d$, this probability is bounded by

$$\begin{aligned} \frac{|\Delta \cap W_m|}{|\Delta|} &\leq \frac{CdN^{1+2\epsilon}}{N^3} \\ &= \frac{Cd}{N^{2-2\epsilon}} \\ &\leq \frac{1}{d} \end{aligned}$$

provided $d \ll N^{1-\epsilon}$. Thus the probability that S has nullity at least k is bounded by the probability that for d independent events with probability $\frac{1}{d}$ at least k occur.

The probability that exactly k events from d independent events with probability $\frac{1}{d}$ occur is exactly

$$g(k, d) = \binom{d}{k} \frac{(d - 1)^{d-k}}{d^d}.$$

The numbers $g(k, d)$ decrease by a factor of more than 2 as k is increased by 1 as long as $k > 2$. This completes the proof of the lemma □

Now that we know our random subset is likely to have full rank, we estimate the number of $2m$ -tuples it contains in the case it does not have full rank. Given a set S with nullity k we will bound the number of possible additive $2m$ -tuplets between elements of S . Specifically:

Lemma 5.7. *A set S of size d and nullity k has $E_{2m}(S) \lesssim C_m k^{2m}$.*

Proof. We write a list E of all equations among elements of S which involve $2m$ or fewer elements of S . Because the nullity is k , the span of these equations has dimension at most k . We pick a basis B for E , and the equations in B involve

at most $2mk$ elements of S . Thus all of the equations of E involve at most $2mk$ elements of S . Thus there are at most

$$h(m, k) = 2^m(2m)! \binom{2mk}{2m}$$

additive $2m$ -tuplets from S . We refer to $h(m, k)$ as the number of possible m -tuplets in S . Note that $h(m, k)$ is a polynomial of degree $2m$ in k . □

Proof of Lemma 5.4 . Now let S be a random selection of d elements from Δ . Then by Lemma 5.6, the probability that S has nullity k is $\lesssim 2^{-k}$. Thus the expected value of the number of possible $2m$ -tuples $\sum_{k \geq 0} h(m, k)$ is $\lesssim_m \sum_{k \geq 0} 2^{-k} k^{2m} \lesssim 1$.

Now we will show that we have defined additive smoothing so that the expected number of $2m$ -tuples is $\gg 1$. This will give us a contradiction.

We know that $d \gtrsim N^{1-\epsilon}$. Thus our selection S will be expected to have $\gg 1$ nontrivial $2m$ -tuples, whenever $E_{2m}(\Delta) \gg N^{4m+2m\epsilon}$. (We simply calculate the probability that an individual $2m$ -tuple involves only elements of S .) Thus we may assume that

$$E_{2m}(\Delta) \lesssim N^{4m+8m\epsilon}.$$

Using the fact that $|\Delta| \lesssim N^{3+3\epsilon}$ and the second part of Lemma 5.2, we get that

$$E_8(\Delta) \lesssim N^{\frac{15m+27m\epsilon-1-\epsilon}{m-1}}.$$

Choosing m sufficiently large gives

$$E_8(\Delta) \lesssim N^{15+27\epsilon}.$$

The choice of m and hence the constants depends on ϵ but not on N . □

6. STRUCTURE OF ROBUST ADDITIVELY NONSMOOTHING SETS

In this section, the only properties of the spectrum Δ which we shall use are its size, its additive structure, and its nonadditive smoothing. Consequently the results can be stated in somewhat more generality. We leave intact, however, the numerology coming from the case of a spectrum of cap sets.

Definition 6.1. We will say that a symmetric set $\Delta \subset F_3^N$ is a *robust additively nonsmoothing set of strength δ* provided that we know its size:

$$(6.1) \quad N^3 \lesssim |\Delta| \lesssim N^{3+\delta},$$

that we know how many additive quadruples can be made from any large subset of it, namely that if $\Delta' \subset \Delta$ with $|\Delta'| \geq \frac{3}{5}|\Delta|$ and Δ' symmetric, we have

$$(6.2) \quad E_4(\Delta') \gtrsim N^{7-\delta}$$

and that we have additive nonsmoothing, namely

$$(6.3) \quad E_8(\Delta) \lesssim N^{15+\delta}$$

and moreover that for each element $a \in \Delta$, there are at most $N^{4+\delta}$ quadruples of the form

$$\pm a \pm b = \pm c \pm d$$

with $b, c, d \in \Delta$.

Let us pause to consider the case of Δ , the spectrum of a cap set with no strong increments. We know that the number of $a \in \Delta$ participating in more than $N^{4+O(\epsilon)}$ quadruplets

$$\pm a \pm b = \pm c \pm d$$

is smaller than $\frac{1}{10}|\Delta|$ since otherwise Δ would have more quadruples and hence more octuples than allowed by Lemma 5.4. Let Δ' be the remaining elements of Δ . Note that by its definition Δ' is still symmetric. Note that any symmetric subset of Δ' containing at least three fifths of its elements must contain at least half the elements of Δ . Thus from Proposition 3.2, Corollary 4.1, and Lemma 5.4 we know that:

Proposition 6.2. *Let Δ be the spectrum of a large capset with no strong increments. There is a subset Δ' of Δ so that Δ' is a robust additively nonsmoothing set of strength $O(\epsilon)$.*

Returning to the setting of robust additively nonsmoothing sets, we let, for the remainder of the section, the set Δ be a robust additively nonsmoothing set of strength δ .

Given a value $x \in \Delta - \Delta$, we define $m(x)$ to be the number of pairs $(a, b) \in \Delta \times \Delta$ so that $a - b = x$. Clearly we have

$$E_4(\Delta) = \sum_{x \in \Delta - \Delta} m(x)^2.$$

Given a robust additively nonsmoothing set Δ of strength δ , for each α , we may define $G_\alpha \subset \Delta \times \Delta$ by

$$G_\alpha = \{(a, b) \in \Delta \times \Delta : N^{1+\alpha} \leq m(a, b) < 2N^{1+\alpha}\}.$$

By the dyadic pigeonhole principle, there is an α so that

$$|G_\alpha| \gtrsim \frac{N^{6-\alpha-\delta}}{\log N}.$$

Moreover, we know that no a in Δ participates in more than $N^{4+\delta}$ quadruples. Thus no element a in Δ participates in more than $N^{3-\alpha+\delta}$ pairs in G_α . Thus there are at least $\frac{N^{3-2\delta}}{\log N}$ elements of Δ , each of which participates in at least $\frac{N^{3-\alpha-2\delta}}{\log N}$ pairs in G_α .

We now forget about optimizing our exponents and consolidate this information in a single definition.

Definition 6.3. We say that (Δ, G, D) is an *additive structure at height α with ambiguity η* if the following hold. We have

$$|\Delta| \leq N^{3+\eta}.$$

We have

$$G \subset \Delta \times \Delta$$

with the property that for each $(a, b) \in G$ we have that $a - b \in D$, and so that each $d \in D$ has $\sim N^{1+\alpha}$ representations as a difference of a pair in G . We have $|G| \sim N^{6-\alpha-\eta}$. Moreover there are at least $N^{3-\eta}$ elements of Δ participating in at least $N^{3-\alpha-\eta}$ sums each. Finally there are no more than $N^{15+\eta}$ additive octuples among elements of Δ .

We summarize what we have shown so far in a proposition.

Proposition 6.4. *Given a robust additively nonsmoothing set Δ of strength δ we may find $G \subset \Delta \times \Delta$, and $D \subset \Delta - \Delta$ and $\alpha \geq 0$ so that (Δ, G, D) is an additive structure at height α and ambiguity $O(\delta)$.*

We now describe a slightly deeper property of additive structures at height α and ambiguity η . Given a structure (Δ, G, D) , for each $x \in D$, we define the set $\Delta_G[x]$ to be the set of $a \in \Delta$ so that there exists $b \in \Delta$ with $(a, b) \in G$ and $a - b = x$. In light of our definitions, we have for each $x \in D$ that $|\Delta_G[x]| \sim N^{1+\alpha}$. We consider the quantity

$$(6.4) \quad K(\Delta, G, D) = \sum_{x \in D} \sum_{y \in D} |\Delta_G[x] \cap \Delta_G[y]|.$$

Clearly $K(\Delta, G, D)$ counts the number of triples (a, x, y) with $a \in \Delta_G[x]$ and $a \in \Delta_G[y]$. Each element in a is contained in exactly as many sets $\Delta_G[x]$ as it participates (in the first position) in pairs in G . Thus we conclude that for (Δ, G, D) an additive structure with height α and ambiguity η that

$$K(\Delta, G, D) \gtrsim N^{9-2\alpha-3\eta}.$$

Now examining (6.4) and dyadically pigeonholing, we observe that we can find β so that there are at least $N^{9-2\alpha-\beta-4\eta}$ pairs (x, y) so that for each such pair, we have $N^\beta \leq |\Delta_G[x] \cap \Delta_G[y]| < 2N^\beta$.

Definition 6.5. We say that the additive structure (Δ, G, D) at height α and with ambiguity η has *comity* μ if we can find the above-mentioned β with $\beta > 1 + \alpha - \mu$.

Lemma 6.6. *Given an additive structure (Δ, G, D) at height α and with ambiguity η either it has comity μ or there is an additive structure (Δ, G', D') with height $\beta - 1 < \alpha - \mu$ and ambiguity $O(\eta)$.*

Proof. We dyadically pigeonhole (6.4) to find β so that there is a set of at least $N^{9-2\alpha-\beta-4\eta}$ pairs (x, y) so that for each such pair, we have $N^\beta \leq |\Delta_G[x] \cap \Delta_G[y]| < 2N^\beta$. If it happens that $\beta - 1 > \alpha - \mu$, then we are done. Otherwise, we will construct an additive structure at height $\beta - 1$.

Now any time that $a \in \Delta_G[x] \cap \Delta_G[y]$, this means that we can write $x = a - b$ and $y = a - c$. Thus we have $x - y = c - b$. We have between N^β and $2N^\beta$ representations of the difference $x - y$. It remains to determine how many such differences there are.

We have two distinct upper bounds on the number of such differences. First there are $\lesssim N^{6-\beta+2\eta}$, since each difference is represented by $\sim N^\beta$ pairs in $\Delta \times \Delta$ and there are only $N^{6+2\eta}$ such pairs. The second estimate is that there are $\lesssim N^{7-2\beta+O(\eta)}$ many such differences, because otherwise $E_4(\Delta)$ would be much larger than N^7 which would make $E_8(\Delta)$ larger than $N^{15+\eta}$. The first upper bound is most effective (ignoring ambiguity) when $\beta < 1$ while the second is most effective when $\beta > 1$. Our plan (modulo ambiguity) is that we shall rule out the case $\beta < 1$ and that we shall show that the second upper bound is tight up to a factor of $N^{O(\eta)}$. Both estimates will follow from the upper bound on $E_8(\Delta)$ and the Cauchy-Schwarz inequality, namely Lemma 2.1.

Since we have $N^{9-2\alpha-\beta-O(\eta)}$ pairs (x, y) with at most $N^{6-\beta+O(\eta)}$ differences, by the Cauchy-Schwarz inequality, there must be at least $N^{12-4\alpha-\beta-O(\eta)}$ additive quadruples in D , namely $x - y = x' - y'$. (Here we let S be the set of pairs (x, y) , we let T be the set of differences with $\sim N^\beta$ representations as difference of Δ and

we let ρ be the difference map, $\rho(x, y) = x - y$. Then we can apply Lemma 2.1.) However since each difference x, y can be represented in $N^{1+\alpha}$ ways as a difference in Δ , we can represent each quadruple in D as an octuple in Δ in $N^{4+4\alpha}$ ways. Thus there are at least $N^{16-\beta-O(\eta)}$ many such octuples which implies $\beta \geq 1 - O(\eta)$.

Thus we are in the regime where the estimate that there are at most $N^{7-2\beta+O(\eta)}$ many differences is most effective. Suppose that there were only $N^{7-2\beta-\gamma}$ many such differences with $\gamma \gg \eta$. Then applying Cauchy-Schwarz again, we would see that there are at least $N^{11-4\alpha-O(\eta)+\gamma}$ many quadruples in D which implies $N^{15-O(\eta)+\gamma}$ octuples in Δ , a contradiction.

Thus taking D' to be the differences $x - y$ obtained from (x, y) so that $N^\beta \leq |\Delta[x] \cap \Delta[y]| < 2N^\beta$ and taking G' to consist of representatives of these differences coming from the intersections, we obtain an additive structure (Δ, G', D') with height $\beta - 1 < \alpha - \mu$ and ambiguity $O(\eta)$. \square

Corollary 6.7. *Given an additive structure (Δ, G, D) at height α and with ambiguity η there is an additive structure (Δ, G, D) at height $\alpha' \leq \alpha$ with ambiguity μ and comity μ with $\mu \lesssim \frac{1}{\log \frac{1}{\eta}}$.*

Proof. We iteratively apply Lemma 6.6 with comity μ fixed by

$$\mu = \frac{K}{\log \frac{1}{\eta}}$$

with K a large constant, and with the ambiguity increasing by a constant factor C in each iteration. Since α decreases by μ each time we don't find comity we need only $\frac{1}{\mu}$ iterations to achieve comity. At this point, we have ambiguity given by $C^{\frac{\log(\frac{1}{\eta})}{K}} \eta \ll \mu$, as long as K was chosen sufficiently large. \square

Now we begin to investigate what we can say about the shape of the set H of all pairs (b, c) in $\Delta \times \Delta$ having the property that $b - c$ has at least $N^{1+\alpha-O(\mu)}$ representations in $\Delta \times \Delta$ for (Δ, G, D) an additive structure with height α and ambiguity and comity μ . We will find that the set H is rather thick in a product set whose projection has size $N^{3-\alpha-O(\mu)}$.

Lemma 6.8. *Let (Δ, G, D) be an additive structure with height α and ambiguity and comity μ . Then there is a subset $B \subset \Delta$ with $|B| \gtrsim N^{3-\alpha-O(\mu)}$ so that there is a set $H \subset B \times B$ with*

$$|H| \gtrsim N^{-O(\mu)} |B|^2,$$

so that for any $(b, c) \in H$, the difference $b - c$ has $N^{1+\alpha-O(\mu)}$ representations in $\Delta \times \Delta$.

Proof. From the hypotheses, we have that

$$\sum_{x \in D} \sum_{y \in D} |\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^{9-2\alpha-O(\mu)},$$

and that there are at least $N^{8-3\alpha-O(\mu)}$ pairs (x, y) for which

$$|\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^{1+\alpha-O(\mu)}.$$

Using the pigeonhole principle, we fix one value of x for which $N^{3-\alpha-O(\mu)}$ choices of y exist such that

$$|\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^{1+\alpha-O(\mu)}.$$

Again using the pigeonhole principle, we find an $a \in \Delta$ and a set $Y \subset D$ so that $a \in \Delta[y]$ for every $y \in Y$, so that $|Y| = N^{3-\alpha-O(\mu)}$ and so that for each $y \in Y$, we have

$$|\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^{1+\alpha-O(\mu)}.$$

We notice that by definition, $a-Y \subset \Delta$. We choose $B = a-Y$. Finally since each $\Delta_G[y]$ has relative density $N^{-O(\mu)}$ in $\Delta_G[x]$, we have by Cauchy-Schwarz (Lemma 2.2) that

$$\sum_{y \in Y} \sum_{y' \in Y} |\Delta_G[y] \cap \Delta_G[y']| \gtrsim |Y|^2 N^{1+\alpha-O(\mu)}.$$

This implies that B satisfies the conclusion of the lemma. The reason is that by Lemma 2.3, we have a set \tilde{Y} of pairs y, y' so that $|\Delta[y] \cap \Delta[y']| \gtrsim N^{1+\alpha-O(\mu)}$. This implies that $y - y'$ has at least $N^{1+\alpha-O(\mu)}$ representatives as a difference of two elements of Δ . \square

Now we are going to use Lemma 6.8 repeatedly to show that for any robust additively nonsmoothing set of size δ we can find an additive structure of ambiguity η with $\eta \lesssim \frac{1}{\log \frac{1}{\delta}}$ which breaks into dense blocks.

Lemma 6.9. *Let Δ be a robust additively nonsmoothing set of strength δ . Choose $\mu \sim \frac{1}{\log \frac{1}{\delta}}$. Then for some $0 \leq \alpha \leq 1$, there is an additive structure (Δ, G, D) of height α and ambiguity μ and disjoint subsets B_1, \dots, B_K of Δ with each B_j satisfying $|B_j| \lesssim N^{3-\alpha+O(\mu)}$ so that*

$$G \subset \bigcup_{j=1}^K B_j \times B_j.$$

Note that since we are requiring that (Δ, G, D) be an additive structure, this requires $|G| \gtrsim N^{6-\alpha-O(\mu)}$, which implies $K \gtrsim N^{\alpha-O(\mu)}$.

Proof. Using Proposition 6.4, Corollary 6.7, and Lemma 6.8. We can find a subset B_1 of Δ so that for some choice of α , it has size at most $N^{3-\alpha+O(\mu)}$ and nevertheless $B_1 \times B_1$ contains at least $N^{6-2\alpha-O(\mu)}$ pairs whose differences have $N^{1+\alpha-O(\mu)}$ representations as differences in Δ .

Having done this, we use the robustness property of Δ to apply the same argument to $\Delta \setminus (B_1 \cup -B_1)$. We continue removing sets from Δ until we have exhausted half of Δ . Now one difficulty is that the disjoint sets B which we chose do not all have the same α . We use dyadic pigeonholing to resolve this for only a small cost in the number of sets. We call these sets B_1, \dots, B_K . Now Lemma 6.8 guarantees us in each $B_j \times B_j$, a subset H_j of cardinality at least $N^{6-2\alpha-O(\mu)}$ so that each difference in H_j is represented in $\Delta \times \Delta$ at least $N^{1+\alpha-O(\mu)}$ times. We denote by D_α the set of differences represented in $\Delta \times \Delta$ at least $N^{1+\alpha-O(\mu)}$ times, and note that

$$|D_\alpha| \lesssim N^{5-2\alpha+O(\mu)},$$

lest there be enough quadruples in Δ to violate the additive nonsmoothing condition. Pigeonholing, we find some $\alpha' \gtrsim \alpha - O(\mu)$ so that at least $N^{5-2\alpha-O(\mu)}$ differences are represented at least $N^{1+\alpha'}$ many times in $\bigcup_j H_j \times H_j$. We denote this set of differences as D'_α . We let D be D'_α and let G be a subset of $\bigcup_j H_j \times H_j$ consisting of $N^{1+\alpha'}$ representatives of each difference in D . \square

Our goal now will be to use Lemma 6.9 to find almost additively closed sets E of size at least $N^{1-f(\mu)}$ inside robust nonadditively smoothing sets of strength δ . Here $f : [0, 1] \rightarrow [0, \infty)$ is some function with $\lim_{t \rightarrow 0} f(t) = 0$. We will be employing such functions from now on in the paper. They, like constants, will change from line to line.

The project of finding additively closed sets will be easiest when we have additive structures of height zero having ambiguity and comity μ . For this reason, we are about to define a stylized structure which generalizes this situation. We will eventually use the generalized version, replacing Δ with the blocks B_j .

Definition 6.10. We will now define a μ -full stylized ρ -structure which is τ -energetic and has ambiguity and comity μ . (The error exponents μ are all the same.) This will be a set (Δ', G, D) , where

$$|\Delta'| \sim N^\rho$$

(hence a ρ -structure), where $G \subset \Delta' \times \Delta'$ with

$$|G| > N^{2\rho-O(\mu)}$$

(this was the μ -fullness), where D is the set of differences in pairs in G and each difference represented $\gtrsim N^\tau$ and $\lesssim N^{\tau+O(\mu)}$ times, hence τ -energetic. Finally we assume that there are at least $N^{3\rho-\tau-O(\mu)}$ pairs $(x, y) \in D \times D$ so that

$$|\Delta'[x] \cap \Delta'[y]| \gtrsim N^{\tau-O(\mu)},$$

which is of course the μ -comity.

Definition 6.11. We shall say that a set K is μ -additively closed provided that

$$|K - K| \lesssim N^{O(\mu)}|K|.$$

Lemma 6.12. *There is a function $f : [0, 1] \rightarrow [0, \infty)$ with*

$$\lim_{t \rightarrow 0} f(t) = 0,$$

so that the following holds. Let (Δ', G, D) be a μ -full stylized ρ -structure which is τ -energetic and has ambiguity and comity μ . Then there is an $f(\mu)$ additively closed set K with

$$|K| \gtrsim N^{\tau-f(\mu)}$$

and a set X so that

$$|X| \lesssim N^{f(\mu)} \frac{|\Delta'|}{|K|},$$

so that

$$|\Delta' \cap (X + K)| \gtrsim N^{\rho-f(\mu)}.$$

Proof. We proceed essentially as in the proof of Lemma 6.8. We find $x \in D$ so that there is a set Y of $y \in D$ with $|Y| \gtrsim N^{\rho-O(\mu)}$ so that

$$|\Delta'_G[x] \cap \Delta'_G[y]| \gtrsim N^{\tau-O(\mu)},$$

for every $y \in Y$. As before, we use the pigeonhole principle to find $a \in \Delta'$ so that there is a subset Y_a of Y so that for each $y \in Y_a$, we have

$$a \in \Delta'_G[y]$$

and so that

$$|Y_a| \gtrsim N^{\rho-O(\mu)}.$$

However $Y_a \subset a - A$. Thus we think of Y_a as a dense part of a translate of $-A$. Now we know that

$$\sum_{y \in Y_a} |\Delta'_G[x] \cap \Delta'_G[y]| \gtrsim N^{\rho+\tau-O(\mu)}.$$

This precisely means that there are $N^{\rho+\tau-O(\mu)}$ triples (b, c, d) with $b, c \in \Delta'$ and $d \in \Delta'[x]$ with $a - b = d - c$ with a still fixed. Applying Cauchy-Schwarz we find $N^{\rho+2\tau-O(\mu)}$ quadruples (d, c, d', c') with $d, d' \in \Delta'[x]$ and $c, c' \in \Delta'$. As it happens, this is precisely the hypothesis of the asymmetric Balog-Szemerédi-Gowers theorem (Theorem 2.5) applied to Δ' and $\Delta_G[x]$. The conclusion follows directly. \square

We are now prepared to state the main result of this section.

Theorem 6.13. *Let Δ be a robust additively nonsmoothing set of strength δ . As before choose $\mu \sim \frac{1}{\log \frac{1}{\delta}}$. There is $f : [0, 1] \rightarrow [0, \infty)$ with*

$$\lim_{t \rightarrow 0} f(t) = 0,$$

so that for some $\gamma \geq 0$, there is an $f(\mu)$ -additively closed set K with

$$|K| \gtrsim N^{1+\gamma-f(\mu)},$$

contained in Δ . In the event that we must have $\gamma = O(f(\mu))$, for some $0 \leq \alpha \leq 1$, we may find pairwise disjoint subsets $B_1, \dots, B_M \subset \Delta$ with $M \gtrsim N^{\alpha-O(\mu)}$ so that for each integer $1 \leq j \leq m$, we have

$$N^{3-\alpha-O(\mu)} \lesssim |B_j| \lesssim N^{3-\alpha+O(\mu)},$$

and moreover we find for each j a μ -additively closed set K_j with

$$N^{1-f(\mu)} \lesssim |K_j| \lesssim N^{1+f(\mu)},$$

together with a set X_j with

$$N^{2-\alpha-f(\mu)} \lesssim |X_j| \lesssim N^{2-\alpha+f(\mu)},$$

so that

$$|B_j \cap (X_j + K_j)| \gtrsim N^{3-\alpha-f(\mu)}.$$

Further, there is a set D with $|D| \gtrsim N^{5-2\alpha+f(\mu)}$ so that each element of D has at least $N^{1+\alpha-f(\mu)}$ representations as a difference of elements of Δ and so that for each j , the set of 4-tuples $Q_j = \{(k_1, d_1, k_2, d_2) : k_1, k_2 \in K_j, d_1, d_2 \in D : k_1 - d_1 = k_2 - d_2\}$ satisfies

$$|Q_j| \gtrsim N^{7-2\alpha-f(\mu)}.$$

Moreover we may choose K_j to be contained in the set of differences having at least $N^{5-2\alpha-f(\mu)}$ representations as a difference between elements of D .

Proof. We apply Lemma 6.9 and restrict our attention to

$$G \subset \bigcup_{j=1}^K B_j \times B_j,$$

where the B_j are the blocks obtained there. Now to G , we apply the argument used in the proof of Lemma 6.6.

That is, for any element $x \in -(G)$, we study

$$\Delta_G[x] = \{a \in \Delta : \exists b \in \Delta : (a, b) \in G; \quad a - b = x\}.$$

Then, we observe

$$\sum_{x,y \in -\langle G \rangle} |\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^{9-2\alpha-O(\mu)},$$

and we observe as before that there is some $1 + \alpha \geq \beta \geq 1 - O(\mu)$ so that there are at least $N^{9-2\alpha-\beta-O(\mu)}$ pairs (x, y) for which $|\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^\beta$. We note that when this happens, for each a in the intersection $\Delta_G[x] \cap \Delta_G[y]$, we have $x = a - b_1$ and $y = a - b_2$. Here if $a \in B_j$, we must have $b_1 \in B_j$ and $b_2 \in B_j$, since $(a, b_1), (a, b_2) \in G$. We argue as in the proof of Lemma 6.6, that there are at least $N^{7-2\beta-O(\mu)}$ differences having N^β representations in

$$\bigcup_{j=1}^K B_j \times B_j,$$

which is impossible unless $\beta \geq \alpha - O(\mu)$ since there are at most $N^{6-\alpha+O(\mu)}$ pairs in

$$\bigcup_{j=1}^K B_j \times B_j.$$

Clearly this gives us $O(\mu)$ comity for G .

Thus we have a set H of pairs x, y for which $|\Delta_G[x] \cap \Delta_G[y]| \gtrsim N^{1+\alpha-O(\mu)}$ and so that $|H| \gtrsim N^{8-3\alpha-O(\mu)}$. Now we use the pairwise disjointness of the blocks B_j to write the identity

$$(6.5) \quad \sum_{(x,y) \in H} |\Delta_G[x] \cap \Delta_G[y]| = \sum_{j=1}^K \sum_{(x,y) \in H} |\Delta_G[x] \cap \Delta_G[y] \cap B_j| \gtrsim N^{9-2\alpha-O(\mu)}.$$

Now we begin to use the comity of G . We first eliminate from the second sum in (6.5) all terms for which the relative density of $\Delta_G[x]$ in $\Delta_G[y] \cap B_j$ or the relative density of $\Delta_G[y]$ in $\Delta_G[x] \cap B_j$ is smaller than $N^{-C\mu}$ for too large a constant C . By choosing C sufficiently large we do not reduce the sum by a factor of more than 2. We dyadically pigeonhole to obtain the largest possible sum from those terms where $N^{1+\gamma} \leq |\Delta_G[x] \cap \Delta_G[y] \cap B_j| \leq 2N^{1+\gamma}$. (We denote this set of (x, y) as $H_{\gamma,j}$.) Thus we have reduced the sum by at most a factor of $\log N$. We keep only those j for which

$$(6.6) \quad \sum_{(x,y) \in H_{\gamma,j}} |\Delta_G[x] \cap \Delta_G[y] \cap B_j| \gtrsim N^{9-3\alpha-O(\mu)}.$$

We observe that for each x , there are at most $N^{3-\alpha+O(\mu)}$ choices of y so that $(x, y) \in H$. The reason is that any $a \in \Delta$ belongs to at most $N^{3-\alpha+O(\mu)}$ sets $\Delta_G[y]$ because G is contained in $\bigcup_j B_j \times B_j$. However if there were more than $N^{3-\alpha+O(\mu)}$ choices of y so that $(x, y) \in H$, then there would be elements $a \in \Delta_G[x]$ which are contained in $\Delta_G[y]$ for more than $N^{3-\alpha+O(\mu)}$ choices of y . Thus, since we have at least $N^{8-2\alpha-\gamma-O(\mu)}$ triples (j, x, y) for which

$$|\Delta_G[x] \cap \Delta_G[y] \cap B_j| \gtrsim N^{1+\gamma},$$

while at the same time

$$|\Delta_G[x] \cap B_j| \lesssim N^{1+\gamma+O(\mu)},$$

it must be that for some values of j , we must have at least $N^{8-3\alpha-\gamma}$ pairs $(x, y) \in H_{\gamma,j}$. This means, fixing one such value of j (since at most $N^{3-\alpha+O(\mu)}$ values of

y are paired with a given x), there are at least $N^{5-2\alpha+\gamma-O(\mu)}$ differences x with $N^{1+\gamma}$ representations in $G \cap (B_j \times B_j)$. We call this set $D_{j,\gamma}$.

Thus $G \cap (B_j \times B_j)$ is μ -full and $(1+\gamma)$ -energetic. Another way of describing the μ -fullness is that $N^{5-2\alpha+\gamma-O(\mu)}$ (up to $N^{O(\mu)}$ factors) is the largest number of such x possible, purely based on the size of $B_j \times B_j$. Thus it must be that for a set of size $N^{5-2\alpha+\gamma-O(\mu)}$ many such x , there are $N^{3-\alpha-O(\mu)}$ such y with $(x, y) \in H_{\gamma,j}$. Thus $(B_j, G \cap (B_j \times B_j), D_{j,\gamma})$ has $O(\mu)$ -comity. Clearly $(B_j, G \cap (B_j \times B_j), D_{j,\gamma})$ is a $3-\alpha$ structure with ambiguity μ . Thus we are in a position to apply Lemma 6.12. This proves the first part of the theorem. (Indeed, since all our estimates were optimal up to $N^{O(\mu)}$ factors, there is a set J of choices of j for which we could apply Lemma 6.12 with $|J| \gtrsim N^{\alpha-O(\mu)}$.)

To prove the second part, we consider in detail the case $\gamma = 0$. We will apply the argument proving Lemma 6.12 to all $j \in J$. This will give us μ -additive sets K_j and sets X_j with appropriate upper and lower bounds since we can assume Δ contains no μ -additive sets with more than $N^{1+f(\mu)}$ elements.

We will allow f to vary from line to line and we will express even quantities that are clearly $O(\mu)$ as $f(\mu)$.

We let D be the set of all differences x for which $|\Delta[x] \cap B_j| \gtrsim N^{1-f(\mu)}$ for at least $N^{\alpha-f(\mu)}$ values of $j \in J$. For each value of $j \in J$, there are at least $N^{8-3\alpha-f(\mu)}$ pairs $(x, y) \in D^2$ with $|\Delta_G[x] \cap \Delta_G[y] \cap B_j| \gtrsim N^{1-f(\mu)}$. Note that we may also restrict D to differences which cannot be represented in more than $N^{1+f(\mu)}$ ways as differences of elements of B_j for more than $N^{\alpha-f(\mu)}$ values in j and so that in each B_j our count of good pairs (x, y) consists only of pairs of differences which cannot be represented as differences in B_j in more than $N^{1+f(\mu)}$ ways. Otherwise, we could choose $\gamma > f(\mu)$.

Now we recall the structure of the argument in Lemma 6.12. We chose an $a \in B_j$ and a set B_a of size $N^{3-\alpha-f(\mu)}$ of the differences in which a participates, and a set K_a which is actually of the form $\Delta_G[x] \cap B_j$ and has size $N^{1-f(\mu)}$. We find $N^{5-\alpha-f(\mu)}$ additive quadruples made up of two elements of B_a and two elements of K_a . We may strip down K_a further to those elements which participate in at least $N^{4-\alpha-f(\mu)}$ of these quadruples and not harm our estimate. Now, we note that since B_a is a large subset of a translate of B_j , it must be that there are $N^{2-f(\mu)}$ pairs $(q_1, q_2) \in K_a^2$ with the property that $q_1 - q_2$ is represented $N^{3-\alpha-f(\mu)}$ times as a difference of elements of B_j . We let $K_{1,j}$ be the set of differences of B_j that can be represented in $N^{3-\alpha-f(\mu)}$ ways as differences in B_j . Because B_j contains no μ -additively closed set of size more than $N^{1+f(\mu)}$, we have that $|K_{1,j}| \leq N^{1+f(\mu)}$. Otherwise we could apply the asymmetric Balog-Szemerédi-Gowers theorem to obtain a μ -additively closed set, contained in B_j , which is larger than Q .

We can replace $N^{5-\alpha-f(\mu)}$ quadruples $q_1 - q_2 = x_1 - x_2$ with $q_1, q_2 \in K_{1,j}$ and $x_1, x_2 \in A_a$ by an equation of the form $q = x_1 - x_2$ with multiplicity at most $N^{1+f(\mu)}$. Thus we obtain at least $N^{4-\alpha-f(\mu)}$ such equations. We rewrite the equation as $x_1 - q = x_2$ and use Cauchy-Schwarz to obtain $N^{5-\alpha-f(\mu)}$ quadruples $x_1 - q = x'_1 - q'$. We see then that without losing more than $N^{f(\mu)}$ factors in our estimates, we can replace K_a by $K_{1,j}$. This is good since we have made it independent of the choice of a . Now we need only show that we can find many

quadruples not only between $K_{1,j}$ and B_a but between $K_{1,j}$ and D . This will give us the desired result.

To do this, we observe that we may delete from B_a a set with relative density $N^{-f(\mu)}$ without harming our estimates on the number of quadruplets between B_a and $K_{1,j}$. Our goal will be to cover a subset D' by a disjoint union of subsets of the form B'_a , where B'_a is a subset of B_a with relative density $1 - N^{-f(\mu)}$. To do this we observe that for any fixed $a \in B_j$,

$$\sum_{a_2} |B_{a_1} \cap B_{a_2}| \lesssim N^{4-\alpha+f(\mu)}.$$

We can do this because the sum counts triples (x, a, a_2) with x a difference and a, a_2 are parts of representations of it. We have assumed that we are only dealing with differences with fewer than $N^{1+O(\mu)}$ representations. Here we are using that we are in the $\gamma = 0$ case,

Now we produce D' as follows. We choose a_1 and keep the set B_{a_1} . We add all elements of B_{a_1} to D' . We choose B_{a_2} to have the minimal possible sized intersection with B_{a_1} and let B'_{a_2} be those elements in B_{a_2} that are not already in D' . We choose B_{a_3} to have minimal possible intersection with $B_{a_1} \cup B_{a_2}$. We continue in this way until we reach a k so that B'_{a_k} no longer has relative density $1 - N^{-f(\mu)}$ in B_{a_k} . Because the average intersection $|B_a \cap B_{a'}|$ is bounded by $N^{1+f(\mu)}$, we get that k is at least $N^{2-\alpha-f(\mu)}$. Thus our set D' has relative density at least $N^{-f(\mu)}$ in D . Thus we have that $K_{1,j}$ has $N^{7-2\alpha-f(\mu)}$ quadruples with D and *a fortiori* with D' .

Now we slightly refine $K_{1,j}$ to $K_{2,j}$ consisting only of differences of elements of $K_{1,j}$ which participate in at least $N^{5-\alpha-f(\mu)}$ of the quadruples with D' . (We perform this refinement so that after we apply Balog-Szemerédi-Gowers to $K_{2,j}$, we will still be guaranteed to have many quadruples between the μ -additively closed set we obtain and D . Since D' is a disjoint union of sets with relative density $N^{-f(\mu)}$ inside translates of B_j , it must be that $K_{2,j}$ still has $N^{5-\alpha-f(\mu)}$ quadruples with B_j . Thus we can apply the asymmetric Balog-Szemerédi-Gowers theorem to find a subset K_j satisfying the conclusions of the theorem. \square

7. STRUCTURE OF SPECTRUM OF LARGE CAP SETS WITH NO STRONG INCREMENTS

In this section, we transfer the results obtained in Theorem 6.13 over to the setting of the spectrum of large cap sets with no strong increments. This turns out to be rather simple. The main ideas which we have not yet taken advantage of are Freiman's theorem and the use of the estimate in Proposition 3.3 which bounds the number of elements of the spectrum in a subspace of dimension d by $dN^{1+2\epsilon}$.

We state the main result of the section.

Theorem 7.1. *Let Δ be the spectrum of a large cap set without strong increments. There is a function $f : [0, 1] \rightarrow [0, \infty]$ with $\lim_{t \rightarrow 0} f(t) = 0$ so that the following holds. There is a subspace H of F_3^N of dimension $f(\epsilon)$ and a set $\Lambda \subset F_3^N$ of size $N^{2-f(\epsilon)}$ so that for each element $\lambda \in \Lambda$, there is a subset $H_\lambda \subset H$ with the properties that*

$$|H_\lambda| \gtrsim N^{1-f(\epsilon)},$$

and the sets $\lambda + H_\lambda$ are pairwise disjoint subsets of Δ . For any subspace $W \subset F_3^N$ of dimension d , we have that W contains at most $dN^{f(\epsilon)}$ elements of Λ .

Proof. As before we allow our function f to vary from line to line until we achieve the desired result.

In light of Theorem 2.6 and Proposition 3.3, any $f(\epsilon)$ -additively closed set in which the spectrum has $N^{-f(\epsilon)}$ relative density, must be bounded in size by $N^{1+f(\epsilon)}$. Therefore, we are in the $\gamma = 0$ case of Theorem 6.13. We know that there is a set Δ' of density $N^{-f(\epsilon)}$ in the spectrum Δ which is contained in

$$\bigcup_{j=1}^K X_j + K_j,$$

with each K_j an $f(\epsilon)$ -additively closed set (of size at least $N^{1-f(\epsilon)}$ and at most $N^{1+f(\epsilon)}$) and with each set X_j of size $N^{2-\alpha\pm f(\epsilon)}$. Moreover, each set K_j lies in the set K of differences having at least $N^{5-2\alpha-f(\epsilon)}$ representations as differences of elements of D , the differences among elements of the spectrum which have at least $N^{1+\alpha-f(\epsilon)}$ representations. In light of the nonadditive smoothing property of Δ , we have that $|K| \lesssim N^{1+f(\epsilon)}$ since there can be at most $N^{11-4\alpha+f(\epsilon)}$ quadruplets among elements of D . We may eliminate all elements q of each K_j for which $\Delta \cap q + X_j$ does not have size at least $N^{2-\alpha-f(\epsilon)}$. Now we let K' be the set of elements of K which appear in at least $N^{\alpha-f(\epsilon)}$ many K_j . We can find some K_j which has intersection of size $N^{1-f(\epsilon)}$ with K' . Then $K' \cap K_j = K''$ is an $f(\epsilon)$ -additively closed set with cardinality at least $N^{1-f(\epsilon)}$. Moreover each element of K'' is contained in $N^{\alpha-f(\epsilon)}$ many K_j . Thus by pigeonholing there are at least $N^{\alpha-f(\epsilon)}$ many K_j so that $|K_j \cap K''| \gtrsim N^{1-f(\epsilon)}$. We only keep these j and replace K_j by $K_j \cap K''$. But by Theorem 2.6, we have that K'' is contained in a subspace of dimension $N^{f(\epsilon)}$, which we call H .

This basically proves the first part of the theorem. We have that a subset of the spectrum of density $N^{-f(\epsilon)}$ is contained in $K'' + X$, where X is the union of the X_j 's. We will pick Λ and the sets H_λ as follows: Find x_1 in X so that at least $N^{1-f(\epsilon)}$ elements of Δ are contained in $x_1 + K''$. Let Δ_1 be the elements of Δ contained in $X + K''$ but not in $x_1 + K''$. Let Δ^1 be those elements of Δ contained in $x_1 + K''$ and let $H_{x_1} = \Delta^1 - x_1$. Note that H_{x_1} is contained in K'' and therefore in H . Now we proceed iteratively. Find $x_j \in X$ so that there are at least $N^{1-f(\epsilon)}$ elements of Δ_{j-1} in $x_j + K''$. When this is no longer possible, we terminate the process. Then we let Δ_j be the elements of Δ_{j-1} not in $x_j + K''$ and we let Δ^j be the ones that are. We let $H_{x_j} = \Delta^j - x_j$. We let $\Lambda = \{x_j\}$ after the iteration has terminated.

To prove the second part of the theorem, let S be any subset of Λ with some cardinality M . But $S + H$ contains at least $MN^{1-f(\epsilon)}$ elements of Δ . This contradicts Proposition 3.3 unless the span of S has dimension at least $MN^{-f(\epsilon)}$. \square

8. CONTRADICTION

The goal of this section is to obtain a contradiction from the existence of large cap sets without strong increments by using the result of Theorem 7.1. We begin by recording some easy consequences of Plancherel's identity for the interaction between the Fourier transform of the characteristic function of a set and the Fourier transforms of its fibers over a subspace.

For any set $A \subset F_3^N$, we define its Fourier transform

$$\hat{A}(x) = \frac{1}{3^N} \sum_{a \in A} e(a \cdot x).$$

We state Plancherel's identity:

Proposition 8.1.

$$\sum_{x \in F_3^N} |\hat{A}(x)|^2 = 3^{-N} |A|.$$

We let H be a subspace of F_3^N and we let H^\perp be its annihilator. We let V be a subspace of the same dimension as H which is transverse to H^\perp . We define the fiber $A_{H,v}$ for $v \in V$ by

$$A_{H,v} = A \cap (H^\perp + v).$$

If we have $h \in H$, then

$$\hat{A}(h) = 3^{-N} \sum_{v \in V} e(h \cdot v) |A_{H,v}|.$$

Thus we arrive at another form of Plancherel:

Proposition 8.2.

$$\sum_{h \in H} |\hat{A}(h)|^2 = |H| 3^{-2N} \sum_{v \in V} |A_{H,v}|^2.$$

Moreover

$$\sum_{h \neq 0 \in H} |\hat{A}(h)|^2 = |H| 3^{-2N} \sum_{v \in V} (|A_{H,v}| - \frac{|A|}{|H|})^2.$$

Next, we consider the situation where we have a subspace $H \subset F_3^N$ and a larger subspace K with $H \subset K \subset F_3^N$. We let V be a subspace transverse to H^\perp as before and we would like to consider the Fourier transforms of the fibers of A , namely the sets $A_{H,v}$. We can think of each fiber $A_{H,v}$ as being identified with a subset of H^\perp (by translation by v) and of course H^\perp can be identified with $F_3^N \setminus H$. That is, we define functions $A_{H,v} : F_3^N \setminus H \rightarrow \mathbf{C}$ by

$$\hat{A}_{H,v}(w) = \frac{|H|}{3^N} \sum_{a \in A_{H,v}} e((a - v) \cdot w).$$

The function $\hat{A}_{H,v}(w)$ is well defined on $F_3^N \setminus H$ since $a - v$ is in H^\perp . Next, we write down a version of Proposition 8.2 which shows how the L^2 norms of the Fourier transforms of the fibers on $K \setminus H$ compare with the L^2 norm of the Fourier transform on K . We let W be a subspace transverse to K^\perp with $V \subset W$.

Proposition 8.3. *With H, K, V , and W as above,*

$$\sum_{k \neq 0 \in K} |\hat{A}(k)|^2 = \sum_{h \neq 0 \in H} |\hat{A}(h)|^2 + \frac{1}{|H|} \sum_{v \in V} \sum_{k \neq 0 \in K/H} |\hat{A}_{H,v}(k)|^2.$$

Proof. Since clearly we have $K^\perp \subset H^\perp$, there is a unique subspace $W' \subset W$ with $W' + K^\perp = H^\perp$. We have $V + W' = W$.

We consider the following function on W :

$$g(w) = |A_{K,w}| - \frac{|A|}{|K|}.$$

Clearly, in light of the second part of Proposition 8.2, the left-hand side of the identity we are trying to prove is the normalized square of the L^2 norm of the function $g(w)$.

We now break up g as the sum of a function g_0 which is constant on translates of W' and functions g_v with v running over V having mean zero and supported on the translate $v + W'$ of W' . Clearly the functions g_0 and $\{g_v\}$ are pairwise orthogonal. The first term on the right-hand side of the identity is the normalized square L^2 norm of the function g_0 . The second term on the right-hand side represents the sum over v of the normalized square L^2 norm of the functions g_v . The identity is then an application of the Pythagorean theorem. \square

(We remark that Proposition 8.3 can simply be thought of as Plancherel for a “local Fourier transform” of A . Here, we localize to the translates of H^\perp .)

Now we are prepared to apply Proposition 8.3 to the setting in which A is a large cap set without strong increments and H is the subspace given to us by Theorem 7.1.

We let A be a large cap set with no strong increments. As usual, f will be a function taking $[0, 1]$ to $[0, \infty]$ with $\lim_{t \rightarrow 0} f(t) = 0$. We will vary f from line to line.

Then there is a subspace H with dimension $N^{f(\epsilon)}$ and a set Λ of size $N^{2-f(\epsilon)}$ so that for each $\lambda \in \Lambda$ there is a subset H_λ of H , so that $|H_\lambda| > N^{1-f(\epsilon)}$ so that for each $h \in H_\lambda$, we have that

$$|\hat{A}(h + \lambda)| \gtrsim N^{-1-f(\epsilon)} 3^{-N} |A|.$$

We also have that the sets $\lambda + H_\lambda$ are pairwise disjoint.

Note therefore that

$$\sum_{\lambda \in \Lambda} \sum_{h \in H_\lambda} |\hat{A}(h + \lambda)|^2 \gtrsim N^{-f(\epsilon)} N 3^{-2N} |A|^2 \gtrsim N^{-f(\epsilon)} 3^{-N} |A|.$$

Thus the structured elements of the spectrum of A account for a large proportion of the squared L^2 norm of the Fourier transform of A .

Now we would like to consider the fibers $A_{H,v}$, where H is the subspace we've been discussing. Because the cap set A has no strong increments, we know that for each value v , we have

$$|A_{H,v}| \leq \left(\frac{|A|}{|H|}\right)(1 + N^{f(\epsilon)-1}).$$

(We will now momentarily fix the function f .)

However, we don't have a good lower bound on $|A_{H,v}|$ in general. All we know is that the sum of all the positive increments is equal to the sum of all the negative increments. We let V_{bad} be the set of all $v \in V$ for which

$$|A_{H,v}| \leq \left(\frac{|A|}{|H|}\right)(1 - N^{2\sqrt{f(\epsilon)}-1}).$$

(That is, V_{bad} is the set of those $v \in V$ for which the fiber has a bad negative increment.) We know that

$$|V_{bad}| \lesssim N^{-\sqrt{f(\epsilon)}} |V|.$$

We define

$$A_{bad} = \bigcup_{v \in V_{bad}} A_{H,v},$$

and we let

$$A' = A \setminus A_{bad}.$$

We know that

$$|A_{bad}| \lesssim N^{-\sqrt{f(\epsilon)}}|A|.$$

Thus

$$\sum_x |\hat{A}_{bad}(x)|^2 \lesssim N^{-\frac{\sqrt{f(\epsilon)}}{2}} \sum_{\lambda \in \Lambda} \sum_{h \in H_\lambda} |\hat{A}(h + \lambda)|^2.$$

Thus removing A_{bad} does not perturb the large spectrum of A too much.

(In making this precise, we now resume changing the function f from line to line, observing that we may take the next f to be larger than the previous \sqrt{f} .) We may find a set Λ' which is a subset of Λ with $|\Lambda'| \gtrsim N^{2-f(\epsilon)}$ and so that for each $\lambda \in \Lambda'$ there is a subset H'_λ of H so that for each $\lambda \in \Lambda'$ and each $h \in H'_\lambda$ we have

$$|\hat{A}'(h + \lambda)| \gtrsim N^{-1-f(\epsilon)}3^{-N}|A|.$$

Thus from the point of view of the structure of the spectrum, we have that A' is essentially as good as A . However, the set A' has a big advantage over A in that we have good bounds on the Fourier transform of its fibers. This is because the fibers are either empty or close to size $\frac{|A|}{H}$. Empty fibers achieve no increments. On the other hand, fibers which are close to average cannot have an increment too large, or else the set A will have a strong increment on a translate of a codimension 1 subspace of H^\perp . Precisely, the estimate we get is

$$|\hat{A}'_{H,v}(x)| \lesssim 3^{-N}|A|N^{-1+f(\epsilon)} = \rho N^{-1+f(\epsilon)}.$$

(This is because the negative increment on density to pass from A to $A'_{H,v}$ does no more than to reduce the density by a factor of $1 - N^{f(\epsilon)-1}$ and the codimension of H is only $f(\epsilon)$. Should the Fourier transform of $A'_{H,v}$ exceed the above bound, then A will have a strong increment into a codimension one subspace of $H^\perp + v$.)

Moreover, the set $A'_{H,V}$ is a large cap set without strong increments (on subspaces of codimension no more than $\frac{N}{2} - N^{f(\epsilon)}$) but with ϵ replaced by $f(\epsilon)$. Thus we see using Proposition 3.3 that for any subspace L of $F_3^N \setminus H$ with dimension d , we have that

$$\sum_{x \in W} |\hat{A}'_{H,V}(x)|^2 \lesssim \left(\frac{|A|}{3^N}\right)^2 dN^{-1+f(\epsilon)}.$$

This estimate is a version of the bound on the number of elements of the spectrum of $A'_{H,V}$ in W .

From this information, together with the fact that no subspace K of dimension d contains more than $dN^{f(\epsilon)}$ elements of Λ and hence of Λ' , we are ready to achieve a contradiction.

We introduce some measure spaces on which we will apply Lemma 2.2. For each $v \in V$, we define χ_v , a measure on $F_3^N \setminus H - \{0\}$, by

$$\chi_v(X) = \sum_{x \in X} |\hat{A}'_{H,v}(x)|^2.$$

Clearly the total measure of χ_v is bounded by $3^{-N}|A|N^{f(\epsilon)}$. We now give a construction for sets which have large density for many of the measures χ_v .

Let $\Xi \subset \Lambda'$ with $|\Xi| = d \lesssim N^{1-f(\epsilon)}$. Then in light of the fact that $\Xi + H$ contains at least $dN^{1-f(\epsilon)}$ points x where $\hat{A}'(x)$ is large and in light of Proposition 8.3, there

must be $O(N^{-f(\epsilon)}|H|)$ values of v for which $\text{span}(\Xi)$ has density at least $dN^{-2-f(\epsilon)}$ for χ_v . (Here, by slight abuse of notation, the span is taken in $F_3^N \setminus H$.)

We will take $d \sim N^{\frac{2}{3}}$. Now let Π be any collection of $N^{\frac{8}{3}+O(\sqrt{\mu})}$ subsets of Λ' with cardinality $N^{\frac{2}{3}}$. We chose the exponents $\frac{2}{3}$ and $\frac{8}{3}$ somewhat arbitrarily. By pigeonholing, we may find a set of v 's of cardinality $|H|N^{-f(\epsilon)}$ for which at least $N^{\frac{8}{3}-f(\epsilon)}$ of the sets Ξ in Π have the property that $\text{span}(\Xi)$ has density $N^{-\frac{4}{3}-f(\epsilon)}$ for χ_v . From this and Lemma 2.2 we obtain the lower bound

$$(8.1) \quad \sum_{v \in V - V_{bad}} \sum_{\Xi_1 \in \Pi} \sum_{\Xi_2 \in \Pi} \frac{\chi_v(\text{span}(\Xi_1) \cap \text{span}(\Xi_2))}{\chi_V(F_3^N \setminus H)} \gtrsim |H|N^{\frac{8}{3}-f(\epsilon)}.$$

What this says is that modulo $N^{f(\epsilon)}$ terms, the average density (in χ_v measure) of an intersection $\text{span}(\Xi_1) \cap \text{span}(\Xi_2)$ is at least approximately $N^{-\frac{8}{3}}$. This is rather large since the density of any single element of $F_3^N \setminus H$ is only approximately N^{-3} in light of the bound on the Fourier transforms of the fibers.

We will now produce a random construction of Π . We choose each set of Π uniformly at random. We will write down an upper bound on the expected density in any χ_v of the intersection $\text{span}(\Xi_1) \cap \text{span}(\Xi_2)$, which contradicts the lower bound (8.1). We observe that if ρ is the density in χ_v of $\text{span}(\Xi_1) \cap \text{span}(\Xi_2)$, then

$$\rho \leq N^{-3+O(f(\epsilon))}(3^k - 1),$$

where k is the nullity of the set $\Xi_1 \cup \Xi_2$. We recall that there are no more than $N^{\frac{2}{3}+f(\epsilon)}$ elements of Λ' in any subspace of dimension $2N^{\frac{2}{3}}$. Thus as we choose the $2N^{\frac{2}{3}}$ elements of Ξ_1 and Ξ_2 , the probability of introducing nullity at each selection is bounded by $N^{-\frac{2}{3}+f(\epsilon)}$. Thus the probability p_k of having nullity k is bounded by

$$p_k \lesssim (N^{-\frac{2}{3}+f(\epsilon)})^k.$$

Thus the expected density of $\text{span}(\Xi_1) \cap \text{span}(\Xi_2)$ is $O(N^{-\frac{11}{3}+f(\epsilon)})$. This is a contradiction since we have just proved that Δ cannot contain a set as thick in $\Lambda + H$ as required by Theorem 7.1.

Thus we have proved Theorem 1.4 and hence Theorem 1.1.

9. EPILOGUE ON EFFICIENCY

The ϵ which we obtain in Theorem 1.1 is of necessity quite small. Like many arguments in analysis, ours does not aim for great efficiency, and we seem to lose a factor in the size of ϵ in nearly every line of the argument. However, there are a few points in the argument where we lose significantly more. The most notable of these are the use of the asymmetric Balog-Szemerédi-Gowers lemma and the discovery of the additive structure with comity in Corollary 6.7. What these two parts of the argument have in common is that they are iterative. One starts with a structure and seeks a certain property. If the property is lacking, one finds a structure which is in a certain sense better but which has cost us a factor in ϵ .

Iteration can be a powerful thing. It sometimes allows us to prove a very deep result with very few words. The proof doesn't care how many iterates must be calculated to implement it. We have a version of our argument which we will publish elsewhere that adds even another layer of iteration to the argument but spares us some of the difficulties of Theorem 6.13. We have semantic disagreements

about whether the more iterative argument is easier to understand. But certainly, it is less efficient than the argument presented here.

To improve efficiency, it is important to make the argument less iterative. In hopes of doing so, we leave the reader with two conjectures:

Conjecture 9.1. *Let $N > 0$ be a large parameter. (Constants may not depend on N .) Suppose B and C are sets in F_3^N each of whose cardinality is bounded by N^{100} . Suppose the number of quadruplets $b_1 + c_1 = b_2 + c_2$ with $b_1, b_2 \in B$ and $c_1, c_2 \in C$ is at least $|B||C|^2N^{-\sigma}$. Then C is contained in a subspace of dimension $N^{O(\sigma)}$.*

Conjecture 9.2. *Suppose Δ is a subset of F_3^N that is N^σ additively nonsmoothing. (We may assume as in the previous conjecture that $|\Delta| \leq N^{100}$.) Then Δ admits an additive structure at some height with $N^{O(\sigma)}$ comity.*

ACKNOWLEDGEMENTS

The authors thank Izabella Łaba and Olof Sisask for pointing out an error in the statement of the asymmetric Balog-Szemerédi-Gowers theorem in a preliminary version of this paper.

REFERENCES

- [CS11] E. Croot and O. Sisask, *A Probabilistic Technique for Finding Almost-Periods of Convolutions*, *Geom. Funct. Anal.* **20** (2010), 1367–1396. MR2738997
- [G] T. Gowers, *What is difficult about the cap set problem?*, <http://gowers.wordpress.com/2011/01/11/what-is-difficult-about-the-cap-set-problem/>.
- [K1] (M. Bateman and) N.H. Katz, <http://gowers.wordpress.com/2011/01/11/what-is-difficult-about-the-cap-set-problem/#comment-10533>.
- [K2] (M. Bateman and) N.H. Katz, <http://gowers.wordpress.com/2011/01/11/what-is-difficult-about-the-cap-set-problem/#comment-10540>.
- [KK10] N. H. Katz and P. Koester, *On Additive Doubling and Energy*, *SIAM J. Discrete Math.* **24** (2010), 1684–1693. MR2746716
- [M95] R. Meshulam, *On subsets of finite abelian groups with no 3-term arithmetic progressions*, *J. Combin. Theory Ser. A* **71** (1995), 168–172. MR1335785 (96g:20033)
- [PM] Polymath on wikipedia, http://en.wikipedia.org/wiki/Polymath_project#Polymath_Project.
- [PM6] Polymath 6: Improving the bounds for Roth’s theorem, <http://polymathprojects.org/2011/02/05/polymath6-improving-the-bounds-for-roths-theorem/>.
- [R99] I. Ruzsa, *An analog of Freiman’s theorem in groups*, *Structure theory of set addition*, *Astérisque* No. **258** (1999), 323–329. MR1701207 (2000h:11111)
- [S08] T. Sanders, *A note on Freiman’s theorem in vector spaces*, *Combin. Probab. Comput.* **17** (2008), 297–305. MR2396355 (2009a:11024)
- [S10] T. Sanders, *Structure in Sets with Logarithmic Doubling*, Arxiv 1002.1552.
- [S11] T. Sanders, *On Roth’s Theorem on Progressions*, Arxiv 1011.0104.
- [Sch11] T. Schoen, *Near optimal bounds in Freiman’s theorem*, *Duke Math. J.* **158**, Number 1 (2011), 1–12. MR2794366
- [Shk08] I. Shkredov, *On sets of large trigonometric sums*, 2008 *Izv. Ross. Akad. Nauk Ser. Math.* **72**, pp. 161–182. MR2394976 (2009e:11151)
- [TV06] T. Tao and V. Vu, *Additive Combinatorics*, Cambridge Univ. Press, Cambridge, 2006. MR2289012 (2008a:11002)

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CALIFORNIA 90095
E-mail address: bateman@math.ucla.edu

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405-7000
E-mail address: nhkatz@indiana.edu