

## COMMUTATORS ON $L_p$ , $1 \leq p < \infty$

DETELIN DOSEV, WILLIAM B. JOHNSON, AND GIDEON SCHECHTMAN

*Dedicated to the memory of Nigel Kalton*

### 1. INTRODUCTION

When studying derivations on a general Banach algebra  $\mathcal{A}$ , a natural problem that arises is to classify the commutators in the algebra; i.e., elements of the form  $AB - BA$ . The problem as stated is hard to tackle on general Banach algebras. The only known obstruction was proved in 1947 by Wintner ([14]). He proved that the identity in a unital Banach algebra is not a commutator, which immediately implies that no operator of the form  $\lambda I + K$ , where  $K$  belongs to a norm closed (proper) ideal  $\mathcal{I}$  of  $\mathcal{A}$  and  $\lambda \neq 0$ , is a commutator in the Banach algebra  $\mathcal{A}$ . On the other hand, there seem to be no general conditions for checking whether an element of a Banach algebra is a commutator.

The situation changes if instead of an arbitrary Banach algebra we consider the algebra  $\mathcal{L}(\mathcal{X})$  of all bounded linear operators on the Banach space  $\mathcal{X}$ . In this setting, one hopes that the underlying structure of the space  $\mathcal{X}$  will provide enough information about the operators on  $\mathcal{X}$  to allow one to attack the problem successfully. Indeed, this is the case provided the space  $\mathcal{X}$  has some “nice” properties. The first complete classification of the commutators in  $\mathcal{L}(\mathcal{X})$  was given in 1965 by Brown and Percy ([3]) for the case  $\mathcal{X} = \ell_2$ . They proved that the only operators in  $\mathcal{L}(\ell_2)$  that are not commutators have the form  $\lambda I + K$ , where  $K$  is compact and  $\lambda \neq 0$ . In 1972, Apostol proved in [1] that the same classification holds for the commutators on  $\ell_p$ ,  $1 < p < \infty$ , and one year later, he proved that the same classification holds in the case of  $\mathcal{X} = c_0$  ([2]). Apostol had some partial results in [1] and [2] about special classes of operators on  $\ell_1$ ,  $\ell_\infty$ , and  $C([0, 1])$ , but he was unable to obtain a complete classification of the commutators on any of those spaces. A year before Apostol’s results, Schneeberger proved that the compact operators on  $L_p$ ,  $1 < p < \infty$ , are commutators but, as will become apparent later, one needs a stronger result in order to classify the commutators on these spaces.

All of the aforementioned spaces have one common property; namely, if  $\mathcal{X} = \ell_p$ ,  $1 \leq p < \infty$ , or  $\mathcal{X} = c_0$ , then  $\mathcal{X} \simeq (\sum \mathcal{X})_p$  ( $p = 0$  if  $\mathcal{X} = c_0$ ). It turns out that this property plays an important role for proving the classification of the commutators on other spaces. Thirty-five years after Apostol’s result, the first author obtained

---

Received by the editors February 2, 2011 and, in revised form, May 30, 2012.

2010 *Mathematics Subject Classification*. Primary 47B47; Secondary 46E30.

The first author was Young Investigator, NSF Workshop in Analysis and Probability, Texas A&M University.

The second author was supported in part by NSF DMS-1001321 and U.S.-Israel Binational Science Foundation.

The third author was supported in part by U.S.-Israel Binational Science Foundation. Participant NSF Workshop in Analysis and Probability, Texas A&M University.

in [4] a complete classification of the commutators on  $\ell_1$ , which, as one may expect, is the same as the classification of the commutators on  $\ell_2$ . A common feature of all the spaces  $\mathcal{X} = \ell_p$ ,  $1 \leq p < \infty$  and  $\mathcal{X} = c_0$  is that the ideal of compact operators  $\mathcal{K}(\mathcal{X})$  on  $\mathcal{X}$  is the largest nontrivial ideal in  $\mathcal{L}(\mathcal{X})$ . The situation for  $\mathcal{X} = \ell_\infty$  is different. Recall that an operator  $T : \mathcal{X} \rightarrow \mathcal{Y}$  is strictly singular provided the restriction of  $T$  to any infinite-dimensional subspace of  $\mathcal{X}$  is not an isomorphism. On  $\ell_p$ ,  $1 \leq p < \infty$ , and on  $c_0$ , every strictly singular operator is compact, but on  $\mathcal{L}(\ell_\infty)$ , the ideal of strictly singular operators contains noncompact operators (and, incidentally, agrees with the ideal of weakly compact operators). In  $\mathcal{L}(\ell_\infty)$ , the ideal of strictly singular operators is the largest ideal, and it was proved in [5] that all operators on  $\ell_\infty$  that are not commutators have the form  $\lambda I + S$ , where  $\lambda \neq 0$  and  $S$  is strictly singular.

The classification of the commutators on  $\ell_p$ ,  $1 \leq p \leq \infty$ , and on  $c_0$ , as well as partial results on other spaces, suggest the following:

**Conjecture 1.** *Let  $\mathcal{X}$  be a Banach space such that  $\mathcal{X} \simeq (\sum \mathcal{X})_p$ ,  $1 \leq p \leq \infty$  or  $p = 0$  (we say that such a space admits a Pełczyński decomposition). Assume that  $\mathcal{L}(\mathcal{X})$  has a largest ideal  $\mathcal{M}$ . Then every noncommutator on  $\mathcal{X}$  has the form  $\lambda I + K$ , where  $K \in \mathcal{M}$  and  $\lambda \neq 0$ .*

In [13], M. Tarbard constructs a Banach space  $X$  on which there is a strictly singular noncompact operator  $S$  so that every bounded linear operator on  $X$  is of the form  $\lambda I + \alpha S + K$  for some compact operator  $K$  and some scalars  $\lambda$  and  $\alpha$ . On this space the strictly singular operators are the unique maximal ideal in the bounded operators on  $X$ , yet  $S$  is clearly not a commutator. Consequently, some assumption on an infinite-dimensional space is needed to show that the space satisfies the conclusion of Conjecture 1.

Here and elsewhere in this paper, when we refer to an ideal of operators we always mean a nontrivial, norm closed, two-sided ideal. This conjecture is stated in [5]. To verify Conjecture 1 for a given Banach space  $\mathcal{X}$ , one must prove two steps:

*Step 1.* Every operator  $T \in \mathcal{M}$  is a commutator.

*Step 2.* If  $T \in \mathcal{L}(\mathcal{X})$  is not of the form  $\lambda I + K$ , where  $K \in \mathcal{M}$  and  $\lambda \neq 0$ , then  $T$  is a commutator.

The methods for proving **Step 1** in most cases where the complete classification of the commutators on the space  $\mathcal{X}$  is known are based on the fact that if  $T \in \mathcal{M}$ , then for every subspace  $Y \subseteq \mathcal{X}$ ,  $Y \simeq \mathcal{X}$  and every  $\varepsilon > 0$  there exists a complemented subspace  $Y_1 \subseteq Y$ ,  $Y_1 \simeq Y$  such that  $\|T|_{Y_1}\| < \varepsilon$ . Let us just mention that this fact is fairly easy to see if  $T$  is a compact operator on  $c_0$  or  $\ell_p$ ,  $1 \leq p < \infty$  ([4, Lemma 9]; see also [1]). (Throughout this work,  $Y \simeq X$  means that  $X$  and  $Y$  are isomorphic, i.e., linearly homeomorphic, while  $Y \equiv X$  means that the spaces are isometrically isomorphic.)

Showing the second step is usually more difficult than showing **Step 1**. In most cases for which we have a complete characterization of the commutators on  $\mathcal{X}$ , we use the following theorem, which is an immediate consequence of Theorem 3.2 and Theorem 3.3 in [5].

**Theorem 1.1.** *Let  $\mathcal{X}$  be a Banach space such that  $\mathcal{X} \simeq (\sum \mathcal{X})_p$ ,  $1 \leq p \leq \infty$  or  $p = 0$ . Let  $T \in \mathcal{L}(\mathcal{X})$  be such that there exists a subspace  $X \subset \mathcal{X}$  such that  $X \simeq \mathcal{X}$ ,*

$T|_X$  is an isomorphism,  $X + T(X)$  is complemented in  $\mathcal{X}$ , and  $d(X, T(X)) > 0$ . Then  $T$  is a commutator.

In the previous theorem the distance is defined as the distance from  $Y$  to the unit sphere of  $X$ . The basic idea is to prove that if  $T \in \mathcal{L}(\mathcal{X})$  is not of the form  $\lambda I + K$ , where  $K \in \mathcal{M}$  and  $\lambda \neq 0$ , then  $T$  satisfies the assumptions of Theorem 1.1 and hence  $T$  is a commutator. This is not obvious even for the classical sequence spaces  $c_0$  and  $\ell_p$ ,  $1 \leq p < \infty$ , but it suggests what one may try to prove for other classical Banach spaces in order to obtain a complete characterization of the commutators on those spaces.

Following the ideas in [5], for a given Banach space  $\mathcal{X}$  we define the set

$$(1) \quad \mathcal{M}_{\mathcal{X}} = \{T \in \mathcal{L}(\mathcal{X}) : I_{\mathcal{X}} \text{ does not factor through } T\}.$$

(We say that  $S \in \mathcal{L}(\mathcal{X})$  factors through  $T \in \mathcal{L}(\mathcal{X})$  if there are  $A, B \in \mathcal{L}(\mathcal{X})$  such that  $S = ATB$ .) As noted in [5], this set comes naturally from the investigation of the structure of the commutators on several classical Banach spaces. In the cases of  $\mathcal{X} = \ell_p$ ,  $1 \leq p \leq \infty$ , and  $\mathcal{X} = c_0$ , the set  $\mathcal{M}_{\mathcal{X}}$  is the largest ideal in  $\mathcal{L}(\mathcal{X})$  (observe that if  $\mathcal{M}_{\mathcal{X}}$  is an ideal, then it is the largest ideal in  $\mathcal{L}(\mathcal{X})$  and  $\mathcal{M}_{\mathcal{X}}$  is an ideal if and only if it is closed under addition). It is also known that  $\mathcal{M}_{\mathcal{X}}$  is the largest ideal for  $\mathcal{X} = L_p$ ,  $1 \leq p < \infty$ , which we discuss later.

In some special cases of finite sums of Banach spaces we know that the classification of the commutators on the sum depends only on the classification of the commutators on each summand. In particular, this is the case with the space  $\ell_{p_1} \oplus \ell_{p_2} \oplus \cdots \oplus \ell_{p_n}$ , where the first two authors proved in [5] that all noncommutators on  $\ell_{p_1} \oplus \ell_{p_2} \oplus \cdots \oplus \ell_{p_n}$  have the form  $\lambda I + K$  where  $\lambda \neq 0$  and  $K$  belongs to some ideal in  $\mathcal{L}(\ell_{p_1} \oplus \ell_{p_2} \oplus \cdots \oplus \ell_{p_n})$ .

In this paper we always denote  $L_p = L_p([0, 1], \mu)$ , where  $\mu$  is the Lebesgue measure. Our main structural results are:

**Theorem 1.2.** *Let  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < 2$ . If  $T - \lambda I \notin \mathcal{M}_{L_p}$  for all  $\lambda \in \mathbb{C}$ , then there exists a subspace  $X \subset L_p$  such that  $X \simeq L_p$ ,  $T|_X$  is an isomorphism,  $X + T(X)$  is complemented in  $L_p$ , and  $d(X, T(X)) > 0$ .*

**Theorem 1.3.** *Let  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < 2$ . If  $T \in \mathcal{M}_{L_p}$ , then for every  $Y \subseteq L_p$ ,  $Y \simeq L_p$ , there exists a subspace  $X \subset Y$  such that  $X$  is complemented in  $L_p$ ,  $X \simeq L_p$ , and  $T|_X$  is a compact operator.*

Notice that Theorem 1.3 implies that for  $1 \leq p < 2$ ,  $\mathcal{M}_{L_p}$  is closed under addition and hence is the largest ideal in  $\mathcal{L}(L_p)$ . It follows by duality that for  $2 < p < \infty$ ,  $\mathcal{M}_{L_p}$  is closed under addition as well and hence is the largest ideal in  $\mathcal{L}(L_p)$ . This duality argument is indeed needed because Theorem 1.3 is false for  $p > 2$ . To see that Theorem 1.3 is false for  $p > 2$  one can consider  $T = JI_{p,2}$ , where  $I_{p,2}$  is the identity from  $L_p$  into  $L_2$  and  $J$  is an isometric embedding from  $L_2$  into  $L_p$ .

In order to prove Theorem 1.2 for  $1 < p < 2$ , it was necessary to improve [8, Proposition 9.11] for the spaces  $L_p$ ,  $1 < p < 2$ , and the improvement is of independent interest. In Theorem 3.4 we show that for a natural equivalent norm on  $L_p$ ,  $1 < p < 2$ , if  $T$  is an operator on  $L_p$  which is an isomorphism on a copy of  $L_p$ , then some multiple of  $T$  is almost an isometry on an isometric copy of  $L_p$ . The proof of Theorem 3.4, which can be read independently from the rest of this paper, is the most difficult argument in this paper, and we will postpone it till the Appendix.

Using Theorem 1.2 and Theorem 1.3 it is easy to show that Conjecture 1 also holds for  $L_p$ ,  $1 \leq p < 2$ . It follows by duality that Conjecture 1 also holds for  $L_p$ ,  $2 < p < \infty$ .

**Theorem 1.4.** *Let  $\mathcal{M}$  be the largest ideal in  $\mathcal{L}(L_p)$ ,  $1 \leq p < \infty$ . An operator  $T \in \mathcal{L}(L_p)$  is a commutator if and only if  $T - \lambda I \notin \mathcal{M}$  for any  $\lambda \neq 0$ .*

*Proof.* As we already mentioned, we only need to consider the case  $1 \leq p < 2$ , and the case  $2 < p < \infty$  will follow by a duality argument.

If  $T$  is a commutator, from the remarks we made in the introduction it follows that  $T - \lambda I$  cannot be in  $\mathcal{M}$  for any  $\lambda \neq 0$ . For proving the other direction we have to consider two cases:

*Case I.* If  $T \in \mathcal{M}$  ( $\lambda = 0$ ), we first apply Theorem 1.3 to obtain a complemented subspace  $X \subset L_p$  such that  $T|_X$  is a compact operator and then apply [4, Corollary 12], which gives us the desired result.

*Case II.* If  $T - \lambda I \notin \mathcal{M}$  for any  $\lambda \in \mathbb{C}$ , we are in position to apply Theorem 1.2, which combined with Theorem 1.1 implies that  $T$  is a commutator.  $\square$

The rest of this paper is devoted to the proofs of Theorems 1.2 and 1.3. We consider the case  $L_1$  separately since some of the ideas and methods used in this case are quite different from those used for the case  $L_p$ ,  $1 < p < \infty$ .

## 2. NOTATION AND BASIC RESULTS

Throughout this manuscript, if  $\mathcal{X}$  is a Banach space and  $X \subseteq \mathcal{X}$  is complemented, by  $P_X$  we denote a projection from  $\mathcal{X}$  onto  $X$ . For any two subspaces (possibly not closed)  $X$  and  $Y$  of a Banach space  $\mathcal{Z}$  let

$$d(X, Y) = \inf\{\|x - y\| : x \in X, y \in Y\}.$$

A well-known consequence of the open mapping theorem is that for any two closed subspaces  $X$  and  $Y$  of  $\mathcal{Z}$ , if  $X \cap Y = \{0\}$ , then  $X + Y$  is a closed subspace of  $\mathcal{Z}$  if and only if  $d(X, Y) > 0$ . Note also that  $2d(X, Y) \geq d(Y, X) \geq 1/2d(X, Y)$ ; thus  $d(X, Y)$  and  $d(Y, X)$  are equivalent up to a constant factor of 2. The following proposition was proved in [5] and will allow us later to consider only isomorphisms instead of arbitrary operators on  $L_p$ .

**Proposition 2.1** ([5, Proposition 2.1]). *Let  $\mathcal{X}$  be a Banach space and  $T \in \mathcal{L}(\mathcal{X})$  be such that there exists a subspace  $Y \subset \mathcal{X}$  for which  $T$  is an isomorphism on  $Y$  and  $d(Y, TY) > 0$ . Then for every  $\lambda \in \mathbb{C}$ ,  $(T - \lambda I)|_Y$  is an isomorphism and  $d(Y, (T - \lambda I)Y) > 0$ .*

We will also need a result similar to Proposition 2.1, where instead of adding a multiple of the identity we want to add an arbitrary operator. Obviously that cannot be done in general, but if we assume that the operator we add has a sufficiently small norm we can derive the desired conclusion.

**Proposition 2.2.** *Let  $T \in \mathcal{L}(\mathcal{X})$  and let  $Y \subset \mathcal{X}$  be such that  $T$  is an isomorphism on  $Y$ ,  $Y \simeq \mathcal{X}$ ,  $d(Y, TY) > 0$ , and  $Y + TY$  is a complemented subspace of  $\mathcal{X}$  isomorphic to  $\mathcal{X}$ . Then there exists an  $\varepsilon > 0$ , depending only on  $d(Y, TY)$ , the norm of the projection onto  $Y + TY$ , and  $\|T|_Y^{-1}\|$  such that if  $K \in \mathcal{L}(\mathcal{X})$  satisfies  $\|K|_Y\| < \varepsilon$ , then  $d(Y, (T + K)Y) > 0$  and  $Y + (T + K)Y$  is a complemented subspace of  $\mathcal{X}$  isomorphic to  $\mathcal{X}$ .*

*Proof.* First we show that if  $\varepsilon$  is sufficiently small, then  $d(Y, (T + K)Y) > 0$ , provided  $\|K\| < \varepsilon$ . As in [5, Proposition 2.1], we have to show that there exists a constant  $c > 0$  such that for all  $y \in S_Y$ ,  $d((T + K)y, Y) > c$ . From  $d(TY, Y) > 0$  it follows that there exists a constant  $C$ , such that for all  $y \in S_Y$ ,  $d(Ty, Y) > C$ . If  $\|K|_Y\| < \frac{C}{2}$ , then

$$\|(T + K)y - z\| \geq \|Ty - z\| - \|Ky\| \geq d(Ty, Y) - \frac{C}{2} \geq \frac{C}{2}$$

for all  $z \in Y$ ; hence  $d((T + K)Y, Y) > 0$ .

Let  $P$  be the projection onto  $Y + TY$ . To show that  $Y + (T + K)Y$  is complemented in  $\mathcal{X}$  we first define an isomorphism  $S : Y + TY \rightarrow Y + (T + K)Y$  by  $S(y + Tz) = y + (T + K)z$  for every  $y, z \in Y$ . From the definition of  $S$  we have that  $\|S - I\| \leq C(Y, T)\|K|_Y\|$  (where  $C(Y, T) = \frac{\|P\|\|T|_Y^{-1}\|}{d(TY, Y)}$ ); hence if  $\|K|_Y\|$  is small enough the operator  $R = SP + I - P$  is an isomorphism on  $\mathcal{X}$ . Now it is not hard to see that  $RPR^{-1}$  is a projection onto  $Y + (T + K)Y$ .  $\square$

### 3. OPERATORS ON $L_p$ , $1 < p < \infty$

Recall (see (1) in the Introduction) that if  $\mathcal{X}$  is a Banach space,  $\mathcal{M}_{\mathcal{X}} = \{T \in \mathcal{L}(\mathcal{X}) : I_{\mathcal{X}} \text{ does not factor through } T\}$ , then  $T \notin \mathcal{M}_{\mathcal{X}}$  if and only if there exists a subspace  $X$  of  $\mathcal{X}$  so that  $T|_X$  is an isomorphism,  $TX$  is complemented in  $\mathcal{X}$ , and  $TX \simeq \mathcal{X}$ .

As we have already mentioned in the Introduction, the set  $\mathcal{M}_{\mathcal{X}}$  is the largest ideal in  $\mathcal{L}(\mathcal{X})$  if and only if it is closed under addition. Using the fact that if  $p = 1$ , then  $\mathcal{M}_{L_1}$  coincides with the ideal of non- $E$  operators, defined in [6], and if  $1 < p < \infty$ , then  $\mathcal{M}_{L_p}$  coincides with the ideal of non- $A$  operators, defined in [8], it is clear that  $\mathcal{M}_{\mathcal{X}}$  is in fact the largest ideal in those spaces. This fact, as we already mentioned, follows from Theorem 1.3 as well. For a more detailed discussion of the  $E$  and  $A$  operators we refer the reader to [6] and [8, Section 9]. Let us also mention that we are not going to use any of the properties of the  $E$  or  $A$  operators and so do not repeat their definitions here.

In this section we mainly consider operators  $T : L_p \rightarrow L_p$ ,  $1 \leq p < \infty$ , that preserve a complemented copy of  $L_p$ ; that is, there exists a complemented subspace  $X \subseteq L_p$ ,  $X \simeq L_p$  such that  $T|_X$  is an isomorphism. The fact that we can automatically take a complemented subspace isomorphic to  $L_p$  instead of just a subspace isomorphic to  $L_p$  follows from [8, Theorem 9.1] in the case  $p > 1$  and [12, Theorem 1.1] in the case  $p = 1$ . From the definition of  $\mathcal{M}_{\mathcal{X}}$  it is easy to see that  $T \notin \mathcal{M}_{L_p}$  if and only if  $T$  maps a copy of  $L_p$  isomorphically onto a complemented copy of  $L_p$ .

Also, recall that an operator  $T : \mathcal{X} \rightarrow \mathcal{Y}$  is called  $Z$ -strictly singular provided the restriction of  $T$  to any subspace of  $\mathcal{X}$ , isomorphic to  $Z$ , is not an isomorphism. From the remarks above, it is clear that the class of operators from  $L_p$  to  $L_p$  that do not preserve a complemented copy of  $L_p$  coincides with the class of  $L_p$ -strictly singular operators; hence the class of  $L_p$ -strictly singular operators is the largest ideal in  $\mathcal{L}(L_p)$ .

**Definition 3.1.** The sequence of functions  $\{h_{0,0}\} \cup \{h_{n,i}\}_{n=0, i=1}^{2^n}$  defined by  $h_{0,0}(t) \equiv 1$  and, for  $n = 0, 1, \dots$  and  $i = 1, 2, \dots, 2^n$ ,

$$h_{n,i}(t) = \begin{cases} 1 & \text{if } t \in ((2i-2)2^{-(n+1)}, (2i-1)2^{-(n+1)}) \\ -1 & \text{if } t \in ((2i-1)2^{-(n+1)}, 2i2^{-(n+1)}) \\ 0 & \text{otherwise} \end{cases}$$

is called the Haar system on  $[0, 1]$ .

The Haar system, in its natural order, is an unconditional monotone basis of  $L_p[0, 1]$  for every  $1 < p < \infty$  (cf. [10, p.3, p.19]), and we denote by  $C_p$  the unconditional basis constant of the Haar system. As usual, by  $\{r_n\}_{n=0}^\infty$  we denote the Rademacher sequence on  $[0, 1]$ , defined by  $r_n = \sum_{i=1}^{2^n} h_{n,i}$ .

**Definition 3.2.** Let  $\{x_i\}_{i=1}^\infty$  be an unconditional basis for  $L_p$ . For  $x = \sum_{i=1}^\infty a_i x_i$ , the square function of  $x$  with respect to  $\{x_i\}_{i=1}^\infty$  is defined by

$$S(x) = \left( \sum_{i=1}^\infty a_i^2 x_i^2 \right)^{\frac{1}{2}}.$$

The following proposition is well known. We include its proof here for completeness.

**Proposition 3.3.** *Let  $\{a_i\}_{i=1}^\infty$  be a block basis of the Haar basis for  $L_p$ ,  $1 < p < \infty$ , such that  $A = \overline{\text{span}}\{a_i : i = 1, 2, \dots\}$  is a complemented subspace of  $L_p$  via a projection  $P$ . Then there exists a projection onto  $A$  that respects supports with respect to the Haar basis and whose norm depends on  $p$  and  $\|P\|$  only.*

*Proof.* Define  $\sigma_i = \{(k, l) : h_{k,l} \in \text{supp}(a_i)\}$ , where the support is taken with respect to the Haar basis, and denote  $X_i = \overline{\text{span}}\{h_{k,l} : (k, l) \in \sigma_i\}$ . It is clear that all spaces  $X_i$  are  $C_p$  complemented in  $L_p$ , via the natural projections  $P_i$ , as a span of subsequences of the Haar basis. Consider the operator  $P_A = \sum_i P_i P P_i$ . Provided it is bounded, it is easy to check that  $P_A$  is a projection onto  $A$  that respects supports. In order to show that  $P_A$  is bounded consider the formal sum

$$P = \sum_{i,j} P_i P P_j.$$

A simple computation shows that

$$(2) \quad \|P_A\| = \|\mathbb{E} \sum_{i,j} \varepsilon_i \varepsilon_j P_i P P_j\| \leq \mathbb{E} \left\| \sum_{i,j} \varepsilon_i \varepsilon_j P_i P P_j \right\| \leq C_p^2 \|P\|,$$

where  $\varepsilon_i$  is a Rademacher sequence on  $[0, 1]$ , which finishes the proof.  $\square$

The following theorem is the main result of this section. We will postpone its proof till the end since the ideas for proving it deviate from the general ideas of this section and the proof as well as the result are of independent interest. Recall [12] that an operator  $T$  on  $L_p$ ,  $1 \leq p < \infty$ , is called a sign-embedding provided there is a set  $S$  of positive measure and  $\delta > 0$  so that  $\|Tf\| \geq \delta$  whenever  $\int f d\mu = 0$  and  $|f| = 1_S$  almost everywhere.

Recall also, that a bounded linear operator  $T : \mathcal{X} \rightarrow \mathcal{Y}$  between two Banach spaces  $\mathcal{X}$  and  $\mathcal{Y}$  is a  $C$ -isomorphism if  $T$  is an isomorphism on  $\mathcal{X}$  and  $\|T\| \|T|_{T\mathcal{X}}^{-1}\| \leq C$ .

**Theorem 3.4.** *For each  $1 < p < 2$  there is a constant  $K_p$  such that if  $T$  is a sign-embedding operator from  $L_p[0, 1]$  into  $L_p[0, 1]$  (and in particular if it is an isomorphism), then there is a  $K_p$  complemented subspace  $X$  of  $L_p[0, 1]$  which is  $K_p$ -isomorphic to  $L_p[0, 1]$  and such that  $T|_X$  is a  $K_p$ -isomorphism and  $T(X)$  is  $K_p$  complemented in  $L_p$ .*

Moreover, if we consider  $L_p[0, 1]$  with the norm  $\|x\|_p = \|S(x)\|_p$  (with  $S$  being the square function with respect to the Haar system), then, for each  $\varepsilon > 0$ , there is a subspace  $X$  of  $L_p[0, 1]$  which is  $(1 + \varepsilon)$ -isomorphic to  $L_p[0, 1]$  and such that  $T|_X$  is a  $(1 + \varepsilon)$ -isomorphism (and  $X$  and  $T(X)$  are  $K_p$  complemented in  $L_p$ ).

*Remark 3.5.* Note that Theorem 3.4 is also true for  $p = 1$ . This result follows from [12, Theorem 1.2], where it is shown that if  $T \in \mathcal{L}(L_1)$  preserves a copy of  $L_1$ , then given  $\varepsilon > 0$ ,  $X$  can be chosen isometric to  $L_1$  so that  $T|_X$  is a  $(1 + \varepsilon)$ -isomorphism.

Having Remark 3.5 in mind, sometimes we may use Theorem 3.4 for the case  $p = 1$  as well.

Before we continue our study of the operators on  $L_p$  that preserve a copy of  $L_p$  we prove Theorem 1.3 in the case of  $L_p$ ,  $1 < p < 2$ . For this we need two lemmas for non-sign-embeddings and  $L_p$ -strictly singular operators on  $L_p$  that we use both in the next section and later on.

**3.1.  $L_p$  - strictly singular operators.** Lemma 3.6 was proved in [12] for the case  $p = 1$ , and basically the same proof works for general  $p$ ,  $1 \leq p < \infty$ .

**Lemma 3.6.** *Let  $T: L_p \rightarrow L_p$ ,  $1 \leq p < \infty$  be a non-sign-embedding operator. Then for all subsets  $S \subset \mathbb{R}$  with positive measure there exists a subspace  $X \subset L_p(S)$  of  $L_p$ ,  $X \cong L_p$ , such that  $T|_X$  is compact.*

*Proof.* We can choose by induction sets  $A_i$  in  $S$  such that  $A_1 = S$ ,  $A_n = A_{2n} \cup A_{2n+1}$ ,  $A_{2n} \cap A_{2n+1} = \emptyset$ ,  $\mu(A_{2n}) = \frac{1}{2}\mu(A_n)$ , and  $\|Tx_n\| < \frac{\varepsilon}{2^{n+1}}$  where  $x_n = \frac{\mathbf{1}_{A_{2n}} - \mathbf{1}_{A_{2n+1}}}{\mu(A_n)^{\frac{1}{p}}}$ . In order to do that assume that we have  $A_1, A_2, \dots, A_k$ , where  $k$  is an odd number and let  $n = \frac{k+1}{2}$ . Since  $T$  is not a sign-embedding, there exists  $y_n$  such that  $\int y_n = 0$ ,  $|y_n| = \mathbf{1}_{A_n}$ , and  $\|Ty_n\| \leq \frac{\varepsilon}{2^{n+1}}\mu(A_n)^{\frac{1}{p}}$ . Then set  $x_n = \frac{y_n}{\mu(A_n)^{\frac{1}{p}}}$ ,  $A_{2n} = \{x : x_n(x) = 1\}$ , and  $A_{2n+1} = \{x : x_n(x) = -1\}$ . It is not hard to see that  $(x_i)_{i=1}^\infty$  is isometrically equivalent to the usual sequence of Haar functions and hence  $X = \text{span}\{x_i\}$  is isometric to  $L_p$ . From the fact that  $(x_i)$  is a monotone basis for  $X$ , it is easy to deduce that  $T|_X$  is a compact operator.  $\square$

Note that Lemma 3.6 immediately implies that for every complemented subspace  $Y$  of  $L_p$ ,  $Y \cong L_p$ , there exists a complemented subspace  $X \subseteq Y$ ,  $X \cong L_p$ , such that  $T|_X$  is compact. The following lemma, which is also an immediate consequence of Lemma 3.6 and Theorem 3.4, will be used in the sequel.

**Lemma 3.7.** *Let  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < 2$ , be an  $L_p$ -strictly singular operator. Then for any  $X \subseteq L_p$ ,  $X \cong L_p$  and  $\varepsilon > 0$  there exists  $Y \subseteq X$ ,  $Y \cong L_p$  such that  $\|T|_Y\| < \varepsilon$ .*

*Proof.* Again, this result immediately follows from Lemma 3.6 and Theorem 3.4. An  $L_p$ -strictly singular operator cannot be a sign-embedding (Theorem 3.4), and then we use the construction in Lemma 3.6. From the fact that  $(x_i)$  is a monotone basis for  $X$  it follows that  $\|T|_X\| < \varepsilon$ .  $\square$

*Remark 3.8.* Clearly the proof we have for Lemma 3.7 depends heavily on Theorem 3.4, which is the deepest result of this paper. We do not know if an analogue of Lemma 3.7 holds for  $2 < p < \infty$ .

*Proof of Theorem 1.3 in the case of  $L_p$ ,  $1 \leq p < 2$ .* This is a direct consequence of Theorem 3.4 and Lemma 3.6. First we find a complemented subspace  $X' \subset Y$ ,  $X' \simeq L_p$ . Now we observe that an  $L_p$ -strictly singular operator cannot be a sign-embedding operator (Theorem 3.4), and then we use Lemma 3.6 for  $X'$  to find a complemented subspace  $X \subset X'$ ,  $X \simeq L_p$ , such that  $T|_X$  is a compact operator.  $\square$

### 3.2. Operators that preserve a copy of $L_p$ .

**Definition 3.9.** For  $T : L_p \rightarrow L_p$  and  $X \subseteq L_p$  define the following two quantities:

$$(3) \quad f(T, X) = \inf_{x \in \mathcal{S}_X} \|Tx\|,$$

$$(4) \quad g(T, X) = \sup_{\substack{Y \subseteq X \\ Y \simeq L_p}} f(T, Y).$$

Clearly  $f(T, X)$  does not decrease and  $g(T, X)$  does not increase if we pass to subspaces. For an arbitrary subspace  $Z \subset L_p$ ,  $Z \simeq L_p$  note the following two (equivalent) basic facts:

- $T|_{Z'}$  is an isomorphism for some  $Z' \subset Z$ ,  $Z' \simeq L_p$  if and only if  $g(T, Z) > 0$ .
- $T|_Z$  is  $L_p$ -strictly singular if and only if  $g(T, Z) = 0$ .

**Proposition 3.10.** *Let  $S : L_p \rightarrow L_p$ ,  $1 \leq p < 2$ , be an  $L_p$ -strictly singular operator and let  $Z$  be a subspace of  $L_p$  which is also isomorphic to  $L_p$ . Then for every operator  $T \in \mathcal{L}(L_p)$  we have  $g(T + S, Z) = g(T, Z)$ .*

*Proof.* If  $T|_Z$  is  $L_p$ -strictly singular, then  $(T+S)|_Z$  is also  $L_p$ -strictly singular; hence  $g(T, Z) = g(T + S, Z) = 0$ . For the rest of the proof we consider the case where there exists  $Z' \subset Z$ ,  $Z' \simeq L_p$  such that  $T|_{Z'}$  is an isomorphism; hence  $g(T, Z) > 0$ .

Let  $0 < \varepsilon < g(T, Z)/4$  and let  $Y \subseteq Z$ ,  $Y \simeq L_p$  be such that  $g(T, Z) - \varepsilon < f(T, Y)$ . Using Lemma 3.7 we find  $Y_1 \subseteq Y$ ,  $Y_1 \simeq L_p$  such that  $\|S|_{Y_1}\| < \varepsilon$ . Now

$$g(T + S, Z) \geq f(T + S, Y_1) > f(T, Y_1) - \varepsilon \geq f(T, Y) - \varepsilon > g(T, Z) - 2\varepsilon;$$

hence  $g(T + S, Z) \geq g(T, Z) - 2\varepsilon$ . Switching the roles of  $T$  and  $T + S$  (apply the previous argument for  $T + S$  and  $-S$ ) gives us  $g(T, Z) \geq g(T + S, Z) - 2\varepsilon$  and since  $\varepsilon$  was arbitrarily small we conclude that  $g(T + S, Z) = g(T, Z)$ .  $\square$

**Lemma 3.11.** *Let  $X$  and  $Y$  be two subspaces of  $L_p$ ,  $1 \leq p < 2$ , such that  $X \simeq Y \simeq L_p$ . Then there exist subspaces  $X_1 \subseteq X$ ,  $Y_1 \subseteq Y$  such that  $X_1 \simeq Y_1 \simeq L_p$ ,  $d(X_1, Y_1) > 0$ , and  $X_1 + Y_1$  is a complemented subspace of  $L_p$ . Moreover,  $X_1$  and  $Y_1$  can be chosen in such a way that there exists a projection onto  $X_1 + Y_1$  with norm depending only on  $p$ , and  $X_1 + Y_1$  is isomorphic to  $L_p$  with the isomorphism constant depending only on  $p$ .*

*Proof.* Without loss of generality, by passing to a subspace if necessary, we may assume that  $X \simeq Y \simeq L_p$  are two complemented subspaces of  $L_p$ . Our first step is to find two subspaces  $X_1 \subseteq X$  and  $Y_1 \subseteq Y$  which are isomorphic to  $L_p$  and  $d(X_1, Y_1) > 0$ .



Let  $P: L_p \rightarrow X$  and  $Q: L_p \rightarrow Y$  be two onto projections. We consider two cases for the operator  $Q|_X: X \rightarrow Y$ :

*Case 1.*  $Q|_X$  is  $L_p$ -strictly singular. Fix  $\delta > 0$ . Using Lemma 3.7 we find  $X_1 \subseteq X$ ,  $X_1 \simeq L_p$  such that  $\|Q|_{X_1}\| < \delta$ . We are going to show that  $d(X_1, Y) > 0$ .

Let  $x \in S_{X_1}$  and  $y \in Y$  be arbitrary. If  $\|y\| \notin [\frac{1}{2}, 2]$ , then clearly  $\|x - y\| > \frac{1}{2}$ . If not, then

$$\|x - y\| \geq \frac{\|Qx - Qy\|}{\|Q\|} = \frac{\|Qx - y\|}{\|Q\|} \geq \frac{\|y\| - \|Qx\|}{\|Q\|} \geq \frac{1}{\|Q\|} \left(\frac{1}{2} - \delta\right).$$

Since  $x$  and  $y$  were arbitrary we conclude that  $d(X_1, Y) > 0$  (here we take  $Y_1 = Y$ ).

*Case 2.*  $Q|_X$  is not  $L_p$ -strictly singular. Fix  $\delta > 0$ . Then using Theorem 3.4 we find  $X' \subseteq X$  such that  $X'$  is isomorphic to  $L_p$  and  $Q|_{X'}$  is a  $K_p$ -isomorphism. Without loss of generality we may assume that  $X' = L_p(\nu)$  for some nonatomic measure  $\nu$ . Now we find two disjoint,  $\nu$ -measurable sets  $A$  and  $B$  with positive measure and denote  $X_1 = L_p(A) \subseteq L_p(\nu)$ ,  $X_2 = L_p(B) \subseteq L_p(\nu)$ , and  $Y_1 = QX_2$ . Clearly  $d(X_1, X_2) = 1$  and we are going to show that  $d(X_1, QX_2) > 0$ .

Let  $x \in S_{X_1}$  and  $y = Qz \in QX_2$ . Then

$$\|x - y\| \geq \frac{\|Qx - Qz\|}{\|Q\|} \geq \frac{\|x - z\|}{K_p} \geq \frac{1}{K_p}.$$

Having  $X_1$  and  $Y_1$  from our first step, without loss of generality (by passing to a subspace if necessary) we may assume that  $X_1$  and  $Y_1$  are  $K_p$ -complemented in  $L_p$  and  $K_p$  isomorphic to  $L_p$  and, for simplicity of notation, we will use  $X$  and  $Y$  instead of  $X_1$  and  $Y_1$ .

Let  $P: L_p \rightarrow X$  and  $Q: L_p \rightarrow Y$  be two onto projections of norm at most  $K_p$ . It is easy to see that  $(I - Q)|_X$  is not an  $L_p$ -strictly singular operator. If we assume that this is not the case, fixing  $\delta > 0$  and using Lemma 3.7, we find  $X' \subset X$ ,  $X' \simeq L_p$ , such that  $\|(I - Q)|_{X'}\| < \delta$ . But then for  $x' \in S_{X'}$  and  $Qx' \in Y$  we have  $\|x' - Qx'\| < \delta$ , which is a contradiction with  $d(X, Y) > 0$  since  $\delta$  was arbitrary. Similarly, we show that  $(I - P)|_Y$  is not an  $L_p$ -strictly singular operator.

Fix  $\varepsilon > 0$ . Let  $X_1 \subset X$ ,  $X_1 \simeq L_p$ , be such that  $(I - Q)|_{X_1}$  is an isomorphism and  $(I - Q)|_{X_1}$  is a  $K_p$ -isomorphism. By Theorem 3.4 there exists a  $K_p$ -complemented subspace  $X_2 \subset (I - Q)(X_1)$  which is also  $K_p$ -isomorphic to  $L_p$ . Denote  $X' = ((I - Q)|_{X_2})^{-1}(X_2)$ . Now  $X' \simeq L_p$  and it is easy to see that  $X'$  is complemented in  $L_p$  (via the projection  $((I - Q)|_{X_2})^{-1}P_{X_2}(I - Q)$  of norm at most  $K_p^2$ , where  $P_{X_2}$  is a projection of norm  $K_p$  onto  $X_2$ ).

Similarly, we find  $Y' \subseteq Y$  such that  $Y'$  is  $K_p$ -isomorphic to  $L_p$ ,  $K_p^2$ -complemented in  $L_p$ , and  $(I - P)|_{Y'}$  is an isomorphism.

Let  $R_1$  and  $R_2$  be projections onto  $(I - Q)X'$  and  $(I - P)Y'$ , respectively, of norm at most  $K_p^2$ . Denote by  $V_1: (I - Q)X' \rightarrow X'$  the inverse map of  $(I - Q)|_{X'}: X' \rightarrow (I - Q)X'$  and, similarly, denote by  $V_2: (I - P)Y' \rightarrow Y'$  the inverse map of  $(I - P)|_{Y'}: Y' \rightarrow (I - P)Y'$ . Then a basic algebraic computation shows that  $PV_1R_1(I - Q) + QV_2R_2(I - P)$  is a projection onto  $X' + Y'$  of norm at most  $2K_p^4$ .

Consider two onto isomorphisms  $T_1: X' \rightarrow L_p$ ,  $T_2: Y' \rightarrow L_p$  such that  $\|T_i\| = 1$  and  $\|T_i^{-1}\| \leq K_p$  for  $i = 1, 2$  (such exist by our choice of  $X'$  and  $Y'$ ). Then the operator  $T_3: X' + Y' \rightarrow L_p \oplus_p L_p \equiv L_p$  defined by  $T_3(z) = (T_1V_1(I - Q)z, T_2V_2(I - P)z)$ ,

for every  $z \in X' + Y'$ , is an isomorphism with isomorphism constant depending only on  $p$ , which finishes the proof.  $\square$

**Lemma 3.12.** *Let  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < 2$ , and assume that for every  $X \subseteq L_p$ ,  $X \simeq L_p$ , there exists a subspace  $X_1 \subseteq X$ ,  $X_1 \simeq L_p$ , and a constant  $\lambda(X_1)$  such that  $T|_{X_1} = \lambda(X_1)I_{X_1} + S(X_1)$ , where  $S(X_1)$  is an  $L_p$ -strictly singular operator. Then there exists a constant  $\lambda$  and an  $L_p$ -strictly singular operator  $S$  such that  $T = \lambda I + S$ .*

*Proof.* Let  $X$  and  $Y$  be arbitrary subspaces of  $L_p$  which are also isomorphic to  $L_p$  and let  $X_1 \subseteq X$  and  $Y_1 \subseteq Y$  be the subspaces from the statement of the lemma. We will show that  $\lambda(X_1) = \lambda(Y_1)$ . Without loss of generality, using Lemma 3.11, we can assume that  $X_1 \cap Y_1 = \{0\}$  and  $X_1 + Y_1$  is a closed and complemented subspace of  $L_p$ . Let

$$T|_{X_1} = \lambda_1 I_{X_1} + S_1, \quad T|_{Y_1} = \lambda_2 I_{Y_1} + S_2.$$

Let  $\tau: X_1 \rightarrow Y_1$  be an isomorphism and define  $Z = \{x + \tau(x) \mid x \in X_1\}$ . Then

$$(5) \quad T|_Z = T(x + \tau x) = \lambda_1 x + \lambda_2 \tau x + S_1 x + S_2 \tau x.$$

The operator  $S: Z \rightarrow L_p$  defined by  $S(x + \tau x) = S_1 x + S_2 \tau x$  is  $L_p$ -strictly singular as a sum of two such operators. From the assumption of the lemma, there exist  $Z_1 \subseteq Z$  and  $\lambda_3 \in \mathbb{C}$  such that  $Z_1 \simeq L_p$  and

$$(6) \quad T|_{Z_1} = \lambda_3 I_{Z_1} + S_3,$$

where  $S_3$  is  $L_p$ -strictly singular. From (5) and (6) we obtain that the operator  $T_1: Z \rightarrow L_p$  defined by  $T_1(x + \tau x) = \lambda_1 x + \lambda_2 \tau x - \lambda_3(x + \tau x)$  is also  $L_p$ -strictly singular on  $Z_1$ ; i.e.,  $T_1|_{Z_1}$  is  $L_p$  strictly singular. The last conclusion is possible if and only if  $\lambda_1 = \lambda_2 = \lambda_3$ . In fact, if we assume that  $\lambda_1 \neq \lambda_3$ , then the operator  $T_1$  will be an isomorphism on  $Z$  because for every  $x \in S_{X_1}$  we will have

$$\begin{aligned} \|T_1(x + \tau x)\| &= |\lambda_1 - \lambda_3| \left\| x + \frac{\lambda_2 - \lambda_3}{\lambda_1 - \lambda_3} \tau x \right\| \geq |\lambda_1 - \lambda_3| d(X_1, Y_1) \|x\| \\ &\geq \frac{|\lambda_1 - \lambda_3|}{1 + \|\tau\|} d(X_1, Y_1) \|x + \tau x\|. \end{aligned}$$

Let  $\lambda = \lambda(X_1)$  for every subspace  $X_1$  as in the statement of the lemma. Now it easily follows that  $\lambda I - T$  is  $L_p$ -strictly singular. Indeed, if we assume otherwise, then there exists a subspace  $Z \subseteq L_p$ ,  $Z \simeq L_p$  such that  $(\lambda I - T)|_Z$  is an isomorphism. But according to the assumptions of the lemma, there exists  $Z_1 \subset Z$ ,  $Z_1 \simeq L_p$  such that  $(\lambda I - T)|_{Z_1}$  is  $L_p$ -strictly singular, which contradicts the fact that  $(\lambda I - T)|_Z$  is an isomorphism. This finishes the proof.  $\square$

An immediate corollary of Lemma 3.12 is that for an operator  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < \infty$ , not of the form  $\lambda I + S$ , where  $S$  is an  $L_p$ -strictly singular operator, there exists a complemented subspace  $X \subset L_p$ ,  $X \simeq L_p$  such that  $(T - \lambda I)|_X$  preserves a copy of  $L_p$  for every  $\lambda \in \mathbb{C}$ , and this is in fact what we are going to use in the sequel.

**Lemma 3.13.** *Let  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < 2$ , and assume that for every  $X \subseteq L_p$ ,  $X \simeq L_p$ , and every  $\varepsilon > 0$  there exist  $X_1 \subset X$ ,  $X_1 \simeq L_p$ , and  $\lambda = \lambda(X_1)$  such that  $g(\lambda I - T, X_1) < \varepsilon$ . Then for every  $\varepsilon > 0$  there exists  $\lambda_\varepsilon$  such that  $g(\lambda_\varepsilon I - T, L_p) < D_p \varepsilon$ , where  $D_p$  is a constant depending only on  $p$ .*

*Proof.* From the assumption in the statement of the lemma, without loss of generality we may assume that for every  $X \subseteq L_p$ ,  $X \simeq L_p$ , and every  $\varepsilon > 0$  there exist  $X_1 \subset X$ ,  $X_1 \simeq L_p$ , and  $\lambda = \lambda(X_1)$  such that  $g(\lambda I - T, X_1) < \varepsilon$  and  $(\lambda I - T)|_{X_1}$  is an isomorphism. If  $g(\lambda I - T, X_1) > 0$ , this can be achieved by passing to a subspace  $Y_1$  of  $X_1$  for which  $f(\lambda I - T, Y_1) > 0$ . If  $g(\lambda I - T, X_1) = 0$  for each  $X_1 \subset X$ ,  $X_1 \simeq L_p$ , using the fact that  $g$  is a continuous function of  $\lambda$ , we find  $\lambda_0$  such that  $0 < g(\lambda_0 I - T, X) < \varepsilon$  and then find  $X_1 \subset X$ ,  $X_1 \simeq L_p$  such that  $f(\lambda_0 I - T, X_1) > 0$ .

Fix an  $\varepsilon > 0$  and let  $Y_1$  and  $Y_2$  be any two subspaces of  $L_p$  such that  $Y_1 \simeq Y_2 \simeq L_p$ . From our assumptions, there exist complemented subspaces  $Y'_1, Y'_2$  such that  $Y'_i \subseteq Y_i$ ,  $Y'_i \simeq L_p$ , and  $g(\lambda(Y'_i)I - T, Y'_i) < \varepsilon$  for  $i = 1, 2$  and without loss of generality, using Theorem 3.4 and passing to a subspace if necessary, we may assume that  $Y'_1$  and  $Y'_2$  are  $K_p$ -complemented in  $L_p$  and  $((\lambda(Y'_i)I - T)|_{Y'_i})$  is a  $K_p$ -isomorphism for  $i = 1, 2$ . Then we apply Lemma 3.11 to get subspaces  $X_1, X_2$  such that

- $X_i \subseteq Y'_i$ ,  $X_i \simeq L_p$  for  $i = 1, 2$ ,
- $X_1 \cap X_2 = \{0\}$ ,
- $X_1$  and  $X_2$  are  $d_p$ -complemented and  $d_p$ -isomorphic to  $L_p$  via  $\lambda(Y'_1)I - T$  and  $\lambda(Y'_2)I - T$ , respectively, for some constant  $d_p$  depending only on  $p$  (this follows from Lemma 3.11 and our choice of  $Y'_1$  and  $Y'_2$ ),
- $X_1 + X_2$  is a closed and complemented subspace of  $L_p$ , and there exists a projection onto  $X_1 + X_2$  with norm depending only on  $p$ .

Since  $X_i \subseteq Y'_i$ ,  $i = 1, 2$ , we have  $g(\lambda(Y'_i)I - T, X_i) < \varepsilon$  for  $i = 1, 2$ , which in view of our choice of  $X_1$  and  $X_2$  implies that

$$(7) \quad \max(\|(\lambda(Y'_1)I - T)|_{X_1}\|, \|(\lambda(Y'_2)I - T)|_{X_2}\|) < d_p \varepsilon.$$

Our goal is to show that  $|\lambda(Y'_1) - \lambda(Y'_2)| < c_p \varepsilon$  for some constant  $c_p$  independent of  $Y'_1$  and  $Y'_2$ .

Let  $\tau: X_1 \rightarrow X_2$  be an isomorphism such that  $\|\tau\| \leq d_p^2$  and  $\|\tau^{-1}\| = 1$ . Define

$$Z = \{x + \tau x \mid x \in X_1\}.$$

By assumption, there exists  $Z' \subseteq Z$ ,  $Z' \simeq L_p$ , and  $\lambda(Z')$  such that  $0 < g(\lambda(Z')I - T, Z') < \varepsilon$  and  $\lambda(Z')I - T$  is an isomorphism on  $Z'$  (like the argument in the beginning of the proof). Using Theorem 3.4 we find  $Z'' \subseteq Z'$  such that  $Z''$  is  $K_p$  isomorphic to  $L_p$  via  $\lambda(Z')I - T$  (clearly  $g(\lambda(Z')I - T, Z'') < \varepsilon$ ). Let  $U = \lambda(Z')I - T$  and define an operator  $S: Z' \rightarrow L_p$  by  $S(x + \tau x) = \lambda(Y'_1)x + \lambda(Y'_2)\tau x - Tx - T\tau x$ . A simple application of the triangle inequality combined with (7) implies that

$$(8) \quad \|S(x + \tau x)\| \leq d_p^3 \varepsilon (\|x\| + \|\tau x\|).$$

From our choice of  $Z''$  we also have

$$(9) \quad \|\lambda(Z')(x + \tau x) - T(x + \tau x)\| \leq K_p \varepsilon (\|x\| + \|\tau x\|)$$

and combining (8) and (9) gives us

$$(10) \quad \|(U - S)(x + \tau x)\| \leq (K_p + d_p^3)\varepsilon (\|x\| + \|\tau x\|) \leq (K_p + d_p^3)(1 + d_p^2)\varepsilon \|x\|.$$

On the other hand,

$$\begin{aligned}
\|(U - S)(x + \tau x)\| &= \|(\lambda(Z') - \lambda(Y'_1))x + (\lambda(Z') - \lambda(Y'_2))\tau x\| \\
(11) \qquad \qquad \qquad &\geq A_p(|\lambda(Z') - \lambda(Y'_1)|\|x\| + |\lambda(Z') - \lambda(Y'_2)|\|\tau x\|) \\
&\geq A_p(|\lambda(Z') - \lambda(Y'_1)| + |\lambda(Z') - \lambda(Y'_2)|)\|x\| \\
&\geq A_p|\lambda(Y'_2) - \lambda(Y'_1)|\|x\|
\end{aligned}$$

(where  $A_p$  depends on  $d_p$ ,  $K_p$  and the norm of the projection onto  $X_1 + X_2$  which also depends on  $p$  only). Combining (8) and (9) we get  $|\lambda(Y'_1) - \lambda(Y'_2)| < \frac{(K_p + d_p^3)(1 + d_p^2)}{A_p}\varepsilon$ . Now we define  $\lambda_\varepsilon = \lambda(Y'_1)$  and it is not hard to check that  $g(\lambda_\varepsilon I - T, L_p) \leq (1 + \frac{(K_p + d_p^3)(1 + d_p^2)}{A_p})\varepsilon$ .  $\square$

*Remark 3.14.* Let  $1 \leq p < \infty$ . Suppose that there exists a sequence of numbers  $\{\lambda_i\}_{i=1}^\infty$  such that  $g(\lambda_n I - T, L_p) \xrightarrow{n \rightarrow \infty} 0$ . Then there exists  $\lambda$  such that  $g(\lambda I - T, L_p) = 0$ . This is easy to see by noticing that  $g(\lambda I - T, L_p)$  is bounded away from 0 for large  $\lambda$ , so without loss of generality we can assume that  $\lambda_n \xrightarrow{n \rightarrow \infty} \lambda$ . Then, if  $g(\lambda I - T, L_p) = 4\delta > 0$ , there exists  $Y \subseteq L_p$ ,  $Y \simeq L_p$ , such that  $f(\lambda I - T, Y) > 2\delta$ . Now if  $|\lambda - \mu| < \delta$ , then  $f(\mu I - T, Y) > \delta$  and hence  $g(\mu I - T, L_p) > \delta$ , which contradicts our original assumption about the sequence  $\{\lambda_i\}_{i=1}^\infty$ .

From the last remark it trivially follows that if  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < \infty$ , is such that  $\lambda I - T$  preserves a copy of  $L_p$  for every  $\lambda$ , then  $\inf_\lambda g(\lambda I - T, L_p) > 0$ .

**Lemma 3.15.** *Let  $T \in \mathcal{L}(L_p)$ ,  $1 \leq p < 2$ , and assume that  $\lambda I - T$  preserves a copy of  $L_p$  for every  $\lambda \in \mathbb{C}$ . Then there exists  $\varepsilon > 0$  and a subspace  $X \subseteq L_p$ ,  $X \simeq L_p$ , such that for every  $X' \subseteq X$ ,  $X' \simeq L_p$ , and every  $\lambda \in \mathbb{C}$  we have  $g(\lambda I - T, X') > \varepsilon$ .*

*Proof.* The proof is straightforward from Lemma 3.13 and the last remark.  $\square$

The next result of this section is a reduction lemma that will help us later.

**Lemma 3.16.** *Let  $T, P \in \mathcal{L}(L_p)$ ,  $1 \leq p < \infty$ , be such that  $P$  is a projection satisfying  $PL_p \simeq L_p$  and  $(I - P)TP$  is an isomorphism on a subspace  $X \subseteq L_p$ ,  $X \simeq L_p$ . Then there exists a subspace  $Z \subset L_p$  such that  $Z \simeq TZ \simeq L_p$ ,  $d(TZ, Z) > 0$ , and  $Z + TZ$  is a subspace isomorphic to  $L_p$  and complemented in  $L_p$ .*

*Proof.* Denote  $Y = PX$  and note that from the statement we have  $Y \simeq L_p$ . Since  $(I - P)T$  is an isomorphism on  $Y$ , for every  $x \in Y$  we have  $\|(I - P)Tx\| \geq c\|x\| \geq c\frac{\|Tx\|}{\|T\|}$ . Now if  $x, y \in Y$  are arbitrary such that  $Tx \in S_Y$ , then

$$d(TY, Y) \geq \|Tx - y\| \geq \frac{\|(I - P)Tx\|}{\|I - P\|} \geq \frac{c}{\|T\|\|I - P\|} > 0.$$

Without loss of generality we may assume that both  $Y$  and  $TY$  are complemented in  $L_p$ . This can be done, for example, by choosing a complemented subspace  $Y_1 \subseteq TY$ ,  $Y_1 \simeq L_p$  and then considering  $(T|_Y)^{-1}Y_1$  instead of  $Y$ .

Let  $P_Y$  be a projection onto  $Y$  and denote  $T' = (I - P_Y)TP_Y$ . Since  $T$  is an isomorphism on  $Y$  and  $d(Y, TY) > 0$  we have that  $T'$  is an isomorphism on  $Y$ . Now we choose a subspace  $Y_1 \subseteq T'Y$  which is isomorphic to  $L_p$  and complemented in  $L_p$  and denote  $Z = (T'|_Y)^{-1}Y_1$ . Clearly  $Z$  is isomorphic to  $L_p$  and complemented in  $L_p$  via  $(T'|_Y)^{-1}P_{Y_1}T'$ , where  $P_{Y_1} : L_p \rightarrow Y_1$  is a projection onto  $Y_1$ . Let  $V$  be

the inverse of the isomorphism  $T(y) \rightarrow T'(y)$ ,  $y \in Y$ . It is not hard to see that  $P_{TZ} := VP_{Y_1}(I - P_Y)$  is a projection from  $L_p$  onto  $TZ$  such that  $P_{TZ}Z = \{0\}$ , and then it is straightforward to verify that  $P_TZ + P_Z(I - P_{TZ})$  is a projection onto  $Z + TZ$ , which finishes the proof.  $\square$

*Proof of Theorem 1.2 for  $L_p$ ,  $1 < p < 2$ .* In view of Lemma 3.12, we can apply Lemma 3.15 for  $T$  and let  $X$  and  $\varepsilon$  be the ones from Lemma 3.15. Without loss of generality we may assume that  $X$  is complemented (otherwise we may pass to a complemented subspace). Let  $V$  be an isomorphism from  $L_p$  into  $L_p$  such that  $VL_p = X$ .

Fix  $\delta > 0$  to be chosen later. We will build sequences  $\{y_i\}_{i=1}^\infty$  in  $X$ , and  $\{a_i\}_{i=1}^\infty$  and  $\{b_i\}_{i=1}^\infty$  in  $L_p$  such that:

- (1)  $\{a_i\}_{i=1}^\infty$  and  $\{b_i\}_{i=1}^\infty$  are block bases of  $\{h_{0,0}\} \cup \{h_{n,i}\}_{n=0, i=1}^\infty$  such that if we denote  $\sigma_i = \text{supp}\{a_i\} \cup \text{supp}\{b_i\}$ , where the support is with respect to the Haar basis, then  $\{\sigma_i\}_{i=1}^\infty$  is a disjoint sequence of subsets of  $\{h_{0,0}\} \cup \{h_{n,i}\}_{n=0, i=1}^\infty$ .
- (2)  $\{y_i\}_{i=1}^\infty$  is equivalent to the Haar basis for  $L_p$ .
- (3)  $\|y_i - a_i\| < \frac{\delta}{2^i}\|y_i\|$  and  $\|Ty_i - b_i\| < \frac{\delta}{2^i}\|Ty_i\|$  for all  $1 \leq i < \infty$ .

The construction of these sequences is similar to the construction of Lemma 3.11, and we sketch it below for completeness. As before, by  $P_{(k,s)}$  we denote the projection onto the linear span of  $\{h_{n,i}\}_{n=k, i=1}^{2^n}$ .

Let  $y_1 = Sh_{1,1}$ . There exists  $n_1$  such that  $\|y_1 - P_{(1,n_1)}y_1\| < \frac{\delta}{2}\|y_1\|$  and  $\|Ty_1 - P_{(1,n_1)}Ty_1\| < \frac{\delta}{2}\|Ty_1\|$ . Let  $a_1 = P_{(1,n_1)}y_1$  and  $b_1 = P_{(1,n_1)}Ty_1$ . Denote  $A_1^+ = \{x : V^{-1}y_1(x) = 1\}$  and  $A_1^- = \{x : V^{-1}y_1(x) = -1\}$ . Let  $\{z_i\}_{i=1}^\infty$  be the Rademacher sequence on  $A_1^+$  and  $\{z'_i\}_{i=1}^\infty$  be the Rademacher sequence on  $A_1^-$ . Using the fact that the Rademacher sequence is weakly null, we find  $l$  such that

$$\begin{aligned} \|Sz_l - P_{(n_1,\infty)}Sz_l\| &< \frac{\delta}{16}\|Sz_l\|, \\ \|TSz_l - P_{(n_1,\infty)}TSz_l\| &< \frac{\delta}{16}\|TSz_l\|, \\ \|Sz'_l - P_{(n_1,\infty)}Sz'_l\| &< \frac{\delta}{16}\|Sz'_l\|, \\ \|TSz'_l - P_{(n_1,\infty)}TSz'_l\| &< \frac{\delta}{16}\|TSz'_l\|. \end{aligned}$$

Define  $y_2 = Sz_l$ ,  $y_3 = Sz'_l$  and find  $n_2$  such that  $\|y_k - P_{(n_1,n_2)}y_k\| < \frac{\delta}{8}\|y_k\|$  and  $\|Ty_k - P_{(n_1,n_2)}Ty_k\| < \frac{\delta}{8}\|Ty_k\|$  for  $k = 2, 3$ . As before, let  $a_k = P_{(n_1,n_2)}y_k$  and  $b_k = P_{(n_1,n_2)}Ty_k$  for  $k = 2, 3$ .

Continuing this way, we build the other elements of the sequences  $\{y_i\}_{i=1}^\infty$ ,  $\{a_i\}_{i=1}^\infty$  and  $\{b_i\}_{i=1}^\infty$ . From the construction it is clear that  $\{(\text{supp } y_i \cup \text{supp } Ty_i)\}_{i=1}^\infty$  are essentially disjoint. If we denote  $Y = \overline{\text{span}}\{y_i : i = 1, 2, \dots\}$  we have that  $Y$  is a complemented subspace of  $L_p$  which is also isomorphic to  $L_p$ . To see this it is enough to notice that  $V^{-1}Y$  is isometric to  $L_p$  (since it is spanned by a sequence which is isometrically equivalent to the Haar basis) and hence complemented in  $L_p$ . One projection onto  $Y$  is given by  $P_Y = VP_{V^{-1}Y}V^{-1}P_X$ . From now on, without loss of generality we assume that  $X = Y$  (since we can pass to a subspace in the beginning if necessary).

As in the argument in Lemma 3.11, using the principle of small perturbations, it is easy to see that the subspace  $A = \overline{\text{span}}\{a_i : i = 1, 2, \dots\}$  is complemented. A projection onto  $A$  is given by  $P'_A = GP_YG^{-1}$ , where  $G \in \mathcal{L}(L_p)$  is defined by

$$G = I - \sum_i y_i^*(\cdot)(y_i - a_i)$$

(in the definition above,  $\{y_i^*\}$  are the biorthogonal functionals to  $\{y_i\}$ ). Using Proposition 3.3 we find a projection  $P_A$  onto  $A$  that respects supports. Let  $S : A \rightarrow L_p$  be the operator defined by

$$S = I - \sum_i a_i^*(\cdot)(a_i - y_i),$$

where  $\{a_i^*\}$  are the biorthogonal functionals to  $\{a_i\}$ , and note that  $SA = Y(\equiv X)$  and  $\|I - S\| < 4\delta$ . Also let  $S' : A \rightarrow L_p$  be the operator defined by  $S'a_i = Ta_i - b_i$  and let  $T' = T - S'P_A$ . It follows easily that  $\|S'\| < C\delta\|T\|$ , where  $C$  depends only on  $p$ , since

$$\|S'a_i\| = \|Ta_i - b_i\| = \|T(a_i - y_i) + Ty_i - b_i\| < \frac{2\delta\|T\|\|y_i\|}{2^i}.$$

First we show that  $(I - P_A)T'P_A$  preserves a copy of  $L_p$ .

If not, we have that  $(I - P_A)T'P_A$  is  $L_p$ -strictly singular and hence  $P_AT'P_A$  preserves a copy of  $L_p$  (otherwise  $T'|_A$  will be  $L_p$ -strictly singular, which is false). We also have the inequality  $g(P_AT'P_A, A) > \varepsilon/2$ . In order to show it we need to go back to the definitions of  $f(\cdot, \cdot)$  and  $g(\cdot, \cdot)$ . Fix  $A' \subseteq A$  and  $\lambda \in \mathbb{C}$ . We shall show that  $g(\lambda I + T', A') > \varepsilon/2$ . We may assume that  $|\lambda| < \|T'\| + 1$  (otherwise  $f(\lambda I - T', A') > 1$ ). Let  $x \in S_{A'}$  be arbitrary. Then

$$\begin{aligned} \|(\lambda I - T')x\| &\geq \|(\lambda I - T')Sx\| - \|(\lambda I - T')(x - Sx)\| \\ &\geq \|(\lambda I - T)Sx\| - \|S'P_ASx\| - \|(\lambda I - T')(x - Sx)\| \\ &\geq (1 - 4\delta)f((\lambda I - T), A') - C\delta\|T\|\|P_A\|(1 + 4\delta) - 4(2\|T'\| + 1)\delta. \end{aligned}$$

Taking the infimum over the left side we obtain  $f((\lambda I - T'), A') > f((\lambda I - T), A')/2$  for sufficiently small  $\delta$  and hence  $g(\lambda I - T', A') > \varepsilon/2$  for every  $\lambda \in \mathbb{C}$ . Using the fact that  $(I - P_A)TP_A$  is an  $L_p$ -strictly singular operator and Proposition 3.10 we obtain

$$g(P_AT'P_A, A) + g((I - P_A)T'P_A, A) = g(T'P_A, A) > \frac{\varepsilon}{2}.$$

Let  $P_AT'a_i = \lambda_i a_i$  (we can do that since  $P_A$  respects supports). Till the end of this proof it will be convenient to switch the enumeration of the  $\{a_i\}_{i=1}^\infty$  to  $\{a_{n,i}\}_{n=0, i=1}^\infty$ , which is actually how we constructed them. For each  $n$ , using the pigeon-hole principle, we can find a set  $\sigma(n)$  with cardinality at least  $\frac{\varepsilon 2^n}{4\|P_AT\|}$  and a number  $\mu_n$  such that  $|\mu_n - \lambda_i| < \frac{\varepsilon}{4}$  for every  $i \in \sigma(n)$ . Clearly, there exists an infinite subset  $N_1 \subseteq \mathbb{N}$  and a number  $\mu_\varepsilon$  such that

$$\sum_{n \in N_1} |\mu_n - \mu_\varepsilon| < \frac{\varepsilon}{100}.$$

Let  $Z = \overline{\text{span}}\{a_{n,i} : n \in N_1, i \in \sigma(n)\}$ . Using a result of Gamlen and Gaudet (see [7]), we have that  $Z \simeq L_p$  and clearly  $Z$  is complemented in  $L_p$ . Now note that

$$(12) \quad \|(\mu_\varepsilon I - P_A T')|_Z\| < \frac{\varepsilon}{2}; \text{ hence } g(\mu_\varepsilon I - P_A T', Z) < \frac{\varepsilon}{2}.$$

On the other hand,

$$(13) \quad \frac{\varepsilon}{2} < g(\mu_\varepsilon I - T', A) = g((\mu_\varepsilon I - P_A T') - (I - P_A)T', A) = g(\mu_\varepsilon I - P_A T', A)$$

since  $(I - P_A)T'_A$  is  $L_p$ -strictly singular. The equations (12) and (13) lead to a contradiction, which shows that  $(I - P_A)T'P_A$  preserves a copy of  $L_p$ , say  $Z'$ . Now  $\|(I - P_A)T'P_A - (I - P_A)TP_A\| = \|(I - P_A)S'P_A\| < C\delta\|T\|(\|P_A\| + 1)^2$ ; hence, for sufficiently small  $\delta$ , we have that  $(I - P_A)TP_A$  is an isomorphism on  $Z'$  and clearly  $d((I - P_A)TP_A Z', Z') > 0$ . In view of Lemma 3.16, this finishes the proof.  $\square$

#### 4. OPERATORS ON $L_1$

Recall that we have already proved Theorem 1.3 in the case of  $L_1$ . The proof in this case does not involve anything new and can be done only using the ideas found in [12], which we have already mentioned.

Now we switch attention to the operators not of the form  $\lambda I + K$ , where  $\lambda \in \mathbb{C}$  and  $K$  is in  $\mathcal{M}_{L_1}$ . Our investigation will rely on the representation Kalton gave for a general operator on  $L_1$  in [9], but again Rosenthal's paper [12] is a better reference for us. Before we state Kalton's representation we need a few definitions.

**Definition 4.1.** An operator  $T : L_1 \rightarrow L_1$  is called an atom if  $T$  maps disjoint functions to disjoint functions. That is, if  $\mu(\text{supp } f \cap \text{supp } g) = 0$ , then  $\mu(\text{supp } Tf \cap \text{supp } Tg) = 0$ .

Unlike the notation for  $1 < p < \infty$ , here  $\text{supp}$  refers to the support with respect to the interval  $[0, 1]$ . A simple characterization of the atoms is given by the following known structural result.

**Proposition 4.2** ([12, Proposition 1.3]). *An operator  $T : L_1 \rightarrow L_1$  is an atom if and only if there exist measurable functions  $a : (0, 1) \rightarrow \mathbb{R}$  and  $\sigma : (0, 1) \rightarrow (0, 1)$  with  $Tf(x) = a(x)f(\sigma x)$  a.e. for all  $f \in L_1$ . Moreover, a pair of functions  $a$  and  $\sigma$  represents an atom  $T$  if and only if  $K = \sup\{(\mu(s))^{-1} \int_{\sigma^{-1}(S)} |a| d\mu : \mu(s) > 0\} < \infty$ , and then  $\|T\| = K$ .*

In [12] the following definition is given:

**Definition 4.3.** Let  $T : L_1 \rightarrow L_1$  be a given operator.

- (a) Say that  $T$  has *atomic part* if there exists a nonzero atom  $A : L_1 \rightarrow L_1$  with  $0 \leq A \leq |T|$ .
- (b) Say that  $T$  is *purely continuous* if  $T$  has no atomic part.
- (c) Say that  $T$  is *purely atomic* if  $T$  is a strong  $\ell_1$ -sum of atoms.

The condition (c) in the preceding definition simply means that there is a sequence of atoms  $\{T_j\}_{j=1}^\infty$  from  $L_1$  to  $L_1$  and  $K < \infty$  so that for all  $f \in L_1$ ,  $\sum \|T_j f\| \leq K\|f\|$  and  $Tf = \sum T_j f$ .

Here is Kalton's representation theorem for operators on  $L_1$  the way it is stated in [12].

**Theorem 4.4.** *Let  $T : L_1 \rightarrow L_1$  be a given operator. There are unique operators  $T_a, T_c \in \mathcal{L}(L_1)$  so that  $T_a$  is purely atomic,  $T_c$  is purely continuous, and  $T = T_a + T_c$ . Moreover, there exists a sequence of atoms  $\{T_j\}_{j=1}^\infty$  so that  $T_a$  is a strong  $\ell_1$ -sum of  $\{T_j\}_{j=1}^\infty$  and the following four conditions hold:*

- (1)  $\sum_{i=1}^\infty \|T_i f\| \leq \|T_a\| \|f\|$ .
- (2)  $(T_i f)(x) = a_i(x) f(\sigma_i x)$  a.e., where  $a_i : (0, 1) \rightarrow \mathbb{R}$  are measurable functions, and  $\sigma_i : (0, 1) \rightarrow (0, 1)$ .
- (3) For all  $i \neq j$ ,  $\sigma_i(x) \neq \sigma_j(x)$  a.e.
- (4)  $|a_j(x)| \geq |a_{j+1}(x)|$  a.e.

Note that if  $E$  is a set of positive measure such that  $a_j(x) \neq 0$  a.e. on  $E$ , then  $\mu(\sigma_j(E)) > 0$ . Indeed, let  $F \subset E$  be such that  $|a_j(x)| > \alpha > 0$  for every  $x \in F$ . Now  $T_j 1_{\sigma_j(F)} = 1_F a_j$  implies that  $\|T_j 1_{\sigma_j(F)}\| > 0$ ; hence  $\mu(\sigma_j(F)) > 0$ . The power of Kalton's representation theorem is that it reduces many problems about operators on  $L_1$  to measure-theoretic considerations. This is illustrated in the proof of the following proposition.

**Proposition 4.5.** *Let  $T \in \mathcal{L}(L_1)$  be a nonzero atom such that  $T \neq \lambda I$  for any  $\lambda \in \mathbb{C}$ . Then there exists a subspace  $Y \subset L_1$  such that  $Y \equiv L_1$ ,  $d(Y, TY) > 0$ , and  $Y + TY$  is complemented in  $L_1$ .*

*Proof.* By the definition of an atom we have that  $(Tf)(x) = a(x)f(\sigma x)$  for some  $a$  and  $\sigma$ . We consider two possibilities depending on  $\sigma$ .

1. If  $\sigma = id$  a.e. on  $(0, 1)$ , then  $a(x) \not\equiv \text{const}$  a.e. (otherwise  $T = \lambda I$  for some  $\lambda$ ). Then we find two different numbers  $\lambda_1, \lambda_2$  and a positive number  $\delta$  such that:

- $|\lambda_1 - \lambda_2| > 3\delta$ .
- There are closed sets  $\Delta_i \subseteq \{x : |a(x) - \lambda_i| < \delta\}$  so that  $\mu(\Delta_i) > 0$  for  $i = 1, 2$ .

To see this we can consider a good enough approximation of  $a(x)$  with a step function and without loss of generality we may assume that  $\lambda_1 \neq -1$ . Note also that we can choose  $\delta$  as small as we want (independent of  $\lambda_1$  and  $\lambda_2$ ) which choice we leave for later. Clearly,  $\Delta_1 \cap \Delta_2 = \emptyset$  and, since they are closed, by shrinking  $\delta$  we can assume that they are at a distance of at least  $\delta$  apart. From our choice of  $\Delta_i$ ,  $i = 1, 2$ , we also have

$$\|(Tf - \lambda_i f) \mathbf{1}_{\Delta_i}\| < \delta \|f \mathbf{1}_{\Delta_i}\|, \quad i = 1, 2.$$

Let  $S : L_1(\Delta_1) \rightarrow L_1(\Delta_2)$  be an isometry and define

$$Z = \{f_1 + S f_1 : f_1 \in L_1(\Delta_1)\}.$$

Since  $\|f_1 + S f_1\| = 2\|f_1\|$  we immediately have that  $Z \equiv L_1$ . It is also clear that  $T|_Z$  is an isomorphism because  $\max\{|\lambda_1|, |\lambda_2|\} > \frac{3}{2}\delta$  and hence  $\|T(f_1 + S f_1)\| \geq \max\{|\lambda_1 - \delta|, |\lambda_2 - \delta|\} \|f_1\| > (\delta/2) \|f_1\|$  for every  $f_1 \in L_1(\Delta_1)$ .



To show that  $d(Z, TZ) > 0$  assume that  $\|T(g + Sg)\| = 1$  for some  $g \in L_1(\Delta_1)$ . Then for arbitrary  $f \in L_1(\Delta_1)$  we have

$$\begin{aligned} \|f + Sf - Tg - TSg\| &= \|f + Sf - ag - aSg\| = \|f - ag\| + \|Sf - aSg\| \\ &= \|f - \lambda_1 g + (\lambda_1 - a)g\| + \|Sf - \lambda_2 Sg + (\lambda_2 - a)Sg\| \\ &\geq \|f - \lambda_1 g\| - \|(\lambda_1 - a)g\| + \|Sf - \lambda_2 Sg\| - \|(\lambda_2 - a)Sg\| \\ &\geq \|f - \lambda_1 g\| + \|f - \lambda_2 g\| - \delta\|g\| - \delta\|Sg\| \\ &\geq |\lambda_1 - \lambda_2|\|g\| - 2\delta\|g\| \geq \frac{|\lambda_1 - \lambda_2|}{3}\|g\|. \end{aligned}$$

Now we observe that  $\|T(g + Sg)\| = 1$  implies that  $\|g\| \geq \frac{1}{2\|T\|}$ ; hence

$$d(TZ, Z) = \inf_{\substack{\|T(g+Sg)\|=1 \\ f, g \in L_1(\Delta_1)}} \|f + Sf - Tg - TSg\| \geq \frac{|\lambda_1 - \lambda_2|}{6\|T\|}.$$

Define  $T_1 f(x) = \lambda_1 f(x)\mathbf{1}_{\Delta_1}(x) + \lambda_2 f(x)\mathbf{1}_{\Delta_2}(x)$  and let  $K = T - T_1$ . Denote by  $P_1$  the natural, norm one, projection from  $L_1$  onto  $L_1(\Delta_1)$  and let  $P = P_1 + \frac{\lambda_2 + 1}{\lambda_1 + 1}SP_1$ . It is easy to see that  $P$  is an idempotent operator since  $P_1 SP_1 \equiv 0$ . To see that  $P$  is a projection onto  $Z + T_1 Z$ , note that

$$\begin{aligned} Pf &= P_1 f + \frac{\lambda_2 + 1}{\lambda_1 + 1}SP_1 f = \frac{1}{\lambda_1 + 1}((\lambda_1 + 1)P_1 f + (\lambda_2 + 1)SP_1 f) \\ &= \frac{1}{\lambda_1 + 1}(P_1 f + SP_1 f + \lambda_1 P_1 f + \lambda_2 SP_1 f) \in Z + T_1 Z. \end{aligned}$$

Now we observe that  $Z + TZ = Z + (T_1 + K)Z$  and use the fact that  $\|K|_Z\| < \delta$  to conclude that for sufficiently small  $\delta$ , Proposition 2.2 guarantees that the subspace  $Z + TZ$  is complemented.

**2.** If  $\sigma \neq id$  a.e. on  $(0, 1)$ , let  $A = \{x \mid \sigma(x) = x\}$  and denote  $A' = (0, 1) \setminus A$  and  $B = A' \cap \{x \mid a(x) \neq 0\}$ . We have two cases depending on  $\mu(B)$ .

*Case 1.*  $\mu(B) > 0$ . In this case we show that there exists  $\Delta \subset B$  such that  $\mu(\Delta \cap \sigma(\Delta)) = 0$ . Denote  $\alpha_k = \{x : |x - \sigma(x)| > \frac{1}{k}\} \cap B$ . Obviously  $\bigcup_{k=1}^{\infty} \alpha_k = \{x : |x - \sigma(x)| > 0\} \cap B = B$  and the latter set has positive measure by assumption; hence there exists  $k_0$  for which  $\mu(\alpha_{k_0}) > 0$ . Now

$$\alpha_{k_0} = \bigcup_{n=0}^{2k_0-1} \left( \alpha_{k_0} \cap \left[ \frac{n}{2k_0}, \frac{n+1}{2k_0} \right] \right),$$

so there exists  $n_0$  such that if we denote  $\beta' = \alpha_{k_0} \cap \left[ \frac{n_0}{2k_0}, \frac{n_0+1}{2k_0} \right]$ , then  $\mu(\beta') > 0$ . From the way we defined  $\beta'$  it is evident that  $\beta' \cap \sigma(\beta') = \emptyset$  because  $\text{diam}(\beta') < \frac{1}{2k_0}$  and  $|x - \sigma(x)| > \frac{1}{k_0}$  for every  $x \in \beta'$ . Using the proof of [9, Lemma 5.1] we find a measurable set  $\beta \subset \beta'$  such that  $T|_{L_1(\beta)}$  is an isomorphism. It is clear that  $d(L_1(\beta), TL_1(\beta)) = 1$  since  $L_1(\beta)$  and  $TL_1(\beta)$  have disjoint supports. The fact that  $L_1(\beta) + TL_1(\beta)$  is complemented in  $L_1$  follows from the facts that  $L_1(\beta)$  is norm-one complemented,  $T$  is an isomorphism (hence  $TL_1(\beta)$  is complemented), and  $L_1(\beta)$  and  $TL_1(\beta)$  have disjoint supports.

*Case 2.*  $\mu(B) = 0$ . In this case we have  $\mu(A) > 0$  (otherwise  $T$  will be a zero atom). There are two subcases:

- If  $a(x) \neq \text{const}$  a.e. on  $A$ .  
Then we proceed as in the case  $\sigma = id$  a.e. on  $(0, 1)$ , but we consider  $A$  instead of  $(0, 1)$ .
- If  $a(x) = \text{const} = \lambda$  a.e. on  $A$ .  
Note that  $\lambda \neq 0$  since otherwise  $T$  will be a zero atom. Then again we proceed as in the case  $\sigma = id$  a.e. on  $(0, 1)$  considering  $\lambda_1 = \lambda$  and  $\lambda_2 = 0$ . We can do this since  $\mu(A') > 0$  and  $\mu(B) = 0$ ; hence  $\mu(A' \setminus B) > 0$ , and  $Tf(x) = a(x)f(\sigma_1 x)$ , where

$$\sigma_1(x) = \begin{cases} \sigma(x) & \text{if } x \notin A' \setminus B, \\ x & \text{if } x \in A' \setminus B. \end{cases}$$

□

*Remark 4.6.* Note that Proposition 4.5 is also valid when considering operators  $T : L_1(\nu_1) \rightarrow L_1(\nu_2)$ , where  $\nu_1$  and  $\nu_2$  are two nonatomic measures on some  $\sigma$ -subalgebra on  $(0, 1)$  and this is in fact how we are going to use it.

Now we proceed to the proof of Theorem 1.2 in the case  $p = 1$ .

*Proof of Theorem 1.2 in the case  $L_1$ .* First, using Lemma 3.15, we find  $\varepsilon > 0$  and a subspace  $X \subseteq L_1$  such that for every  $X' \subseteq X$ ,  $X' \simeq L_1$ , and every  $\lambda \in \mathbb{C}$  we have  $g(\lambda I - T, X') > \varepsilon$ . Then consider an onto isomorphism  $S : L_1 \rightarrow L_1$  such that  $SX = L_1(\Delta)$ , where  $\Delta$  can be any nonempty open interval. For  $T' = STS^{-1}$  we have  $g(\lambda I - T', X') > \varepsilon'$  for every  $X' \subseteq SX = L_1(\Delta)$ , where  $\varepsilon' = \frac{\varepsilon}{\|S\| \|S^{-1}\|}$ . It is clear that it is enough to prove the theorem for  $T'$ , so without loss of generality we may assume that  $T' = T$  and  $\varepsilon = \varepsilon'$ .

Let  $T = T_a + T_c$  be the Kalton representation for  $T$  and fix  $\delta$ . First, we use Rosenthal's remark before Lemma 2.1 in [12] to find an atom  $V$  and set  $\Delta_1 \subseteq \Delta$  such that  $\|T_a|_{L_1(\Delta_1)} - V\| < \frac{\varepsilon}{10}$ . Since purely continuous operators are not sign-embeddings, we apply [12, Lemma 3.1] to find a norm-one complemented subspace  $X' \subseteq L_1(\Delta_1)$  such that  $X' \equiv L_1$  and  $\|T_c|_{X'}\| < \frac{\varepsilon}{10}$ . From our choice of  $X'$  it follows that

$$g(\lambda I - V, X'') > \frac{\varepsilon}{2} \text{ for every } \lambda \in \mathbb{C} \text{ and every } X'' \subseteq X', X'' \simeq L_1.$$

From the last inequality it is clear that  $V : X' \rightarrow L_1$  is a nonzero atom and  $V|_{X'} \neq \lambda I$ . Now Proposition 4.5 gives us the desired result. □

## 5. APPENDIX

Before we start with the proof of Theorem 3.4 we recall some of the notation we previously used and note some of the properties of  $S(x)$ , the square function defined with respect to the Haar basis (for a general definition of  $S(x)$ , see Definition 3.2).

Recall that unless otherwise noted,  $L_p$  denotes  $L_p([0, 1], \mu)$  where  $\mu$  is the Lebesgue measure. The unconditional basis constant of the usual Haar basis  $\{h_{n,i}\}_{n=0, i=1}^\infty$  in  $L_p$ ,  $1 < p < \infty$ , is denoted by  $C_p$ . Recall also that  $\{r_n\}_{n=0}^\infty$  is the Rademacher sequence on  $[0, 1]$  (defined by  $r_n = \sum_{i=1}^{2^n} h_{n,i}$ ).

Denote by  $\mathcal{E}_n$  the finite algebra generated by the dyadic intervals  $[(i-1)2^{-n}, i2^{-n}]$ ,  $i = 1, 2, \dots, 2^n$ , and by  $\mathcal{E}$ , the union of all these algebras. It is clear that the algebra  $\mathcal{E}_n$  is generated by the supports of  $\{(h_{n,i})\}_{i=1}^{2^n}$ .

If  $f$  and  $g$  are functions in  $L_p$  which have disjoint supports with respect to the Haar basis, then it is obvious that  $S^2(f + g) = S^2(f) + S^2(g)$ . This will be used numerous times. Let  $\{x_k\}_{k=1}^\infty$  be a sequence of functions in  $L_p$ ,  $1 < p < \infty$ , which are disjointly supported with respect to the Haar basis. Using the unconditionality of the Haar basis and Khintchine's inequality we obtain

$$(14) \quad \begin{aligned} \left\| \sum_{k=1}^\infty x_k \right\|_p &\geq C_p^{-1} \left( \int_0^1 \left\| \sum_{k=1}^\infty r_k(u)x_k \right\|_p^p du \right)^{1/p} \\ &= C_p^{-1} \left\| \left( \int_0^1 \left| \sum_{k=1}^\infty r_k(u)x_k \right|^p du \right)^{1/p} \right\|_p \geq C_p^{-1} A_p \left\| \left( \sum_{k=1}^\infty |x_k|^2 \right)^{\frac{1}{2}} \right\|_p, \end{aligned}$$

where  $A_p$  is the constant from Khintchine's inequality. If  $x_k = \sum_i \alpha_{k,i} h_{n_{k,i}}$ ,  $k = 1, 2, \dots$  are disjointly supported vectors with respect to the Haar basis, using (14) for  $\{\alpha_{k,i} h_{n_{k,i}}\}_{k,i}$  we obtain

$$(15) \quad \left\| \sum_{k=1}^\infty x_k \right\|_p \geq C_p^{-1} A_p \left( \int_0^1 \left( \sum_{k=1}^\infty S^2(x_k) \right)^{\frac{p}{2}} \right)^{\frac{1}{p}}.$$

In a similar manner

$$(16) \quad \left\| \sum_{k=1}^\infty x_k \right\|_p \leq C_p B_p \left( \int_0^1 \left( \sum_{k=1}^\infty S^2(x_k) \right)^{\frac{p}{2}} \right)^{\frac{1}{p}}.$$

From the last two inequalities it follows that the norm  $\|\cdot\|_p = \|S(\cdot)\|_p$  is equivalent to the usual norm in  $L_p$ ,  $1 < p < \infty$ . Now we proceed to the main theorem of this section.

**Theorem 3.4.** *For each  $1 < p < 2$  there is a constant  $K_p$  such that if  $T$  is a sign-embedding operator from  $L_p[0, 1]$  into  $L_p[0, 1]$  (and in particular if it is an isomorphism), then there is a  $K_p$ -complemented subspace  $X$  of  $L_p[0, 1]$  which is  $K_p$ -isomorphic to  $L_p[0, 1]$  and such that  $T|_X$  is a  $K_p$ -isomorphism and  $T(X)$  is  $K_p$ -complemented in  $L_p$ .*

Moreover, if we consider  $L_p[0, 1]$  with the norm  $\|x\|_p = \|S(x)\|_p$  (with  $S$  being the square function with respect to the Haar system), then, for each  $\varepsilon > 0$ , there is a subspace  $X$  of  $L_p[0, 1]$  which is  $(1 + \varepsilon)$ -isomorphic to  $L_p[0, 1]$  and such that  $T|_X$  is a  $(1 + \varepsilon)$ -isomorphism (and  $X$  and  $T(X)$  are  $K_p$ -complemented in  $L_p$ ).

*Proof of Theorem 3.4.* Let  $T$  be as in the statement of the theorem. Without loss of generality (see e.g. Lemma 9.10 in [8] and note that only the boundedness of  $T$  is used)  $\{Th_{n,i}\}$  is a block basis of  $\{h_{n,i}\}_{n=0, i=1}^\infty$ . For  $E \in \mathcal{E}_n$  put

$$v_n(E) = S \left( \sum_{h_{n,i} \subseteq E} Th_{n,i} \right),$$

where  $h_{n,i} \subseteq E$  is a shorthand notation for  $\text{supp}(h_{n,i}) \subseteq E$ . Also put  $v_n = v_n([0, 1]) = S(\sum_{i=1}^{2^n} Th_{n,i}) = S(Tr_n)$ .

*Claim 5.1.* The convex hull of  $\{v_n^2\}$  is  $p/2$ -equi-integrable; i.e., the set

$$V = \left\{ \left( \sum \alpha_n^2 v_n^2 \right)^{p/2} : \sum \alpha_n^2 \leq 1 \right\}$$

is equi-integrable.

*Proof.* The proof is a refinement of the argument on page 265 of [8]. Since the convex hull of any finite set in  $L_{p/2}$  is  $p/2$ -equi-integrable, it follows that if the convex hull of  $\{v_n^2\}$  is not  $p/2$ -equi-integrable, then there are  $\varepsilon_0 > 0$ , successive subsets  $\sigma_m \subset \mathbb{N}$ , and disjoint subsets  $\{A_m\}_{m=1}^\infty$  of  $[0, 1]$  such that for  $w_m^2 = \sum_{n \in \sigma_m} \alpha_n^2 v_n^2$ , where  $\sum_{n \in \sigma_m} \alpha_n^2 = 1$  for all  $m$ , we have

$$(17) \quad \left( \int_{A_m} w_m^p \right)^{1/p} \geq \varepsilon_0.$$

Using (15) and (17) (the estimate in (15) we can use since  $\{Tr_n\}$  are disjointly supported with respect to the Haar basis), for all  $\{a_m\}_{m=1}^\infty \in \ell_2$  we have

$$\begin{aligned} \left( \sum_{m=1}^\infty a_m^2 \right)^{1/2} &= \left\| \sum_{m=1}^\infty a_m \sum_{n \in \sigma_m} \alpha_n r_n \right\|_2 \geq \left\| \sum_{m=1}^\infty a_m \sum_{n \in \sigma_m} \alpha_n r_n \right\|_p \\ &\geq \|T\|^{-1} \left\| \sum_{m=1}^\infty a_m \sum_{n \in \sigma_m} \alpha_n Tr_n \right\|_p \\ &\geq \|T\|^{-1} C_p^{-1} A_p \left( \int_0^1 \left( \sum_{m=1}^\infty a_m^2 \sum_{n \in \sigma_m} \alpha_n^2 S^2(Tr_n) \right)^{p/2} \right)^{1/p} \\ &= \|T\|^{-1} C_p^{-1} A_p \left( \int_0^1 \left( \sum_{m=1}^\infty a_m^2 \sum_{n \in \sigma_m} \alpha_n^2 v_n^2 \right)^{p/2} \right)^{1/p} \\ &= \|T\|^{-1} C_p^{-1} A_p \left( \int_0^1 \left( \sum_{m=1}^\infty a_m^2 w_m^2 \right)^{p/2} \right)^{1/p} \\ &\geq \|T\|^{-1} C_p^{-1} A_p \left( \sum_{m=1}^\infty |a_m|^p \int_{A_m} w_m^p \right)^{1/p} \\ &\geq \|T\|^{-1} C_p^{-1} A_p \varepsilon_0 \left( \sum_{m=1}^\infty |a_m|^p \right)^{1/p}, \end{aligned}$$

which leads us to a contradiction since  $p < 2$ . □

**Proposition 5.2.** *There is an additive  $L_{p/2}^+$ -valued measure,  $\Lambda$ , on  $\mathcal{E}$  and there are successive convex combinations  $u_m(\cdot)$  of  $\{v_n^2(\cdot)\}$  such that for all  $E \in \mathcal{E}$  we have  $u_m(E) \rightarrow \Lambda(E)$  almost surely and in  $L_{p/2}$ . Moreover, for any sequence  $\varepsilon_n \rightarrow 0$ , there are measurable sets  $D_n \subset [0, 1]$  with  $\mu(D_n) > 1 - \varepsilon_n$  and such that for all  $E \in \mathcal{E}$  and all  $n$ ,  $u_m(E) \mathbf{1}_{D_n} \rightarrow \Lambda(E) \mathbf{1}_{D_n}$  as  $m \rightarrow \infty$  also in  $L_1$ .*

*Proof.* We start as in the proof of Lemma 6.4 in [8]: Since the set  $V$  from Claim 5.1 is bounded in  $L_{p/2}$ , by a result of Nikišin [11], for each  $\varepsilon > 0$  there exists a set

$D = D_\varepsilon \subset [0, 1]$  of measure larger than  $1 - \varepsilon$  such that

$$\sup_{v \in V} \int_D v d\mu < \infty.$$

Note that we may assume that  $D_{1/n} \subset D_{1/(n+1)}$  for  $n = 2, 3, \dots$ . As in the proof of [8, Lemma 6.4], using the weak compactness of  $V|_{D_{1/2}} \subset L_1$ , we can find successive convex combinations  $u_m(\cdot)$  of the  $v_n^2(\cdot)$  such that  $u_m(E)\mathbf{1}_{D_{1/2}}$  converges pointwise and in  $L_1$  to  $\Lambda_1(E)\mathbf{1}_{D_{1/2}}$  for every  $E \in \mathcal{E}$ , where  $\Lambda_1\mathbf{1}_{D_{1/2}}$  is  $L_1^+$ -valued additive measure. Now we can find successive convex combinations  $w_m(\cdot)$  of the  $u_m(\cdot)$  such that  $w_m(E)\mathbf{1}_{D_{1/3}}$  converges pointwise and in  $L_1$  to  $\Lambda_2(E)\mathbf{1}_{D_{1/3}}$  for every  $E \in \mathcal{E}$ , where  $\Lambda_2\mathbf{1}_{D_{1/3}}$  is  $L_1^+$ -valued additive measure. Note that  $\Lambda_2\mathbf{1}_{D_{1/2}} = \Lambda_1\mathbf{1}_{D_{1/2}}$ . Continuing in this manner and taking a diagonal sequence of the sequences of successive convex combinations we get a sequence, which we still denote  $u_m$ , of successive convex combinations of the  $v_n^2$  and an  $L_0^+$ -valued additive measure  $\Lambda$  such that for every  $n$ ,  $\Lambda\mathbf{1}_{D_{1/n}}$  is  $L_1^+$ -valued and  $u_m(E)\mathbf{1}_{D_{1/n}}$  converges, as  $m \rightarrow \infty$ , pointwise and in  $L_1$  to  $\Lambda(E)\mathbf{1}_{D_{1/n}}$  for every  $E \in \mathcal{E}$ .

It remains to show that the convergence is also in  $L_{p/2}$  (on the whole interval). Since for each  $E$ ,  $\{u_m(E)\}$  is  $p/2$ -equi-integrable, it follows that, given any  $\delta > 0$ , if  $n$  is large enough  $\int_{D_n^c} u_m(E)^{p/2} < \delta$  for all  $m$ . Consequently, we also have  $\int_{D_n^c} \Lambda(E)^{p/2} \leq \delta$  and

$$\begin{aligned} \limsup_{m \rightarrow \infty} \int |u_m(E) - \Lambda(E)|^{p/2} &\leq \limsup_{m \rightarrow \infty} \int_{D_n} |u_m(E) - \Lambda(E)|^{p/2} + 2\delta \\ &\leq \limsup_{m \rightarrow \infty} \|(u_m(E) - \Lambda(E))\mathbf{1}_{D_n}\|_1^{p/2} + 2\delta = 2\delta. \end{aligned}$$

Since this is true for any  $\delta$  we get the desired result.  $\square$

Note first that if we denote  $C = (C_p^2 B_p A_p^{-1} \|T\|)^p$ , then for all  $m$  and all  $E \in \mathcal{E}$  we have  $\int u_m(E)^{p/2} \leq C\mu(E)$ , where  $\{u_m\}$  are from Proposition 5.2. Indeed, let  $u_m = \sum_{k \in \sigma_m} \alpha_{m,k} v_k^2$ , where  $\{\sigma_m\}_{m=0}^\infty$  are successive subsets of  $\mathbb{N}$  and  $\{\alpha_{m,k}\}_{m=0, k \in \sigma_m}^\infty$  is a sequence of nonnegative numbers such that  $\sum_{k \in \sigma_m} \alpha_{m,k} = 1$ . Then using (15), (16) and the unconditionality of the Haar basis we get

$$\begin{aligned} \int u_m(E)^{p/2} &= \int \left( \sum_{k \in \sigma_m} \alpha_{m,k} v_k^2(E) \right)^{p/2} = \int \left( \sum_{k \in \sigma_m} \alpha_{m,k} S^2 \left( \sum_{h_{k,i} \subseteq E} T h_{k,i} \right) \right)^{p/2} \\ &\leq (C_p A_p^{-1})^p \left\| \sum_{k \in \sigma_m} \alpha_{m,k}^{1/2} \left( \sum_{h_{k,i} \subseteq E} T h_{k,i} \right) \right\|_p^p \\ &\leq (C_p A_p^{-1} \|T\|)^p \left\| \sum_{k \in \sigma_m} \alpha_{m,k}^{1/2} \left( \sum_{h_{k,i} \subseteq E} h_{k,i} \right) \right\|_p^p \\ &\leq (C_p^2 B_p A_p^{-1} \|T\|)^p \left\| \left( \sum_{k \in \sigma_m} \alpha_{m,k} S^2 \left( \sum_{h_{k,i} \subseteq E} h_{k,i} \right) \right)^{1/2} \right\|_p^p \\ &\leq (C_p^2 B_p A_p^{-1} \|T\|)^p \|\mathbf{1}_E\|_p^p = (C_p^2 B_p A_p^{-1} \|T\|)^p \mu(E). \end{aligned}$$

This implies that for all  $E \in \mathcal{E}$ ,  $\int \Lambda(E)^{p/2} \leq C\mu(E)$ . Now consider the linear operator  $R$  defined on the functions of the form  $f = \sum_{i=1}^r a_i \mathbf{1}_{E_i}$  by  $Rf = \sum_{i=1}^r a_i \Lambda(E_i)$ , where the  $E_i$ 's are disjoint sets in  $\mathcal{E}$ . Then  $R$  is bounded as an operator from a

subspace of  $L_{p/2}$  to  $L_{p/2}$ . Indeed,

$$\int |Rf|^{p/2} \leq \int \sum_{i=1}^r |a_i|^{p/2} \Lambda(E_i)^{p/2} \leq C \int \sum_{i=1}^r |a_i|^{p/2} \mu(E_i) = C \|f\|_{p/2}^{p/2}.$$

Consequently,  $R$  can be extended to all of  $L_{p/2}$  and then we define  $\Lambda(E) = R\mathbf{1}_E$  for all  $E$  in the Borel  $\sigma$ -algebra.

*Remark 5.3.* From the comments above it follows that  $\Lambda$  can be extended to an  $L_{p/2}^+$ -valued measure satisfying  $\int \Lambda(E)^{p/2} \leq C\mu(E)$  for some constant  $C < \infty$  and for all  $E$  in the Borel  $\sigma$ -algebra. For each  $n$ ,  $\Lambda(E)\mathbf{1}_{D_n}$  is an  $L_1^+$ -valued measure. Note also that, since  $T$  is a sign-embedding,  $\Lambda$  is not identically zero.

**Lemma 5.4.** *Let  $\Lambda$  be a nonzero  $L_{p/2}^+$ -valued measure on the Borel  $\sigma$ -algebra  $\mathcal{B}$  satisfying  $\int \Lambda(A)^{p/2} \leq C\mu(A)$  for some  $C < \infty$  and all  $A \in \mathcal{B}$ . Then for all  $\varepsilon > 0$  there exist a set  $A_0$  and a number  $c$ ,  $0 < c \leq C$ , such that  $\int \Lambda(A_0) > 0$  and*

$$c\mu(A) \leq \int \Lambda(A)^{p/2} \leq c(1 + \varepsilon)\mu(A)$$

for all  $A \subseteq A_0$ .

*Proof.* Fix an  $\varepsilon > 0$  and denote  $m = \sup\{\int \Lambda(A)^{p/2}/\mu(A) : A \in \mathcal{B}\}$ . Let  $B_0 \in \mathcal{B}$  be such that  $\frac{\int \Lambda(B_0)^{p/2}}{\mu(B_0)} \geq \frac{m}{1+\varepsilon}$ . Also let  $\mathcal{C}$  be a maximal collection of disjoint Borel subsets of  $B_0$  of positive measure satisfying  $\frac{\int \Lambda(B)^{p/2}}{\mu(B)} < \frac{m}{1+\varepsilon}$ . The collection  $\mathcal{C}$  is necessarily countable and if we assume that  $A_0 = B_0 \setminus \bigcup_{B \in \mathcal{C}} B$  has measure 0, then we have

$$\begin{aligned} \frac{m}{1+\varepsilon}\mu(B_0) &\leq \int \Lambda(B_0)^{p/2} = \int (\sum_{B \in \mathcal{C}} \Lambda(B))^{p/2} \\ &\leq \int \sum_{B \in \mathcal{C}} \Lambda(B)^{p/2} < \frac{m}{1+\varepsilon} \sum_{B \in \mathcal{C}} \mu(B) \\ &= \frac{m}{1+\varepsilon}\mu(B_0), \end{aligned}$$

which is a contradiction. Therefore,  $A_0$  satisfies the conclusion of the lemma with  $c = \frac{m}{1+\varepsilon}$ .  $\square$

**Lemma 5.5.** *Let  $\Lambda$  be an  $L_{p/2}^+$ -valued measure and suppose that  $A_0$  is such that for all  $A \subseteq A_0$  and some constant  $c > 0$ ,*

$$c\mu(A) \leq \int \Lambda(A)^{p/2} \leq c(1 + \varepsilon)\mu(A).$$

Then for any measurable partition  $A_0 = \bigcup_{i=1}^n F_i$ ,

$$\int \max_{1 \leq i \leq n} \Lambda(F_i)^{p/2} \geq (1 + \varepsilon)^{-p/(2-p)} c\mu(A_0) \geq (1 + \varepsilon)^{-2/(2-p)} \int \Lambda(A_0)^{p/2}.$$

*Proof.*

$$\begin{aligned} c\mu(A_0) &\leq \int \Lambda(A_0)^{p/2} = \int (\sum_{i=1}^n \Lambda(F_i))^{p/2} \\ &\leq \int (\sum_{i=1}^n \Lambda(F_i)^{p/2})^{p/2} \max_{1 \leq i \leq n} \Lambda(F_i)^{(1-p/2)p/2} \\ &\leq (\int \sum_{i=1}^n \Lambda(F_i)^{p/2})^{p/2} (\int \max_{1 \leq i \leq n} \Lambda(F_i)^{p/2})^{(1-p/2)} \\ &\leq (1 + \varepsilon)^{p/2} c^{p/2} (\mu(A_0))^{p/2} (\int \max_{1 \leq i \leq n} \Lambda(F_i)^{p/2})^{(1-p/2)}. \end{aligned}$$

Consequently,

$$\int \max_{1 \leq i \leq n} \Lambda(F_i)^{p/2} \geq (1 + \varepsilon)^{-p/(2-p)} c\mu(A_0) \geq (1 + \varepsilon)^{-2/(2-p)} \int \Lambda(A_0)^{p/2}.$$

$\square$

Fix an  $\varepsilon > 0$  and let  $A_0 \in \mathcal{B}$  and  $c$  be as in Lemma 5.5 so that for any partition  $A_0 = \bigcup_{i=1}^n F_i$ ,

$$(18) \quad \int \max_{1 \leq i \leq n} \Lambda(F_i)^{p/2} > (1 + \varepsilon)^{-p/(2-p)} c\mu(A_0).$$

Approximating by a set from  $\mathcal{E}$ , we may assume that the set  $A_0$  satisfying (18) is in  $\mathcal{E}$ . Let  $\{E_{n,i}\}_{n=0, i=1}^\infty$  be a dyadic tree of sets in  $\mathcal{E}$  with  $E_{0,1} = A_0$  and let

$$M_n = \max_{1 \leq i \leq 2^n} \Lambda(E_{n,i}).$$

$M_n$  is a nonincreasing sequence of functions in  $L_{p/2}^+$ . Denote by  $M$  its limit (in  $L_{p/2}$  or, equivalently, almost everywhere). Clearly,

$$\int M^{p/2} \geq (1 + \varepsilon)^{-p/(2-p)} c\mu(A_0).$$

We now define a sequence of functions  $\varphi_n : [0, 1] \rightarrow A_0$  which will play a role similar to the one played by the sequence with the same name in [8, Lemma 9.8]. For each  $n$  order the set  $\{1, 2, \dots, 2^n\}$  according to the order of the leftmost points in  $\{\overline{E_{n,i}}\}$ ; i.e.,  $i \prec j$  if  $\min\{t \in \overline{E_{n,i}}\} < \min\{t \in \overline{E_{n,j}}\}$ . Let  $\varphi_n : [0, 1] \rightarrow A_0$  be defined by  $\varphi_n(t) = \min\{t \in \overline{E_{n,i}}\}$  if  $1 \leq i \leq 2^n$  is the first, in the order  $\prec$ , such that  $\Lambda(E_{n,i})(t) \geq M(t)$ . For each  $t$ ,  $\{\varphi_n(t)\}$  is a nondecreasing and thus a converging sequence. Let  $\varphi(t)$  denote its limit. Notice that

$$\mathbf{1}_{\varphi^{-1}(A)}(t)M(t) \leq \Lambda(A)(t)$$

for every  $t$  and every  $A$  which is a union of the interiors of the  $E_{n,i}$ -s. Indeed, it is enough to prove this for  $A = E_{n,i}^\circ$  for some  $n$  and  $i$ . But if  $t \in \varphi^{-1}(E_{n,i}^\circ)$ , then for  $k$  large enough,  $E_{k,i(k)} \subset E_{n,i}$  (where  $\varphi_k(t)$  is the leftmost point of  $\overline{E_{k,i(k)}}$ ) and

$$\Lambda(E_{n,i})(t) \geq \Lambda(E_{k,i(k)})(t) \geq M(t).$$

Fix a sequence  $\{\varepsilon_n\}_{n=1}^\infty$ ,  $\varepsilon_n \rightarrow 0$ , to be chosen later. Consider the vector measure

$$m(A) = (\mu(A), \int_{\varphi^{-1}(A)} M^{p/2}), \quad A \subseteq A_0$$

and notice that it is nonatomic and even absolutely continuous with respect to Lebesgue measure. Indeed, for  $A$  in the algebra generated by the  $E_{n,i}$ 's,

$$\int_{\varphi^{-1}(A)} M^{p/2} \leq \int_{\varphi^{-1}(A)} \Lambda(A)^{p/2} \leq \int \Lambda(A)^{p/2} \leq C\mu(A).$$

The inequality clearly extends to all  $A \in \mathcal{B}$ ,  $A \subseteq A_0$ .

By Lyapunov's theorem one can find a partition of  $A_0$  into two sets,  $\tilde{F}_{1,1}$  and  $\tilde{F}_{1,2}$ , of equal  $m$  measure. For any  $\varepsilon_1 > 0$ , we can perturb  $\tilde{F}_{1,1}$  and  $\tilde{F}_{1,2}$  slightly to get  $F_{1,1}, F_{1,2}$  in the algebra generated by the  $E_{n,i}$ 's which satisfy

$$\mu(F_{1,1}) = \mu(F_{1,2}) = \frac{1}{2}\mu(A_0)$$

and

$$\frac{(1 - \varepsilon_1)}{2} \int M^{p/2} \leq \int_{\varphi^{-1}(F_{1,1})} M^{p/2}, \int_{\varphi^{-1}(F_{1,2})} M^{p/2} \leq \frac{(1 + \varepsilon_1)}{2} \int M^{p/2}.$$

Now we partition each of  $F_{1,1}$  and  $F_{1,2}$  in a similar manner and then continue the process. This way, for every positive sequence  $\{\varepsilon_n\}_{n=1}^\infty$ ,  $\varepsilon_n \downarrow 0$ , we construct a dyadic tree  $\{F_{n,i}\}_{n=0, i=1}^\infty$  of subsets of  $A_0$  such that the elements of the tree

$\{F_{n,i}\}_{n=0}^{\infty} \}_{i=1}^{2^n}$ , are in the algebra generated by the  $E_{n,i}$ 's and for all  $n = 0, 1, \dots$ ,  $i = 1, \dots, 2^n$ , we have

$$(19) \quad \mu(F_{n,i}) = 2^{-n} \mu(A_0)$$

and

$$(20) \quad 2^{-n} \prod_{j=1}^n (1 - \varepsilon_j) \int M^{p/2} \leq \int_{\varphi^{-1}(F_{n,i})} M^{p/2} \leq 2^{-n} \prod_{j=1}^n (1 + \varepsilon_j) \int M^{p/2}.$$

Define  $G_{n,i} = \varphi^{-1}(F_{n,i})$  for  $n = 0, 1, \dots$  and  $i = 1, 2, \dots, 2^n$ . Fix  $1 - \delta = (\prod_{j=1}^{\infty} (1 - \varepsilon_j))(1 + \varepsilon)^{-2/(2-p)}$ . The main remaining ingredient in the proof of the theorem is the following claim.

*Claim 5.6.* There exist  $a$  and  $b$ ,  $0 < a < b = (1 - \delta)^{-1}a$ , such that for all  $N$  and all coefficients  $\{a_{n,i}\}_{n=0}^N \}_{i=1}^{2^n}$ ,

$$a \|S(\sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i})\|_p^p \leq \| \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 \Lambda(F_{n,i}) \|_{p/2}^{p/2} \leq b \|S(\sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i})\|_p^p.$$

*Proof.* We shall use the shorthand notation  $h_{n,i} \supseteq h_{N,j}$  for  $\text{supp}(h_{n,i}) \supseteq \text{supp}(h_{N,j})$ . The first equality below follows by expressing  $\Lambda(F_{n,i})$  in terms of the  $\Lambda(F_{N,j})$ 's and changing the order of summation. We also use Lemma 5.4 and (19). Then

$$\begin{aligned} \left\| \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 \Lambda(F_{n,i}) \right\|_{p/2}^{p/2} &= \int \left( \sum_{j=1}^{2^N} \Lambda(F_{N,j}) \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} \\ &\leq \int \sum_{j=1}^{2^N} \Lambda(F_{N,j})^{p/2} \left( \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} \\ &\leq c(1 + \varepsilon) \mu(A_0) 2^{-N} \int \sum_{j=1}^{2^N} \left( \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} \\ &= c(1 + \varepsilon) \mu(A_0) \|S(\sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i})\|_p^p. \end{aligned}$$

For the other direction we use Lemma 5.5, (20), and the fact that  $\Lambda(F_{n,i}) \geq M$  on  $G_{n,i} = \varphi^{-1}(F_{n,i})$ . Then

$$\begin{aligned} \int \left( \sum_{j=1}^{2^N} \Lambda(F_{N,j}) \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} &\geq \int \left( \sum_{j=1}^{2^N} \mathbf{1}_{G_{N,j}} M \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} \\ &= \sum_{j=1}^{2^N} \int_{G_{N,j}} M^{p/2} \left( \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} \\ &\geq \left( \prod_{n=1}^N (1 - \varepsilon_n) \right) 2^{-N} \int M^{p/2} \sum_{j=1}^{2^N} \left( \sum_{(n,i); h_{n,i} \supseteq h_{N,j}} a_{n,i}^2 \right)^{p/2} \\ &\geq \left( \prod_{n=1}^N (1 - \varepsilon_n) \right) (1 + \varepsilon)^{\frac{-p}{2-p}} c \mu(A_0) \|S(\sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i})\|_p^p. \end{aligned}$$



It is clear that the claim follows with  $a = (\prod_{n=1}^\infty (1 - \varepsilon_n))(1 + \varepsilon)^{\frac{-p}{2-p}} c\mu(A_0)$ .  $\square$

Now we continue as in the proof of [8, Theorem 9.1, case  $1 < p < 2$ ]. From the fact that the  $\|\cdot\|_p$  is equivalent to the usual norm in  $L_p$ ,  $1 < p < \infty$ , it is enough to prove only the “moreover” part of Theorem 3.4. The fact that  $T(X)$  is  $K_p$ -complemented in  $L_p$  will follow from [8]. (The norm of the projection there depends only on the isomorphism constant and on  $p$ ; see Lemma 9.6 and the proof of Theorem 9.1 in the case  $1 < p < 2$  there.)

Let  $\{\beta_{m,j}\}_{m=0, j=1}^\infty, 2^m$  be a sequence of positive numbers such that  $\sum_{n,i} \beta_{n,i} = \delta$ . Since we obtained  $\Lambda$  as a limit of successive convex combinations of  $\{v_n^2(\cdot)\}$ , there exists a sequence of disjoint finite sets  $\{\sigma_{m,j}\}_{m=0, j=1}^\infty, 2^m \subseteq \mathbb{N}$  with  $\sigma_{m,j} > \inf\{l : F_{m,j} \in \mathcal{E}_l\}$  and a sequence of nonnegative numbers  $\{\alpha_n\}_{n=1}^\infty$  such that  $\sum_{n \in \sigma_{m,j}} \alpha_n = 1$ ,  $m = 0, 1, \dots$ ,  $j = 1, 2, \dots, 2^m$ , and

$$\int \left| \sum_{n \in \sigma_{m,j}} \alpha_n v_n^2(F_{m,j}) - \Lambda(F_{m,j}) \right|^{p/2} < \beta_{m,j} \int \Lambda(F_{m,j})^{p/2}$$

for all  $m = 0, 1, \dots$ ,  $j = 1, 2, \dots, 2^m$ . Put  $u_{m,j} = \sum_{n \in \sigma_{m,j}} \alpha_n v_n^2$ .

As in [8, Theorem 9.1], we define a Gaussian Haar system by

$$k_{m,j} = \sum_{n \in \sigma_{m,j}} \alpha_n^{1/2} \sum_{h_{n,i} \subseteq F_{m,j}} Th_{n,i}$$

for all  $m = 0, 1, \dots$ ,  $j = 1, 2, \dots, 2^m$ .

Set

$$X = \overline{\text{span}}\left\{ \sum_{n \in \sigma_{m,j}} \alpha_n^{1/2} \sum_{h_{n,i} \subseteq F_{m,j}} h_{n,i} \right\}_{m=0, j=1}^\infty, 2^m$$

and

$$Y = \overline{\text{span}}\{k_{m,j}\}_{m=0, j=1}^\infty, 2^m.$$

We first show that some multiple of the sequence  $\{k_{m,j}\}$  is almost isometrically equivalent to the Haar basis in the norm  $\|\cdot\|_p$ .

For all coefficients  $\{a_{n,i}\}_{n=0, i=1}^N, 2^n$  we have

$$\begin{aligned} (21) \quad & \left\| \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 (\Lambda(F_{n,i}) - u_{n,i}(F_{n,i})) \right\|_{p/2}^{p/2} = \int \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 (\Lambda(F_{n,i}) - u_{n,i}(F_{n,i})) \right)^{p/2} \\ & \leq \int \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^p |\Lambda(F_{n,i}) - u_{n,i}(F_{n,i})|^{p/2} \\ & \leq \int \sum_{n=0}^N \sum_{i=1}^{2^n} \beta_{n,i} a_{n,i}^p |\Lambda(F_{n,i})|^{p/2} \\ & \leq \delta \int \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 \Lambda(F_{n,i}) \right)^{p/2} \\ & = \delta \left\| \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 \Lambda(F_{n,i}) \right\|_{p/2}^{p/2} \end{aligned}$$

and using Claim 5.6 we immediately get

$$(22) \quad a(1-\delta) \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i} \right) \right\|_p^p \leq \left\| \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 u_{n,i}(F_{n,i}) \right\|_{p/2}^{p/2} \\ \leq b(1+\delta) \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i} \right) \right\|_p^p.$$

Since  $\{Th_{n,i}\}$  are disjointly supported with respect to the Haar basis, it follows that

$$S^2(k_{m,j}) = \sum_{n \in \sigma_{m,j}} \alpha_n S^2 \left( \sum_{h_{n,i} \subseteq F_{m,j}} Th_{n,i} \right) = \sum_{n \in \sigma_{m,j}} \alpha_n v_n^2(F_{m,j}) = u_{m,j}(F_{m,j})$$

and now using the fact that  $\{k_{n,i}\}$  are disjointly supported with respect to the Haar basis we get

$$(23) \quad S^2 \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} k_{n,i} \right) = \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 S^2(k_{n,i}) = \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i}^2 u_{n,i}(F_{n,i}).$$

Now we just have to observe that for any  $x \in L_p$  we have  $\|S(x)\|_p^p = \|S^2(x)\|_{p/2}^{p/2}$  and combining this with (22) and (23) gives us

$$(24) \quad a(1-\delta) \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i} \right) \right\|_p^p \leq \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} k_{n,i} \right) \right\|_p^p \\ \leq b(1+\delta) \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i} \right) \right\|_p^p.$$

The last estimate shows that some multiple of the sequence  $\{k_{n,i}\}$  is almost isometrically equivalent to the Haar basis with respect to  $\|\cdot\|_p$ . We must mention that (24) also implies that some multiple of  $T$  is almost an isometry on  $X$ . This follows from the fact that  $S^2(\sum_{n \in \sigma_{m,j}} \alpha_n^{1/2} \sum_{h_{n,i} \subseteq F_{m,j}} h_{n,i}) = \mathbf{1}_{F_{m,j}}$ ; hence

$$\mu(A_0) \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} h_{n,i} \right) \right\|_p^p = \left\| S \left( \sum_{n=0}^N \sum_{i=1}^{2^n} a_{n,i} \left( \sum_{m \in \sigma_{n,i}} \alpha_m^{1/2} \sum_{h_{m,j} \subseteq F_{n,i}} h_{m,j} \right) \right) \right\|_p^p.$$

□

## REFERENCES

- [1] C. Apostol, *Commutators on  $\ell_p$  spaces*, Rev. Roum. Math. Appl. **17** (1972), 1513–1534. MR0336432 (49:1206)
- [2] C. Apostol, *Commutators on  $c_0$ -spaces and on  $\ell_\infty$ -spaces*. Rev. Roum. Math. Pures Appl. **18** (1973), 1025–1032. MR0336433 (49:1207)
- [3] A. Brown, C. Pearcy, *Structure of commutators of operators*, Ann. of Math. (2) **82** (1965), 112–127. MR0178354 (31:2612)
- [4] D. Dosev, *Commutators on  $\ell_1$* , J. of Funct. Analysis **256** (2009), 3490–3509. MR2514050 (2010f:47067)
- [5] D. Dosev, W. B. Johnson, *Commutators on  $\ell_\infty$* , Bull. London Math. Soc. **42** (2010), 155–169. MR2586976 (2011d:47084)
- [6] P. Enflo, T. W. Starbird, *Subspaces of  $L_1$  containing  $L_1$* , Studia Math. **65** (1979), 203–225. MR557491 (81j:46034)
- [7] J. L. B. Gamlen, R. J. Gaudet, *On subsequences of the Haar system in  $L_p[0, 1]$ ,  $1 < p < \infty$* , Israel J. Math. **15** (1973), 404–413. MR0328575 (48:6917)

- [8] W. B. Johnson, B. Maurey, G. Schechtman, L. Tzafriri, *Symmetric structures in Banach spaces*. Mem. Amer. Math. Soc. **19** (1979), no. 217, v+298 pp. MR527010 (82j:46025)
- [9] N. J. Kalton, *The Endomorphisms of  $L_p$*  ( $0 \leq p \leq 1$ ), Indiana Univ. Math. J. **27** (1978), no. 3, 353–381. MR0470670 (57:10416)
- [10] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces. I*. Ergebnisse der Mathematik und ihrer Grenzgebiete, **92**, Springer-Verlag, Berlin-New York, 1977. MR0500056 (58:17766)
- [11] E. M. Nikišin, *Resonance theorems and superlinear operators* (Russian) Uspehi Mat. Nauk **25** (1970), no. 6(156), 129–191. MR0296584 (45:5643)
- [12] H. P. Rosenthal, *Embeddings of  $L_1$  in  $L_1$* , Contemp. Math. **26** (1984), 335–349. MR737412 (86k:46027)
- [13] M. Tarbard, *Hereditarily indecomposable, separable  $\mathcal{L}_\infty$  spaces with  $\ell_1$  dual having few operators, but not very few operators*, J. London Math. Soc. (2012) : jdr066v1-jdr066
- [14] A. Wintner, *The unboundedness of quantum-mechanical matrices*, Phys. Rev. **71** (1947), 738–739. MR0020724 (8:589c)

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, ISRAEL  
*E-mail address*: `dosevd@weizmann.ac.il`

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843  
*E-mail address*: `johnson@math.tamu.edu`

DEPARTMENT OF MATHEMATICS, WEIZMANN INSTITUTE OF SCIENCE, REHOVOT, ISRAEL  
*E-mail address*: `gideon@weizmann.ac.il`