

SOBOLEV EXTENSION BY LINEAR OPERATORS

CHARLES L. FEFFERMAN, ARIE ISRAEL, AND GARVING K. LULI

CONTENTS

1. Introduction	69
2. Notation and preliminaries	76
3. Plan for the proof	80
4. The inductive hypothesis	88
5. The CZ decomposition	92
6. Paths to keystone cubes I	94
7. Paths to keystone cubes II	103
8. Representatives	106
9. A partition of unity	109
10. Local extension operators	111
11. Extension near the keystone cubes	113
12. Estimates for auxiliary polynomials	117
13. Estimates for local extensions	117
14. The jet of a near optimal extension I	121
15. A constrained problem	127
16. The jet of a near optimal extension II	129
17. The extension operator	132
18. Proofs of the main theorems for finite E	135
19. Passage to infinite E	137
Acknowledgments	144
References	144

1. INTRODUCTION

Fix positive integers m, n and let f be a real-valued function defined on an (arbitrary) given subset $E \subset \mathbb{R}^n$. How can we tell whether f extends to a C^m function F on the whole \mathbb{R}^n ? If such an F exists, then how small can we take its C^m -norm? What can we say about the derivatives $\partial^\alpha F(x)$ at a given point x ? Can we take F to depend linearly on f ?

Received by the editors May 11, 2012 and, in revised form, November 12, 2012.

2010 *Mathematics Subject Classification*. Primary 42B99.

The first author was partially supported by NSF and ONR grants DMS 09-01040 and N00014-08-1-0678.

The second author was partially supported by NSF postdoctoral fellowship DMS-1103978.

The third author was partially supported by NSF and ONR grants DMS 09-01040 and N00014-08-1-0678.

Suppose E is finite. Can we compute an extension F whose C^m -norm has the least possible order of magnitude? How many computer operations does it take?

The above questions were answered in [8–12, 14, 15], building on earlier work of H. Whitney, G. Glaeser, Y. Brudnyi, P. Shvartsman, E. Bierstone, P. Milman and W. Pawłucki [2–4, 16, 25–27].

Now we want to address the analogous questions for Sobolev spaces in place of C^m . Important first steps were taken by P. Shvartsman [20–22] and A. Israel [17]; we discuss these papers later in the introduction.

To state our results, we set up notation. We work in the Sobolev space $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$ ($n < p < \infty$) with seminorm

$$\|F\|_{\mathbb{X}} = \max_{|\alpha|=m} \left(\int_{\mathbb{R}^n} |\partial^\alpha F(x)|^p dx \right)^{1/p}$$

or else in $\mathbb{X} = C^m(\mathbb{R}^n)$ with norm

$$\|F\|_{\mathbb{X}} = \max_{|\alpha| \leq m} \sup_{x \in \mathbb{R}^n} |\partial^\alpha F(x)|.$$

Given $E \subset \mathbb{R}^n$, we write $\mathbb{X}(E)$ for the space of restrictions to E of functions in \mathbb{X} . The space $\mathbb{X}(E)$ has a natural seminorm

$$\|f\|_{\mathbb{X}(E)} = \inf\{\|F\|_{\mathbb{X}} : F \in \mathbb{X}, F = f \text{ on } E\}.$$

When $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$, our standing assumption $n < p < \infty$ guarantees by the Sobolev theorem that any $F \in \mathbb{X}$ has continuous derivatives up to order $(m-1)$. Consequently, F may be restricted to E , and our definition of $\mathbb{X}(E)$ makes sense.

When $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$, then c, C, C' , etc. denote constants depending only on m, n, p . Similarly, when $\mathbb{X} = C^m(\mathbb{R}^n)$, then c, C, C' , etc. denote constants depending only on m, n . These symbols may denote different constants in different occurrences.

The simplest form of our results is as follows. Let $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$ until further notice.

Theorem 1 (Existence of linear extension operators). *Given $E \subset \mathbb{R}^n$, there exists a linear map $T: \mathbb{X}(E) \rightarrow \mathbb{X}$ such that*

$$Tf = f \text{ on } E \quad \text{and} \quad \|Tf\|_{\mathbb{X}} \leq C \|f\|_{\mathbb{X}(E)} \quad \text{for all } f \in \mathbb{X}(E).$$

Theorem 2 (Computation of the norm). *Suppose $E \subset \mathbb{R}^n$ is finite; let $N = \#(E)$ be the number of points in E . Then there exist linear functionals $\xi_1, \dots, \xi_L: \mathbb{X}(E) \rightarrow \mathbb{R}$ such that $L \leq CN$ and*

$$c \sum_{\ell=1}^L |\xi_\ell(f)|^p \leq \|f\|_{\mathbb{X}(E)}^p \leq C \sum_{\ell=1}^L |\xi_\ell(f)|^p \quad \text{for all } f \in \mathbb{X}(E).$$

For finite $E \subset \mathbb{R}^n$, we can say more about the linear map T in Theorem 1 and the linear functionals ξ_1, \dots, ξ_L in Theorem 2; they have “assisted bounded depth”, a notion that we explain below. We expect that this will be useful when we look for algorithms analogous to those in [14, 15].

To motivate the idea of assisted bounded depth, let us compute the variance of N given real numbers x_1, \dots, x_N . Two standard formulas are

$$(Var 1) \quad \sigma^2 = \frac{1}{2N^2} \sum_{i,j=1}^N (x_i - x_j)^2 \quad \text{and}$$

$$(Var 2) \quad \sigma^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \bar{x})^2, \quad \text{where} \quad \bar{x} = \frac{1}{N} \sum_{j=1}^N x_j.$$

Note that it takes $\sim N^2$ computer operations to apply formula (Var 1), but only $\sim N$ operations to apply (Var 2). By precomputing \bar{x} , we save a lot of work. We will later return to the problem of computing a variance.

Now we explain the notion of assisted bounded depth. Let $E \subset \mathbb{R}^n$ be finite, and let $N = \#(E)$. Any linear functional ω on the space $\mathbb{X}(E)$ can be written in the form

$$(1) \quad \omega(f) = \sum_{x \in E} \lambda(x) f(x), \quad \text{for coefficients } \lambda(x) \text{ independent of } f.$$

We write $\text{dp}(\omega)$ (the ‘‘depth’’ of ω) to denote the number of non-zero coefficients $\lambda(x)$. A collection of functionals $\Omega = \{\omega_1, \dots, \omega_S\}$ on $\mathbb{X}(E)$ will be said to be a set of ‘‘assists’’ provided we have

$$\sum_{s=1}^S \text{dp}(\omega_s) \leq CN.$$

The significance of this condition is that for any given $f \in \mathbb{X}(E)$, it takes at most CN computer operations to calculate all the values $\omega_1(f), \dots, \omega_S(f)$. The assists $\omega_s(f)$ ($s = 1, \dots, S$) will play the role of the mean \bar{x} from our earlier remarks on computing a variance.

Let $\Omega = \{\omega_1, \dots, \omega_S\}$ be a collection of assists for $\mathbb{X}(E)$. A linear functional $\xi : \mathbb{X}(E) \rightarrow \mathbb{R}$ is said to have ‘‘ Ω -assisted bounded depth’’ if it can be expressed in the form

$$(2) \quad \xi(f) = \sum_{y \in E} \lambda(y) f(y) + \sum_{s=1}^S \mu_s \omega_s(f), \quad \text{for all } f \in \mathbb{X}(E),$$

where the $\lambda(y), \mu_s \in \mathbb{R}$ are independent of f and where at most C of the coefficients $\lambda(y), \mu_s$ are non-zero.

Similarly, a linear map $T : \mathbb{X}(E) \rightarrow \mathbb{X}$ is said to have Ω -assisted bounded depth if it can be expressed in the form

$$(3) \quad Tf(x) = \sum_{y \in E} \lambda(x, y) f(y) + \sum_{s=1}^S \mu_s(x) \omega_s(f), \quad \text{for all } f \in \mathbb{X}(E), x \in \mathbb{R}^n,$$

where $\lambda(x, y), \mu_s(x) \in \mathbb{R}$ are independent of f and where for each $x \in \mathbb{R}^n$, at most C of the $\lambda(x, y), \mu_s(x)$ are non-zero.

An operator or functional is said to have ‘‘bounded depth’’ if it has Ω -assisted bounded depth with Ω equal to the empty set (no assists). See [11, 18].

We will prove the following result.

Theorem 3. *Suppose $E \subset \mathbb{R}^n$ is finite. Then there exist assists $\Omega = \{\omega_1, \dots, \omega_S\}$ such that the operator T in Theorem 1 and the functionals ξ_1, \dots, ξ_L in Theorem 2 can be taken to have Ω -assisted bounded depth.*

If an oracle told us the coefficients in (1) for the assists $\omega_1, \dots, \omega_S$, and in (3) for the operator T in Theorems 1 and 3, then we could efficiently compute Tf for any given $f \in \mathbb{X}(E)$. We could first precompute the assists $\omega_1(f), \dots, \omega_S(f)$ using at most CN computer operations by applying (1). After that we could compute $Tf(x)$ at any given point $x \in \mathbb{R}^n$ using at most C operations by applying (3).

Similarly, if an oracle told us the coefficients in (1) for the assists $\omega_1, \dots, \omega_S$, and in (2) for the functionals ξ_1, \dots, ξ_L in Theorems 2 and 3, then we could compute the order of magnitude of $\|f\|_{\mathbb{X}(E)}$ for any given f by first computing the assists $\omega_1(f), \dots, \omega_S(f)$ and then computing $\xi_1(f), \dots, \xi_L(f)$ in Theorems 2 and 3. This would require at most CN computer operations.

We hope that the coefficients arising in our formula for the assists $\omega_1, \dots, \omega_S$ and the functionals ξ_1, \dots, ξ_L can be computed using at most $CN \log N$ operations and that after the one-time work $CN \log N$ we can compute the coefficients relevant to $Tf(x)$ at a given query point x using at most $C \log N$ operations. This would provide an efficient algorithm for interpolation of data in the Sobolev space $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$ ($n < p < \infty$), analogous to the algorithms in Fefferman-Klartag [14, 15] for $C^m(\mathbb{R}^n)$.

Let us compare our present results to what we know about $C^m(\mathbb{R}^n)$. Switching over to $\mathbb{X} = C^m(\mathbb{R}^n)$, we recall the following results [10, 12, 14].

Theorem 4. *For any $E \subset \mathbb{R}^n$, there exists a linear map $T : \mathbb{X}(E) \rightarrow \mathbb{X}$ such that*

$$Tf = f \text{ on } E \text{ and } \|Tf\|_{\mathbb{X}} \leq C\|f\|_{\mathbb{X}(E)} \text{ for all } f \in \mathbb{X}(E).$$

Moreover, if E is finite, then T has bounded depth.

Theorem 5. *Let $E \subset \mathbb{R}^n$ be finite, and let $N = \#(E)$.*

Then there exist subsets $S_1, \dots, S_K \subset E$, with $K \leq CN$ and with $\#(S_k) \leq C$ for each k such that

$$\|f\|_{\mathbb{X}(E)} \leq C \cdot \max_{1 \leq k \leq K} \|(f|_{S_k})\|_{\mathbb{X}(S_k)} \text{ for all } f \in \mathbb{X}(E).$$

Corollary 1. *Let $E \subset \mathbb{R}^n$ be finite, and let $N = \#(E)$. Then there exist linear functionals $\xi_1, \dots, \xi_L : \mathbb{X}(E) \rightarrow \mathbb{R}$ such that $L \leq CN$, each ξ_ℓ has bounded depth, and*

$$c \cdot \max_{1 \leq \ell \leq L} |\xi_\ell(f)| \leq \|f\|_{\mathbb{X}(E)} \leq C \cdot \max_{1 \leq \ell \leq L} |\xi_\ell(f)| \text{ for all } f \in \mathbb{X}(E).$$

For the rest of the introduction, we again take $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$ ($n < p < \infty$). Motivated by Theorems 4, 5 and Corollary 1, one might wonder whether we can dispense with the assists in Theorem 3 and take T, ξ_1, \dots, ξ_L to have bounded depth. An optimist might even conjecture that $\|f\|_{\mathbb{X}(E)}^p$ is comparable to $\sum_{\ell=1}^L \lambda_\ell \cdot \|(f|_{S_\ell})\|_{\mathbb{X}(S_\ell)}^p$ for subsets $S_1, \dots, S_L \subset E$ and coefficients $\lambda_1, \dots, \lambda_L$ independent of f , with $L \leq CN$ and $\#(S_\ell) \leq C$ for each ℓ .

In fact, a counterexample [13] shows that the extension operator T in Theorem 3 cannot be taken to have bounded depth. On the other hand, the remarkable work of Batson-Spielman-Srivastava [1] on “sparsification” gives hope that bounded-depth ξ ’s may exist. We illustrate the result of [1] by returning to the computation of the variance of real numbers x_1, \dots, x_N .

Given $\epsilon > 0$, there exist coefficients $\lambda_1, \dots, \lambda_L > 0$ and integers $i_1, \dots, i_L, j_1, \dots, j_L \in \{1, \dots, N\}$ such that $L \leq C(\epsilon)N$ and such that the variance of x_1, \dots, x_N differs by at most a factor of $(1 + \epsilon)$ from

$$\sum_{\ell=1}^L \lambda_\ell (x_{i_\ell} - x_{j_\ell})^2,$$

for any real numbers x_1, \dots, x_N . Here, $C(\epsilon)$ depends only on ϵ . Thus, one can compute a variance to within, say, a one percent error by using $O(N)$ functionals without assists. The results in [1] sparsify an arbitrary sum of squares of linear forms, not just a variance. See also [23, 24].

Dealing with sums of p^{th} powers ($p > 2$) is more difficult than dealing with sums of squares. We don't know whether sparsification applies to our problems or whether the functionals ξ_1, \dots, ξ_L in Theorems 2 and 3 can be taken to have bounded depth.

Theorems 1, 2, and 3 deal with the homogeneous Sobolev space $L^{m,p}(\mathbb{R}^n)$. It is easy to pass from these results to analogous theorems for the inhomogeneous Sobolev space $W^{m,p}(\mathbb{R}^n)$; see section 18.

So far, we have looked for functions $F \in L^{m,p}(\mathbb{R}^n)$ that agree perfectly with a given $f : E \rightarrow \mathbb{R}$. More generally, we can look for functions F that agree approximately with a given f . To do so (for E finite), we specify a weight function $\mu : E \rightarrow (0, \infty)$ along with our usual $f : E \rightarrow \mathbb{R}$. We then look for $F \in L^{m,p}(\mathbb{R}^n)$ and $M \geq 0$ such that

$$\sum_{x \in E} \mu(x) |F(x) - f(x)|^p \leq M^p \quad \text{and} \quad \|F\|_{L^{m,p}(\mathbb{R}^n)} \leq M,$$

with M having the smallest possible order of magnitude.

We believe that Theorems 1, 2, 3, and the algorithms to which (we hope) they give rise can be extended to solve this more general problem. Analogous results for $C^m(\mathbb{R}^n)$ are given in [8, 12, 14, 15].

Let us sketch the proof of Theorem 1 for the case of finite sets E . The general case (infinite E) follows by taking a Banach limit, and the proofs of Theorems 2 and 3 arise by carefully examining our proof of Theorem 1. We will oversimplify the discussion so that the spirit of the ideas comes through.

We first set up a bit more notation. We will write Q, Q' , etc. to denote cubes in \mathbb{R}^n (with sides parallel to the coordinate axes). We write δ_Q to denote the sidelength of the cube Q . For any real number $A > 0$, we write AQ to denote the cube having the same center as Q and having sidelength $A \cdot \delta_Q$. If F is a C^{m-1} function on a neighborhood of a point $x \in \mathbb{R}^n$, then we write $J_x(F)$ (the “jet” of F at x) to denote the $(m-1)^{\text{st}}$ degree Taylor polynomial of F at x . Thus, $J_x(F)$ belongs to \mathcal{P} , the vector space of real $(m-1)^{\text{st}}$ degree polynomials on \mathbb{R}^n (“jets”).

Next, we introduce a local variant of the problem addressed by Theorem 1. Suppose we are given the following data:

- A cube Q .
- A finite set $E \subset 3Q$.
- A function $f : E \rightarrow \mathbb{R}$.
- A point $x \in E$.
- A jet $P \in \mathcal{P}$.

The local interpolation problem $LIP(Q, E, f, x, P)$ is the problem of finding a function $F \in \mathbb{X}(3Q)$ such that

$$(4) \quad F = f \text{ on } E, \quad J_x(F) = P,$$

and

$$(5) \quad \|F\|_{\mathbb{X}(3Q)} \text{ is as small as possible, up to a factor } C.$$

To prove Theorem 1, we will solve an arbitrary local interpolation problem $LIP(Q, E, f, x, P)$ in such a way that the interpolant F depends linearly on the data (f, P) for fixed Q, E, x .

Once we give such a linear solution, then Theorem 1 for finite E follows easily by taking Q to be a large enough cube containing E .

We will measure the difficulty of a local interpolation problem by assigning to it a “label” \mathcal{A} . Here, \mathcal{A} is any subset of the set \mathcal{M} of all multi-indices $\alpha = (\alpha_1, \dots, \alpha_n)$ of order $|\alpha| = \alpha_1 + \dots + \alpha_n \leq m - 1$. Labels \mathcal{A} come with a (total) order relation $<$. Roughly speaking, whenever $\mathcal{A} < \mathcal{A}'$, a local interpolation problem that carries the label \mathcal{A} is easier than one that carries the label \mathcal{A}' . In our ordering, the empty set \emptyset is maximal and the set \mathcal{M} is minimal. Accordingly, \emptyset labels the hardest interpolation problems and \mathcal{M} labels the easiest problems.

To assign a label \mathcal{A} to a local interpolation problem, we introduce the convex set

$$(6) \quad \sigma(x, E) = \{J_x(F) : F \in \mathbb{X}, \|F\|_{\mathbb{X}} \leq 1, F = 0 \text{ on } E\} \subset \mathcal{P}.$$

This set is clearly relevant to the problem of finding $F \in \mathbb{X}$ such that $F = f$ on E ; it measures our freedom of action in assigning $J_x(F)$ for such an interpolant.

Roughly speaking, a local interpolation problem $LIP(Q, E, f, x, P)$ carries a label $\mathcal{A} \subset \mathcal{M}$ if for each $\alpha \in \mathcal{A}$, the monomial

$$(7) \quad P_\alpha(y) = \delta_Q^{m-|\alpha|-n/p} (y-x)^\alpha \text{ belongs to } \sigma(x, E).$$

(Recall that δ_Q is the sidelength of Q and that $\mathbb{X} = L^{m,p}(\mathbb{R}^n)$.)

Thus, we have defined the notion of a local interpolation problem that carries a given label \mathcal{A} . We list a few relevant properties of labels.

- (8) Any problem that carries a label \mathcal{A} also carries the label \mathcal{A}' for any $\mathcal{A}' \subset \mathcal{A}$. If $\mathcal{A}' \subset \mathcal{A}$, then $\mathcal{A} < \mathcal{A}'$, where $<$ is the order relation used to rate the difficulty of a local interpolation problem.
- (9) Every local interpolation problem carries the label \emptyset , since we then make no requirement that any monomial P_α belongs to $\sigma(x, E)$.
- (10) On the other hand, a local interpolation problem carries the label \mathcal{M} only when E is the empty set.
- (11) The assignment of a label to $LIP(Q, E, f, x, P)$ depends only on (Q, E, x) ; the function f and the jet P play no role.

For each label \mathcal{A} , we will prove the following

Main Lemma for \mathcal{A} . *Any local interpolation problem $LIP(Q, E, f, x, P)$ that carries the label \mathcal{A} has a solution F that depends linearly on the data (f, P) .*

In particular, the MAIN LEMMA FOR $\mathcal{A} = \emptyset$ tells us that every local interpolation problem admits a solution F depending linearly on the data (f, P) . (See (9).) Consequently, the proof of Theorem 1 reduces to the task of proving the MAIN

LEMMA for every label \mathcal{A} . We proceed by induction on \mathcal{A} with respect to the order $<$.

In the BASE CASE, we take $\mathcal{A} = \mathcal{M}$, the smallest possible label under $<$. The MAIN LEMMA FOR \mathcal{M} holds trivially. For any local interpolation problem $LIP(Q, E, f, x, P)$ with label \mathcal{M} , we simply take our interpolant to be $F = P$. We have $F = f$ on E vacuously, since E is empty, and have $\|F\|_{\mathbb{X}} = 0$. (See (10).)

For the INDUCTION STEP, we fix a label $\mathcal{A} \neq \mathcal{M}$, and make the following inductive assumption:

$$(12) \quad \text{The MAIN LEMMA FOR } \mathcal{A}' \text{ holds for all } \mathcal{A}' < \mathcal{A}.$$

Under this assumption, we will prove the MAIN LEMMA FOR \mathcal{A} .

Thus, let

$$(13) \quad \begin{array}{l} LIP^\circ = LIP(Q^\circ, E^\circ, f^\circ, x^\circ, P^\circ) \text{ be a local interpolation problem} \\ \text{that carries the label } \mathcal{A}. \end{array}$$

Our task is to find an interpolant $F \in \mathbb{X}(3Q^\circ)$ that solves the interpolation problem LIP° and depends linearly on the data (f°, P°) . To do so, we proceed in several steps, as follows.

Step 1. We partition $3Q^\circ$ into ‘‘Calderón-Zygmund’’ subcubes Q_ν ($\nu = 1, \dots, \nu_{\max}$). Each Q_ν is such that $E_\nu := E \cap 3Q_\nu$ is non-empty. We pick a point $x_\nu \in E_\nu$ for each ν .

Step 2. For each ν , we pick a jet P_ν , depending linearly on the data (f°, P°) . We then set $f_\nu := f|_{E_\nu}$ and consider the local interpolation problem

$$(14) \quad LIP_\nu = LIP(Q_\nu, E_\nu, f_\nu, x_\nu, P_\nu).$$

Step 3. The partition of $3Q^\circ$ into Calderón-Zygmund cubes Q_ν has been defined to guarantee that each local interpolation problem LIP_ν carries a label $\mathcal{A}' < \mathcal{A}$ (except in the trivial case in which E_ν contains at most one point). Hence, by our inductive assumption (12), we can solve each problem LIP_ν , obtaining an interpolant $F_\nu \in \mathbb{X}(3Q_\nu)$ that depends linearly on (f_ν, P_ν) . Since f_ν and P_ν also depend linearly on (f°, P°) (see STEP 2), it follows that each F_ν depends linearly on (f°, P°) .

Step 4. We introduce a partition of unity,

$$1 = \sum_{\nu} \theta_{\nu} \quad \text{on } 3Q^\circ, \text{ with each } \theta_{\nu} \text{ supported in } 3Q_{\nu}.$$

Using that partition of unity, we patch together the local interpolants F_ν into a single interpolant F by taking

$$F = \sum_{\nu} \theta_{\nu} F_{\nu}.$$

Thus, we obtain a function $F \in \mathbb{X}(3Q^\circ)$ that satisfies $F = f^\circ$ on E° , and $J_{x^\circ}(F) = P^\circ$. It remains to show that the seminorm of F in $\mathbb{X}(3Q^\circ)$ is as small as possible up to a factor C .

This has a chance only if the jets P_ν in STEP 2 are carefully picked to be mutually consistent. Arranging this consistency is the hardest part of our proof and is the main difference between our arguments here and those in [8–11] for $C^m(\mathbb{R}^n)$ interpolation.

To assign the jets P_ν in STEP 2 and achieve their needed mutual consistency, we first pick out from among the Calderón-Zygmund cubes Q_ν the subcollection of keystone cubes. A given Q_ν is a keystone cube if every $Q_{\nu'}$ that meets $100Q_\nu$ is at least as big as Q_ν .

We carry out STEP 2 by first assigning jets P_ν to the keystone cubes. Each keystone cube may be treated separately, without worrying about mutual consistency. For any Calderón-Zygmund cube Q_ν other than a keystone cube, we carefully select a nearby keystone cube $Q_{\kappa(\nu)}$ and then simply set

$$P_\nu = P_{\kappa(\nu)}; \text{ the right-hand side has already been defined.}$$

The above procedure associates jets P_ν to all the Calderón-Zygmund cubes Q_ν in such a way that the required mutual consistency is guaranteed.

This concludes our sketch of the proof of Theorem 1. We again warn the reader that it is oversimplified. Even the notation and definitions in the subsequent sections differ from those presented in the introduction.

Throughout this paper, we study $L^{m,p}(\mathbb{R}^n)$ only for $n < p < \infty$. It would be natural to investigate the more general case $\frac{n}{m} < p < \infty$, since then any $F \in L^{m,p}(\mathbb{R}^n)$ is continuous and may therefore be restricted to an arbitrary subset $E \subset \mathbb{R}^n$.

We briefly review the earlier work on Sobolev extensions. The first breakthrough was the discovery by P. Shvartsman [20] that Theorem 1 for the Sobolev space $L^{1,p}(\mathbb{R}^n)$ (i.e., $m = 1$) holds with T given by the classical proof of the Whitney extension theorem [25]. Shvartsman gave a formula [20] for the order of magnitude of the seminorm $\|f\|_{\mathbb{X}(E)}$ when $\mathbb{X} = L^{1,p}(\mathbb{R}^n)$ and $E \subset \mathbb{R}^n$ is arbitrary (possibly infinite). See also [21]. The proof of Theorem 1 for the Sobolev space $L^{m,p}(\mathbb{R})$ (i.e., $n = 1$) was given by G.K. Luli [19] when $E \subset \mathbb{R}$ is finite and by P. Shvartsman [21] in the general case.

The next significant progress was the proof by A. Israel of Theorems 1 and 2 for the case $\mathbb{X} = L^{2,p}(\mathbb{R}^2)$ (with the bound $L \leq CN^2$ in place of $L \leq CN$ in Theorem 2). See [17]. The proof in [17] makes explicit use of the keystone cubes.

Our main technical achievement here is to combine the ideas used previously for interpolation problems with labels and those used to exploit keystone cubes. In particular, we have to dispense with the convex sets called $\Gamma(x, M)$ in [14, 15]; these sets played a crucial role in our earlier analysis of $C^m(\mathbb{R}^n)$.

P. Shvartsman has lectured at several workshops on results and ideas that appear closely related to ours. He is now writing up his results, and we look forward to studying them; see also [22]. It would be interesting to study the relationship between our keystone cubes and the “important” cubes from [22].

2. NOTATION AND PRELIMINARIES

2.1. Notation. Fix integers $m, n \geq 1$ and a real number $p > n$. Unless we say otherwise, constants written c, c', C, C' , etc. depend only on m, n , and p and are always positive. We refer to such constants as “universal constants”. The lowercase letters denote small (universal) constants, while the uppercase letters denote large (universal) constants.

The notation regarding constants is somewhat different in section 6. The reader may refer there for further details.

For non-negative quantities A, B , we write $A \simeq B$, $A \lesssim B$, or $A \gtrsim B$ to indicate (respectively) that $cB \leq A \leq CB$, $A \leq CB$, or $A \geq cB$.

This paper is divided into sections. The label (p.q) refers to formula (q) in section p. Within section p, we abbreviate (p.q) to (q).

A *cube* $Q \subset \mathbb{R}^n$ is any subset of the form

$$Q = \{a\} + (-\delta, \delta]^n \quad (a \in \mathbb{R}^n, \delta > 0).$$

The *sidelength* of Q is denoted $\delta_Q = 2\delta$, while the *center* of Q is the point a . For $A > 0$ let AQ denote the cube having the same center as Q but with sidelength $A\delta_Q$. A *dyadic* cube $Q \subset \mathbb{R}^n$ is any cube of the form

$$Q = (j_1 \cdot 2^k, (j_1 + 1) \cdot 2^k] \times (j_2 \cdot 2^k, (j_2 + 1) \cdot 2^k] \times \cdots \times (j_n \cdot 2^k, (j_n + 1) \cdot 2^k] \\ (j_1, j_2, \dots, j_n, k \in \mathbb{Z}).$$

To *bisect* a cube $Q \subset \mathbb{R}^n$ is to partition it into 2^n disjoint subcubes of sidelength $\frac{1}{2}\delta_Q$. These subcubes are called the *children* of Q . If $Q \subsetneq Q'$ are dyadic cubes, we say that Q' is an *ancestor* of Q . Every dyadic cube Q has a smallest ancestor called its *parent*, which we denote by Q^+ .

We use the following notation:

$$\begin{aligned} |x| &:= |x|_\infty = \max\{|x_1|, \dots, |x_n|\} & (x = (x_1, \dots, x_n) \in \mathbb{R}^n); \\ \text{dist}(x, \Omega) &:= \inf\{|x - y| : y \in \Omega\} & (\Omega \subset \mathbb{R}^n, x \in \mathbb{R}^n); \\ \text{dist}(\Omega', \Omega) &:= \inf\{|x - y| : x \in \Omega', y \in \Omega\} & (\Omega, \Omega' \subset \mathbb{R}^n); \\ B(\Omega, R) &:= \{x \in \mathbb{R}^n : \text{dist}(x, \Omega) \leq R\} & (\Omega \subset \mathbb{R}^n, R > 0); \text{ and} \\ \text{diam}(S) &:= \max\{|x - y| : x, y \in S\} & (S \subset \mathbb{R}^n \text{ finite}). \end{aligned}$$

The analogous objects defined with respect to the Euclidean norm $|x|_2 = (|x_1|^2 + \cdots + |x_n|^2)^{1/2}$ are denoted by $\text{dist}_2(x, \Omega)$, $\text{dist}_2(\Omega', \Omega)$, $B_2(\Omega, R)$ and $\text{diam}_2(S)$.

We write \mathcal{M} for the collection of all multi-indices α of order $|\alpha| \leq m - 1$. If α and β are multi-indices, then $\delta_{\alpha\beta}$ denotes the Kronecker delta.

If F is a C^{m-1} function on a neighborhood of a point $y \in \mathbb{R}^n$, then we write $J_y(F)$ (the ‘‘jet’’ of F at y) for the $(m - 1)^{\text{st}}$ degree Taylor polynomial of F at y .

Let $U \subset \mathbb{R}^n$ be a domain. The homogeneous Sobolev space $L^{m,p}(U)$ consists¹ of all functions $F : U \rightarrow \mathbb{R}$ with finite seminorm

$$\|F\|_{L^{m,p}(U)} := \max_{|\alpha|=m} \left(\int_U |\partial^\alpha F(x)|^p dx \right)^{1/p}.$$

Similarly, the inhomogenous Sobolev space $W^{m,p}(U)$ consists of all functions with finite norm

$$\|F\|_{W^{m,p}(U)} := \max_{|\alpha| \leq m} \left(\int_U |\partial^\alpha F(x)|^p dx \right)^{1/p}.$$

For $0 < s < 1$, the homogeneous Hölder space $\dot{C}^{m-1,s}(U)$ consists of all functions $F : U \rightarrow \mathbb{R}$ with finite seminorm

$$\|F\|_{\dot{C}^{m-1,s}(U)} := \max \left\{ \frac{|\partial^\alpha F(x) - \partial^\alpha F(y)|}{|x - y|^s} : x, y \in U, |\alpha| = m - 1 \right\}.$$

¹Here the derivatives $\partial^\alpha F$ are defined as distributions. Strictly speaking, $F \in L^{m,p}$ is defined only up to a set of measure zero. Under our assumption $n < p < \infty$, the Sobolev theorem allows us to regard any $F \in L^{m,p}$ as a function in $C_{loc}^{m-1, 1-n/p}$. For a discussion of these technicalities, see [7].

Likewise, the inhomogeneous Hölder space $C^{m-1,s}(U)$ consists of all functions with finite norm:

$$\|F\|_{C^{m-1,s}(U)} := \max \left\{ \frac{|\partial^\alpha F(x) - \partial^\alpha F(y)|}{|x-y|^s} : x, y \in U, |\alpha| = m-1 \right\} \\ + \max \{ |\partial^\alpha F(x)| : x \in U, |\alpha| \leq m-1 \}.$$

Let \mathcal{P} denote the space of real-valued $(m-1)^{\text{st}}$ degree polynomials on \mathbb{R}^n . Then \mathcal{P} is a vector space of dimension $D := \dim(\mathcal{P})$. For each $x \in \mathbb{R}^n$, the multiplication \odot_x on \mathcal{P} is defined by

$$P \odot_x Q := J_x(P \cdot Q) \quad (P, Q \in \mathcal{P}).$$

Given a finite subset $E' \subset \mathbb{R}^n$, the space of Whitney fields on E' is denoted

$$Wh(E') := \{ (P_x)_{x \in E'} : P_x \in \mathcal{P} \text{ for all } x \in E' \}.$$

For every function F that is C^{m-1} on a neighborhood of E' , we define the Whitney field $J_{E'}(F) := (J_x(F))_{x \in E'} \in Wh(E')$ (the “jet” of F on E').

Given an arbitrary subset $E \subset \mathbb{R}^n$ and a finite subset $E' \subset \mathbb{R}^n$, we define the trace space

$$L^{m,p}(E; E') := \left\{ (f, \vec{P}) : \vec{P} \in Wh(E'), f : E \rightarrow \mathbb{R}, \right. \\ \left. \exists F \in L^{m,p}(\mathbb{R}^n) \text{ with } F|_E = f \text{ and } J_{E'}(F) = \vec{P} \right\}.$$

This space comes equipped with the natural seminorm

$$\|(f, \vec{P})\|_{L^{m,p}(E; E')} := \inf \{ \|F\|_{L^{m,p}(\mathbb{R}^n)} : F|_E = f \text{ and } J_{E'}(F) = \vec{P} \}.$$

If E' is empty, then $L^{m,p}(E; E')$ is simply $L^{m,p}(E)$, and we take $\vec{P} \equiv 0$. Similarly, if E is empty, then $L^{m,p}(E; E') = Wh(E')$ (equipped with the obvious seminorm), and we take $f \equiv 0$. For $z \in \mathbb{R}^n$, we often write $L^{m,p}(E; z)$ in place of $L^{m,p}(E; \{z\})$.

A function $F \in L^{m,p}(\mathbb{R}^n)$ is called an *extension* of $(f, \vec{P}) \in L^{m,p}(E; E')$ if

$$F = f \text{ on } E \text{ and } J_{E'}(F) = \vec{P}.$$

By a *near optimal* extension of (f, \vec{P}) , we mean an extension that satisfies

$$\|F\|_{L^{m,p}(\mathbb{R}^n)} \leq C \|(f, \vec{P})\|_{L^{m,p}(E; E')} \text{ for some universal constant } C.$$

An *extension operator* is a linear map $T : L^{m,p}(E; E') \rightarrow L^{m,p}(\mathbb{R}^n)$ such that $T(f, \vec{P})$ is an extension of (f, \vec{P}) for each $(f, \vec{P}) \in L^{m,p}(E; E')$. We call T *bounded* provided that

$$\|T\| := \sup \left\{ \|T(f, \vec{P})\|_{L^{m,p}(\mathbb{R}^n)} : \|(f, \vec{P})\|_{L^{m,p}(E; E')} \leq 1 \right\} \leq C.$$

Let λ be a linear map from $L^{m,p}(E; E')$ into $V = \mathbb{R}$ or $V = \mathcal{P}$. The *depth* of λ (denoted $\text{dp}(\lambda)$) is the smallest non-negative integer d such that there exist points $x_1, \dots, x_r \in E$ and $y_1, \dots, y_s \in E'$ with $r + s = d$, multi-indices $\alpha_1, \dots, \alpha_s \in \mathcal{M}$, and elements $v_1, \dots, v_d \in V$ such that

$$\lambda(f, \vec{P}) = f(x_1) \cdot v_1 + \dots + f(x_r) \cdot v_r + \partial^{\alpha_1} P_{y_1}(y_1) \cdot v_{r+1} + \dots + \partial^{\alpha_s} P_{y_s}(y_s) \cdot v_{r+s} \\ \text{for all } (f, \vec{P}) \in L^{m,p}(E; E').$$

We now introduce the notion of linear maps with “assisted bounded depth”.

Definition 1. Let $V = \mathbb{R}$ or $V = \mathcal{P}$. Let a collection of linear functionals $\Omega \subset [L^{m,p}(E)]^*$ and an integer $d \geq 0$ be given. We say that a linear map $\lambda : L^{m,p}(E; E') \rightarrow V$ has Ω -assisted depth d if there exist $\omega_1, \dots, \omega_d \in \Omega$, elements $v_1, \dots, v_d \in V$, and a linear map $\tilde{\lambda} : L^{m,p}(E; E') \rightarrow V$ with $\text{dp}(\tilde{\lambda}) \leq d$ such that

$$\lambda(f, \vec{P}) = \omega_1(f) \cdot v_1 + \dots + \omega_d(f) \cdot v_d + \tilde{\lambda}(f, \vec{P}) \quad \text{for all } (f, \vec{P}) \in L^{m,p}(E; E').$$

Let $U \subset \mathbb{R}^n$ be a domain. We say that a linear map $T : L^{m,p}(E; E') \rightarrow L^{m,p}(U)$ has Ω -assisted depth d if the linear map

$$(f, \vec{P}) \in L^{m,p}(E; E') \mapsto J_x(T(f, \vec{P})) \in \mathcal{P} \quad \text{has } \Omega\text{-assisted depth } d, \text{ for all } x \in U.$$

If T (resp. λ) has Ω -assisted depth d for some constant d that depends only on m, n and p , we say that T (resp. λ) has Ω -assisted bounded depth.

For a finite subset $E \subset \mathbb{R}^n$ and E' empty, we have introduced the notion of an Ω -assisted bounded depth map $T : L^{m,p}(E) \rightarrow L^{m,p}(U)$ above and also in the introduction. It is not very difficult to see that these definitions are equivalent; we leave this as an exercise for the interested reader.

Let $E \subset \mathbb{R}^n$, and let $x \in \mathbb{R}^n$. Then we define

$$\sigma(x, E) := \{P \in \mathcal{P} : \exists \phi \in L^{m,p}(\mathbb{R}^n) \text{ with } J_x \phi = P, \phi|_E = 0, \text{ and } \|\phi\|_{L^{m,p}(\mathbb{R}^n)} \leq 1\}.$$

Note that $\sigma(x, E)$ is a symmetric ($P \in \mathcal{P} \implies -P \in \mathcal{P}$) convex subset of \mathcal{P} .

2.2. Preliminaries. For each $x \in \mathbb{R}^n$ and $\delta > 0$, we define a norm on \mathcal{P} by

$$(1) \quad |P|_{x,\delta} := \left(\sum_{|\alpha| \leq m-1} |\partial^\alpha P(x)|^p \delta^{n+(|\alpha|-m)p} \right)^{1/p} \quad (P \in \mathcal{P}).$$

For $z \in \mathbb{R}^n$, we have the Taylor expansion

$$\partial^\alpha P(x) = \sum_{|\gamma| \leq m-1-|\alpha|} \frac{1}{\gamma!} \partial^{\alpha+\gamma} P(z) \cdot (x-z)^\gamma \quad (|\alpha| \leq m-1).$$

Thus, the norms defined in (1) satisfy the inequality

$$(2) \quad |P|_{x,\delta} \leq C' |P|_{z,\delta} \quad (x, z \in \mathbb{R}^n, |x-z| \leq C\delta).$$

We will consider the following result in several settings.

The Sobolev Inequality for U : For every $F \in L^{m,p}(U)$, we have

$$(3) \quad |\partial^\alpha (J_y F - F)(x)| \leq C_U |x-y|^{m-|\alpha|-n/p} \|F\|_{L^{m,p}(U)} \quad (x, y \in U, |\alpha| \leq m-1).$$

Note that the norms in (1) are naturally related with the Sobolev Inequality. For instance, (3) can be seen (through (2)) to be equivalent to

$$|J_y F - J_x F|_{z,\delta} \leq C'_U \|F\|_{L^{m,p}(U)} \quad (x, y, z \in U, \delta \geq \text{Diameter}\{x, y, z\}).$$

For appropriate domains U , the constant C_U in (3) will be universal. The usual proof of the Sobolev Inequality establishes (3) when U is any cube. By applying a linear transformation, we then obtain (3) when U is a non-degenerate axis-parallel rectangular box (i.e., whose sidelengths differ by at most a universal constant factor).

Finally, we consider the case when $U = R_1 \cup R_2$ is the union of two such rectangular boxes; we assume that R_1 and R_2 have an interior point in common. Take $x, y \in U$ and fix some function $F \in L^{m,p}(U)$. If x and y lie in the same box R_i ,

then (3) follows from the Sobolev Inequality for R_i . Thus, we may assume that x and y lie in different boxes. Let $z \in R_1 \cap R_2$ be any point that satisfies

$$|x - z| \leq |x - y| \quad \text{and} \quad |y - z| \leq |x - y|.$$

(Here, we exploit the axis-parallel structure of R_1 and R_2 .) For any α with $|\alpha| \leq m - 1$, since $|\alpha| + n/p - m < 0$ we obtain

$$\begin{aligned} |\partial^\alpha (J_y F - J_x F)(z)| \cdot |x - y|^{|\alpha| + n/p - m} &\leq |\partial^\alpha (J_y F - J_z F)(z)| \cdot |y - z|^{|\alpha| + n/p - m} \\ &\quad + |\partial^\alpha (J_x F - J_z F)(z)| \cdot |x - z|^{|\alpha| + n/p - m}, \end{aligned}$$

which is bounded by $C' \|F\|_{L^{m,p}(R_1 \cup R_2)}$, thanks to the Sobolev Inequality for R_i ($i = 1, 2$). Inserting this inequality on the right-hand side of (2), where we have set $P = J_y F - J_x F$ and $\delta = |x - y|$, we obtain (3). This completes the proof of the Sobolev Inequality for $U = R_1 \cup R_2$.

3. PLAN FOR THE PROOF

Let $E \subset \mathbb{R}^n$ be finite, and let $z \in \mathbb{R}^n$. Our main goal is to prove the following theorem.

Extension Theorem for (E, z) . *There exist a linear map $T : L^{m,p}(E; z) \rightarrow L^{m,p}(\mathbb{R}^n)$, a map $M : L^{m,p}(E; z) \rightarrow \mathbb{R}_+$, and a collection of linear functionals $\Omega \subset [L^{m,p}(E)]^*$ that satisfy the following properties:*

(E1) *T is an extension operator.*

(E2) $\|(f, P)\|_{L^{m,p}(E; z)} \leq \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)} \leq C \cdot \|(f, P)\|_{L^{m,p}(E; z)}$, and

(E3) $C^{-1} \cdot M(f, P) \leq \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)} \leq C \cdot M(f, P)$,

for each $(f, P) \in L^{m,p}(E; z)$.

(E4) $\sum_{\omega \in \Omega} dp(\omega) \leq C \cdot \#(E)$.

(E5) *T has Ω -assisted bounded depth.*

(E6) *There exists a collection of linear functionals $\Xi \subset [L^{m,p}(E; z)]^*$ such that*

(a) *each functional in Ξ has Ω -assisted bounded depth,*

(b) $\#(\Xi) \leq C \cdot \#(E)$, and

(c) $M(f, P) = \left(\sum_{\xi \in \Xi} |\xi(f, P)|^p \right)^{1/p}$ for each $(f, P) \in L^{m,p}(E; z)$.

Here, C depends only on m , n and p .

Theorem 1 (for finite E) and Theorems 2, 3 follow easily from this extension theorem. (See section 18.)

3.1. Order relation on labels. To prove the Extension Theorem for (E, z) we proceed by induction on the “shape” of the convex subsets $\sigma(x, E) \subset \mathcal{P}$, where x ranges through E . The shape of a single convex subset $\sigma \subset \mathcal{P}$ will be defined in terms of a label $\mathcal{A} \subset \mathcal{M}$. We use the following order relation on labels.

Definition 2. Given distinct elements $\alpha = (\alpha_1, \dots, \alpha_n), \beta = (\beta_1, \dots, \beta_n) \in \mathcal{M}$, let $k \in \{1, \dots, n\}$ be maximal subject to the condition $\sum_{i=1}^k \alpha_i \neq \sum_{i=1}^k \beta_i$. We write $\alpha < \beta$ if

$$\sum_{i=1}^k \alpha_i < \sum_{i=1}^k \beta_i.$$

Given distinct labels $\mathcal{A}, \overline{\mathcal{A}} \subset \mathcal{M}$, we write $\mathcal{A} < \overline{\mathcal{A}}$ if the minimal element of the symmetric difference $\mathcal{A} \Delta \overline{\mathcal{A}}$ (under the order $<$ on elements) lies in \mathcal{A} .

The next lemma is immediate from the above definition.

Lemma 3.1. *The following properties hold:*

- If $\alpha, \beta \in \mathcal{M}$ and $|\alpha| < |\beta|$, then $\alpha < \beta$.
- If $\alpha + \gamma \in \mathcal{M}$ and $\beta + \gamma \in \mathcal{M}$, then $\alpha < \beta \iff \alpha + \gamma < \beta + \gamma$.
- If $\mathcal{A} \subsetneq \overline{\mathcal{A}} \subset \mathcal{M}$, then $\overline{\mathcal{A}} < \mathcal{A}$. In particular, the empty set is maximal and \mathcal{M} is minimal under the order on labels.

3.2. Polynomial bases. In this section, we introduce the notion of a labeled basis for a symmetric convex subset $\sigma \subset \mathcal{P}$.

Definition 3. Given $\mathcal{A} \subset \mathcal{M}$, $x \in \mathbb{R}^n$, $\epsilon > 0$ and $\delta > 0$, we say that $(P_\alpha)_{\alpha \in \mathcal{A}} \subset \mathcal{P}$ forms an $(\mathcal{A}, x, \epsilon, \delta)$ -basis for σ if

- (B1) $P_\alpha \in \epsilon \cdot \delta^{n/p+|\alpha|-m} \cdot \sigma$, for $\alpha \in \mathcal{A}$;
- (B2) $\partial^\beta P_\alpha(x) = \delta_{\alpha\beta}$, for $\alpha, \beta \in \mathcal{A}$; and
- (B3) $|\partial^\beta P_\alpha(x)| \leq \epsilon \cdot \delta^{|\alpha|-|\beta|}$, for $\alpha \in \mathcal{A}$, $\beta \in \mathcal{M}$ with $\beta > \alpha$.

Remark 1. The above definition is monotone in the parameters (ϵ, δ) in the following sense: Suppose that $(P_\alpha)_{\alpha \in \mathcal{A}}$ forms an $(\mathcal{A}, x, \epsilon, \delta)$ -basis for σ . Let $\epsilon' \geq \epsilon$ and $0 < \delta' \leq \delta$. Then $(P_\alpha)_{\alpha \in \mathcal{A}}$ forms an $(\mathcal{A}, x, \epsilon', \delta')$ -basis for σ .

Remark 2. The above definition provides a natural scale-invariant measure for the shape of a symmetric convex subset. Indeed, for $x \in \mathbb{R}^n$ and $\delta > 0$, consider the scaling map

$$(\tau_{x,\delta} F)(y) := \delta^{n/p-m} F(\delta(y-x) + x), \text{ defined for } F \in L^{m,p}(\mathbb{R}^n).$$

This scaling preserves the $L^{m,p}(\mathbb{R}^n)$ -norm. It is straightforward to verify that

(P_α) forms an $(\mathcal{A}, x, \epsilon, \delta)$ -basis for σ

$$\iff (\delta^{-|\alpha|} P_\alpha(\delta \cdot)) \text{ forms an } (\mathcal{A}, x, \epsilon, 1)\text{-basis for } \tau_{x,\delta} \sigma.$$

In the case when $\sigma = \sigma(x, E)$ for some $E \subset \mathbb{R}^n$, we have

$$\tau_{x,\delta} [\sigma(x, E)] = \sigma(x, \delta^{-1}(E-x) + x).$$

These comments imply that the hypotheses of the Main Lemma introduced in the next section are scale-invariant (see the first few paragraphs in section 4, where we reduce to the unit lengthscale).

3.3. The main lemma. Fix a collection of multi-indices $\mathcal{A} \subset \mathcal{M}$. By induction on \mathcal{A} , we prove the following:

Main Lemma for \mathcal{A} . *There exists a constant $\epsilon = \epsilon(\mathcal{A})$, depending only on \mathcal{A} , m , n and p , such that the following holds. Let $E \subset \mathbb{R}^n$ and $z \in \mathbb{R}^n$ satisfy $2 \leq \#(E \cup \{z\}) < \infty$, and assume that*

$$(1) \quad \begin{aligned} \sigma(x, E) \text{ contains an } (\mathcal{A}, x, \epsilon, \delta_{E,z})\text{-basis for every } x \in E, \\ \text{where } \delta_{E,z} := 10 \cdot \text{diam}(E \cup \{z\}). \end{aligned}$$

Then the Extension Theorem for (E, z) holds.

Note that (1) from the Main Lemma for \emptyset holds vacuously. Thus, the Main Lemma for $\mathcal{A} = \emptyset$ implies the Extension Theorem for (E, z) , whenever $\#(E \cup \{z\}) \geq 2$. The Extension Theorem for (E, z) obviously holds when $\#(E \cup \{z\}) \leq 1$ (see Lemma 3.2). Thus, we have reduced the proof of the Extension Theorem to the task of proving the Main Lemma for \mathcal{A} for every $\mathcal{A} \subset \mathcal{M}$.

We proceed by induction and establish the following.

Base Case: The Main Lemma for \mathcal{M} holds.

Induction Step: Let $\mathcal{A} \subset \mathcal{M}$ with $\mathcal{A} \neq \mathcal{M}$. Assume that the Main Lemma for \mathcal{A}' holds for each $\mathcal{A}' \subset \mathcal{A}$. Then the Main Lemma for \mathcal{A} holds.

To prove the Main Lemma for \mathcal{M} , we define $\epsilon(\mathcal{M}) = 1$. Let $(P_\alpha^x)_{\alpha \in \mathcal{M}}$ be an $(\mathcal{M}, x, 1, \delta)$ -basis for $\sigma(x, E)$. For each $x \in E$, by taking $\alpha = \beta = 0$ in **(B2)** we see that $P_0^x(x) = 1$. On the other hand, from **(B1)** we have $P_0^x(x) = 0$. This contradiction shows that E must be empty, hence we cannot have $\#(E \cup \{z\}) \geq 2$. Thus, the Main Lemma for \mathcal{M} holds vacuously. This proves the Base Case of the induction.

3.4. Small extension problems.

Lemma 3.2. *Suppose that $\#(E \cup \{z\}) \leq 2$. Then the Extension Theorem for (E, z) holds.*

Proof. If E is empty, then the Extension Theorem for (E, z) holds with (T, M, Ω, Ξ) defined by

$$T(P) = P \text{ and } M(P) = 0 \text{ for each } P \in \mathcal{P}; \quad \Omega = \emptyset \text{ and } \Xi = \emptyset.$$

Suppose that E is non-empty. Since the Extension Theorem for (E, z) is equivalent to the Extension Theorem for $(E \setminus \{z\}, z)$, it suffices to assume that $E = \{\hat{x}\}$ for some $\hat{x} \in \mathbb{R}^n$ and that $z \in \mathbb{R}^n \setminus E$.

Fix data $f : E \rightarrow \mathbb{R}$ and $P \in \mathcal{P}$. Define the polynomial $R \in \mathcal{P}$ by

$$(2) \quad R(\hat{x}) = f(\hat{x}), \text{ and } \partial^\alpha R(\hat{x}) = \partial^\alpha P(\hat{x}) \text{ for all } \alpha \in \mathcal{M} \setminus \{0\}.$$

(Note that R is determined uniquely and linearly from (f, P) .) Take a cutoff function $\theta \in C^\infty(\mathbb{R}^n)$ that satisfies

$$(3) \quad \theta \text{ is supported on } B(\hat{x}, |z - \hat{x}|/4);$$

$$(4) \quad \theta \equiv 1 \text{ in a neighborhood of } \hat{x}; \text{ and}$$

$$(5) \quad |\partial^\alpha \theta| \leq C|z - \hat{x}|^{-|\alpha|} \text{ for } |\alpha| \leq m.$$

(Note that θ necessarily vanishes in a neighborhood of the point z .) Define the linear map $T : L^{m,p}(E; z) \rightarrow L^{m,p}(\mathbb{R}^n)$ by

$$T(f, P) = P + \theta(R - P), \text{ for each } (f, P).$$

Then (2), (3) and (4) imply that T is an extension operator, and hence **(E1)** holds.

Since $T(f, P)$ agrees with an $(m-1)^{\text{st}}$ degree polynomial on $\mathbb{R}^n \setminus B(\hat{x}, |z - \hat{x}|/2)$, the Leibniz rule and (5) show that

$$\begin{aligned} \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}^p &= \|T(f, P)\|_{L^{m,p}(B(\hat{x}, |\hat{x}-z|/2))}^p \\ &\lesssim \sum_{|\alpha| \leq m} |\partial^\alpha (P - R)(\hat{x})|^p \cdot |z - \hat{x}|^{n+(|\alpha|-m)p} \\ (6) \qquad \qquad \qquad &\stackrel{(2)}{=} |P(\hat{x}) - f(\hat{x})|^p \cdot |z - \hat{x}|^{n-mp}. \end{aligned}$$

From the Sobolev Inequality, we have

$$(7) \quad |P(\hat{x}) - f(\hat{x})|^p \cdot |z - \hat{x}|^{n-mp} \leq C \cdot \|F\|_{L^{m,p}(\mathbb{R}^n)}^p,$$

for every $F \in L^{m,p}(\mathbb{R}^n)$ that satisfies $F(\hat{x}) = f(\hat{x})$ and $J_z(F) = P$.

Taking the infimum with respect to F in (7) yields

$$|P(\hat{x}) - f(\hat{x})|^p \cdot |z - \hat{x}|^{n-mp} \leq C \cdot \|(f, P)\|_{L^{m,p}(E;z)}^p.$$

Since T is an extension operator, the definition of the trace seminorm shows that

$$\|(f, P)\|_{L^{m,p}(E;z)} \leq \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}.$$

The last two inequalities and (6) suffice to prove **(E2)**. If we define $M(f, P) = |P(\hat{x}) - f(\hat{x})| \cdot |z - \hat{x}|^{n/p-m}$, then the same inequalities establish **(E3)**. Finally, **(E4)**, **(E5)**, and **(E6)** follow easily upon taking $\Omega = \emptyset$ and $\Xi = \{\xi\}$, where $\xi(f, P) = [P(\hat{x}) - f(\hat{x})] \cdot |z - \hat{x}|^{\frac{n}{p}-m}$. This concludes the proof of Lemma 3.2. \square

3.5. Technical lemmas. In this section we present two technical lemmas used in our proof of the Induction Step. The proofs involve nothing but elementary linear algebra, though they are somewhat involved (the reader may wish to omit the proofs on first reading). Recall that D stands for the dimension of the space of polynomials \mathcal{P} .

Lemma 3.3. *There exist universal constants $c_1 \in (0, 1)$ and $C_1 \geq 1$ so that the following holds. Suppose we are given the following data:*

- Real numbers $\epsilon_1 \in (0, c_1]$ and $\epsilon_2 \in (0, \epsilon_1^{2D+2}]$.
- A lengthscale $\delta > 0$.
- A collection of multi-indices $\mathcal{A} \subset \mathcal{M}$.
- Two finite subsets $\emptyset \neq E_1 \subset E_2 \subset \mathbb{R}^n$, with $\text{diam}(E_2) \leq 10\delta$.
- A family of polynomials $(\tilde{P}_\alpha^x)_{\alpha \in \mathcal{A}}$ that form an $(\mathcal{A}, x, \epsilon_2, \delta)$ -basis for $\sigma(x, E_1)$, for each $x \in E_2$, and satisfy

$$(8) \quad \max\{|\partial^\beta \tilde{P}_\alpha^x(x)| \delta^{|\beta|-|\alpha|} : x \in E_2, \alpha \in \mathcal{A}, \beta \in \mathcal{M}\} \geq \epsilon_1^{-D-1}.$$

Then there exists $\bar{\mathcal{A}} < \mathcal{A}$ so that $\sigma(x, E_1)$ contains an $(\bar{\mathcal{A}}, x, C_1 \cdot \epsilon_1, \delta)$ -basis for every $x \in E_2$.

Proof. By rescaling we may assume that $\delta = 1$. Let c_1 be a sufficiently small constant to be determined later. Our hypothesis tells us that $\epsilon_1 \leq c_1$, $\epsilon_2 \leq \epsilon_1^{2D+2}$ and that $(\tilde{P}_\alpha^x)_{\alpha \in \mathcal{A}}$ forms an $(\mathcal{A}, x, \epsilon_2, 1)$ -basis for $\sigma(x, E_1)$, for each $x \in E_2$. That

is,

$$(9) \quad \tilde{P}_\alpha^x \in \epsilon_2 \cdot \sigma(x, E_1);$$

$$(10) \quad \partial^\beta \tilde{P}_\alpha^x(x) = \delta_{\alpha\beta} \quad (\alpha, \beta \in \mathcal{A}); \text{ and}$$

$$(11) \quad |\partial^\beta \tilde{P}_\alpha^x(x)| \leq \epsilon_2 \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}, \beta > \alpha).$$

For each $\alpha \in \mathcal{A}$, we define $Z_\alpha = \max \left\{ |\partial^\beta \tilde{P}_\alpha^x(x)| : x \in E_2, \beta \in \mathcal{M} \right\}$. Then hypothesis (8) is equivalent to $\max \{ Z_\alpha : \alpha \in \mathcal{A} \} \geq \epsilon_1^{-D-1}$. Let $\bar{\alpha} \in \mathcal{A}$ be the minimal index with $Z_{\bar{\alpha}} \geq \epsilon_1^{-D-1}$. Thus,

$$(12) \quad Z_\alpha < \epsilon_1^{-D-1}, \text{ for all } \alpha \in \mathcal{A} \text{ with } \alpha < \bar{\alpha},$$

and there exist $x_0 \in E_2$ and $\beta_0 \in \mathcal{M}$ with

$$(13) \quad \epsilon_1^{-D-1} \leq Z_{\bar{\alpha}} = |\partial^{\beta_0} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)|.$$

Thus, (10) and (11) imply that $\beta_0 \neq \bar{\alpha}$ and $\beta_0 \leq \bar{\alpha}$, respectively. Therefore, $\beta_0 < \bar{\alpha}$. By definition of $Z_{\bar{\alpha}}$, we also have

$$(14) \quad |\partial^\beta \tilde{P}_{\bar{\alpha}}^y(y)| \leq |\partial^{\beta_0} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)|, \text{ for all } y \in E_2 \text{ and } \beta \in \mathcal{M}.$$

Let the elements of \mathcal{M} between β_0 and $\bar{\alpha}$ be ordered as follows:

$$\beta_0 < \beta_1 < \dots < \beta_k = \bar{\alpha}.$$

Note that $k+1 \leq \#\mathcal{M} = D$. Define

$$a_i = |\partial^{\beta_i} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)|, \text{ for } i = 0, \dots, k.$$

Then, (10) and (13) imply that $a_k = 1$ and $a_0 \geq \epsilon_1^{-D-1}$. Choose $r \in \{0, \dots, k\}$ with $a_r \epsilon_1^{-r} = \max \{ a_l \epsilon_1^{-l} : 0 \leq l \leq k \}$. Note that $a_0 \geq \epsilon_1^{-D-1} > a_k \epsilon_1^{-k}$, which implies $r \neq k$. Moreover, we have

$$(15) \quad a_r \geq \epsilon_1^D a_0 \text{ and } a_r \geq \epsilon_1^{-1} a_i \text{ for } i = r+1, \dots, k.$$

Define $\bar{\beta} = \beta_r \in \mathcal{M}$. Then, (15) states that

$$(16) \quad |\partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \geq \epsilon_1^D |\partial^{\beta_0} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \stackrel{(13)}{\geq} \epsilon_1^{-1}.$$

Also, (11) and (16) imply that

$$|\partial^\beta \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \leq \epsilon_2 \leq 1 \leq \epsilon_1 |\partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \quad (\beta \in \mathcal{M}, \beta > \bar{\alpha}).$$

Meanwhile, for $\bar{\beta} < \beta \leq \bar{\alpha}$, (15) implies that $|\partial^\beta \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \leq \epsilon_1 |\partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)|$. Thus, we have

$$(17) \quad |\partial^\beta \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \leq \epsilon_1 |\partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| \quad (\beta \in \mathcal{M}, \beta > \bar{\beta}).$$

By (16) we have $|\partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| > 1$. Hence, (10) and (11) imply that

$$(18) \quad \bar{\beta} < \bar{\alpha} \text{ and } \bar{\beta} \notin \mathcal{A}.$$

Define $\tilde{P}_{\bar{\beta}}^{x_0} = \tilde{P}_{\bar{\alpha}}^{x_0} / \partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)$, which satisfies

$$(19) \quad \tilde{P}_{\bar{\beta}}^{x_0} \in \epsilon_2 \cdot \sigma(x_0, E_1), \quad \text{thanks to (9) and } |\partial^{\bar{\beta}} \tilde{P}_{\bar{\alpha}}^{x_0}(x_0)| > 1;$$

$$(20) \quad |\partial^{\beta} \tilde{P}_{\bar{\beta}}^{x_0}(x_0)| \leq \epsilon_1 \quad (\beta \in \mathcal{M}, \beta > \bar{\beta}), \quad \text{thanks to (17);}$$

$$(21) \quad |\partial^{\beta} \tilde{P}_{\bar{\beta}}^{x_0}(x_0)| \leq \epsilon_1^{-D} \quad (\beta \in \mathcal{M}), \quad \text{thanks to (14) and (16); and}$$

$$(22) \quad \partial^{\bar{\beta}} \tilde{P}_{\bar{\beta}}^{x_0}(x_0) = 1.$$

By (19), there exists $\varphi_{\bar{\beta}} \in L^{m,p}(\mathbb{R}^n)$ with

$$(23) \quad J_{x_0} \varphi_{\bar{\beta}} = \tilde{P}_{\bar{\beta}}^{x_0};$$

$$(24) \quad \varphi_{\bar{\beta}}|_{E_1} = 0; \text{ and}$$

$$(25) \quad \|\varphi_{\bar{\beta}}\|_{L^{m,p}(\mathbb{R}^n)} \leq \epsilon_2.$$

Fix an arbitrary point $y \in E_2$ and define $\tilde{P}_{\bar{\beta}}^y := J_y \varphi_{\bar{\beta}}$. Then (24)-(25) and the definition of $\sigma(\cdot, \cdot)$ imply that

$$(26) \quad \tilde{P}_{\bar{\beta}}^y \in \epsilon_2 \cdot \sigma(y, E_1).$$

Since $\tilde{P}_{\bar{\beta}}^y = J_y \varphi_{\bar{\beta}}$, we have

$$(27) \quad |\partial^{\beta} \tilde{P}_{\bar{\beta}}^y(y) - \delta_{\beta \bar{\beta}}| \leq |\partial^{\beta} \varphi_{\bar{\beta}}(y) - \partial^{\beta} \tilde{P}_{\bar{\beta}}^{x_0}(y)| \\ + \left| \sum_{|\gamma| \leq m-1-|\beta|} \frac{1}{\gamma!} \partial^{\beta+\gamma} \tilde{P}_{\bar{\beta}}^{x_0}(x_0) \cdot (y-x_0)^{\gamma} - \delta_{\beta \bar{\beta}} \right| \quad (\beta \in \mathcal{M}).$$

Since $x_0, y \in E_2$ and $\text{diam}(E_2) \leq 10\delta = 10$, we have $|x_0 - y| \leq 10$. Therefore, the first term on the right-hand side of (27) is bounded by $C\|\varphi_{\bar{\beta}}\|_{L^{m,p}(\mathbb{R}^n)}$; this uses (23) and the Sobolev Inequality (2.3). In turn this is bounded by $C\epsilon_2$, thanks to (25). If $\beta > \bar{\beta}$, then $\beta + \gamma > \bar{\beta}$ for each multi-index γ with $|\gamma| \leq m-1-|\beta|$. In this case (20) and $|x_0 - y| \leq 10$ imply that the second term on the right-hand side of (27) is controlled by $C\epsilon_1$. If $\beta = \bar{\beta}$, then (22) implies that the second term from (27) equals

$$\left| \sum_{0 < |\gamma| \leq m-1-|\beta|} \frac{1}{\gamma!} \partial^{\beta+\gamma} \tilde{P}_{\bar{\beta}}^{x_0}(x_0) \cdot (y-x_0)^{\gamma} \right|,$$

which is bounded by $C\epsilon_1$, again by (20) and $|x_0 - y| \leq 10$. Thus we have argued that

$$(28) \quad |\partial^{\beta} \tilde{P}_{\bar{\beta}}^y(y) - \delta_{\beta \bar{\beta}}| \leq C\epsilon_1 \quad (\beta \in \mathcal{M}, \beta \geq \bar{\beta}).$$

Finally, if $\beta \in \mathcal{M}$, then (21) and $|x_0 - y| \leq 10$ imply that the second term on the right-hand side of (27) is controlled by $C\epsilon_1^{-D}$. Therefore,

$$(29) \quad |\partial^{\beta} \tilde{P}_{\bar{\beta}}^y(y)| \leq |\partial^{\beta} \tilde{P}_{\bar{\beta}}^y(y) - \delta_{\beta \bar{\beta}}| + 1 \\ \leq C\epsilon_1 + C\epsilon_1^{-D} + 1 \leq C'\epsilon_1^{-D} \quad (\beta \in \mathcal{M}).$$

Define $\overline{\mathcal{A}} = \{\alpha \in \mathcal{A} : \alpha < \overline{\beta}\} \cup \{\overline{\beta}\}$. Then (18) implies that the minimal element of $\mathcal{A} \Delta \overline{\mathcal{A}}$ is $\overline{\beta}$. Thus, we have $\overline{\mathcal{A}} < \mathcal{A}$. For each $\alpha \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}$, we have $\alpha < \overline{\beta}$; hence, (18) implies that $\alpha < \overline{\alpha}$. Thus, (12) and the definition of Z_α imply that

$$(30) \quad |\partial^\beta \tilde{P}_\alpha^y(y)| \leq \epsilon_1^{-D-1} \quad (\alpha \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}, \beta \in \mathcal{M}).$$

Define

$$P_\beta^y := \tilde{P}_\beta^y - \sum_{\gamma \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}} \partial^\gamma \tilde{P}_\beta^y(y) \tilde{P}_\gamma^y.$$

Notice that $\overline{\mathcal{A}} \setminus \{\overline{\beta}\} \subset \mathcal{A}$; thus (10) implies that

$$\partial^\alpha P_\beta^y(y) = \partial^\alpha \tilde{P}_\beta^y(y) - \sum_{\gamma \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}} \partial^\gamma \tilde{P}_\beta^y(y) \delta_{\alpha\gamma} = 0 \quad (\alpha \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}).$$

Also, (26), (29) and (9) imply that

$$P_\beta^y \in (\epsilon_2 + C\epsilon_1^{-D}\epsilon_2) \cdot \sigma(y, E_1) \subseteq C\epsilon_1^{-D}\epsilon_2 \cdot \sigma(y, E_1).$$

Since $\overline{\beta}$ is the maximal element of $\overline{\mathcal{A}}$, it follows that for any $\beta \geq \overline{\beta}$ and any $\gamma \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}$, we have $\beta > \gamma$. Thus, (28), (29) and (11) imply that

$$(31) \quad |\partial^\beta P_\beta^y(y) - \delta_{\beta\overline{\beta}}| \leq C(\epsilon_1 + \epsilon_1^{-D}\epsilon_2) \quad (\beta \in \mathcal{M}, \beta \geq \overline{\beta}).$$

Finally, (29) and (30) imply that

$$|\partial^\beta P_\beta^y(y)| \leq C\epsilon_1^{-2D-1} \quad (\beta \in \mathcal{M}).$$

Recall that $\epsilon_2 \leq \epsilon_1^{D+1}$ and $\epsilon_1 \leq c_1$. We now fix c_1 to be a small universal constant so that (31) yields $\partial^{\overline{\beta}} P_\beta^y(y) \in [1/2, 2]$. We then define $\hat{P}_\beta^y = P_\beta^y / \partial^{\overline{\beta}} P_\beta^y(y)$. The above four lines give that

$$(32) \quad \hat{P}_\beta^y \in C\epsilon_1^{-D}\epsilon_2 \cdot \sigma(y, E_1);$$

$$(33) \quad \partial^\beta(\hat{P}_\beta^y)(y) = \delta_{\beta\overline{\beta}} \quad (\beta \in \overline{\mathcal{A}});$$

$$(34) \quad |\partial^\beta \hat{P}_\beta^y(y)| \leq C(\epsilon_1 + \epsilon_1^{-D}\epsilon_2) \quad (\beta \in \mathcal{M}, \beta > \overline{\beta}); \text{ and}$$

$$(35) \quad |\partial^\beta \hat{P}_\beta^y(y)| \leq C\epsilon_1^{-2D-1} \quad (\beta \in \mathcal{M}).$$

For each $\alpha \in \overline{\mathcal{A}} \setminus \{\overline{\beta}\}$ we define $\hat{P}_\alpha^y = \tilde{P}_\alpha^y - \left[\partial^{\overline{\beta}}(\tilde{P}_\alpha^y)(y) \right] \hat{P}_\beta^y$. Note that $\alpha < \overline{\beta}$, and hence $|\partial^{\overline{\beta}}(\tilde{P}_\alpha^y)(y)| \leq \epsilon_2 \leq 1$, thanks to (11). From (9) and (32), we now obtain

$$(36) \quad \hat{P}_\alpha^y \in C'\epsilon_1^{-D}\epsilon_2 \cdot \sigma(y, E_1).$$

From (11) and (35), we have

$$(37) \quad \begin{aligned} \left| \partial^\beta(\hat{P}_\alpha^y)(y) \right| &\leq \left| \partial^\beta(\tilde{P}_\alpha^y)(y) \right| + \left| \partial^{\overline{\beta}}(\tilde{P}_\alpha^y)(y) \right| \cdot \left| \partial^\beta(\hat{P}_\beta^y)(y) \right| \leq \epsilon_2 + \epsilon_2 \cdot C\epsilon_1^{-2D-1} \\ &\leq C'\epsilon_1^{-2D-1}\epsilon_2 \quad (\beta \in \mathcal{M}, \beta > \alpha). \end{aligned}$$

Recall that $\bar{\mathcal{A}} = \{\alpha \in \mathcal{A} : \alpha < \bar{\beta}\} \cup \{\bar{\beta}\}$. From (10) and (33), we have

$$\begin{aligned}
 \partial^\beta \widehat{P}_\alpha^y &= \partial^\beta (\widetilde{P}_\alpha^y)(y) - \left[\partial^{\bar{\beta}} (\widetilde{P}_\alpha^y)(y) \right] \partial^\beta (\widehat{P}_{\bar{\beta}}^y)(y) \\
 &= \begin{cases} \delta_{\alpha\beta} - \left[\partial^{\bar{\beta}} (\widetilde{P}_\alpha^y)(y) \right] \delta_{\bar{\beta}\bar{\beta}} & = \delta_{\alpha\beta} \quad \text{for } \beta \in \mathcal{A}, \beta < \bar{\beta}, \\ \partial^{\bar{\beta}} (\widetilde{P}_\alpha^y)(y) - \left[\partial^{\bar{\beta}} (\widetilde{P}_\alpha^y)(y) \right] \delta_{\bar{\beta}\bar{\beta}} = 0 & \text{for } \beta = \bar{\beta} \end{cases} \\
 (38) \quad &= \delta_{\alpha\beta} \quad (\beta \in \bar{\mathcal{A}}).
 \end{aligned}$$

By now varying the point $y \in E_2$, we deduce from (32)-(34) and (36)-(38) that $\sigma(y, E_1)$ contains an $(\bar{\mathcal{A}}, y, C \cdot [\epsilon_1 + \epsilon_1^{-2D-1}\epsilon_2], 1)$ -basis for each $y \in E_2$. Since $\epsilon_2 \leq \epsilon_1^{2D+2}$, the conclusion of Lemma 3.3 is immediate. \square

Definition 4 (Near-triangular matrices). Let $S \geq 1$, $\epsilon \in (0, 1)$ and $\mathcal{A} \subset \mathcal{M}$ be given. A matrix $B = (B_{\alpha\beta})_{\alpha, \beta \in \mathcal{A}}$ is called (S, ϵ) near-triangular if

$$\begin{aligned}
 |B_{\alpha\beta} - \delta_{\alpha\beta}| &\leq \epsilon \quad (\alpha, \beta \in \mathcal{A}, \alpha \leq \beta); \text{ and} \\
 |B_{\alpha\beta}| &\leq S \quad (\alpha, \beta \in \mathcal{A}).
 \end{aligned}$$

Lemma 3.4. *Given $R \geq 1$, there exist constants $c_2 > 0$, $C_2 \geq 1$ depending only on R, m, n , such that the following holds. Suppose we are given $\epsilon_2 \in (0, c_2]$, $x \in \mathbb{R}^n$, a symmetric convex subset $\sigma \subset \mathcal{P}$ and a family of polynomials $(P_\alpha)_{\alpha \in \mathcal{A}} \subset \mathcal{P}$ such that*

$$(39a) \quad P_\alpha \in \epsilon_2 \cdot \sigma \quad (\alpha \in \mathcal{A});$$

$$(39b) \quad |\partial^\beta P_\alpha(x) - \delta_{\alpha\beta}| \leq \epsilon_2 \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}, \beta \geq \alpha); \text{ and}$$

$$(39c) \quad |\partial^\beta P_\alpha(x)| \leq R \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}).$$

Then there exists a $(C_2, C_2\epsilon_2)$ near-triangular matrix $B = (B_{\alpha\beta})_{\alpha, \beta \in \mathcal{A}}$, for which

$$(40) \quad \widehat{P}_\alpha := \sum_{\beta \in \mathcal{A}} B_{\alpha\beta} P_\beta \quad (\alpha \in \mathcal{A})$$

forms an $(\mathcal{A}, x, C_2\epsilon_2, 1)$ -basis for σ . Furthermore, $|\partial^\beta \widehat{P}_\alpha(x)| \leq C_2$ for every $\alpha \in \mathcal{A}$, $\beta \in \mathcal{M}$.

Proof. Let $c_2 \in (0, 1)$ and $C_2 \geq 1$ be constants depending only on R, m and n , that will be determined later. Define the matrix D by setting $D_{\alpha\beta} = \partial^\beta P_\alpha(x)$ for $\alpha, \beta \in \mathcal{A}$. From (39b) and (39c) it follows that D is (R, ϵ_2) near-triangular, where $\epsilon_2 \leq c_2$. Taking c_2 sufficiently small determined by R, m and n , we can ensure that D is invertible and that $B := D^{-1} = (B_{\alpha\beta})_{\alpha, \beta \in \mathcal{A}}$ is $(C', C'\epsilon_2)$ near-triangular, for some constant C' determined by R, m and n . This follows by a standard compactness argument.

For each $\alpha \in \mathcal{A}$ we define \widehat{P}_α as in (40). Since σ is symmetric and convex, from $|B_{\alpha\beta}| \leq C'$ and (39a) we deduce that

$$\widehat{P}_\alpha \in C''\epsilon_2 \cdot \sigma, \quad \text{where } C'' = C''(R, m, n).$$

Since $B = D^{-1}$, we have

$$\partial^\gamma \widehat{P}_\alpha(x) = \sum_{\beta \in \mathcal{A}} B_{\alpha\beta} D_{\beta\gamma} = \delta_{\alpha\gamma} \quad (\text{all } \gamma \in \mathcal{A}).$$

Finally, for each $\gamma \in \mathcal{M}$ with $\gamma > \alpha$, we write

$$|\partial^\gamma \widehat{P}_\alpha(x)| \leq \left| \sum_{\substack{\beta \in \mathcal{A} \\ \beta \leq \alpha}} B_{\alpha\beta} \partial^\gamma P_\beta(x) \right| + \left| \sum_{\substack{\beta \in \mathcal{A} \\ \beta > \alpha}} B_{\alpha\beta} \partial^\gamma P_\beta(x) \right|.$$

From (39b), (39c) and the fact that B is $(C', C'\epsilon_2)$ near-triangular, each summand above is dominated by $C'R\epsilon_2$. Hence,

$$|\partial^\gamma \widehat{P}_\alpha(x)| \leq \widehat{C}\epsilon_2 \quad (\text{all } \gamma \in \mathcal{M}, \gamma > \alpha), \quad \text{where } \widehat{C} = \widehat{C}(R, m, n).$$

Taking C_2 sufficiently large, determined by R , m and n , we have shown that $(\widehat{P}_\alpha)_{\alpha \in \mathcal{A}}$ forms an $(\mathcal{A}, x, C_2\epsilon_2, 1)$ -basis for σ . Finally, (39c) and $|B_{\alpha\beta}| \leq C'$ imply the last conclusion of the lemma. \square

4. THE INDUCTIVE HYPOTHESIS

Let $\mathcal{A} \subsetneq \mathcal{M}$. Here, we start on the proof of the Induction Step. We make the inductive assumption that the Main Lemma for \mathcal{A}' holds for each $\mathcal{A}' < \mathcal{A}$. Put

$$\epsilon_0 = \min\{\epsilon(\mathcal{A}') : \mathcal{A}' < \mathcal{A}\},$$

with $\epsilon(\mathcal{A}')$ as in the Main Lemma for \mathcal{A}' . From Remark 1 in section 3.2, we have:

(IH) Let $\widehat{\mathcal{A}} < \mathcal{A}$. Let $\widehat{E} \subset \mathbb{R}^n$ and $\widehat{z} \in \mathbb{R}^n$ satisfy $2 \leq \#(\widehat{E} \cup \{\widehat{z}\}) < \infty$, and assume that

$$(1) \quad \sigma(x, \widehat{E}) \text{ contains an } (\widehat{\mathcal{A}}, x, \epsilon_0, \delta_{\widehat{E}, \widehat{z}})\text{-basis for every } x \in \widehat{E},$$

$$\text{where } \delta_{\widehat{E}, \widehat{z}} = 10 \cdot \text{diam}(\widehat{E} \cup \{\widehat{z}\}).$$

Then the Extension Theorem for $(\widehat{E}, \widehat{z})$ holds.

Let us start on the proof of the Main Lemma for \mathcal{A} . The value of the universal constant $\epsilon(\mathcal{A})$ is determined later in the paper. We now assume that $\epsilon = \epsilon(\mathcal{A})$ is less than a small enough universal constant (the “small ϵ assumption”).

Fix $E \subset \mathbb{R}^n$ and $z \in \mathbb{R}^n$ with $2 \leq \#(E \cup \{z\}) < \infty$ and such that condition (3.1) from the Main Lemma for \mathcal{A} holds. By rescaling and translating E and $\{z\}$, we can arrange that

$$(2) \quad \delta_{E,z} := 10 \cdot \text{diam}(E \cup \{z\}) = 1 \text{ and}$$

$$E \cup \{z\} \subset \frac{1}{8}Q^\circ, \text{ where } Q^\circ := (0, 1]^n.$$

(Note that Q° is a dyadic cube as per our notation.)

Suppose that there exists $\overline{\mathcal{A}} < \mathcal{A}$ such that $\sigma(x, E)$ contains an $(\overline{\mathcal{A}}, x, \epsilon_0, 1)$ -basis for every $x \in E$. Then the Extension Theorem for (E, z) holds in view of **(IH)**. Having reached the conclusion of the Main Lemma for \mathcal{A} in this case, we now assume that

$$(3) \quad \text{for every } \overline{\mathcal{A}} < \mathcal{A}, \text{ there exists } x \in E$$

$$\text{such that } \sigma(x, E) \text{ does not contain an } (\overline{\mathcal{A}}, x, \epsilon_0, 1)\text{-basis.}$$

If $\#(E) \leq 1$, then the Extension Theorem for (E, z) holds (see Lemma 3.2). Thus, we may assume that

$$(4) \quad \#(E) \geq 2.$$

The assumptions (2)-(4) on (E, z) will remain in place until the end of section 18, when we prove Theorem 1 for finite E and establish Theorems 2, 3.

4.1. Auxiliary polynomials. Placing $\delta_{E;z} = 1$ into condition (3.1) from the Main Lemma for \mathcal{A} , we obtain $(\tilde{P}_\alpha^x)_{\alpha \in \mathcal{A}}$ that forms an $(\mathcal{A}, x, \epsilon, 1)$ -basis for $\sigma(x, E)$, for every $x \in E$. The goal of this subsection is to exhibit a similar basis for $\sigma(x, E)$ at every other point $x \in Q^\circ$.

As a consequence of (3), we will show that

$$(5) \quad |\partial^\beta \tilde{P}_\alpha^x(x)| \leq C \quad (x \in E, \alpha \in \mathcal{A}, \beta \in \mathcal{M}).$$

To prove (5), we fix a universal constant $\epsilon_1 \leq \min\{c_1, \epsilon_0/C_1\}$, where c_1 and C_1 are the constants from Lemma 3.3. For the sake of contradiction, suppose that (5) fails to hold with $C = \epsilon_1^{-D-1}$:

$$(6) \quad \max\{|\partial^\beta \tilde{P}_\alpha^x(x)| : x \in E, \alpha \in \mathcal{A}, \beta \in \mathcal{M}\} > \epsilon_1^{-D-1}.$$

We may assume that $\epsilon \leq \epsilon_1^{2D+2}$. Then the hypotheses of Lemma 3.3 hold with parameters

$$\left(\epsilon_1, \epsilon_2, \delta, \mathcal{A}, E_1, E_2, (\tilde{P}_\alpha^x)_{\alpha \in \mathcal{A}, x \in E_2}\right) := \left(\epsilon_1, \epsilon, 1, \mathcal{A}, E, E, (\tilde{P}_\alpha^x)_{\alpha \in \mathcal{A}, x \in E}\right).$$

(Here, the left-hand side denotes the local parameters of Lemma 3.3 and the right-hand side is defined as in the paragraph above. Notice that $\text{diam}(E) \leq 10$ and $E \neq \emptyset$ follow from (2) and (4), respectively.) Thus we find $\bar{\mathcal{A}} < \mathcal{A}$ so that $\sigma(x, E)$ contains an $(\bar{\mathcal{A}}, x, C_1\epsilon_1, 1)$ -basis for each $x \in E$. Since $C_1\epsilon_1 \leq \epsilon_0$, this contradicts (3), which concludes our proof of (5).

Fix $x_0 \in E$ and $\alpha \in \mathcal{A}$. Since $\tilde{P}_\alpha^{x_0} \in \epsilon \cdot \sigma(x_0, E)$, there exists $\varphi_\alpha \in L^{m,p}(\mathbb{R}^n)$ with

$$(7) \quad \varphi_\alpha = 0 \text{ on } E;$$

$$(8) \quad J_{x_0}(\varphi_\alpha) = \tilde{P}_\alpha^{x_0}; \text{ and}$$

$$(9) \quad \|\varphi_\alpha\|_{L^{m,p}(\mathbb{R}^n)} \leq \epsilon.$$

In view of (5), as well as **(B2)** and **(B3)** from the definition of $(\tilde{P}_\alpha^{x_0})_{\alpha \in \mathcal{A}}$ being an $(\mathcal{A}, x_0, \epsilon, 1)$ -basis, we have

$$(10) \quad |\partial^\beta \varphi_\alpha(x_0)| = |\partial^\beta \tilde{P}_\alpha^{x_0}(x_0)| \leq C \quad \text{for } \beta \in \mathcal{M};$$

$$(11) \quad \partial^\alpha \varphi_\alpha(x_0) = \partial^\alpha \tilde{P}_\alpha^{x_0}(x_0) = 1; \text{ and}$$

$$(12) \quad |\partial^{\beta+\gamma} \varphi_\alpha(x_0)| = |\partial^{\beta+\gamma} \tilde{P}_\alpha^{x_0}(x_0)| \leq \epsilon \quad \text{for } \beta > \alpha, |\gamma| \leq m-1-|\beta| \text{ and} \\ \text{for } \beta = \alpha, 0 < |\gamma| \leq m-1-|\beta|$$

(since in either case we have $\beta + \gamma > \alpha$).

For each $x \in Q^\circ$, define $\hat{P}_\alpha^x = J_x(\varphi_\alpha)$. Then (7) and (9) yield

$$(13) \quad \hat{P}_\alpha^x \in \epsilon \cdot \sigma(x, E).$$

For each $\beta \in \mathcal{M}$, by the Sobolev Inequality (notice that $|x_0 - x| \leq \text{diam}(Q^\circ) = 1$), we have

$$\begin{aligned} \left| \partial^\beta \widehat{P}_\alpha^x(x) - \delta_{\alpha\beta} \right| &= \left| \partial^\beta \varphi_\alpha(x) - \delta_{\alpha\beta} \right| \leq \left| \partial^\beta J_{x_0}(\varphi_\alpha)(x) - \delta_{\alpha\beta} \right| \\ &\quad + \left| \partial^\beta [J_{x_0}(\varphi_\alpha) - \varphi_\alpha](x) \right| \\ &\lesssim \left| \sum_{|\gamma| \leq m-1-|\beta|} \frac{1}{\gamma!} \partial^{\beta+\gamma} \varphi_\alpha(x_0)(x-x_0)^\gamma - \delta_{\alpha\beta} \right| + \|\varphi_\alpha\|_{L^{m,p}(\mathbb{R}^n)}. \end{aligned}$$

Thus, (9), (11), and (12) show that

$$(14) \quad \left| \partial^\beta \widehat{P}_\alpha^x(x) - \delta_{\alpha\beta} \right| \leq \widehat{C}\epsilon \quad (\beta \in \mathcal{M}, \beta \geq \alpha),$$

while (9) and (10) imply that

$$(15) \quad \left| \partial^\beta \widehat{P}_\alpha^x(x) \right| \leq \left| \partial^\beta \widehat{P}_\alpha^x(x) - \delta_{\alpha\beta} \right| + 1 \leq \widehat{C} + \epsilon + 1 \quad (\beta \in \mathcal{M}).$$

It follows from (13), (14), and (15) that (3.39a), (3.39b), and (3.39c) hold for $(\widehat{P}_\alpha^x)_{\alpha \in \mathcal{A}}$, with R equal to a universal constant, $\epsilon_2 = \widehat{C}\epsilon$ and $\sigma = \sigma(x, E)$. We may assume that $\widehat{C}\epsilon \leq c_2$, with the constant c_2 as in Lemma 3.4. We have verified the hypotheses of Lemma 3.4; hence for each $x \in Q^\circ$ there exists a $(C', C'\epsilon)$ near-triangular matrix $A^x = (A_{\alpha\beta}^x)_{\alpha, \beta \in \mathcal{A}}$ such that

$$P_\alpha^x := \sum_{\beta \in \mathcal{A}} A_{\alpha\beta}^x \cdot J_x(\varphi_\beta) \text{ satisfies:}$$

$$(16) \quad \begin{aligned} (P_\alpha^x)_{\alpha \in \mathcal{A}} \text{ is an } (\mathcal{A}, x, C'\epsilon, 1)\text{-basis for } \sigma(x, E), \text{ and} \\ \left| \partial^\beta P_\alpha^x(x) \right| \leq C', \text{ for every } \beta \in \mathcal{M}. \end{aligned}$$

That is,

$$(17a) \quad P_\alpha^x \in C'\epsilon \cdot \sigma(x, E) \quad (x \in Q^\circ, \alpha \in \mathcal{A});$$

$$(17b) \quad \partial^\beta P_\alpha^x(x) = \delta_{\alpha\beta} \quad (x \in Q^\circ, \alpha, \beta \in \mathcal{A});$$

$$(17c) \quad \left| \partial^\beta P_\alpha^x(x) \right| \leq C'\epsilon \quad (x \in Q^\circ, \alpha \in \mathcal{A}, \beta \in \mathcal{M}, \beta > \alpha); \text{ and}$$

$$(17d) \quad \left| \partial^\beta P_\alpha^x(x) \right| \leq C' \quad (x \in Q^\circ, \alpha \in \mathcal{A}, \beta \in \mathcal{M}).$$

Here and elsewhere, $\partial^\beta P_\alpha^x(x)$ denotes the value of $\partial_y^\beta P_\alpha^x(y)$ at $y = x$, not the β^{th} derivative of the function $x \mapsto P_\alpha^x(x)$.

4.2. Reduction to monotonic \mathcal{A} . The notion of monotonic labels played a key role in the study of C^m extension problems. It continues to be crucial for us here.

Definition 5 (Monotonic labels). A collection of multi-indices $\mathcal{A} \subset \mathcal{M}$ is monotonic if

$$\alpha \in \mathcal{A} \text{ and } |\gamma| \leq m-1-|\alpha| \implies \alpha + \gamma \in \mathcal{A}.$$

If the above property fails, we say that \mathcal{A} is non-monotonic.

In this section we deduce the monotonicity of \mathcal{A} using assumption (3) and condition (3.1) from the Main Lemma for \mathcal{A} .

We proceed by contradiction and assume that \mathcal{A} is non-monotonic. In this setting, we prove

⊛ There exists $\overline{\mathcal{A}} < \mathcal{A}$ so that $\sigma(x, E)$ contains an $(\overline{\mathcal{A}}, x, \epsilon_0, 1)$ -basis for each $x \in E$.

This contradicts (3); thus, our proof of the Induction Step will be reduced to the case of monotonic \mathcal{A} .

In order to construct $\overline{\mathcal{A}}$, we exploit the non-monotonicity of \mathcal{A} and choose $\alpha_0 \in \mathcal{A}$ and $\gamma \in \mathcal{M}$ with

$$(18) \quad 0 < |\gamma| \leq m - 1 - |\alpha_0| \quad \text{and} \quad \overline{\alpha} := \alpha_0 + \gamma \in \mathcal{M} \setminus \mathcal{A}.$$

We then define $\overline{\mathcal{A}} = \mathcal{A} \cup \{\overline{\alpha}\}$. Note that the minimal (only) element of $\overline{\mathcal{A}}\Delta\mathcal{A}$ is $\overline{\alpha}$, which is a member of $\overline{\mathcal{A}}$. Hence, $\overline{\mathcal{A}} < \mathcal{A}$ by definition of the order. Also, note that $\alpha_0 < \overline{\alpha}$.

For each $y \in Q^\circ$, we have defined $(P_\alpha^y)_{\alpha \in \mathcal{A}}$ that satisfy (16)–(17d). For each $y \in E$, we now define

$$(19) \quad P_{\overline{\alpha}}^y = P_{\alpha_0}^y \odot_y q^y, \quad \text{where} \quad q^y(x) := \frac{\alpha_0!}{\overline{\alpha}!} (x - y)^\gamma.$$

Expanding out this product, we have

$$P_{\overline{\alpha}}^y(x) = \frac{\alpha_0!}{\overline{\alpha}!} \sum_{|\omega| \leq m-1-|\gamma|} \frac{1}{\omega!} \partial^\omega P_{\alpha_0}^y(y) (x - y)^{\omega+\gamma}.$$

Note that $\omega = \alpha_0$ arises in the sum above, thanks to (18). Also, the terms with $\omega + \gamma > \overline{\alpha} = \alpha_0 + \gamma$ correspond precisely to $\omega > \alpha_0$ by definition of the order. The following properties are now immediate:

$$(20a) \quad \partial^{\overline{\alpha}} P_{\overline{\alpha}}^y(y) = 1, \quad \text{thanks to (17b) with } \alpha = \beta = \alpha_0.$$

$$(20b) \quad |\partial^\beta P_{\overline{\alpha}}^y(y)| \leq C'\epsilon \quad (\beta \in \mathcal{M}, \beta > \overline{\alpha}), \quad \text{thanks to (17c) with } \alpha = \alpha_0.$$

$$(20c) \quad |\partial^\beta P_{\overline{\alpha}}^y(y)| \leq C' \quad (\beta \in \mathcal{M}), \quad \text{thanks to (17d)}.$$

Next, we establish that

$$(20d) \quad P_{\overline{\alpha}}^y \in C'\epsilon \cdot \sigma(y, E).$$

From (17a) (with $\alpha = \alpha_0$), we find a function $\varphi \in L^{m,p}(\mathbb{R}^n)$ with

$$(21) \quad \varphi = 0 \text{ on } E;$$

$$(22) \quad J_y(\varphi) = P_{\alpha_0}^y; \text{ and}$$

$$(23) \quad \|\varphi\|_{L^{m,p}(\mathbb{R}^n)} \leq C\epsilon.$$

Consider $x \in Q^\circ$ and $\beta \in \mathcal{M}$ with $\beta > \alpha_0$. Since x and y belong to the unit cube Q° , we have $|x - y| \leq \delta_{Q^\circ} = 1$. Thus, (22) and the Sobolev Inequality yield

$$\begin{aligned} |\partial^\beta \varphi(x)| &\leq |\partial^\beta P_{\alpha_0}^y(x)| + |\partial^\beta \varphi(x) - \partial^\beta P_{\alpha_0}^y(x)| \leq |\partial^\beta P_{\alpha_0}^y(x)| + C\|\varphi\|_{L^{m,p}(\mathbb{R}^n)} \\ &= \left| \sum_{|\gamma| \leq m-1-|\beta|} \frac{1}{\gamma!} \partial^{\beta+\gamma} P_{\alpha_0}^y(y) (x - y)^\gamma \right| + C\|\varphi\|_{L^{m,p}(\mathbb{R}^n)}. \end{aligned}$$

Note that $\beta > \alpha_0 \implies \beta + \gamma > \alpha_0$. Thus, (17c),(23) imply that

$$(24) \quad |\partial^\beta \varphi(x)| \leq C'\epsilon \quad (x \in Q^\circ, \beta \in \mathcal{M}, \beta > \alpha_0).$$

Choose a cutoff function $\theta \in C_0^\infty(Q^\circ)$ with

$$(25) \quad \theta \equiv 1 \text{ in a neighborhood of } E \text{ and}$$

$$(26) \quad |\partial^\alpha \theta| \lesssim 1 \text{ when } |\alpha| \leq m.$$

(This cutoff exists in view of (2).) Define

$$\widehat{\varphi} = P_{\overline{\alpha}}^y + \theta(q^y \varphi - P_{\overline{\alpha}}^y).$$

Since $y \in E$, by (21), (22), and (25), we have

$$(27) \quad J_y \widehat{\varphi} = P_{\alpha_0}^y \odot_y q^y = P_{\alpha}^y \quad \text{and} \quad \widehat{\varphi}|_E = 0.$$

Because θ is compactly supported on Q° , the function $\widehat{\varphi}$ agrees with an $(m-1)^{\text{st}}$ degree polynomial outside $\text{supp}(\theta) \subset Q^\circ$. Also, since the m^{th} order derivatives of $P_{\alpha}^y \in \mathcal{P}$ vanish, we find that

$$\begin{aligned} \|\widehat{\varphi}\|_{L^{m,p}(\mathbb{R}^n)} &= \|\widehat{\varphi}\|_{L^{m,p}(Q^\circ)} = \|\theta \cdot (q^y \varphi - P_{\alpha}^y)\|_{L^{m,p}(Q^\circ)} \\ &\stackrel{(19),(22)}{=} \|\theta \cdot (q^y \varphi - J_y[q^y \varphi])\|_{L^{m,p}(Q^\circ)}. \end{aligned}$$

Thus, by (26) and the Sobolev Inequality, we have

$$(28) \quad \|\widehat{\varphi}\|_{L^{m,p}(\mathbb{R}^n)} \lesssim \sum_{|\alpha| \leq m} \|\partial^\alpha (q^y \varphi - J_y[q^y \varphi])\|_{L^p(Q^\circ)} \lesssim \|q^y \varphi\|_{L^{m,p}(Q^\circ)}.$$

Consider $x \in Q^\circ$ and some multi-index β with $|\beta| = m$. Since q^y is a polynomial of degree $|\gamma|$, we have

$$(29) \quad \partial^\beta [q^y \varphi](x) = \sum_{\omega + \omega' = \beta} \frac{\beta!}{\omega! \omega'!} \partial^\omega q^y(x) \partial^{\omega'} \varphi(x) = \sum_{\substack{\omega + \omega' = \beta \\ |\omega| \leq |\gamma|}} \frac{\beta!}{\omega! \omega'!} \partial^\omega q^y(x) \partial^{\omega'} \varphi(x).$$

If $\omega + \omega' = \beta$ and $|\omega| \leq |\gamma|$, then $|\omega'| \geq |\beta| - |\gamma| = m - |\gamma|$. Thus, from (18) we have $|\omega'| > |\alpha_0|$. By definition of the order on \mathcal{M} , either $[\omega = 0 \text{ and } \omega' = \beta]$ or $[\omega' \in \mathcal{M} \text{ and } \omega' > \alpha_0]$. Using either $\|q^y\|_{L^\infty(Q^\circ)} \leq C$ (see the definition of q^y in (19)) or (24) to bound the corresponding summand from (29), we obtain

$$|\partial^\beta [q^y \varphi](x)| \leq C' \cdot |\partial^\beta \varphi(x)| + C' \cdot \epsilon \quad (x \in Q^\circ, |\beta| = m).$$

Taking p^{th} powers, integrating over Q° and maximizing with respect to the multi-indices β with $|\beta| = m$, we obtain

$$\|\widehat{\varphi}\|_{L^{m,p}(\mathbb{R}^n)}^p \stackrel{(28)}{\lesssim} \|q^y \varphi\|_{L^{m,p}(Q^\circ)}^p \lesssim \|\varphi\|_{L^{m,p}(Q^\circ)}^p + \epsilon^p \stackrel{(23)}{\lesssim} \epsilon^p.$$

Together with (27), this proves (20d).

Now (17a)-(17d) and (20a)-(20d) imply that $(P_{\alpha}^y)_{\alpha \in \overline{\mathcal{A}}}$ satisfies (3.39a)-(3.39c) with R equal to a universal constant, $\epsilon_2 = C' \epsilon$ and $\sigma = \sigma(y, E)$. We may assume that $C' \epsilon \leq c_2$, where the constant c_2 comes from Lemma 3.4. Then the hypotheses of Lemma 3.4 hold, and hence $\sigma(y, E)$ contains an $(\overline{\mathcal{A}}, y, C'' \epsilon, 1)$ -basis for some universal constant $C'' \geq 1$.

Finally, we may assume that $\epsilon \leq \epsilon_0 / C''$, in which case $\sigma(y, E)$ contains an $(\overline{\mathcal{A}}, y, \epsilon_0, 1)$ -basis. Since $y \in E$ was arbitrary, this concludes the proof of \circledast . As already mentioned, this contradicts (3). Thus we may assume that $\mathcal{A} \subsetneq \mathcal{M}$ is a monotonic set. This property will be called upon much later to prove Proposition 3 (but nowhere else).

5. THE CZ DECOMPOSITION

To start, we make two definitions.

Definition 6 (OK cubes). A dyadic cube $Q \subset Q^\circ$ is OK if either $\#(3Q \cap E) \leq 1$ or there exists $\overline{\mathcal{A}} < \mathcal{A}$ such that $\sigma(x, E \cap 3Q)$ contains an $(\overline{\mathcal{A}}, x, \epsilon_0, 30\delta_Q)$ -basis for every $x \in E \cap 3Q$.

Definition 7 (Calderón-Zygmund cubes). A dyadic cube $Q \subset Q^\circ$ is CZ if Q is OK and all dyadic cubes $Q' \subset Q^\circ$ that properly contain Q are not OK.

The collection of CZ cubes will be denoted by

$$CZ^\circ = \{Q_1, \dots, Q_{\nu_{\max}}\}.$$

Lemma 5.1. *The Calderón-Zygmund cubes CZ° form a non-trivial finite partition of Q° into dyadic cubes. In particular, each $Q \in CZ^\circ$ has a (unique) dyadic parent $Q^+ \subset Q^\circ$.*

Proof. Recall our assumption that $E \subset \mathbb{R}^n$ is finite with $\#(E) \geq 2$ (see (4.4)). Therefore,

$$\delta := \inf \{|x - y| : x, y \in E \text{ distinct}\} \in (0, \infty).$$

Let $Q \subset Q^\circ$ be some dyadic cube with sidelength $\delta_Q \leq \delta/4$. It follows that $\#(E \cap 3Q) \leq 1$, and hence Q is OK. Since there are finitely many dyadic subcubes of Q° with sidelength greater than or equal to $\delta/4$, it follows that CZ° is a finite partition of Q° .

Suppose that Q° is OK. Since $E \subset Q^\circ$, this means that either

- (a) $\#(E) \leq 1$ or
- (b) $\sigma(x, E)$ contains an $(\mathcal{A}', x, \epsilon_0, 30\delta_{Q^\circ})$ -basis for each $x \in E$ and some $\mathcal{A}' < \mathcal{A}$.

Since $\delta_{Q^\circ} = 1$, we can replace $30\delta_{Q^\circ}$ by 1 in (b), while retaining validity (see Remark 1 in section 3.2). Therefore, (a) contradicts $\#(E) \geq 2$, while (b) contradicts (4.3).

It follows that Q° is not OK. Therefore, CZ° is not the trivial partition $\{Q^\circ\}$. \square

Two cubes $Q_\nu, Q_{\nu'} \in CZ^\circ$ are called “neighbors” if their closures satisfy $\text{cl}(Q_\nu) \cap \text{cl}(Q_{\nu'}) \neq \emptyset$. (In particular, any CZ cube is neighbors with itself.) We denote this property by $Q_\nu \leftrightarrow Q_{\nu'}$ or $\nu \leftrightarrow \nu'$.

Suppose that $\#(E \cap 3Q^+) \leq 1$ for some $Q \in CZ^\circ$. Then by definition Q^+ is OK, which contradicts $Q \in CZ^\circ$. Thus, we have

$$(1) \quad \#(E \cap 9Q) \geq \#(E \cap 3Q^+) \geq 2 \text{ for each } Q \in CZ^\circ.$$

(Here, we are using the fact that $Q^+ \subset 3Q$ for any cube $Q \subset \mathbb{R}^n$.)

Lemma 5.2 (Good geometry). *If $Q, Q' \in CZ^\circ$ satisfy $Q \leftrightarrow Q'$, then $\frac{1}{2}\delta_Q \leq \delta_{Q'} \leq 2\delta_Q$.*

Proof. For the sake of contradiction, suppose that $Q, Q' \in CZ^\circ$ satisfy

$$\text{cl}(Q) \cap \text{cl}(Q') \neq \emptyset \text{ and } 4\delta_Q \leq \delta_{Q'}.$$

Since Q^+, Q'^+ are dyadic cubes, it follows that $3Q^+ \subset 3Q'$. Since Q' is OK, either

- (a) $\#(3Q' \cap E) \leq 1$ or
- (b) $\sigma(x, E \cap 3Q')$ contains an $(\overline{\mathcal{A}}, x, \epsilon_0, 30\delta_{Q'})$ -basis for each $x \in 3Q' \cap E$ and some $\overline{\mathcal{A}} < \mathcal{A}$.

Note that (a) implies that $\#(3Q^+ \cap E) \leq 1$, which contradicts (1). It remains to consider (b).

By definition of $\sigma(\cdot, \cdot)$, we have $\sigma(x, E \cap 3Q') \subset \sigma(x, E \cap 3Q^+)$ for every $x \in E \cap 3Q^+$. Therefore, (b) implies that $\sigma(x, E \cap 3Q^+)$ contains an $(\overline{\mathcal{A}}, x, \epsilon_0, 30\delta_{Q'})$ -basis for each $x \in 3Q^+ \cap E$. Since $\delta_{Q^+} \leq \delta_{Q'}$, we may replace $\delta_{Q'}$ by δ_{Q^+} in the previous statement, while retaining validity. It follows that Q^+ is OK, which contradicts $Q \in CZ^\circ$. \square

Lemma 5.3 (More good geometry). *For each $Q \in CZ^\circ$, the following properties hold:*

- *If $Q' \in CZ^\circ$ is such that $(1.3)Q' \cap (1.3)Q \neq \emptyset$, then $Q \leftrightarrow Q'$. Consequently, each point $x \in \mathbb{R}^n$ belongs to at most $C(n)$ of the cubes $(1.3)Q$ with $Q \in CZ^\circ$.*
- *If $Q' \in CZ^\circ$ is such that $(1.1)Q' \cap 10Q \neq \emptyset$, then $\delta_{Q'} \leq 50\delta_Q$.*
- *If $Q' \in CZ^\circ$ is such that $(1.1)Q' \cap 100Q \neq \emptyset$, then $\delta_{Q'} \leq 10^3\delta_Q$.*
- *If $\text{cl}(Q) \cap \partial Q^\circ \neq \emptyset$, then $\delta_Q \geq \frac{1}{20}\delta_{Q^\circ}$.*

Proof. Let $Q \in CZ^\circ$ be fixed. We start with the first bullet point. Suppose that $Q' \in CZ^\circ$ does not neighbor Q . Put $\delta = \max\{\delta_Q/2, \delta_{Q'}/2\}$. Then Lemma 5.2 implies that $\text{dist}(Q, Q') \geq \delta$ (the CZ cubes that neighbor the larger cube have sidelength at least δ , providing a buffer between Q and Q' of this width). Thus,

$$(1.3)Q \cap (1.3)Q' \subset B(Q, (0.3)\delta) \cap B(Q', (0.3)\delta) = \emptyset.$$

This completes the proof of the first statement of the first bullet point. The second statement is immediate from Lemma 5.2. We pass to the second bullet point.

For the sake of contradiction, suppose that there exists $Q' \in CZ^\circ$ with $(1.1)Q' \cap 10Q \neq \emptyset$ and $\delta_{Q'} > 50\delta_Q$. Note that

$$(1.1)Q' \cap 10Q \neq \emptyset \implies \text{dist}(Q, Q') \leq (0.05)\delta_{Q'} + (4.5)\delta_Q.$$

Since $\delta_{Q'} > 50\delta_Q$, it follows that $\text{dist}(Q, Q') < (0.05 + .09)\delta_{Q'} < (0.15)\delta_{Q'}$. Therefore, $(1.3)Q \cap (1.3)Q' \neq \emptyset$. However, this contradicts $\delta_{Q'} > 50\delta_Q$ in view of the first bullet point and Lemma 5.2. This completes the proof of the second bullet point.

The proof of the third bullet point is analogous to the above.

Finally, we prove the fourth bullet point. For the sake of contradiction, suppose that $\text{cl}(Q) \cap \partial Q^\circ \neq \emptyset$ and $\delta_Q < \delta_{Q^\circ}/20$. Therefore, $9Q \subset \mathbb{R}^n \setminus (1/8)Q^\circ$. Because $E \subset (1/8)Q^\circ$, we have $9Q \cap E = \emptyset$. However, this contradicts (1). This proves the fourth bullet point and completes the proof of Lemma 5.3 \square

6. PATHS TO KEYSTONE CUBES I

In this section, constants called c_G, c, c', C, C' , etc. depend only on the dimension n . In this section we refer to such constants as “controlled constants”. The lowercase letters denote small (controlled) constants while the uppercase letters denote large (controlled) constants. A is a constant to be picked later. We assume A is greater than a large enough controlled constant (“large A assumption”).

We derive our main proposition in the following setting:

- We are given a CZ decomposition: \mathbb{R}^n is partitioned into a collection CZ of dyadic cubes, such that $Q, Q' \in CZ$ and $(1 + 10c_G)Q \cap (1 + 10c_G)Q' \neq \emptyset \implies \frac{1}{64}\delta_Q \leq \delta_{Q'} \leq 64\delta_Q$ (“good geometry”).
- We are given a finite set $E \subset \mathbb{R}^n$ with cardinality $N = \#(E) \geq 2$.
- The CZ cubes are related to the set E as follows: $\#(E \cap 9Q) \geq 2$ for each $Q \in CZ$.

Definition 8. A cube $Q \in CZ$ is called a keystone cube if for any $Q' \in CZ$ with $Q' \cap 100Q \neq \emptyset$, we have $\delta_{Q'} \geq \delta_Q$.

This section is devoted to a proof of the following proposition:

Proposition 1. *We can find a subset $CZ_{\text{special}} \subset CZ$ and an assignment to each $Q \in CZ$ of a finite sequence $\mathcal{S}_Q = (Q_1, Q_2, \dots, Q_L)$ of CZ cubes (with length L depending on Q) such that the following properties hold:*

- (i) CZ_{special} contains at most $C \cdot N$ distinct cubes.
- (ii) $Q_1 = Q$, Q_L is a keystone cube, $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$ for every $1 \leq l < L$, and

$$\delta_{Q_k} \leq C \cdot (1 - c)^{k-1} \delta_{Q_l} \text{ for } 1 \leq l \leq k \leq L.$$

- (iii) Let $Q, Q' \in CZ \setminus CZ_{\text{special}}$, and let $\mathcal{S}_Q = (Q_1, \dots, Q_L)$, $\mathcal{S}_{Q'} = (Q'_1, \dots, Q'_{L'})$. If $(1 + c_G)Q \cap (1 + c_G)Q' \neq \emptyset$, then $Q_L = Q'_{L'}$.

We begin the proof of Proposition 1.

Definition 9 (Clusters). A subset $S \subset E$ is a cluster if it satisfies $\#(S) \geq 2$ and $\text{dist}(S, E \setminus S) \geq A^3 \text{diam}(S)$, where the left-hand side $= \infty$ if $E \setminus S = \emptyset$.

Definition 10 (Representatives). For each cluster S we pick a representative $x(S) \in S$. We write $Q(S)$ to denote the CZ cube containing $x(S)$.

Definition 11 (Halos). For each cluster S , we define the halo of S to be

$$H(S) = \{x \in \mathbb{R}^n : A \cdot \text{diam}(S) < |x - x(S)| < A^{-1} \text{dist}(S, E \setminus S)\}.$$

Again, $\text{dist}(S, E \setminus S) = \infty$ if $S = E$.

Definition 12 (Interstellar cubes). $Q \in CZ$ is interstellar if $\text{diam}(A^{10}Q \cap E) \leq A^{-10} \delta_Q$ and $(1 + 3c_G)Q \cap E = \emptyset$.

We recall the Well Separated Pairs Decomposition [5].

Theorem 6 (Well Separated Pairs Decomposition). *Let $E \subset \mathbb{R}^n$ be finite. There exists a list of non-empty Cartesian products $E'_1 \times E''_1, E'_2 \times E''_2, \dots, E'_{\nu_{\max}} \times E''_{\nu_{\max}}$, each contained in $E \times E \setminus \{(x, x) : x \in E\}$, such that the following properties hold:*

- For each $\nu = 1, \dots, \nu_{\max}$, we have $\text{diam}(E'_\nu) + \text{diam}(E''_\nu) \leq 10^{-6} \text{dist}(E'_\nu, E''_\nu)$.
- Each pair $(x', x'') \in E \times E$ with $x' \neq x''$ belongs to precisely one $E'_\nu \times E''_\nu$.
- $\nu_{\max} \leq CN$.

Fix a representative $(x'_\nu, x''_\nu) \in E'_\nu \times E''_\nu$ for each $\nu = 1, \dots, \nu_{\max}$. Then for any $(x', x'') \in E \times E$ with $x' \neq x''$, we have $(x', x'') \in E'_\nu \times E''_\nu$ for some ν . For that ν , we have

$$\begin{aligned} |x'_\nu - x'| + |x''_\nu - x''| &\leq \text{diam}(E'_\nu) + \text{diam}(E''_\nu) \\ &\leq 10^{-6} \text{dist}(E'_\nu, E''_\nu) \leq 10^{-6} |x' - x''|, \end{aligned}$$

and similarly

$$|x'_\nu - x'| + |x''_\nu - x''| \leq 10^{-6} |x'_\nu - x''_\nu|.$$

Using the Well Separated Pairs Decomposition, we prove the following.

Lemma 6.1. *The number of non-interstellar cubes $Q \in CZ$ is at most $C(A) \cdot N$; here $C(A)$ depends only on A and on the dimension n .*

Lemma 6.2. *The number of distinct clusters is at most CN .*

Proof of Lemma 6.1. The non-interstellar cubes are $Q \in CZ$ such that $(1+3c_G)Q \cap E \neq \emptyset$ or $\text{diam}(A^{10}Q \cap E) > A^{-10}\delta_Q$.

For each $x \in E$, there are at most C distinct CZ cubes Q such that $x \in (1+3c_G)Q$. Therefore, there are at most CN distinct CZ cubes Q such that $(1+3c_G)Q \cap E \neq \emptyset$.

Next, we consider $Q \in CZ$ such that $\text{diam}(A^{10}Q \cap E) > A^{-10}\delta_Q$. For such Q , we can find $x', x'' \in E \cap A^{10}Q$ such that $|x' - x''| > A^{-10}\delta_Q$. For some ν , we have

$$|x'_\nu - x'| + |x''_\nu - x''| \leq 10^{-6} |x' - x''| \leq A^{10}\delta_Q.$$

Hence, $|x'_\nu - x''_\nu| \geq \frac{1}{2}A^{-10}\delta_Q$, yet $x'_\nu, x''_\nu \in CA^{10}Q$.

Therefore, the number of $Q \in CZ$ such that $\text{diam}(A^{10}Q \cap E) > A^{-10}\delta_Q$ is at most the sum over all $\nu = 1, \dots, \nu_{\max}$ of

$$\left[\begin{array}{l} \text{the number of distinct dyadic cubes } Q \text{ such that } x'_\nu, x''_\nu \in CA^{10}Q \\ \text{and } |x'_\nu - x''_\nu| \geq \frac{1}{2}A^{-10}\delta_Q. \end{array} \right]$$

For each fixed ν , the quantity in the square brackets is at most $C(A)$, and the number of distinct $\nu = 1, \dots, \nu_{\max}$ is at most CN .

Thus, there are at most $C(A)N$ distinct $Q \in CZ$ such that $\text{diam}(A^{10}Q \cap E) > A^{-10}\delta_Q$.

The proof of Lemma 6.1 is complete. \square

Proof of Lemma 6.2. Let S be a cluster. Fix $x', x'' \in S$ such that $|x' - x''| = \text{diam}(S)$. Then any point $y \in E \setminus S$ satisfies $|x' - y| \geq A^3|x' - x''|$. Fix ν such that

$$|x'_\nu - x'| + |x''_\nu - x''| \leq 10^{-6} |x' - x''|$$

and

$$|x'_\nu - x'| + |x''_\nu - x''| \leq 10^{-6}|x'_\nu - x''_\nu|.$$

Then

$$(1 - 10^{-6})|x' - x''| \leq |x'_\nu - x''_\nu| \leq (1 + 10^{-6}) |x' - x''|.$$

Any $y \in S$ satisfies

$$\begin{aligned} |y - x'_\nu| &\leq |y - x'| + |x' - x'_\nu| \leq \text{diam}(S) + 10^{-6}|x' - x''| \\ &= |x' - x''| + 10^{-6}|x' - x''| \leq 2|x'_\nu - x''_\nu|. \end{aligned}$$

On the other hand, any $y \in E \setminus S$ satisfies

$$\begin{aligned} |y - x'_\nu| &\geq |y - x'| - |x'_\nu - x'| \geq A^3|x' - x''| - 10^{-6}|x' - x''| \\ &> A|x'_\nu - x''_\nu|. \end{aligned}$$

Consequently, $S = E \cap B(x'_\nu, 10|x'_\nu - x''_\nu|)$ for some ν .

Thus, every cluster S arises as $E \cap B(x'_\nu, 10|x'_\nu - x''_\nu|)$ for some $\nu = 1, \dots, \nu_{\max}$. Since $\nu_{\max} \leq CN$, there can be at most CN distinct clusters. The proof of Lemma 6.2 is complete. \square

Lemma 6.3. *If Q is an interstellar cube, then $(1+c_G)Q \subset H(S)$ for some cluster S .*

Proof. Let $S = E \cap A^{10}Q$. Then $\#(S) \geq \#(E \cap 9Q) \geq 2$; also $\text{diam}(S) \leq A^{-10}\delta_Q$. Since S intersects $9Q$, it follows that $S \subseteq 13Q$. On the other hand, $E \setminus S \subseteq \mathbb{R}^n \setminus A^{10}Q$. Consequently, $\text{dist}(S, E \setminus S) \geq \text{dist}(13Q, \mathbb{R}^n \setminus A^{10}Q) \geq cA^{10}\delta_Q$. Thus, $\text{dist}(S, E \setminus S) \geq cA^{20}\text{diam}(S)$, proving that S is a cluster.

Next, let $x \in (1+c_G)Q$. We know that $(1+3c_G)Q \cap E = \emptyset$ since Q is interstellar. Hence, $|x - x(S)| \geq c\delta_Q$ since $x(S) \in S \subseteq E$. Therefore, $|x - x(S)| \geq cA^{10}\text{diam}(S)$. On the other hand, $x(S) \in S \subseteq 13Q$ and $x \in (1+c_G)Q$. It follows that $|x - x(S)| \leq C\delta_Q \leq C'A^{-10}\text{dist}(S, E \setminus S)$.

Thus, $cA^{10}\text{diam}(S) \leq |x - x(S)| \leq C'A^{-10}\text{dist}(S, E \setminus S)$. We have shown that each $x \in (1+c_G)Q$ belongs to $H(S)$. \square

Lemma 6.4. *Let S be a cluster. Let $Q \in CZ$, and let $x \in (1+c_G)Q \cap H(S)$. Then*

$$\frac{1}{2}A^{-1} \left[|x - x(S)| + \delta_{Q(S)} \right] \leq \delta_Q \leq A \left[|x - x(S)| + \delta_{Q(S)} \right].$$

Proof. We know that $A \cdot \text{diam}(S) < |x - x(S)| < A^{-1}\text{dist}(S, E \setminus S)$. In particular, $S \subset B(x, |x - x(S)| + \text{diam}(S)) \subset B(x, [1 + A^{-1}] \cdot |x - x(S)|) \subset B(x, 2|x - x(S)|)$ and

$$\text{dist}(x, S) \geq |x - x(S)| - \text{diam}(S) > [1 - A^{-1}] \cdot |x - x(S)| \geq \frac{1}{2}|x - x(S)|.$$

Also,

$$\begin{aligned} \text{dist}(x, E \setminus S) &\geq \text{dist}(x(S), E \setminus S) - |x - x(S)| \geq \text{dist}(S, E \setminus S) - |x - x(S)| \\ &> [A - 1] \cdot |x - x(S)|. \end{aligned}$$

In particular,

$$\text{dist}(x, E \setminus S) > [A - 1] \cdot |x - x(S)| > 2|x - x(S)| \geq \text{dist}(x, S).$$

Therefore, $\text{dist}(x, E) = \text{dist}(x, S) \geq \frac{1}{2}|x - x(S)|$. On the other hand, $E \cap 9Q \neq \emptyset$ since $Q \in CZ$, and $x \in (1+c_G)Q$. Therefore, $\text{dist}(x, E) \leq C\delta_Q$. It follows that $|x - x(S)| \leq 2\text{dist}(x, E) \leq C'\delta_Q$.

Next, we check that $\delta_{Q(S)} \leq A\delta_Q$. In fact, suppose that $\delta_{Q(S)} > A\delta_Q$. Then also $\delta_{Q(S)} > cA|x - x(S)|$. Since $x(S) \in Q(S)$ by definition, it follows that $x \in (1+c_G)Q(S)$. On the other hand, $x \in (1+c_G)Q$. Therefore, $\delta_{Q(S)}$ and δ_Q differ by at most a factor of 64, thanks to the good geometry of the CZ cubes. This contradicts our assumption that $\delta_{Q(S)} > A\delta_Q$, completing the proof that $\delta_{Q(S)} \leq A\delta_Q$.

We now know that $\frac{1}{2}A^{-1}[\delta_{Q(S)} + |x - x(S)|] \leq \delta_Q$.

Next, we show that $\delta_Q \leq A[\delta_{Q(S)} + |x - x(S)|]$. Indeed, suppose that

$$\delta_Q > A[\delta_{Q(S)} + |x - x(S)|].$$

Since $x \in (1+c_G)Q$ and $A^{-1}\delta_Q > |x - x(S)|$, it follows that $x(S) \in (1+2c_G)Q$. On the other hand, $x(S) \in Q(S)$. By the good geometry of the CZ cubes, the side lengths δ_Q and $\delta_{Q(S)}$ can differ by at most a factor of 64. This contradicts our assumption that $\delta_Q > A\delta_{Q(S)}$, completing the proof that $\delta_Q \leq A[\delta_{Q(S)} + |x - x(S)|]$. \square

Lemma 6.5. *For any two distinct clusters S, S' , the halos $H(S), H(S')$ are disjoint.*

Proof. Suppose $x \in H(S) \cap H(S')$, with S and S' distinct clusters. Then

$$A \cdot \text{diam}(S) < |x - x(S)| < A^{-1} \text{dist}(S, E \setminus S).$$

Let $R_S = 2|x - x(S)|$. Then

$$S \subset B(x, |x - x(S)| + \text{diam}(S)) \subset B(x, [1 + A^{-1}] \cdot |x - x(S)|) \subset B(x, R_S).$$

For $y \in E \setminus S$, we have

$$\begin{aligned} |y - x| &\geq \text{dist}(y, S) - \text{dist}(x, S) \geq \text{dist}(E \setminus S, S) - \text{dist}(x, S) \\ &\geq \text{dist}(E \setminus S, S) - |x - x(S)| \\ &\geq [A - 1] \cdot |x - x(S)| > 2|x - x(S)| = R_S. \end{aligned}$$

Therefore, $(E \setminus S) \cap B(x, R_S) = \emptyset$. Since we observed that $S \subset B(x, R_S)$ and since $S \subset E$, we conclude that $S = B(x, R_S) \cap E$.

Similarly, $S' = B(x, R_{S'}) \cap E$, where $R_{S'} = 2|x - x(S')|$. It follows that $S \subset S'$ or $S' \subset S$. Without loss of generality, we may suppose $S \subset S'$. Since S and S' are distinct, we can find $y \in S' \setminus S \subset E \setminus S$.

Note that $\text{diam}(S') \geq |y - x(S)| \geq \text{dist}(E \setminus S, S)$. Since $x \in H(S')$, we have $|x - x(S')| > A \cdot \text{diam}(S')$, and hence

$$\begin{aligned} |x - x(S)| &> A \cdot \text{diam}(S') - |x(S) - x(S')| \geq A \cdot \text{diam}(S') - \text{diam}(S') \\ &\quad (\text{since } x(S) \in S \subset S' \text{ and } x(S') \in S') \\ &\geq \frac{1}{2}A \cdot \text{diam}(S'). \end{aligned}$$

On the other hand, since $x \in H(S)$, we have

$$|x - x(S)| \leq A^{-1} \text{dist}(S, E \setminus S) \leq A^{-1} \text{dist}(S, S' \setminus S) \leq A^{-1} \text{diam}(S').$$

Thus, $|x - x(S)| \geq \frac{1}{2}A \cdot \text{diam}(S')$ and $|x - x(S)| \leq A^{-1} \text{diam}(S')$. This contradiction shows that we cannot have $x \in H(S) \cap H(S')$. \square

Lemma 6.6. *Let S be a cluster. Let $x, x' \in H(S)$, and let $Q, Q' \in CZ$, with $x \in Q$ and $x' \in Q'$. If $|x - x'| \leq A^{-2}|x - x(S)|$, then $(1 + c_G)Q \cap (1 + c_G)Q' \neq \emptyset$.*

Proof. By Lemma 6.4, we have $|x' - x| \leq A^{-2} [|x - x(S)| + \delta_{Q(S)}] \leq 2A^{-1} \delta_Q$. Since $x \in Q$, it follows that $x' \in (1 + c_G)Q$. Also, $x' \in Q' \subset (1 + c_G)Q'$. \square

Lemma 6.7. *Fix a cluster S . Let $x, x' \in H(S)$, with $|x - x(S)| \geq |x' - x(S)|$. Then there exist a finite sequence of points $x_1, x_2, \dots, x_L \in H(S)$ and a positive integer L_* with the following properties:*

- $x_1 = x$ and $x_L = x'$.
- $|x_{l+1} - x(S)| \leq |x_l - x(S)|$ for $l = 1, \dots, L - 1$.
- $|x_l - x_{l+1}| \leq A^{-2}|x_l - x(S)|$ for $l = 1, \dots, L - 1$.
- $|x_{l+L_*} - x(S)| \leq (1 - A^{-3})|x_l - x(S)|$ for $1 \leq l \leq L - L_*$.
- $L_* \leq A^3$.

Proof. Define a point $\tilde{x} \in \mathbb{R}^n$ such that

$$\tilde{x} - x(S) = |x - x(S)| \cdot \frac{x' - x(S)}{|x' - x(S)|}.$$

Then $\tilde{x} \in H(S)$ since $|\tilde{x} - x(S)| = |x - x(S)|$ and $x \in H(S)$. We have

$$(1) \quad \tilde{x} - x(S) = T \cdot [x' - x(S)] \text{ for some } T \geq 1.$$

We pick points x_1, \dots, x_{L_1} in $\partial B(x(S), |x - x(S)|)$ such that

(2)

$$x_1 = x; \quad x_{L_1} = \tilde{x};$$

$$|x_{l+1} - x_l| \leq A^{-2}|x - x(S)| = A^{-2}|x_l - x(S)| \quad \text{for } 1 \leq l < L_1; \quad \text{and } L_1 \leq \frac{1}{2}A^3.$$

We then pick positive real numbers $T_{L_1}, T_{L_1+1}, \dots, T_{L-1}, T_L$ with the properties:

$$T_{L_1} = T \quad \text{as in (1); } T_L = 1; \quad T_{l+1} = (1 - A^{-3})T_l \quad \text{for } L_1 \leq l \leq L - 2; \quad \text{and}$$

$$(1 - A^{-3})T_{L-1} \leq T_L \leq T_{L-1}.$$

Define the points $x_{L_1+1}, \dots, x_L \in \mathbb{R}^n$ by setting

(3)

$$x_l - x(S) = T_l \cdot [x' - x(S)] \quad \text{for } l = L_1 + 1, \dots, L.$$

Note that (3) also holds for $l = L_1$ and that $|x_l - x(S)| = T_l \cdot |x' - x(S)|$. We have $1 \leq T_l \leq T$ for each $l = L_1, \dots, L$; therefore $|x' - x(S)| \leq |x_l - x(S)| \leq |x - x(S)|$ for each such l . Since $x, x' \in H(S)$, it follows that each x_l also belongs to $H(S)$.

We have

(4)

$$|x_l - x_{l+1}| \leq A^{-3}|x_l - x(S)| \quad \text{for } L_1 \leq l \leq L - 1;$$

$$|x_{l+1} - x(S)| \leq |x_l - x(S)| \quad \text{for } L_1 \leq l \leq L - 1; \quad \text{and}$$

$$|x_{l+1} - x(S)| = (1 - A^{-3})|x_l - x(S)| \quad \text{for } L_1 \leq l \leq L - 2.$$

We have now defined x_1, \dots, x_L . Note that $x_1 = x$ and $x_L = x'$, which is the first bullet point in Lemma 6.7.

We know that $|x_{l+1} - x(S)| = |x_l - x(S)|$ for $1 \leq l \leq L_1 - 1$ and $|x_{l+1} - x(S)| \leq |x_l - x(S)|$ for $L_1 \leq l \leq L - 1$; therefore $|x_{l+1} - x(S)| \leq |x_l - x(S)|$ for $1 \leq l \leq L - 1$. This establishes the second bullet point of Lemma 6.7. Also, $|x_{l+1} - x(S)| \leq (1 - A^{-3})|x_l - x(S)|$ for $L_1 \leq l \leq L - 2$. The last two estimates together show that $|x_{l+(L_1+3)} - x(S)| \leq (1 - A^{-3})|x_l - x(S)|$ for $1 \leq l \leq L - (L_1 + 3)$. Here, $L_1 + 3 \leq \frac{1}{2}A^3 + 3 < A^3$. Thus, we have proven the last two bullet points of Lemma 6.7. The third bullet point in Lemma 6.7 is immediate from (2) and (4). We have verified all the conclusions of Lemma 6.7. \square

Lemma 6.8. *Fix a cluster S . Let $Q, Q' \in CZ$, and let $x, x' \in H(S)$, with $x \in Q$ and $x' \in Q'$. Assume*

$$|x - x(S)| \geq |x' - x(S)| \geq \delta_{Q(S)}.$$

Then there exist cubes $Q_1, Q_2, \dots, Q_L \in CZ$ with the following properties:

- $Q_1 = Q$ and $Q_L = Q'$.
- $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$ for all $l = 1, 2, \dots, L - 1$.
- $\delta_{Q_k} \leq C(A)(1 - c(A))^{k-l}\delta_{Q_l}$ for $1 \leq l \leq k \leq L$. Here, $0 < c(A) < 1$ and $C(A) > 0$ are constants depending only on A and on the dimension n .

Proof. Pick a sequence $x_1, x_2, \dots, x_L \in H(S)$ and an integer L_* , as in Lemma 6.7. For each l , let Q_l be the CZ cube containing x_l . In particular, $Q_1 = Q$ and $Q_L = Q'$, since $x_1 = x \in Q$ and $x_L = x' \in Q'$. For each $l = 1, 2, \dots, L - 1$, we have $|x_{l+1} - x_l| \leq A^{-2}|x_l - x(S)|$. Also, since $x_l \in Q_l$ and $x_{l+1} \in Q_{l+1}$, Lemma 6.6 tells us that $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$. In particular, $\delta_{Q_{l+1}}$ and δ_{Q_l} differ by at most a factor of 64.

Lemma 6.7 gives $|x_{l+1} - x(S)| \leq |x_l - x(S)|$ for $l = 1, 2, \dots, L - 1$; hence, $|x_l - x(S)| \geq |x' - x(S)| \geq \delta_{Q(S)}$. Hence, by Lemma 6.4 (and the fact that $x_l \in Q_l$),

δ_{Q_l} differs by at most a factor of $2A$ from $\delta_{Q(S)} + |x_l - x(S)|$, which in turn differs by at most a factor of 2 from $|x_l - x(S)|$.

Therefore,

$$\frac{1}{4}A^{-1}|x_l - x(S)| \leq \delta_{Q_l} \leq 4A|x_l - x(S)|, \text{ for each } l = 1, 2, \dots, L.$$

The fourth bullet point of Lemma 6.7 now gives

$$\delta_{Q_{L_*j}} \leq (4A)^2 \cdot (1 - A^{-3})^{j-i} \delta_{Q_{L_*i}} \text{ for } 1 \leq L_*i \leq L_*j \leq L.$$

Also, since $\delta_{Q_{l+1}}$ and δ_{Q_l} differ by at most a factor of 64 ($1 \leq l \leq L-1$) and since $1 \leq L_* \leq A^3$, it follows that

$$\delta_{Q_k} \leq C(A) \cdot (1 - c(A))^{k-l} \delta_{Q_l} \text{ for } 1 \leq l \leq k \leq L.$$

□

Fix a cluster S . For each $Q \in CZ$ such that $Q \cap H(S) \neq \emptyset$, we fix a point $x(Q, S) \in Q \cap H(S)$.

Lemma 6.9. *Given $R \geq \delta_{Q(S)}$, there are at most $CA^{2n} (R/\delta_{Q(S)})^n$ distinct cubes $Q \in CZ$ such that $Q \cap H(S) \neq \emptyset$ and $|x(Q, S) - x(S)| \leq R$.*

Proof. For each Q as in the statement of the lemma, we know from Lemma 6.4 that

$$\delta_Q \leq A \cdot [|x(Q, S) - x(S)| + \delta_{Q(S)}] \leq A \cdot [2R],$$

and therefore $Q \subset B(x(Q, S), CAR) \subset B(x(S), C'AR)$.

On the other hand, the CZ cubes are pairwise disjoint, and each CZ cube such that $Q \cap H(S) \neq \emptyset$ has volume at least $(2A)^{-n} \delta_{Q(S)}^n$, by Lemma 6.4. The conclusion of Lemma 6.9 follows at once. □

We say that $Q \in CZ$ is *privileged* for the cluster S (or *S-privileged*), provided that $Q \cap H(S) \neq \emptyset$ and $|x(Q, S) - x(S)| \leq \delta_{Q(S)}$. According to Lemma 6.9, there are at most CA^{2n} privileged cubes for a given cluster S .

Moreover, Lemma 6.9 shows that if there are CZ cubes Q such that

$$(5) \quad Q \cap H(S) \neq \emptyset \text{ and } |x(Q, S) - x(S)| > \delta_{Q(S)},$$

then there exists $\hat{Q}_S \in CZ$ such that $\hat{Q}_S \cap H(S) \neq \emptyset$, $|x(\hat{Q}_S, S) - x(S)| > \delta_{Q(S)}$, and $|x(\hat{Q}_S, S) - x(S)| \leq |x(Q, S) - x(S)|$ for any $Q \in CZ$ for which (5) holds.

For each such cluster S , we pick such a \hat{Q}_S .

Lemma 6.10. *Let S be a cluster, and let $Q \in CZ$. Suppose $Q \cap H(S) \neq \emptyset$, and suppose Q is not privileged for S . Then there exists a finite sequence of cubes $Q_1, Q_2, \dots, Q_L \in CZ$ with the following properties:*

- $Q_1 = Q$ and $Q_L = \hat{Q}_S$.
- $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$ for all $l = 1, 2, \dots, L-1$.
- $\delta_{Q_k} \leq C(A) \cdot (1 - c(A))^{k-l} \delta_{Q_l}$ for $1 \leq l \leq k \leq L$. Here, $0 < c(A) < 1$ and $C(A) > 0$ are constants depending only on A and on the dimension n .

Proof. Lemma 6.8 applies, with Q as in the present lemma, $Q' = \hat{Q}_S$, $x = x(Q, S)$, and $x' = x(\hat{Q}_S, S)$. □

Lemma 6.11. *Let $Q, Q' \in CZ$ be interstellar cubes, and suppose $(1 + c_G)Q \cap (1 + c_G)Q' \neq \emptyset$. Then there exists a cluster S such that $(1 + c_G)Q \subset H(S)$ and $(1 + c_G)Q' \subset H(S)$, and for any cluster $S' \neq S$, the cubes $(1 + c_G)Q$ and $(1 + c_G)Q'$ are both disjoint from $H(S')$.*

Proof. Immediate from Lemmas 6.3 and 6.5. \square

Lemma 6.12. *Let $Q \in CZ$. Then there exists a finite sequence Q_1, Q_2, \dots, Q_L of CZ cubes such that*

- $Q_1 = Q$ and Q_L is a keystone cube.
- $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$ for $l = 1, 2, \dots, L - 1$.
- $\delta_{Q_k} \leq C \cdot (1 - c)^{k-l} \delta_{Q_l}$ for $1 \leq l \leq k \leq L$.

Proof. Let $x \in \mathbb{R}^n$, and let Q^x be the CZ cube containing x . Then any CZ cube that meets $(1 + c_G)Q^x$ is a dyadic cube of sidelength at least $\delta_{Q^x}/64$. Hence, x has a neighborhood that meets only finitely many CZ cubes. Consequently, every compact set meets only finitely many CZ cubes.

If Q is a keystone cube, then the conclusion of the lemma holds with $L = 1$ and $Q_1 = Q$.

Suppose Q is not a keystone cube. Then there exist cubes $Q' \in CZ$ with

$$(6) \quad Q' \cap 100Q \neq \emptyset \text{ and } \delta_{Q'} \leq \frac{1}{2}\delta_Q.$$

There are only finitely many such Q' . Pick $Q^1 \in CZ$ that satisfies (6) and has minimal distance to Q among cubes satisfying (6). Let $\mathfrak{s} : [0, 1] \rightarrow \mathbb{R}^n$ be an affine map with

$$(7) \quad \mathfrak{s}(0) \in \text{cl}(Q), \quad \mathfrak{s}(1) \in \text{cl}(Q^1) \text{ and } |\mathfrak{s}(1) - \mathfrak{s}(0)| = \text{dist}(Q^1, Q).$$

Since Q^1 meets $100Q$ and the point $\mathfrak{s}(1) \in \text{cl}(Q^1)$ has a minimal distance to Q , it follows that $\mathfrak{s}(1) \in \text{cl}(100Q)$. Also, since $\mathfrak{s}(0) \in \text{cl}(Q)$, we have $\mathfrak{s}((0, 1)) \subset 100Q$.

The bounded set $\mathfrak{s}((0, 1))$ meets only finitely many CZ cubes $Q^{1,1}, \dots, Q^{1,K}$. Thus,

$$(8) \quad Q^{1,k} \cap 100Q \neq \emptyset \text{ and}$$

$$(9) \quad \text{dist}(Q^{1,k}, Q) < \text{dist}(Q^1, Q) \text{ for } k = 1, \dots, K.$$

(Here, we use (7) to prove (9).) Reordering the cubes $Q^{1,1}, \dots, Q^{1,K}$ if necessary, we can arrange that

$$(10) \quad \begin{aligned} &(1 + c_G)Q^{1,1} \cap (1 + c_G)Q \neq \emptyset \text{ and } (1 + c_G)Q^{1,K} \cap (1 + c_G)Q^1 \neq \emptyset; \text{ and} \\ &(1 + c_G)Q^{1,k} \cap (1 + c_G)Q^{1,k+1} \neq \emptyset \text{ for } k = 1, \dots, K - 1. \end{aligned}$$

From (9) and the definition of Q^1 , each $Q^{1,k}$ must not satisfy (6); hence $\delta_{Q^{1,k}} \geq \delta_Q$, thanks to (8). Since each $Q^{1,k}$ meets $100Q$, good geometry shows that $\delta_{Q^{1,k}} \leq C'\delta_Q$, as in the proof of Lemma 5.3. Therefore,

$$(11) \quad \delta_{Q^{1,k}} \simeq \delta_Q \text{ for } k = 1, \dots, K, \text{ and } K \leq C.$$

We call the cube $Q^1 \in CZ$ a *junior partner* of Q . Any sequence $(Q^{1,1}, \dots, Q^{1,K})$ of CZ cubes that satisfies (10) and (11) is said to *join* Q with Q^1 . Since Q^1 meets $100Q$ and satisfies $\delta_{Q^1} \leq \frac{1}{2}\delta_Q$, we have

$$C'' \cdot Q^1 \subset C'' \cdot Q \text{ whenever } Q^1 \text{ is a junior partner to } Q.$$

Now, either Q^1 is a keystone cube or it has a junior partner Q^2 . In the latter case, either Q^2 is a keystone cube or it has a junior partner Q^3 . We continue in this way, either forever or until we arrive at a keystone cube.

If the above process continued indefinitely, then we would have a sequence of CZ cubes Q^1, Q^2, Q^3, \dots with each Q^{j+1} being a junior partner to Q^j . That would imply that $\delta_{Q^{j+1}} \leq \frac{1}{2}\delta_{Q^j}$ and $C'' \cdot Q^{j+1} \subset C'' \cdot Q^j$ for each j . Thus, the cubes Q^j would shrink to a single point as $j \rightarrow \infty$. However, this contradicts the fact that every point has a neighborhood that meets only finitely many CZ cubes. Thus, the above process of successively passing to junior partners must stop after finitely many steps. Accordingly, starting from any $Q \in CZ$, we obtain a finite sequence Q^1, Q^2, \dots, Q^J of CZ cubes such that Q^{j+1} is a junior partner of Q^j for $1 \leq j \leq J-1$, and Q^J is a keystone cube. We now join each Q^j with Q^{j+1} through a sequence of CZ cubes. Concatenating these sequences, we obtain a sequence satisfying the conclusions of Lemma 6.12. \square

Lemma 6.13. *There exists a set CZ_{special} , consisting of at most $C(A) \cdot N$ distinct CZ cubes, for which the following holds. We can associate to each $Q \in CZ$ a finite sequence $\mathcal{S}_Q = (Q_1, Q_2, \dots, Q_L)$ of CZ cubes, with the following properties:*

- $Q_1 = Q$ and Q_L is a keystone cube.
- $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$ for $l = 1, 2, \dots, L-1$.
- $\delta_{Q_k} \leq C(A) \cdot (1 - c(A))^{k-l} \delta_{Q_l}$ for $1 \leq l \leq k \leq L$.
- Let $Q, Q' \in CZ \setminus CZ_{\text{special}}$, and suppose $(1 + c_G)Q \cap (1 + c_G)Q' \neq \emptyset$. Let $\mathcal{S}_Q = (Q_1, \dots, Q_L)$ and $\mathcal{S}_{Q'} = (Q'_1, \dots, Q'_{L'})$ be the finite sequences of CZ cubes associated to Q and to Q' , respectively. Then $Q_L = Q'_{L'}$.

Proof. First, we define the collection CZ_{special} . It consists of all non-interstellar cubes $Q \in CZ$, together with all $Q \in CZ$ that are privileged for some cluster S . If $Q \in CZ_{\text{special}}$, then we say that Q is “special”. We have seen that there are at most $C(A)N$ non-interstellar cubes and at most CN distinct clusters (see Lemmas 6.1 and 6.2). Since there are at most $C(A)$ privileged cubes for each given cluster, it follows that CZ_{special} consists of at most $C(A)N$ distinct CZ cubes.

Next, we define the sequence $\mathcal{S}_Q = (Q_1, Q_2, \dots, Q_L)$ for each CZ cube Q . If $Q \in CZ_{\text{special}}$, we just pick any finite sequence as in Lemma 6.12. Then \mathcal{S}_Q satisfies the first three bullet points in the statement of Lemma 6.13. It remains to define the \mathcal{S}_Q for all $Q \in CZ \setminus CZ_{\text{special}}$ and to prove that our \mathcal{S}_Q and CZ_{special} have the properties asserted in Lemma 6.13.

For each cluster S such that there exist cubes $Q \in CZ$ that are not privileged for S but that meet $H(S)$, we have picked out a cube \hat{Q}_S in the discussion following Lemma 6.9. Applying Lemma 6.12 to \hat{Q}_S , we obtain a finite sequence $\hat{Q}_S^1, \hat{Q}_S^2, \dots, \hat{Q}_S^{L(S)}$ of CZ cubes such that

- $\hat{Q}_S^1 = \hat{Q}_S$, $\hat{Q}_S^{L(S)}$ is a keystone cube.
- $(1 + c_G)\hat{Q}_S^l \cap (1 + c_G)\hat{Q}_S^{l+1} \neq \emptyset$ for $l = 1, 2, \dots, L(S)-1$.
- $\delta_{\hat{Q}_S^\nu} \leq C(A) \cdot (1 - c(A))^{\nu-\mu} \delta_{\hat{Q}_S^\mu}$ for $1 \leq \mu \leq \nu \leq L(S)$.

For each $Q \in CZ$ that is not privileged for S but that meets $H(S)$, we define a sequence Q_1, Q_2, \dots, Q_L as in Lemma 6.10. Thus, $Q_1 = Q$, $Q_L = \hat{Q}_S$, $(1 + c_G)Q_l \cap (1 + c_G)Q_{l+1} \neq \emptyset$ for $1 \leq l \leq L-1$, and $\delta_{Q_k} \leq C(A) \cdot (1 - c(A))^{k-l} \delta_{Q_l}$ for $1 \leq l \leq k \leq L$.

Unless Q is special (in which case we have already defined $\mathcal{S}(Q)$), we then define $\mathcal{S}(Q)$ to be the sequence

$$\mathcal{S}(Q) = \left(Q_1, Q_2, \dots, Q_L, \hat{Q}_S^1, \hat{Q}_S^2, \dots, \hat{Q}_S^{L(S)} \right).$$

This is well defined, since each $Q \in CZ \setminus CZ_{\text{special}}$ is as above for one and only one S (see Lemma 6.11 with $Q' = Q$).

Since $Q_L = \hat{Q}_S = \hat{Q}_S^1$, one checks easily that $\mathcal{S}(Q)$ satisfies the first three bullet points in the statement of Lemma 6.13.

Moreover, if Q and Q' are any two non-special CZ cubes that meet $H(S)$, then the finite sequences $\mathcal{S}(Q)$ and $\mathcal{S}(Q')$ both end with the finite sequence $\hat{Q}_S^1, \hat{Q}_S^1, \dots, \hat{Q}_S^{L(S)}$. In particular, the sequences $\mathcal{S}(Q)$ and $\mathcal{S}(Q')$ both end with the same cube, namely $\hat{Q}_S^{L(S)}$.

The above observation applies to any $Q, Q' \in CZ \setminus CZ_{\text{special}}$ such that $(1+c_G)Q \cap (1+c_G)Q' \neq \emptyset$. Indeed, Lemma 6.11 gives a cluster S such that $(1+c_G)Q, (1+c_G)Q' \subset H(S)$. The cluster S admits non-special cubes Q'' that intersect $H(S)$; indeed, we may take $Q'' = Q$ or Q' . Hence $\mathcal{S}(Q)$ and $\mathcal{S}(Q')$ end with the same cube, by the observation in the preceding paragraph.

The conclusions of Lemma 6.13 are now obvious. \square

Proof of Proposition 1. We simply take A in Lemma 6.13 to be a large enough constant determined by the dimension n . \square

7. PATHS TO KEYSTONE CUBES II

We place ourselves back in the setting of section 5. In particular, CZ° is a dyadic decomposition of the cube $Q^\circ = (0, 1]^n$. We define the collection of keystone cubes for CZ° by

$$(1) \quad CZ_{\text{key}}^\circ = \{Q \in CZ^\circ : \delta_{Q'} \geq \delta_Q \text{ for every } Q' \in CZ^\circ \text{ that meets } 10Q\},$$

which will also be denoted by

$$CZ_{\text{key}}^\circ = \{Q_1^\#, \dots, Q_{\mu_{\text{max}}}^\#\}.$$

Lemma 7.1. *For each $\mu = 1, \dots, \mu_{\text{max}}$, there are at most C' indices $\mu' \in \{1, \dots, \mu_{\text{max}}\}$ with $10Q_{\mu'}^\# \cap 10Q_\mu^\# \neq \emptyset$.*

Proof. Suppose that $Q^\#, \hat{Q}^\# \in CZ_{\text{key}}^\circ$ satisfy $10Q^\# \cap 10\hat{Q}^\# \neq \emptyset$. Without loss of generality, we may assume that $\delta_{Q^\#} \geq \delta_{\hat{Q}^\#}$. Therefore, $100Q^\# \cap \hat{Q}^\# \neq \emptyset$. By the definition of keystone cubes, we have $\delta_{\hat{Q}^\#} \geq \delta_{Q^\#}$. Thus, $\delta_{Q^\#} = \delta_{\hat{Q}^\#}$ whenever $10Q^\# \cap 10\hat{Q}^\# \neq \emptyset$. The conclusion of Lemma 7.1 follows immediately. \square

By applying Proposition 1 to the current setting we prove the following.

Proposition 2. *To each cube $Q_\nu \in CZ^\circ$ ($\nu = 1, \dots, \nu_{\text{max}}$), we can assign a finite sequence $\mathcal{S}_\nu = (Q_{\nu,1}, Q_{\nu,2}, \dots, Q_{\nu,L_\nu})$ of cubes from CZ° such that the following properties hold:*

(K1) $Q_{\nu,1} = Q_\nu$ and Q_{ν,L_ν} is a keystone cube for CZ° , $Q_{\nu,l} \leftrightarrow Q_{\nu,l+1}$ ($1 \leq l \leq L_\nu - 1$), and

$$\delta_{Q_{\nu,k}} \leq C \cdot (1-c)^{k-l} \delta_{Q_{\nu,l}} \quad (1 \leq l \leq k \leq L_\nu).$$

(K2) Let $\mathcal{K} : CZ^\circ \rightarrow CZ^\circ$ be defined by $\mathcal{K}(Q_\nu) = Q_{\nu, L_\nu}$. Then

$$\#\{(\nu, \nu') \in \{1, \dots, \nu_{\max}\}^2 : Q_\nu \leftrightarrow Q_{\nu'} \text{ and } \mathcal{K}(Q_\nu) \neq \mathcal{K}(Q_{\nu'})\} \leq C \cdot N.$$

(K3) $\mathcal{K}(Q_\nu) = Q_\nu$ for any $Q_\nu \in CZ^\circ_{\text{key}}$.

Proof. First we embed CZ° (a dyadic decomposition of Q°) into a dyadic decomposition CZ^+ of the whole \mathbb{R}^n .

Define the collection of dyadic cubes

$$(2) \quad \overline{CZ} = \left[\{Q \subset \mathbb{R}^n \text{ dyadic} : \delta_Q = 1, 3Q^+ \supset Q^\circ\} \cup \{Q \subset \mathbb{R}^n \text{ dyadic} : \delta_Q \geq 2, 3Q^+ \supset Q^\circ, 3Q \not\supset Q^\circ\} \right] \setminus \{Q^\circ\}.$$

We establish the following claims.

Claim 1. \overline{CZ} partitions $\mathbb{R}^n \setminus Q^\circ$ into dyadic cubes.

Claim 2. If $Q \in \overline{CZ}$, then $9Q \supset Q^\circ$.

Claim 3. If $Q, Q' \in \overline{CZ}$ satisfy $\text{cl}(Q) \cap \text{cl}(Q') \neq \emptyset$, then $1/2 \leq \delta_Q/\delta_{Q'} \leq 2$.

Claim 4. If $Q \subset \mathbb{R}^n$ is dyadic and satisfies $\delta_Q = 1$, $\text{cl}(Q) \cap \partial Q^\circ \neq \emptyset$ and $Q \neq Q^\circ$, then $Q \in \overline{CZ}$.

Proof of Claim 1. For each $x \in \mathbb{R}^n \setminus Q^\circ$, let $Q \subset \mathbb{R}^n$ be the smallest dyadic cube with $x \in Q$, $3Q^+ \supset Q^\circ$ and $\delta_Q \geq 1$. Then $Q \neq Q^\circ$, since $x \notin Q^\circ$. If $\delta_Q = 1$, then $Q \in \overline{CZ}$. On the other hand, if $\delta_Q \geq 2$, then $3Q \not\supset Q^\circ$ since Q is minimal, hence also $Q \in \overline{CZ}$. In either case, $Q \in \overline{CZ}$ and $x \in Q$. Thus, \overline{CZ} covers $\mathbb{R}^n \setminus Q^\circ$.

We now prove that the collection \overline{CZ} is pairwise disjoint. For the sake of contradiction, suppose that $Q, Q' \in \overline{CZ}$ are distinct with $Q \cap Q' \neq \emptyset$. Since Q, Q' are dyadic, either $Q \subsetneq Q'$ or $Q' \subsetneq Q$. Without loss of generality, $Q \subsetneq Q'$. Therefore, $Q^+ \subset Q'^+$. It follows that $\delta_{Q'} \geq 2\delta_Q \geq 2$. Also, since $Q \in \overline{CZ}$, we have $3Q^+ \supset Q^\circ$, hence $3Q' \supset Q^\circ$. Thus, $Q' \notin \overline{CZ}$, yielding the desired contradiction.

Obviously, each $Q \in \overline{CZ}$ is disjoint from Q° , which completes the proof of Claim 1.

Proof of Claim 2. Let $Q \in \overline{CZ}$. Then $9Q \supset 3Q^+ \supset Q^\circ$.

Proof of Claim 3. For the sake of contradiction, suppose that $Q, Q' \in \overline{CZ}$ satisfy $\text{cl}(Q) \cap \text{cl}(Q') \neq \emptyset$ and $\delta_{Q'} \geq 4\delta_Q$. It follows that $3Q^+ \subset 3Q'^+$. Therefore, since $Q^\circ \subset 3Q^+$, we have $Q^\circ \subset 3Q'$. Note that $\delta_{Q'} \geq 4$. Therefore, $Q' \notin \overline{CZ}$, yielding the desired contradiction.

Proof of Claim 4. For any cube Q that satisfies $\text{cl}(Q) \cap \partial Q^\circ \neq \emptyset$ and $\delta_Q \geq 1$, we have $3Q^+ \supset Q^\circ$. If Q is also dyadic with $\delta_Q = 1$ and $Q \neq Q^\circ$, then we have $Q \in \overline{CZ}$ by definition.

Let us define

$$CZ^+ = CZ^\circ \cup \overline{CZ}.$$

From Claim 1 and the fact that CZ° is a dyadic decomposition of Q° , it follows that CZ^+ is a dyadic decomposition of \mathbb{R}^n .

The following is immediate from Claims 3, 4, the good geometry of CZ° (Lemma 5.2) and the last bullet point in Lemma 5.3:

$$\text{If } Q, Q' \in CZ^+ \text{ satisfy } \text{cl}(Q) \cap \text{cl}(Q') \neq \emptyset, \text{ then } \delta_Q/\delta_{Q'} \in [1/64, 64].$$

Therefore,

(3) if $Q, Q' \in CZ^+$ satisfy $(1+10^{-5})Q \cap (1+10^{-5})Q' \neq \emptyset$, then $\text{cl}(Q) \cap \text{cl}(Q') \neq \emptyset$.

Thus, by the last two lines,

(4)

if $Q, Q' \in CZ^+$ satisfy $(1+10^{-5})Q \cap (1+10^{-5})Q' \neq \emptyset$, then $\delta_Q/\delta_{Q'} \in [1/64, 64]$.

It follows that CZ^+ satisfies the first bullet point at the beginning of section 6 with $c_G := 10^{-6}$.

From (5.1), we have $\#(E \cap 9Q) \geq 2$ for every $Q \in CZ^\circ$. Also, from Claim 2 and $E \subset Q^\circ$ we have $\#(E \cap 9Q) = \#(E) \geq 2$ for every $Q \in \overline{CZ}$. Thus, CZ^+ and $E \subset \mathbb{R}^n$ satisfy the second and third bullet points at the beginning of section 6.

Define the keystone cubes for CZ^+ as in section 6:

$$CZ_{\text{key}}^+ = \{Q \in CZ^+ : \delta_{Q'} \geq \delta_Q \text{ for every } Q' \in CZ^+ \text{ that meets } 100Q\}.$$

Since each $Q \in \overline{CZ}$ satisfies $\delta_Q \geq 1$ and $3Q^+ \supset Q^\circ$, there must exist cubes $Q' \in CZ^\circ$ with $Q' \cap 100Q \neq \emptyset$ and $\delta_{Q'} \leq \frac{1}{2}\delta_Q$. (Here, we also use the fact that CZ° partitions Q° into cubes of sidelength $\leq \frac{1}{2}$.) Consequently,

$$(5) \quad CZ_{\text{key}}^+ \subset CZ_{\text{key}}^\circ.$$

From Proposition 1, we find a subcollection $CZ_{\text{special}}^+ \subset CZ^+$ and an assignment to each $Q_\nu \in CZ^\circ$ ($\nu = 1, \dots, \nu_{\max}$) of a sequence $\widehat{\mathcal{S}}_\nu = (\widehat{Q}_{\nu,1}, \dots, \widehat{Q}_{\nu,L_\nu})$ of cubes such that the following properties hold:

(a) CZ_{special}^+ contains at most $C \cdot N$ distinct cubes.

(b) $\widehat{Q}_{\nu,1} = Q_\nu$ and $\widehat{Q}_{\nu,L_\nu} \in CZ_{\text{key}}^+$; each $\widehat{Q}_{\nu,k}$ belongs to CZ^+ ;

(6) $(1+10^{-6})\widehat{Q}_{\nu,l} \cap (1+10^{-6})\widehat{Q}_{\nu,l+1} \neq \emptyset$ for every $1 \leq l < L_\nu$; and

(7) $\delta_{\widehat{Q}_{\nu,k}} \leq C \cdot (1-c)^{k-l} \delta_{\widehat{Q}_{\nu,l}}$ for $1 \leq l \leq k \leq L_\nu$.

(c) If $Q_\nu, Q_{\nu'} \in CZ^\circ \setminus CZ_{\text{special}}^+$ satisfy $(1+10^{-6})Q_\nu \cap (1+10^{-6})Q_{\nu'} \neq \emptyset$, then

$$\widehat{Q}_{\nu,L_\nu} = \widehat{Q}_{\nu',L_{\nu'}}.$$

Suppose that $Q_\nu \in CZ^\circ \setminus CZ_{\text{key}}^\circ$ is such that $\widehat{\mathcal{S}}_\nu$ contains at least one cube from \overline{CZ} . In this case, we define $a(\nu)$ to be the first index $a \in \{1, \dots, L_\nu\}$ such that $\widehat{Q}_{\nu,a} \in \overline{CZ}$ and we define $b(\nu)$ to be the last index $b \in \{1, \dots, L_\nu\}$ such that $\widehat{Q}_{\nu,b} \in \overline{CZ}$.

Note that $\widehat{Q}_{\nu,1} = Q_\nu \in CZ^\circ$ and $\widehat{Q}_{\nu,L_\nu} \in CZ^\circ$, thanks to (b) and (5). Therefore, $1 < a(\nu) \leq b(\nu) < L_\nu$. Since $\widehat{Q}_{\nu,a(\nu)} \in \overline{CZ}$ and $\widehat{Q}_{\nu,a(\nu)-1} \in CZ^\circ$, by (3) and (6) it follows that the cube $\widehat{Q}_{\nu,a(\nu)-1}$ must touch the boundary of Q° . Likewise, $\widehat{Q}_{\nu,b(\nu)+1} \in CZ^\circ$ must touch the boundary of Q° .

In view of the last bullet point in Lemma 5.3, we may choose a sequence of CZ° cubes $\mathcal{E}_\nu = (Q_{\nu,a(\nu)-1}, \dots, Q_{\nu,s(\nu)+1})$ of bounded length (i.e., $s(\nu) - a(\nu) \leq C$) that connects $\widehat{Q}_{\nu,a(\nu)-1}$ with $\widehat{Q}_{\nu,b(\nu)+1}$ such that each of the $Q_{\nu,k}$ touches the boundary

of Q° . That is, we can arrange for

$$(8) \quad Q_{\nu, a(\nu)-1} = \widehat{Q}_{\nu, a(\nu)-1} \text{ and } Q_{\nu, s(\nu)+1} = \widehat{Q}_{\nu, b(\nu)+1};$$

$$(9) \quad (1 + 10^{-6})Q_{\nu, k} \cap (1 + 10^{-6})Q_{\nu, k+1} \neq \emptyset \text{ for all } a(\nu) - 1 \leq k \leq s(\nu);$$

$$(10) \quad Q_{\nu, k} \in CZ^\circ \text{ for each } k = a(\nu) - 1, \dots, s(\nu) + 1; \text{ and}$$

$$(11) \quad s(\nu) - a(\nu) \leq C.$$

Define the sequence of CZ° cubes:

$$\mathcal{S}_\nu = (\widehat{Q}_{\nu, 1}, \widehat{Q}_{\nu, 2}, \dots, \widehat{Q}_{\nu, a(\nu)-2}, Q_{\nu, a(\nu)-1}, \dots, Q_{\nu, s(\nu)+1}, \widehat{Q}_{\nu, b(\nu)+2}, \dots, \widehat{Q}_{\nu, L_\nu}).$$

Note that the starting cube $\widehat{Q}_{\nu, 1}$ and the terminating cube \widehat{Q}_{ν, L_ν} of the sequence \mathcal{S}_ν have not been changed, thanks to (8). Also, note that consecutive cubes from \mathcal{S}_ν are neighbors (they have intersecting closures), due to (3), (6), (8), and (9). Thanks to (7), (11) and the good geometry of the CZ° cubes, the sequence \mathcal{S}_ν satisfies the inequality from **(K1)**. Thus, we have defined \mathcal{S}_ν for each $Q_\nu \in CZ^\circ \setminus CZ_{\text{key}}^\circ$ such that $\widehat{\mathcal{S}}_\nu$ contains cubes from \overline{CZ} , and we have proven that \mathcal{S}_ν satisfies **(K1)**.

For any cube $Q_\nu \in CZ^\circ \setminus CZ_{\text{key}}^\circ$ such that the path $\widehat{\mathcal{S}}_\nu$ contains no cubes from \overline{CZ} , we simply take $\mathcal{S}_\nu = \widehat{\mathcal{S}}_\nu$. Then **(K1)** holds in view of (b).

For any cube $Q_\nu \in CZ_{\text{key}}^\circ$, we take $L_\nu = 1$ and $\mathcal{S}_\nu = (Q_{\nu, 1}) = (Q_\nu)$. Clearly, **(K1)** holds in this case.

Thus, we have defined \mathcal{S}_ν for each ν and established **(K1)** in all cases. Also, **(K3)** clearly holds. Thanks to Proposition 1 and the fact that each Q_ν has boundedly many neighbors, we obtain

$$\#\{(Q_\nu, Q_{\nu'}) \in [CZ^\circ \setminus CZ_{\text{key}}^\circ]^2 : Q_\nu \leftrightarrow Q_{\nu'}, Q_{\nu, L_\nu} \neq Q_{\nu', L_{\nu'}}\} \leq C \cdot \#(CZ_{\text{special}}^+).$$

Note that $\#(CZ_{\text{special}}^+) \leq C \cdot N$ by definition. Similarly, we have

$$\#\{(Q_\nu, Q_{\nu'}) \in CZ_{\text{key}}^\circ \times CZ^\circ : Q_\nu \leftrightarrow Q_{\nu'}\} \leq C \cdot \#(CZ_{\text{key}}^\circ).$$

Thanks to (5.1), to each $Q \in CZ_{\text{key}}^\circ$ we may assign a point $y_Q \in E \cap 10Q$. Lemma 7.1 shows that the preimage of each $y \in E$ has cardinality at most a universal constant. Thus, $\#(CZ_{\text{key}}^\circ) \leq C \cdot N$, which together with the previous two lines establishes **(K2)**.

This completes the proof of Proposition 2. \square

Recall that $\{Q_1^\sharp, \dots, Q_{\mu_{\max}}^\sharp\}$ denotes the keystone cubes. Using the map $\mathcal{K} : CZ^\circ \rightarrow CZ_{\text{key}}^\circ$ from Proposition 2 we produce a map on indices $\kappa : \{1, \dots, \nu_{\max}\} \rightarrow \{1, \dots, \mu_{\max}\}$ defined by $\mathcal{K}(Q_\nu) = Q_{\kappa(\nu)}^\sharp$.

8. REPRESENTATIVES

Since Q_ν is OK, the subset $E \cap 3Q_\nu$ lies on the zero set of a non-degenerate smooth function with small norm. Using this fact we prove the next result.

Lemma 8.1. *For each $\nu = 1, \dots, \nu_{\max}$, there exists $\widehat{x}_\nu \in \frac{1}{2}Q_\nu$ such that $\text{dist}(\widehat{x}_\nu, E) \geq c'\delta_{Q_\nu}$.*

Proof. Fix $\nu \in \{1, \dots, \nu_{\max}\}$. If $E \cap (1/4)Q_\nu = \emptyset$, then we take \widehat{x}_ν to be the center of Q_ν and reach the desired conclusion. Thus, we may assume that $E \cap (1/4)Q_\nu \neq \emptyset$. Let $y_\nu \in E \cap (1/4)Q_\nu$ be fixed.

If $\#(E \cap 3Q_\nu) \leq 1$, then the conclusion of the lemma is obvious. Thus we may assume that $\#(E \cap 3Q_\nu) \geq 2$. Thus, because Q_ν is OK there exists an $(\bar{\mathcal{A}}, y_\nu, \epsilon_0, 30\delta_{Q_\nu})$ -basis for $\sigma(y_\nu, E \cap 3Q_\nu)$, for some $\bar{\mathcal{A}} < \mathcal{A}$. Since $\bar{\mathcal{A}} < \mathcal{A}$ and the empty set is maximal under the order $<$, we may choose a multi-index $\alpha_0 \in \bar{\mathcal{A}}$. From the definition of an $(\bar{\mathcal{A}}, y_\nu, \epsilon_0, 30\delta_{Q_\nu})$ -basis (see **(B1)** and **(B2)**), we have a polynomial

$$P_{\alpha_0} \in C\epsilon_0\delta_{Q_\nu}^{n/p+|\alpha_0|-m} \cdot \sigma(y_\nu, E \cap 3Q_\nu) \text{ with } \partial^{\alpha_0} P_{\alpha_0}(y_\nu) = 1.$$

Choose $\alpha' \in \mathcal{M}$ with

$$|\partial^{\alpha'} P_{\alpha_0}(y_\nu)|\delta_{Q_\nu}^{|\alpha'|} \geq |\partial^\beta P_{\alpha_0}(y_\nu)|\delta_{Q_\nu}^{|\beta|} \text{ for all } \beta \in \mathcal{M}.$$

Placing $\beta = \alpha_0$ in the inequality above, we obtain

$$|\partial^{\alpha'} P_{\alpha_0}(y_\nu)| \geq |\partial^{\alpha_0} P_{\alpha_0}(y_\nu)|\delta_{Q_\nu}^{|\alpha_0|-|\alpha'|} = \delta_{Q_\nu}^{|\alpha_0|-|\alpha'|}.$$

Define $P = \left[\partial^{\alpha'} P_{\alpha_0}(y_\nu) \right]^{-1} \cdot P_{\alpha_0}$. Then the above three lines imply that

$$P \in C\epsilon_0\delta_{Q_\nu}^{n/p+|\alpha'|-m} \cdot \sigma(y_\nu, E \cap 3Q_\nu); \partial^{\alpha'} P(y_\nu) = 1; \text{ and}$$

$$|\partial^\beta P(y_\nu)| \leq \delta_{Q_\nu}^{|\alpha'|-|\beta|} \text{ for all } \beta \in \mathcal{M}.$$

By definition of $\sigma(y_\nu, E \cap 3Q_\nu)$, there exists $\varphi \in L^{m,p}(\mathbb{R}^n)$ with

$$(1a) \quad \varphi = 0 \text{ on } E \cap 3Q_\nu \text{ and } J_{y_\nu}(\varphi) = P;$$

$$(1b) \quad |\partial^\beta \varphi(y_\nu)| \leq \delta_{Q_\nu}^{|\alpha'|-|\beta|} \text{ for all } \beta \in \mathcal{M}; \text{ and}$$

$$(1c) \quad \|\varphi\|_{L^{m,p}(\mathbb{R}^n)} \leq C\epsilon_0\delta_{Q_\nu}^{n/p+|\alpha'|-m}.$$

Applying Bernstein's inequality to the polynomial $P \in \mathcal{P}$, we have

$$\max_{B(y_\nu, \delta)} |\partial^{\alpha'} P| \lesssim \delta^{-|\alpha'|} \max_{B(y_\nu, \delta)} |P| \quad (\delta > 0).$$

Since $\partial^{\alpha'} P(y_\nu) = 1$, the left-hand side is bounded from below by 1. Therefore, for every $\delta > 0$ there exists $x_\delta \in B(y_\nu, \delta)$ with $|P(x_\delta)| \geq c \cdot \delta^{|\alpha'|}$. By the Sobolev Inequality,

$$\begin{aligned} |(\varphi - P)(x_\delta)| &= |(\varphi - J_{y_\nu}(\varphi))(x_\delta)| \lesssim \|\varphi\|_{L^{m,p}(\mathbb{R}^n)} |x_\delta - y_\nu|^{m-n/p} \\ &\lesssim \delta_{Q_\nu}^{n/p+|\alpha'|-m} \delta^{m-n/p}, \text{ thanks to (1c) and } |x_\delta - y_\nu| \leq \delta. \end{aligned}$$

Thus,

$$\begin{aligned} |\varphi(x_\delta)| &\geq |P(x_\delta)| - |(\varphi - P)(x_\delta)| \geq c \cdot \delta^{|\alpha'|} - C\delta_{Q_\nu}^{n/p+|\alpha'|-m} \delta^{m-n/p} \\ &= \delta^{|\alpha'|} \cdot \left[c - C \left(\frac{\delta}{\delta_{Q_\nu}} \right)^{m-|\alpha'|-n/p} \right]. \end{aligned}$$

We now set $\delta = c_0\delta_{Q_\nu}$ for some small universal constant $c_0 < 1/8$ so that $|\varphi(x_\delta)| \geq c'' \cdot \delta_{Q_\nu}^{|\alpha'|}$ and

$$x_\delta \in B(y_\nu, \delta) \subset B(y_\nu, \delta_{Q_\nu}/8) \subset (1/2)Q_\nu, \text{ since } y_\nu \in (1/4)Q_\nu.$$

Also, (1b), (1c) and the Sobolev Inequality imply that $|\nabla \varphi| \leq C\delta_{Q_\nu}^{|\alpha'|-1}$ on $3Q_\nu$. Since $\varphi = 0$ on $E \cap 3Q_\nu$, we have

$$c'' \cdot \delta_{Q_\nu}^{|\alpha'|} \leq |\varphi(x_\delta)| = |\varphi(x_\delta) - \varphi(x)| \leq C\delta_{Q_\nu}^{|\alpha'|-1} \cdot |x_\delta - x|, \text{ for every } x \in E \cap 3Q_\nu.$$

Hence, $\text{dist}(x_\delta, E \cap 3Q_\nu) \geq c' \cdot \delta_{Q_\nu}$. Thus the conclusion of Lemma 8.1 holds with $\widehat{x}_\nu = x_\delta$. \square

We have indexed the Calderón-Zygmund cubes and the keystone cubes in an arbitrary manner. Without loss of generality, we may put in place several new indexing assumptions. First, we may assume that Q_1 contains the point $z \in (1/8)Q^\circ$. If Q_1 happens to be keystone, we also assume that $Q_1^\sharp = Q_1$. To summarize, we have

$$(2) \quad z \in Q_1, \text{ and if } Q_1 \text{ is keystone, then } Q_1 = Q_1^\sharp.$$

We make no further indexing assumptions on Q_μ^\sharp and Q_ν . We now define $\mu_{\min} = 2$ if Q_1 is keystone and $\mu_{\min} = 1$ otherwise. Thus,

$$\{Q_{\mu_{\min}}^\sharp, \dots, Q_{\mu_{\max}}^\sharp\} = \{Q_1^\sharp, \dots, Q_{\mu_{\max}}^\sharp\} \setminus \{Q_1\}.$$

We define representative basepoints for the CZ cubes:

$$x_1 = z; \text{ and } x_\nu = \widehat{x}_\nu \text{ for } \nu = 2, \dots, \nu_{\max}.$$

Similarly, we define representative basepoints for the keystone cubes:

$$x_\mu^\sharp = x_\nu \text{ for each } \mu = 1, \dots, \mu_{\max}, \text{ where } \nu \text{ is such that } Q_\mu^\sharp = Q_\nu.$$

Denote the collection of basepoints $E' := \{x_1, \dots, x_{\nu_{\max}}\}$ and the collection of keystone basepoints $E^\sharp := \{x_1^\sharp, \dots, x_{\mu_{\max}}^\sharp\}$.

Lemma 8.2. *The following properties hold:*

- $x_1 \in Q_1$ and $x_\nu \in (1/2)Q_\nu$ for $\nu = 2, \dots, \nu_{\max}$.
- $\text{dist}(x_\nu, E) \gtrsim \delta_{Q_\nu}$ for $\nu = 2, \dots, \nu_{\max}$.
- $x_\nu \in 0.99Q^\circ$ for $\nu = 1, \dots, \nu_{\max}$.
- $|x_\nu - x_{\nu'}| \geq \delta_{Q_\nu}/8$ for $\nu, \nu' = 1, \dots, \nu_{\max}$.

Proof. The first and second bullet points follow immediately from Lemma 8.1. Now we pass to the third bullet point. From Lemma 5.3, we have $\delta_{Q_\nu} \geq \delta_{Q^\circ}/20$ for any Q_ν that touches the boundary of Q° , and hence $(1/2)Q_\nu \subset 0.99Q^\circ$ for such cubes.

On the other hand, if $\text{cl}(Q_{\nu'}) \cap \partial Q^\circ = \emptyset$, then $\text{dist}(Q_{\nu'}, \partial Q^\circ) \geq \delta_{Q^\circ}/20$. (The cubes Q_ν that touch the boundary of Q° provide a buffer between $Q_{\nu'}$ and Q° of width $\delta_{Q^\circ}/20$.) Therefore, $Q_{\nu'} \subset 0.99Q^\circ$ for such cubes. Thus, $x_\nu \in (1/2)Q_\nu \subset 0.99Q^\circ$ for each $\nu = 2, \dots, \nu_{\max}$. Since $x_1 = z \in (1/8)Q^\circ$ by definition, we have established the third bullet point.

Finally, the fourth bullet point is an easy consequence of the first bullet point and the good geometry of CZ° . This concludes the proof of Lemma 8.2. \square

Henceforth a polynomial written P_ν , R_ν , etc. will always denote a jet at x_ν . Similarly, for any subcollection $E'' = \{x_{\nu_1}, \dots, x_{\nu_s}\} \subset E'$, we naturally identify tuples of polynomials $(P_{\nu_1}, \dots, P_{\nu_s})$, $(R_{\nu_1}, \dots, R_{\nu_s})$, etc. with Whitney fields on E'' .

For $P \in \mathcal{P}$, recall the norm defined in (2.1):

$$|P|_{x, \delta} = \left(\sum_{|\alpha| \leq m-1} |\partial^\alpha P(x)|^p \cdot \delta^{n+(|\alpha|-m)p} \right)^{1/p} \quad (x \in \mathbb{R}^n, \delta > 0).$$

For each $\nu = 1, \dots, \nu_{\max}$, we denote

$$|P|_\nu := |P|_{x_\nu, \delta_{Q_\nu}}.$$

Note that $|x_{\kappa(\nu)}^\# - x_\nu| \leq C'\delta_{Q_\nu}$, since $x_\nu \in Q_\nu$, $x_{\kappa(\nu)}^\# \in Q_{\kappa(\nu)}^\#$ and **(K1)** from Proposition 2 holds (recall that $\mathcal{K}(Q_\nu) = Q_{\kappa(\nu)}^\#$). Thus, (2.2) yields

$$(3) \quad |P|_\nu \simeq |P|_{x_{\kappa(\nu)}^\#, \delta_{Q_\nu}} = \left(\sum_{|\alpha| \leq m-1} |\partial^\alpha P(x_{\kappa(\nu)}^\#)|^p \cdot \delta_{Q_\nu}^{n+(|\alpha|-m)p} \right)^{1/p}.$$

Similarly, if $\nu' \in \{1, \dots, \nu_{\max}\}$ is such that $|x_{\nu'} - x_\nu| \leq C\delta_{Q_\nu}$ and $\delta_{Q_\nu}/\delta_{Q_{\nu'}} \in [C^{-1}, C]$, then by (2.2) we have

$$(4) \quad |P|_\nu \simeq |P|_{\nu'}.$$

In particular, (4) holds if $\nu \leftrightarrow \nu'$.

9. A PARTITION OF UNITY

Thanks to the properties of $Q_1, \dots, Q_{\nu_{\max}}$ and $x_1, \dots, x_{\nu_{\max}}$ established in Lemmas 5.2, 5.3 and 8.2, there exists a partition of unity $\{\theta_\nu\}_{\nu=1}^{\nu_{\max}} \subset C^\infty(Q^\circ)$ that satisfies the following properties. (We leave the construction as an exercise for the interested reader.)

(POU1) $0 \leq \theta_\nu \leq 1$.

(POU2) θ_ν vanishes on $Q^\circ \setminus (1.1)Q_\nu$.

(POU3) $|\partial^\alpha \theta_\nu| \leq C\delta_{Q_\nu}^{-|\alpha|}$ whenever $|\alpha| \leq m$.

(POU4) $\theta_\nu \equiv 1$ near x_ν and $\theta_\nu \equiv 0$ near $\{x_{\nu'} : \nu' = 1, \dots, \nu_{\max}, \nu' \neq \nu\}$.

(POU5) $\sum_{\nu=1}^{\nu_{\max}} \theta_\nu \equiv 1$ on Q° .

Lemma 9.1 (Patching estimate). *Let $G_\nu \in L^{m,p}(1.1Q_\nu)$ be given for each $\nu = 1, \dots, \nu_{\max}$, and define*

$$G(x) := \sum_{\nu=1}^{\nu_{\max}} G_\nu(x)\theta_\nu(x) \quad (x \in Q^\circ).$$

Then

$$\|G\|_{L^{m,p}(Q^\circ)}^p \lesssim \sum_{\nu=1}^{\nu_{\max}} \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p + \sum_{\nu \leftrightarrow \nu'} |J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'})|_\nu^p.$$

Proof. Fix $\nu' \in \{1, \dots, \nu_{\max}\}$. Since $\sum \theta_\nu = 1$ on Q° , for each $x \in Q_{\nu'}$ we have

$$G(x) = G(x) - G_{\nu'}(x) \left[\sum_{\nu=1}^{\nu_{\max}} \theta_\nu(x) \right] + G_{\nu'}(x) = \sum_{\nu=1}^{\nu_{\max}} (G_\nu - G_{\nu'})(x)\theta_\nu(x) + G_{\nu'}(x).$$

For each $x \in Q_{\nu'}$ there are at most a bounded number of θ_ν that do not vanish in a neighborhood of x (because $\text{supp}(\theta_\nu) \subset (1.1)Q_\nu$ and from the good geometry of the CZ cubes). Thus, by taking m^{th} derivatives and integrating p^{th} powers over $Q_{\nu'}$, we obtain

(1)

$$\|G\|_{L^{m,p}(Q_{\nu'})}^p \lesssim \|G_{\nu'}\|_{L^{m,p}(Q_{\nu'})}^p + \sum_{|\alpha+\beta|=m} \sum_{\nu=1}^{\nu_{\max}} \int_{Q_{\nu'}} |\partial^\beta (G_\nu - G_{\nu'})(x)|^p |\partial^\alpha \theta_\nu(x)|^p dx.$$

First we consider a term from the sum in (1) when ν is fixed, $|\beta| = m$ and $\alpha = 0$. Since $|\theta_\nu| \leq 1$ and θ_ν is supported in $(1.1)Q_\nu$, we have

$$(2) \quad \int_{Q_{\nu'}} |\partial^\beta (G_\nu - G_{\nu'})(x)|^p |\theta_\nu(x)|^p dx \lesssim \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p + \|G_{\nu'}\|_{L^{m,p}(Q_{\nu'})}^p.$$

The remaining terms in the sum arise when $|\beta| \leq m - 1$ and $|\alpha| = m - |\beta| \geq 1$. Since $|\partial^\alpha \theta_\nu| \lesssim \delta_{Q_\nu}^{-|\alpha|}$ and $\partial^\alpha \theta_\nu$ is supported in $(1.1)Q_\nu$, the sum of these terms is controlled by

$$(3) \quad C \sum_{\substack{1 \leq \nu \leq \nu_{\max} \\ 1.1Q_\nu \cap Q_{\nu'} \neq \emptyset}} \sum_{|\beta| \leq m-1} \sup_{x \in 1.1Q_\nu \cap Q_{\nu'}} |\partial^\beta (G_\nu - G_{\nu'})(x)|^p \delta_{Q_\nu}^n \delta_{Q_\nu}^{-(m-|\beta|)p}.$$

Because $|x - x_\nu| \lesssim \delta_{Q_\nu}$ whenever $x \in (1.1)Q_\nu$ and $|x - x_{\nu'}| \lesssim \delta_{Q_{\nu'}}$ whenever $x \in Q_{\nu'}$, the Sobolev Inequality (2.3) implies that

$$(4) \quad \begin{aligned} |\partial^\beta (G_\nu - G_{\nu'})(x)| &\lesssim |\partial^\beta (J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'}))(x)| + \|G_\nu\|_{L^{m,p}(1.1Q_\nu)} \delta_{Q_\nu}^{m-|\beta|-n/p} \\ &+ \|G_{\nu'}\|_{L^{m,p}(Q_{\nu'})} \delta_{Q_{\nu'}}^{m-|\beta|-n/p} \quad (\beta \in \mathcal{M}). \end{aligned}$$

Now summing (4) over all ν with $\nu \leftrightarrow \nu'$, and using good geometry of the CZ cubes, we find that (3) is bounded by

$$C' \sum_{\substack{1 \leq \nu \leq \nu_{\max} \\ \nu \leftrightarrow \nu'}} \left[\sup_{x \in 1.1Q_\nu \cap Q_{\nu'}} \sum_{|\beta| \leq m-1} \delta_{Q_\nu}^{n-(m-|\beta|)p} |\partial^\beta (J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'}))(x)|^p + \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p \right].$$

(Recall that $\nu' \leftrightarrow \nu'$ and thus the term from (4) that contains $\|G_{\nu'}\|_{L^{m,p}(1.1Q_{\nu'})}$ appears above as is required.) Applying (2.2) (recall that $|\cdot|_\nu = |\cdot|_{x_\nu, \delta_{Q_\nu}}$), we may bound the above by

$$(5) \quad C'' \sum_{\substack{1 \leq \nu \leq \nu_{\max} \\ \nu \leftrightarrow \nu'}} \left[|J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'})|_\nu^p + \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p \right].$$

Hence, by inserting the bounds (2) and (5) in (1) and by using the fact that each CZ cube has boundedly many neighbors, we have

$$(6) \quad \|F\|_{L^{m,p}(Q_{\nu'})}^p \lesssim \sum_{\substack{1 \leq \nu \leq \nu_{\max} \\ \nu \leftrightarrow \nu'}} \left[|J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'})|_\nu^p + \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p \right].$$

We now sum (6) over $\nu' = 1, \dots, \nu_{\max}$. Since the CZ cubes partition Q° and every CZ cube has a bounded number of neighbors, we have

$$\begin{aligned} \|F\|_{L^{m,p}(Q^\circ)}^p &\lesssim \sum_{\nu \leftrightarrow \nu'} \left[|J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'})|_\nu^p + \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p \right] \\ &\lesssim \sum_{\nu \leftrightarrow \nu'} |J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'})|_\nu^p + \sum_{\nu=1}^{\nu_{\max}} \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p, \end{aligned}$$

as required. \square

Lemma 9.2. *Let $0 < a < 1$, a cube $Q \subset \mathbb{R}^n$ and $F \in L^{m,p}(Q)$ be given. Then there exists $G \in L^{m,p}(\mathbb{R}^n)$ which depends linearly on F and satisfies*

$$G = F \text{ on } aQ \text{ and } \|G\|_{L^{m,p}(\mathbb{R}^n)} \leq C \cdot (1-a)^{-m} \cdot \|F\|_{L^{m,p}(Q)}.$$

Moreover, suppose that $F = T(f, \vec{P})$ for some linear map $T : L^{m,p}(E_1; E_2) \rightarrow L^{m,p}(Q)$ with Ω -assisted bounded depth, for some $\Omega \subset [L^{m,p}(E_1)]^$. Then one may take $G = T'(f, \vec{P})$ for some linear map $T' : L^{m,p}(E_1; E_2) \rightarrow L^{m,p}(\mathbb{R}^n)$ with Ω -assisted bounded depth.*

Proof. Let y denote the center of Q , and fix a cutoff function $\theta \in C_0^\infty(Q)$ that satisfies

$$(7) \quad \theta \equiv 1 \text{ on } aQ \text{ and}$$

$$(8) \quad |\partial^\alpha \theta| \lesssim ((1-a)\delta_Q)^{-|\alpha|} \text{ when } |\alpha| \leq m.$$

Define $G = \theta F + (1-\theta) \cdot J_y(F) \in L^{m,p}(\mathbb{R}^n)$. Then G depends linearly on F , while (7) implies that $G = F$ on aQ .

Since G matches an $(m-1)$ st degree polynomial on the complement of Q , we have

$$\begin{aligned} \|G\|_{L^{m,p}(\mathbb{R}^n)}^p &= \|G\|_{L^{m,p}(Q)}^p \lesssim \|F\|_{L^{m,p}(Q)}^p + \|(1-\theta) \cdot (J_y(F) - F)\|_{L^{m,p}(Q)}^p \\ &\lesssim \|F\|_{L^{m,p}(Q)}^p + \sum_{|\beta| \leq m-1} \delta_Q^n \cdot ((1-a)\delta_Q)^{-(m-|\beta|)p} \\ &\quad \cdot \sup_{x \in Q} |\partial^\beta (F - J_y(F))(x)|^p. \end{aligned}$$

(In the last inequality, the m^{th} order derivatives that fall on $(J_y(F) - F)$ have been raised to the p^{th} power and integrated over Q ; these terms are incorporated into $\|F\|_{L^{m,p}(Q)}^p$.) Thus, by the Sobolev Inequality, $\|G\|_{L^{m,p}(\mathbb{R}^n)} \lesssim (1-a)^{-m} \cdot \|F\|_{L^{m,p}(Q)}$.

It remains to analyze this construction from the perspective of assisted bounded depth linear maps. Suppose that $F = T(f, \vec{P})$ depends linearly on some data $(f, \vec{P}) \in L^{m,p}(E_1; E_2)$, where T has Ω -assisted bounded depth for some $\Omega \subset [L^{m,p}(E_1)]^*$. Then the function G is given by the linear map $T'(f, \vec{P}) := \theta T(f, \vec{P}) + (1-\theta) \cdot J_y(T(f, \vec{P}))$. Thus, for each point $x \in \mathbb{R}^n$ there exist linear maps $\psi_x, \phi_x : \mathcal{P} \rightarrow \mathcal{P}$ such that

$$(9) \quad J_x[T'(f, \vec{P})] = \begin{cases} \psi_x(J_x[T(f, \vec{P})]) + \phi_x(J_y[T(f, \vec{P})]) & : x \in Q, \\ \phi_x(J_y[T(f, \vec{P})]) & : x \notin Q. \end{cases}$$

These maps are defined by $\psi_x(P) = J_x[\theta P]$ and $\phi_x(P) = J_x[(1-\theta)P]$ for each $P \in \mathcal{P}$. Note that (9) follows from the containment $\text{supp}(\theta) \subset Q$ and the observation that $J_x[\theta H] = \psi_x(J_x H)$ and $J_x[(1-\theta)H] = \phi_x(J_x H)$ for each function H that is C^{m-1} on a neighborhood of x .

Because T has Ω -assisted bounded depth, (9) implies that T' has Ω -assisted bounded depth. This concludes the proof of Lemma 9.2. \square

10. LOCAL EXTENSION OPERATORS

In this section we apply the inductive hypothesis **(IH)** to construct local extension operators for functions defined on subsets of $E \cap 3Q_\nu$.

Fix $\nu \in \{1, \dots, \nu_{\max}\}$ and $\overline{E}_\nu \subset E \cap 3Q_\nu$.

First suppose that $\#(\overline{E}_\nu \cup \{x_\nu\}) \geq 3$. Note that $10 \operatorname{diam}(\overline{E}_\nu \cup \{x_\nu\}) \leq 30\delta_{Q_\nu}$ because $\overline{E}_\nu \subset 3Q_\nu$ and $x_\nu \in Q_\nu$. Also note that $\#(E \cap 3Q_\nu) \geq \#(\overline{E}_\nu) \geq 2$. By definition, since Q_ν is OK and $\#(E \cap 3Q_\nu) \geq 2$, there exists $\mathcal{A}_\nu < \mathcal{A}$ such that for all $x \in E \cap 3Q_\nu$ we have that

$$\begin{aligned} & \sigma(x, E \cap 3Q_\nu) \text{ contains an } (\mathcal{A}_\nu, x, \epsilon_0, 30\delta_{Q_\nu})\text{-basis} \\ \implies & \sigma(x, \overline{E}_\nu) \text{ contains an } (\mathcal{A}_\nu, x, \epsilon_0, 30\delta_{Q_\nu})\text{-basis} \\ & \text{(because } \sigma(x, E \cap 3Q_\nu) \subset \sigma(x, \overline{E}_\nu)\text{)} \\ \implies & \sigma(x, \overline{E}_\nu) \text{ contains an } (\mathcal{A}_\nu, x, \epsilon_0, 10 \operatorname{diam}(\overline{E}_\nu \cup \{x_\nu\}))\text{-basis.} \end{aligned}$$

(For the last implication, we use Remark 1 in section 3.2.) Hence, (4.1) holds with $\widehat{E} = \overline{E}_\nu$, $\widehat{x} = x_\nu$ and $\widehat{\mathcal{A}} = \mathcal{A}_\nu$. Thus, by the inductive hypothesis **(IH)**, the Extension Theorem for $(\overline{E}_\nu, x_\nu)$ holds.

On the other hand, suppose that $\#(\overline{E}_\nu \cup \{x_\nu\}) \leq 2$. Then Lemma 3.2 implies the Extension Theorem for $(\overline{E}_\nu, x_\nu)$.

In any case, for each $\nu \in \{1, \dots, \nu_{\max}\}$ and every $\overline{E}_\nu \subset E \cap 3Q_\nu$, there exists $(T_\nu, M_\nu, \Omega_\nu)$ with the following properties:

- (L1)** $T_\nu : L^{m,p}(\overline{E}_\nu; x_\nu) \rightarrow L^{m,p}(\mathbb{R}^n)$ is a linear extension operator.
- (L2)** $\|(f_\nu, P_\nu)\|_{L^{m,p}(\overline{E}_\nu, x_\nu)} \leq \|T_\nu(f_\nu, P_\nu)\|_{L^{m,p}(\mathbb{R}^n)} \leq C\|(f_\nu, P_\nu)\|_{L^{m,p}(\overline{E}_\nu, x_\nu)}$, and
- (L3)** $C^{-1}M_\nu(f_\nu, P_\nu) \leq \|T_\nu(f_\nu, P_\nu)\|_{L^{m,p}(\mathbb{R}^n)} \leq CM_\nu(f_\nu, P_\nu)$ for each (f_ν, P_ν) .
- (L4)** $\sum_{\omega \in \Omega_\nu} \operatorname{dp}(\omega) \leq C \cdot \#(\overline{E}_\nu)$.
- (L5)** T_ν has Ω_ν -assisted bounded depth.
- (L6)** There exists a collection of linear functionals $\Xi_\nu \subset L^{m,p}(\overline{E}_\nu; x_\nu)^*$ such that
 - (a)** each functional in Ξ_ν has Ω_ν -assisted bounded depth,
 - (b)** $\#(\Xi_\nu) \leq C \cdot \#(\overline{E}_\nu)$, and
 - (c)** $M_\nu(f_\nu, P_\nu) = \left(\sum_{\xi \in \Xi_\nu} |\xi(f_\nu, P_\nu)|^p \right)^{1/p}$ for each (f_ν, P_ν) .

By patching together local extension operators that satisfy **(L1)-(L6)**, in section 15, we construct a bounded linear extension operator for the trace space $L^{m,p}(E; E')$. Along the way, we find an explicit expression for the $L^{m,p}(E; E')$ trace seminorm. The one remaining task is to determine a jet $\vec{R} \in Wh(E')$ that essentially minimizes this expression.

We now summarize the remaining content of the paper.

- In section 11: we construct keystone components of the jet \vec{R} , i.e. polynomial jets associated with the keystone representative points. Later in section 16 we set the remaining components of \vec{R} equal to certain associated keystone components. This association is determined through the paths introduced in section 7.
- In section 12: we provide local estimates on the auxiliary polynomials defined in section 4.

- In section 13: we estimate the size/shape of $\sigma(x, S)$ for various localized sets $S \subset E$. (These estimates are stated in terms of the dual trace seminorm $\|(0|_S, \cdot)\|_{L^{m,p}(S;x)\cdot}$)
- In section 14: we prove a Sobolev-type inequality which bounds the variance of an $L^{m,p}$ function along the paths introduced in section 7. Using this inequality and the local estimates from section 12, we demonstrate the existence of near-optimal extensions with two additional properties: *Coherence* and *Constancy Along Paths*.
- In section 15: we construct an extension operator for the trace space $L^{m,p}(E; E')$ and estimate the corresponding trace seminorm.
- In section 16: we prove that the jet $\vec{R} \in Wh(E')$ nearly minimizes the $L^{m,p}(E; E')$ trace seminorm. We use results from sections 13 and 14, as well as the estimation of the trace seminorm from section 15.
- In section 17: we prove the Extension Theorem for (E, z) .
- In section 18: we prove Theorem 1 for finite E , and Theorems 2 and 3; we also prove an extension theorem for the inhomogeneous Sobolev space.
- In section 19: we deduce Theorem 1 for infinite sets from the finite case using a Banach limit.

Until the end of section 18, we fix some arbitrary data $(f, P) \in L^{m,p}(E; z)$.

11. EXTENSION NEAR THE KEYSTONE CUBES

We begin with a simple lemma concerning the minimization of the ℓ^p -norm subject to linear constraints. We will work in \mathbb{R}^{N_0+k} and denote a vector in \mathbb{R}^{N_0+k} by (w, w') with $w \in \mathbb{R}^{N_0}$, $w' \in \mathbb{R}^k$.

Lemma 11.1. *Given integers $N_0, k \geq 0$ and linear functionals $\lambda_1, \dots, \lambda_L$ on \mathbb{R}^{N_0+k} , there exists a linear map $\xi : \mathbb{R}^{N_0} \rightarrow \mathbb{R}^k$ so that for each $w \in \mathbb{R}^{N_0}$ we have*

$$\sum_{i=1}^L |\lambda_i(w, \xi(w))|^p \leq C \inf_{w' \in \mathbb{R}^k} \left\{ \sum_{i=1}^L |\lambda_i(w, w')|^p \right\}, \quad \text{with } C = C(k, p).$$

Proof. It suffices to assume that $k = 1$, because by iteratively applying this result we obtain the full version of the lemma. Write $\lambda_i(w, w') = \widehat{\lambda}_i(w) - a_i w'$, where $\widehat{\lambda}_i : \mathbb{R}^{N_0} \rightarrow \mathbb{R}$ is a linear functional and $a_i \in \mathbb{R}$ for each $i = 1, \dots, L$. The expression to be minimized is

$$\sum_{i=1}^L \left| \widehat{\lambda}_i(w) - a_i w' \right|^p.$$

By removing those terms that are independent of w' , we may assume that $a_i \neq 0$ for all $i = 1, \dots, L$, and rewrite the expression as

$$(1) \quad \sum_{i=1}^L \left| \frac{\widehat{\lambda}_i(w)}{a_i} - w' \right|^p |a_i|^p.$$

If $L \neq 0$, then a standard application of Hölder's inequality shows that

$$w' = \xi(w) := \frac{\widehat{\lambda}_1(w)|a_1|^p/a_1 + \dots + \widehat{\lambda}_L(w)|a_L|^p/a_L}{|a_1|^p + \dots + |a_L|^p}$$

minimizes (1) to within a factor of 2^{p+1} . If $L = 0$, we simply take $w' = 0$. \square

Lemma 11.2. *Let Q_μ^\sharp be a keystone cube. Set $f_\mu^\sharp = f|_{E \cap 9Q_\mu^\sharp}$. Then there exists $R_\mu^\sharp \in \mathcal{P}$ with the following properties:*

$$(2) \quad R_\mu^\sharp \text{ depends linearly on } f_\mu^\sharp \text{ and } P;$$

$$(3) \quad \partial^\alpha R_\mu^\sharp(x_\mu^\sharp) = \partial^\alpha P(x_\mu^\sharp) \text{ for every } \alpha \in \mathcal{A};$$

and

$$(4) \quad \|(f_\mu^\sharp, R_\mu^\sharp)\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)} \\ \leq C \inf_{R' \in \mathcal{P}} \left\{ \|(f_\mu^\sharp, R')\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)} : \partial^\alpha R'(x_\mu^\sharp) = \partial^\alpha P(x_\mu^\sharp) \text{ for every } \alpha \in \mathcal{A} \right\}.$$

Local extension problems on 9-dilated cubes play an important role in our proof. For instance, these problems have unique solutions modulo harmless errors once \mathcal{A} -derivatives are specified. (See Lemma 13.1.) Note that 9-dilates of keystone cubes are covered by a controlled number of CZ cubes. This fact is central to the proof of Lemma 11.2, which explains why we restrict our attention to keystone cubes.

Proof. The main content of the proof consists of showing that the trace seminorm

$$(5) \quad \|(f_\mu^\sharp, R')\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)}$$

is given up to a universal constant factor by an expression of the type

$$(6) \quad \left(|\lambda_1(f_\mu^\sharp, R')|^p + \cdots + |\lambda_L(f_\mu^\sharp, R')|^p \right)^{1/p},$$

where $\lambda_1, \dots, \lambda_L$ are some linear functionals.

Let $CZ_\mu^\sharp = \{Q \in CZ^\circ : Q \cap 100Q_\mu^\sharp \neq \emptyset\}$. We write $CZ_\mu^\sharp = \{Q_{\nu_1}, \dots, Q_{\nu_K}\}$, where $Q_{\nu_1} = Q_\mu^\sharp$. Hence also $x_{\nu_1} = x_\mu^\sharp$. By definition of the keystone cubes and Lemma 5.3, we have

$$(7) \quad \delta_{Q_\mu^\sharp} \leq \delta_{Q_{\nu_i}} \leq 10^3 \delta_{Q_\mu^\sharp} \text{ for } i = 1, \dots, K, \\ \text{and hence } K \leq C'.$$

Let $R' \in \mathcal{P}$ and $P_{\nu_i} \in \mathcal{P}$ for $i = 2, \dots, K$ be given (arbitrary) polynomials. Set $P_{\nu_1} = R'$. For each $i = 1, \dots, K$, we define the following objects:

- The subset $S_{\nu_i} = E \cap 9Q_\mu^\sharp \cap 3Q_{\nu_i}$ and the function $f_{\nu_i} = f|_{S_{\nu_i}}$.
- The map $M_{\nu_i} : L^{m,p}(S_{\nu_i}; x_{\nu_i}) \rightarrow \mathbb{R}_+$ that satisfies **(L1)**-**(L6)** for $S_{\nu_i} \subset E \cap 3Q_{\nu_i}$ and the representative $x_{\nu_i} \in Q_{\nu_i}$ (together with some linear map $T_{\nu_i} : L^{m,p}(S_{\nu_i}; x_{\nu_i}) \rightarrow L^{m,p}(\mathbb{R}^n)$ and some collection of linear functionals $\Omega_{\nu_i} \subset [L^{m,p}(S_{\nu_i})]^*$).

From **(L2)**, **(L3)**, and **(L6)**, for any $i = 1, \dots, K$ we have

$$(8) \quad M_{\nu_i}(f_{\nu_i}, P_{\nu_i}) = \left(\sum_{j=1}^{L_i} |\lambda_i^j(f_{\nu_i}, P_{\nu_i})|^p \right)^{1/p} \simeq \|(f_{\nu_i}, P_{\nu_i})\|_{L^{m,p}(S_{\nu_i}; x_{\nu_i})}$$

for some linear functionals $\lambda_i^1, \dots, \lambda_i^{L_i} \in [L^{m,p}(S_{\nu_i}; x_{\nu_i})]^*$.

Thus, there exists $F_{\nu_i} \in L^{m,p}(\mathbb{R}^n)$ with

$$(9) \quad F_{\nu_i} = f_{\nu_i} \text{ on } S_{\nu_i} \text{ and } J_{x_{\nu_i}}(F_{\nu_i}) = P_{\nu_i}; \text{ and}$$

$$(10) \quad \|F_{\nu_i}\|_{L^{m,p}(\mathbb{R}^n)} \simeq \|(f_{\nu_i}, P_{\nu_i})\|_{L^{m,p}(S_{\nu_i}; x_{\nu_i})} \simeq M_{\nu_i}(f_{\nu_i}, P_{\nu_i}).$$

Suppose that $Q \in CZ^\circ$ satisfies $Q \cap 100Q_\mu^\sharp = \emptyset$ and $(1.1)Q \cap 9Q_\mu^\sharp \neq \emptyset$. Hence, $\delta_Q \geq 100\delta_{Q_\mu^\sharp}$. However, this contradicts the second bullet point in Lemma 5.3. Therefore, $Q \notin CZ_\mu^\sharp \implies (1.1)Q \subset \mathbb{R}^n \setminus 9Q_\mu^\sharp$. Thus, by **(POU2)** we have

$$(11) \quad \theta_\nu \equiv 0 \text{ on } 9Q_\mu^\sharp \cap Q^\circ \text{ for } \nu \in \{1, \dots, \nu_{\max}\} \setminus \{\nu_1, \dots, \nu_K\}.$$

Define $F^\circ \in L^{m,p}(Q^\circ)$ by

$$F^\circ(x) := \sum_{i=1}^K F_{\nu_i}(x)\theta_{\nu_i}(x) + \sum_{\nu \notin \{\nu_1, \dots, \nu_K\}} R'(x)\theta_\nu(x) \quad (x \in Q^\circ).$$

From **(POU2)**, (9) and the definition of S_{ν_i} , we have $F_{\nu_i} = f_{\nu_i}$ on $\text{supp}(\theta_{\nu_i}) \cap E \cap 9Q_\mu^\sharp$. Hence, from **(POU5)** and (11), we have $F^\circ = f_\mu^\sharp$ on $E \cap 9Q_\mu^\sharp$ (recall that $E \subset (1/8)Q^\circ$).

From **(POU4)** and (9), we find that $J_{x_{\nu_i}}(F^\circ) = P_{\nu_i}$ for each $i = 1, \dots, K$. To summarize,

$$(12) \quad F^\circ = f_\mu^\sharp \text{ on } E \cap 9Q_\mu^\sharp \text{ and } J_{x_{\nu_i}}(F^\circ) = P_{x_{\nu_i}} \text{ for } i = 1, \dots, K.$$

We prepare to estimate the seminorm $\|F^\circ\|_{L^{m,p}(Q^\circ)}$ using Lemma 9.1.

Let $G_\nu := F_\nu$ when $\nu \in \{\nu_1, \dots, \nu_K\}$ and $G_\nu := R'$ when $\nu \in \{1, \dots, \nu_{\max}\} \setminus \{\nu_1, \dots, \nu_K\}$. Lemma 9.1 provides the estimate

$$\|F^\circ\|_{L^{m,p}(Q^\circ)}^p \lesssim \sum_{\nu=1}^{\nu_{\max}} \|G_\nu\|_{L^{m,p}(1.1Q_\nu)}^p + \sum_{\nu \leftrightarrow \nu'} |J_{x_\nu}(G_\nu) - J_{x_{\nu'}}(G_{\nu'})|_\nu^p.$$

The terms with $\nu \notin \{\nu_1, \dots, \nu_K\}$ and $\nu, \nu' \notin \{\nu_1, \dots, \nu_K\}$ in the first and second sums vanish, respectively. In view of (9), the remaining terms in the second sum are of the form $|P_\nu - P_{\nu'}|_\nu^p$ and $|P_\nu - R'|_\nu^p$ ($\nu, \nu' \in \{\nu_1, \dots, \nu_K\}$). (Here, we also use (8.4).) Thus,

$$\begin{aligned} \|F^\circ\|_{L^{m,p}(Q^\circ)}^p &\lesssim \sum_{i=1}^K \|F_{\nu_i}\|_{L^{m,p}(1.1Q_{\nu_i})}^p + \sum_{i,j=1}^K |P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p \\ &\lesssim \sum_{i=1}^K M_{\nu_i}(f_{\nu_i}, P_{\nu_i})^p + \sum_{i,j=1}^K |P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p \quad (\text{thanks to (10)}). \end{aligned}$$

Lemma 9.2 yields $F \in L^{m,p}(\mathbb{R}^n)$ such that $F = F^\circ$ on $0.99Q^\circ$ and $\|F\|_{L^{m,p}(\mathbb{R}^n)} \leq C\|F^\circ\|_{L^{m,p}(Q^\circ)}$. Therefore,

$$(13) \quad \|F\|_{L^{m,p}(\mathbb{R}^n)}^p \lesssim \sum_{i=1}^K M_{\nu_i}(f_{\nu_i}, P_{\nu_i})^p + \sum_{i,j=1}^K |P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p.$$

Recall that $E \cup E' \subset 0.99Q^\circ$ (see (4.2) and Lemma 8.2). Thus, (12) yields

$$(14) \quad F = f_\mu^\sharp \text{ on } E \cap 9Q_\mu^\sharp \text{ and } J_{x_{\nu_i}}(F) = P_{\nu_i} \text{ for } i = 1, \dots, K.$$

Let $H \in L^{m,p}(\mathbb{R}^n)$ be arbitrary with

$$(*) \quad H = f_\mu^\sharp \text{ on } E \cap 9Q_\mu^\sharp \text{ and } J_{x_{\nu_i}}(H) = P_{\nu_i}, \text{ for } i = 1, \dots, K.$$

From (10) and the definition of the trace seminorm, we have

$$M_{\nu_i}(f_{\nu_i}, P_{\nu_i})^p \simeq \|(f_{\nu_i}, P_{\nu_i})\|^p \leq \|H\|_{L^{m,p}(\mathbb{R}^n)}^p, \quad \text{for } i = 1, \dots, K,$$

while the Sobolev Inequality implies that

$$|P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p = \left| J_{x_{\nu_i}}(H) - J_{x_{\nu_j}}(H) \right|_{\nu_i}^p \lesssim \|H\|_{L^{m,p}(\mathbb{R}^n)}^p, \quad \text{for } i, j = 1, \dots, K.$$

(Here, we use (7) and $x_{\nu_i} \in Q_{\nu_i}$.) Summing these two estimates over $i, j \in \{1, \dots, K\}$ and using (7) provides a bound on the right-hand side of (13) by $C\|H\|_{L^{m,p}(\mathbb{R}^n)}^p$. We now take $E'_\mu := \{x_{\nu_1}, \dots, x_{\nu_K}\}$. Taking the infimum in the previous estimate, with respect to $H \in L^{m,p}(\mathbb{R}^n)$ that satisfy $(*)$, we have

$$\sum_{i=1}^K M_{\nu_i}(f_{\nu_i}, P_{\nu_i})^p + \sum_{i,j=1}^K |P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p \lesssim \|(f_\mu^\sharp, (P_{\nu_i})_{i=1}^K)\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; E'_\mu)}^p.$$

Together with (13) and (14), this tells us that

$$\sum_{i=1}^K M_{\nu_i}(f_{\nu_i}, P_{\nu_i})^p + \sum_{i,j=1}^K |P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p \simeq \|(f_\mu^\sharp, (P_{\nu_i})_{i=1}^K)\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; E'_\mu)}^p.$$

For every $R' \in \mathcal{P}$, the definition of the trace seminorm gives

$$\begin{aligned} & \|(f_\mu^\sharp, R')\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)}^p \\ &= \inf \left\{ \|(f_\mu^\sharp, (P_{\nu_i})_{i=1}^K)\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; E'_\mu)}^p : (P_{\nu_i})_{i=1}^K \in Wh(E'_\mu), P_{\nu_1} = R' \right\}. \end{aligned}$$

Thus, by the last two lines and (8), we have

$$\begin{aligned} & \|(f_\mu^\sharp, R')\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)}^p \\ & \simeq \inf \left\{ \sum_{i=1}^K \sum_{j=1}^{L_i} |\lambda_i^j(f_{\nu_i}, P_{\nu_i})|^p + \sum_{i,j=1}^K |P_{\nu_i} - P_{\nu_j}|_{\nu_i}^p : \right. \\ & \left. (P_{\nu_i})_{i=1}^K \in Wh(E'_\mu), P_{\nu_1} = R' \right\}. \end{aligned}$$

By an application of Lemma 11.1, we solve for $(P_{\nu_i})_{i=2}^K$ which depends linearly on (f_μ^\sharp, R') and minimizes the expression inside the infimum to within the multiplicative factor $C(p, k)$. Here we solve for $k := \dim(\mathcal{P}) \cdot (K - 1)$ real-valued variables. In view of (7), the constant $C(p, k)$ is universal, and hence

$$(15) \quad \|(f_\mu^\sharp, R')\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)}^p \simeq \sum_{i=1}^L |\lambda_i(f_\mu^\sharp, R')|^p,$$

for some integer L and some linear functionals $\lambda_1, \dots, \lambda_L$.

We apply Lemma 11.1 once again, solving for $R' = R'_\mu \in \mathcal{P}$ that depends linearly on (f_μ^\sharp, P) and minimizes the right-hand side of (15) to within a multiplicative factor $C(p, k')$, subject to the condition $[\partial^\alpha R'_\mu(x_\mu^\sharp) = \partial^\alpha P(x_\mu^\sharp)]$ for all $\alpha \in \mathcal{A}$. In this case, we solve for the remaining $k' := \dim(\mathcal{P}) - \#(\mathcal{A})$ coefficients of the polynomial R'_μ . Again, the constant $C(k', p)$ is universal. It follows that R'_μ satisfies (3) and (4). This concludes the proof of Lemma 11.2. \square

12. ESTIMATES FOR AUXILIARY POLYNOMIALS

Recall that for each $x \in Q^\circ$ we defined polynomials $(P_\alpha^x)_{\alpha \in \mathcal{A}}$ that satisfy (4.16)-(4.17d).

Lemma 12.1. *Let $Q \in CZ^\circ$ be fixed. Then, the following estimate holds:*

$$|\partial^\beta P_\alpha^y(y)| \leq C \delta_Q^{|\alpha| - |\beta|} \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}, y \in Q).$$

Proof. Let $E^+ := (E \cap 3Q^+) \cup \{y\}$, and let $x \in E^+$ be given. From (4.17a)-(4.17d), it follows that the collection $(P_\alpha^x)_{\alpha \in \mathcal{A}}$ forms an $(\mathcal{A}, x, C\epsilon, 1)$ -basis for $\sigma(x, E)$. Set $C' = 30^m C$. By definition of $(\mathcal{A}, x, C\epsilon, 1)$ -bases, $(P_\alpha^x)_{\alpha \in \mathcal{A}}$ also forms an $(\mathcal{A}, x, C'\epsilon, 30)$ -basis for $\sigma(x, E)$. Since $Q^+ \subset Q^\circ$ satisfies $\delta_{Q^+} \leq \delta_{Q^\circ} = 1$, we have that $(P_\alpha^x)_{\alpha \in \mathcal{A}}$ forms an $(\mathcal{A}, x, C'\epsilon, 30\delta_{Q^+})$ -basis for $\sigma(x, E)$, hence also for $\sigma(x, E \cap 3Q^+)$. (See Remark 1 in section 3.2.)

Define $\epsilon_1 = \min\{c_1, \epsilon_0/C_1\}$, where c_1 and C_1 are the constants from Lemma 3.3. For the sake of contradiction, suppose that

$$\max\{|\partial^\beta P_\alpha^x(x)| \cdot (30\delta_{Q^+})^{|\beta| - |\alpha|} : x \in E^+, \alpha \in \mathcal{A}, \beta \in \mathcal{M}\} > \epsilon_1^{-D-1}.$$

We assume that $\epsilon \leq \epsilon_1^{2D+2}/C'$. Then the hypotheses of Lemma 3.3 hold with parameters

$$\begin{aligned} & \left(\epsilon_1, \epsilon_2, \delta, \mathcal{A}, E_1, E_2, (\tilde{P}_\alpha^x)_{\alpha \in \mathcal{A}, x \in E_2} \right) \\ & := \left(\epsilon_1, C'\epsilon, 30\delta_{Q^+}, \mathcal{A}, E \cap 3Q^+, E^+, (P_\alpha^x)_{\alpha \in \mathcal{A}, x \in E^+} \right). \end{aligned}$$

Thus, there exists $\bar{\mathcal{A}} < \mathcal{A}$ such that $\sigma(x, E \cap 3Q^+)$ contains an $(\bar{\mathcal{A}}, x, C_1\epsilon_1, 30\delta_{Q^+})$ -basis for each $x \in E^+$. Thus, since $C_1\epsilon_1 \leq \epsilon_0$, we deduce that Q^+ is OK. This contradicts the fact that $Q \in CZ^\circ$ and completes the proof of Lemma 12.1. \square

13. ESTIMATES FOR LOCAL EXTENSIONS

Lemma 13.1. *Let $Q \in CZ^\circ \setminus \{Q_1\}$ and let x_Q be the representative point for Q . Then for every $P \in \mathcal{P}$,*

$$(1) \quad \|(0|_{E \cap 1.1Q}, P)\|_{L^{m,p}(E \cap 1.1Q; x_Q)} \leq \|(0|_E, P)\|_{L^{m,p}(E; x_Q)} \lesssim |P|_{x_Q, \delta_Q}.$$

Moreover, if $P \in \mathcal{P}$ satisfies $\partial^\alpha P(x_Q) = 0$ for all $\alpha \in \mathcal{A}$, then

$$(2) \quad \|(0|_{E \cap 9Q}, P)\|_{L^{m,p}(E \cap 9Q; x_Q)} \simeq |P|_{x_Q, \delta_Q}.$$

Proof. Since $Q \neq Q_1$ we have that $x_Q \neq x_1 = z$. Thus, by the properties of representative points listed in Lemma 8.2, there exists $\theta \in C_0^\infty(Q)$ with

$$(3) \quad \theta \equiv 0 \text{ in a neighborhood of } E;$$

$$(4) \quad \theta \equiv 1 \text{ in a neighborhood of } x_Q; \text{ and}$$

$$(5) \quad |\partial^\alpha \theta| \lesssim \delta_Q^{-|\alpha|} \text{ when } |\alpha| \leq m.$$

Define $H := \theta P$. By a straightforward calculation using $\text{supp}(\theta) \subset Q$ and (5), we have

$$\|H\|_{L^{m,p}(\mathbb{R}^n)}^p \lesssim \sum_{|\alpha| \leq m-1} |\partial^\alpha P(x_Q)|^p \delta_Q^{n - (m - |\alpha|)p} = \left[|P|_{x_Q, \delta_Q} \right]^p.$$

Moreover, (3) and (4) show that H is a competitor in the infimum that defines

$$\|(0|_E, P)\|_{L^{m,p}(E; x_Q)} := \inf \{ \|G\|_{L^{m,p}(\mathbb{R}^n)} : G|_E = 0, J_{x_Q} G = P \},$$

and therefore

$$\|(0|_E, P)\|_{L^{m,p}(E;x_Q)} \leq \|H\|_{L^{m,p}(\mathbb{R}^n)} \lesssim |P|_{x_Q, \delta_Q}.$$

Finally, by definition of the trace seminorm we have

$$(6) \quad \begin{aligned} \|(0|_{E \cap (1.1)Q}, P)\|_{L^{m,p}(E \cap 1.1Q;x_Q)} &\leq \|(0|_{E \cap 9Q}, P)\|_{L^{m,p}(E \cap 9Q;x_Q)} \\ &\leq \|(0|_E, P)\|_{L^{m,p}(E;x_Q)}. \end{aligned}$$

This completes the proof of (1).

To prove (2), we fix $P \in \mathcal{P}$ with

$$(7) \quad \partial^\alpha P(x_Q) = 0 \text{ for all } \alpha \in \mathcal{A},$$

and establish the reverse inequality

$$(8) \quad \|(0|_{E \cap 9Q}, P)\|_{L^{m,p}(E \cap 9Q;x_Q)} \geq c|P|_{x_Q, \delta_Q}.$$

From the definition of $\sigma(\cdot, \cdot)$ and the definition of the trace seminorm, we have

$$(9) \quad \{P \in \mathcal{P} : \|(0|_{E \cap 9Q}, P)\|_{L^{m,p}(E \cap 9Q;x_Q)} \leq 1/2\} \subset \sigma(x_Q, E \cap 9Q).$$

Let $\epsilon_1 > 0$ be some small universal constant to be determined by the end of the proof. For the sake of contradiction, suppose that (8) fails with $c = \epsilon_1^{D+1}$ for some $P \in \mathcal{P}$ that satisfies (7). Upon rescaling P and using (9), we see that

$$(10) \quad P \in C\epsilon_1^{D+1} \cdot \sigma(x_Q, E \cap 9Q) \text{ and}$$

$$(11) \quad \max_{\beta \in \mathcal{M}} \left\{ |\partial^\beta P(x_Q)| \delta_Q^{n/p+|\beta|-m} \right\} = 1.$$

For each integer $\ell \geq 0$, we define

$$\Delta_\ell = \left\{ \alpha \in \mathcal{M} : |\partial^\alpha P(x_Q)| \delta_Q^{n/p+|\alpha|-m} \in (\epsilon_1^\ell, 1] \right\}.$$

Note that $\Delta_\ell \subset \Delta_{\ell+1}$ for each $\ell \geq 0$ and that $\Delta_\ell \neq \emptyset$ for $\ell \geq 1$ by (11). Since \mathcal{M} contains D elements, there exists $0 \leq \ell_* \leq D$ with $\Delta_{\ell_*} = \Delta_{\ell_*+1} \neq \emptyset$. Let $\bar{\alpha} \in \mathcal{M}$ be the maximal element of Δ_{ℓ_*} . Thus,

$$(12) \quad |\partial^{\bar{\alpha}} P(x_Q)| \delta_Q^{n/p+|\bar{\alpha}|-m} \geq \epsilon_1^{\ell_*}.$$

Moreover, for every $\beta \in \mathcal{M}$ with $\beta > \bar{\alpha}$, we have $\beta \notin \Delta_{\ell_*} = \Delta_{\ell_*+1}$, and hence

$$(13) \quad |\partial^\beta P(x_Q)| \delta_Q^{n/p+|\beta|-m} \leq \epsilon_1^{\ell_*+1} \quad (\beta \in \mathcal{M}, \beta > \bar{\alpha}).$$

Thanks to (12), we may define $P_{\bar{\alpha}} := [\partial^{\bar{\alpha}} P(x_Q)]^{-1} P$. Hence,

$$(14) \quad \partial^{\bar{\alpha}} P_{\bar{\alpha}}(x_Q) = 1,$$

while (10), (12) and $\ell_* \leq D$ imply that

$$P_{\bar{\alpha}} \in C\epsilon_1 \delta_Q^{|\bar{\alpha}|+n/p-m} \cdot \sigma(x_Q, E \cap 9Q).$$

Thus, there exists $\varphi_{\bar{\alpha}} \in L^{m,p}(\mathbb{R}^n)$ with

$$(15) \quad \varphi_{\bar{\alpha}} = 0 \text{ on } E \cap 9Q;$$

$$(16) \quad J_{x_Q}(\varphi_{\bar{\alpha}}) = P_{\bar{\alpha}}; \text{ and}$$

$$(17) \quad \|\varphi_{\bar{\alpha}}\|_{L^{m,p}(\mathbb{R}^n)} \leq C\epsilon_1 \delta_Q^{|\bar{\alpha}|+n/p-m}.$$

From (11) and (12) and from (12) and (13), we obtain

$$(18) \quad |\partial^\beta P_{\bar{\alpha}}(x_Q)| \leq \epsilon_1^{-D} \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}) \text{ and}$$

$$(19) \quad |\partial^\beta P_{\bar{\alpha}}(x_Q)| \leq \epsilon_1 \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}, \beta > \bar{\alpha}),$$

while (7) and (12) yield

$$(20) \quad \bar{\alpha} \notin \mathcal{A}.$$

Let $x \in E \cap 9Q$ be arbitrary. Define $P_{\bar{\alpha}}^x = J_x(\varphi_{\bar{\alpha}})$. Thus,

$$(21) \quad \begin{aligned} & \left| \partial^\beta P_{\bar{\alpha}}^x(x) - \sum_{|\gamma| \leq m-1-|\beta|} \frac{1}{\gamma!} \partial^{\beta+\gamma} P_{\bar{\alpha}}(x_Q) \cdot (x-x_Q)^\gamma \right| = |\partial^\beta P_{\bar{\alpha}}^x(x) - \partial^\beta P_{\bar{\alpha}}(x)| \\ & = |\partial^\beta \varphi_{\bar{\alpha}}(x) - \partial^\beta J_{x_Q}(\varphi_{\bar{\alpha}})(x)| \leq C' \epsilon_1 \cdot \delta_Q^{|\bar{\alpha}+n/p-m} \cdot |x-x_Q|^{m-n/p-|\beta|} \\ & \leq C'' \epsilon_1 \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}). \end{aligned}$$

(Here, we have used (16), (17) and the standard Sobolev Inequality, and $|x-x_Q| \leq C\delta_Q$.) By (15) and (17), by inserting (14) and (19) into (21), and by inserting (18) into (21), we have

$$(22) \quad P_{\bar{\alpha}}^x \in C_3 \epsilon_1 \cdot \delta_Q^{|\bar{\alpha}+n/p-m} \cdot \sigma(x, E \cap 9Q);$$

$$(23) \quad |\partial^\beta P_{\bar{\alpha}}^x(x) - \delta_{\beta\bar{\alpha}}| \leq C_3 \epsilon_1 \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}, \beta \geq \bar{\alpha}); \text{ and}$$

$$(24) \quad |\partial^\beta P_{\bar{\alpha}}^x(x)| \leq C_3 \epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}).$$

We recall the defining properties (4.17a)-(4.17d) of the polynomials P_α^x :

$$(25a) \quad P_\alpha^x \in C' \epsilon \cdot \sigma(x, E) \quad (\alpha \in \mathcal{A});$$

$$(25b) \quad \partial^\beta P_\alpha^x(x) = \delta_{\alpha\beta} \quad (\alpha, \beta \in \mathcal{A});$$

$$(25c) \quad |\partial^\beta P_\alpha^x(x)| \leq C' \epsilon \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}, \beta > \alpha); \text{ and}$$

$$(25d) \quad |\partial^\beta P_\alpha^x(x)| \leq C' \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}).$$

Let $Q' \in CZ^\circ$ be such that $x \in Q'$. Then $9Q \cap Q' \neq \emptyset$, since $x \in 9Q$. Thus $\delta_{Q'} \leq 50\delta_Q$, thanks to the second bullet point in Lemma 5.3. Also, $|\partial^\beta P_\alpha^x(x)| \leq C \delta_Q^{|\alpha|-|\beta|}$ for $\alpha \in \mathcal{A}$, $\beta \in \mathcal{M}$, thanks to Lemma 12.1. Thus, we have $|\partial^\beta P_\alpha^x(x)| \leq C' \delta_Q^{|\alpha|-|\beta|}$ for every $\alpha \in \mathcal{A}$ and $\beta \in \mathcal{M}$ with $|\beta| \leq |\alpha|$. In combination with (25d), this gives

$$(26) \quad |\partial^\beta P_\alpha^x(x)| \leq C' \delta_Q^{|\alpha|-|\beta|} \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}).$$

(Here, we use the fact that $\delta_Q \leq 1$ since $Q \subset Q^\circ = (0, 1]^n$.)

Define the polynomial

$$\tilde{P}_{\bar{\alpha}}^x = P_{\bar{\alpha}}^x - \sum_{\alpha \in \mathcal{A}, \alpha < \bar{\alpha}} [\partial^\alpha P_{\bar{\alpha}}^x(x)] \cdot P_\alpha^x.$$

From (22), (24), and (25a), we obtain

$$(27) \quad \begin{aligned} \tilde{P}_{\bar{\alpha}}^x & \in \left[C_3 \epsilon_1 \cdot \delta_Q^{|\bar{\alpha}+n/p-m} + \sum_{\alpha \in \mathcal{A}, \alpha < \bar{\alpha}} \left[C_3 \epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\alpha|} \right] \cdot C' \epsilon \right] \cdot \sigma(x, E \cap 9Q) \\ & \subset C_4 \left[\epsilon_1 + \epsilon \epsilon_1^{-D} \right] \cdot \delta_Q^{|\bar{\alpha}+n/p-m} \cdot \sigma(x, E \cap 9Q). \end{aligned}$$

(Here, we have also used the fact that $\sigma(x, E) \subset \sigma(x, E \cap 9Q)$ and $|\alpha| \leq m-1 < m-n/p$.)

For each $\beta \in \mathcal{A}$, $\beta < \bar{\alpha}$, by (25b) we have

$$(28) \quad \partial^\beta \tilde{P}_\alpha^x(x) = \partial^\beta P_\alpha^x(x) - \partial^\beta P_{\bar{\alpha}}^x(x) = 0.$$

Let $\beta \in \mathcal{M}$ with $\beta \geq \bar{\alpha}$ be given. Note that if $\alpha \in \mathcal{A}$ and $\alpha < \bar{\alpha}$, then $\alpha < \beta$. We estimate

$$(29) \quad \begin{aligned} |\partial^\beta \tilde{P}_\alpha^x(x) - \delta_{\beta\bar{\alpha}}| &\leq |\partial^\beta P_\alpha^x(x) - \delta_{\beta\bar{\alpha}}| + \sum_{\alpha \in \mathcal{A}, \alpha < \bar{\alpha}} |\partial^\alpha P_\alpha^x(x)| \cdot |\partial^\beta P_\alpha^x(x)| \\ &\leq C_3 \epsilon_1 \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} + \sum_{\alpha \in \mathcal{A}, \alpha < \bar{\alpha}} \left[C_3 \epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\alpha|} \right] \cdot C' \epsilon \\ &\quad \text{(thanks to (23), (24), (25c))} \\ &\leq C_4 [\epsilon_1 + \epsilon_1^{-D} \epsilon] \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad \text{(since } \alpha < \beta \implies |\alpha| \leq |\beta|). \end{aligned}$$

On the other hand, consider an arbitrary $\beta \in \mathcal{M}$. We estimate

$$(30) \quad \begin{aligned} |\partial^\beta \tilde{P}_\alpha^x(x)| &\leq |\partial^\beta P_\alpha^x(x)| + \sum_{\alpha \in \mathcal{A}, \alpha < \bar{\alpha}} |\partial^\alpha P_\alpha^x(x)| \cdot |\partial^\beta P_\alpha^x(x)| \\ &\leq C_3 \epsilon_1^{-D} \delta_Q^{|\bar{\alpha}|-|\beta|} + \sum_{\alpha \in \mathcal{A}, \alpha < \bar{\alpha}} \left[C_3 \epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\alpha|} \right] \cdot C' \delta_Q^{|\alpha|-|\beta|} \\ &\quad \text{(thanks to (24) and (26))} \\ &\leq C_4 \epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\beta|}. \end{aligned}$$

Define $\bar{\mathcal{A}} = \{\alpha \in \mathcal{A} : \alpha < \bar{\alpha}\} \cup \{\bar{\alpha}\}$. From (20) and the definition of the order on multi-index sets, we have

$$(31) \quad \bar{\mathcal{A}} < \mathcal{A}.$$

For small enough ϵ and ϵ_1 such that $\epsilon \leq \epsilon_1^{D+1}$, by (29) we have $|\partial^{\bar{\alpha}} \tilde{P}_\alpha^x(x)| \geq 1/2$. Thus, the following polynomial is well defined:

$$\hat{P}_\alpha^x = \left[\partial^{\bar{\alpha}} \tilde{P}_\alpha^x(x) \right]^{-1} \cdot \tilde{P}_\alpha^x.$$

Thanks to (27)-(30), the following properties are immediate:

$$(32a) \quad \hat{P}_\alpha^x \in 2C_4 \cdot [\epsilon_1 + \epsilon_1^{-D}] \cdot \delta_Q^{|\bar{\alpha}+n/p-m} \sigma(x, E \cap 9Q);$$

$$(32b) \quad \partial^\beta \hat{P}_\alpha^x(x) = \delta_{\beta\bar{\alpha}} \quad (\beta \in \bar{\mathcal{A}});$$

$$(32c) \quad |\partial^\beta \hat{P}_\alpha^x(x)| \leq 2C_4 \cdot [\epsilon_1 + \epsilon_1^{-D} \epsilon] \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}, \beta > \bar{\alpha}); \text{ and}$$

$$(32d) \quad |\partial^\beta \hat{P}_\alpha^x(x)| \leq 2C_4 \cdot \epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\beta \in \mathcal{M}).$$

For each $\alpha \in \bar{\mathcal{A}} \setminus \{\bar{\alpha}\}$, we define the polynomial

$$\hat{P}_\alpha^x = P_\alpha^x - [\partial^{\bar{\alpha}} P_\alpha^x(x)] \cdot \hat{P}_\alpha^x.$$

Then, we have

$$(33) \quad \partial^{\bar{\alpha}} \hat{P}_\alpha^x(x) = \partial^{\bar{\alpha}} P_\alpha^x(x) - [\partial^{\bar{\alpha}} P_\alpha^x(x)] \cdot \partial^{\bar{\alpha}} \hat{P}_\alpha^x(x) = 0 \quad \text{(thanks to (32b))}.$$

For $\beta \in \bar{\mathcal{A}} \setminus \{\bar{\alpha}\}$, we have

$$(34) \quad \partial^\beta \hat{P}_\alpha^x(x) = \partial^\beta P_\alpha^x(x) - [\partial^{\bar{\alpha}} P_\alpha^x(x)] \cdot \partial^\beta \hat{P}_\alpha^x(x) = \delta_{\beta\alpha} \quad \text{(thanks to (25b), (32b))}.$$

Note that $\alpha < \bar{\alpha}$, because $\alpha \in \bar{\mathcal{A}} \setminus \{\bar{\alpha}\}$. Thus, for each $\beta \in \mathcal{M}$ with $\beta > \alpha$, we obtain

$$(35) \quad \begin{aligned} |\partial^\beta \widehat{P}_\alpha^x(x)| &\leq C'\epsilon + [C'\epsilon] \cdot 2C_4\epsilon_1^{-D} \cdot \delta_Q^{|\bar{\alpha}|-|\beta|} \quad (\text{thanks to (25c) and (32d)}) \\ &\leq C_5\epsilon \cdot \epsilon_1^{-D} \cdot \delta_Q^{|\alpha|-|\beta|} \quad (\text{since } |\alpha| \leq |\bar{\alpha}|, |\alpha| \leq |\beta|). \end{aligned}$$

Finally, from (25a), (25d), and (32a), we obtain

$$\widehat{P}_\alpha^x \in \left[C'\epsilon + [C'] \cdot 2C_4 \cdot (\epsilon_1 + \epsilon\epsilon_1^{-D}) \cdot \delta_Q^{|\bar{\alpha}|+n/p-m} \right] \cdot \sigma(x, E \cap 9Q).$$

We may assume that $\epsilon \leq \epsilon_1$. Note that $|\alpha| \leq |\bar{\alpha}|$, $|\alpha| + n/p - m \leq 0$ and $\delta_Q \leq 1$. Therefore,

$$(36) \quad \widehat{P}_\alpha^x \in C_5 \cdot [\epsilon_1 + \epsilon\epsilon_1^{-D}] \cdot \delta_Q^{|\alpha|+n/p-m} \cdot \sigma(x, E \cap 9Q).$$

Recall that $x \in E \cap 9Q$ was arbitrary. Thus, from (32a), (32b), (32c) and (33)-(36), it follows that $(\widehat{P}_\alpha^x)_{\alpha \in \bar{\mathcal{A}}}$ forms an $(\bar{\mathcal{A}}, x, C_6 \cdot [\epsilon_1 + \epsilon_1^{-D}\epsilon], \delta_{Q^+})$ -basis for $\sigma(x, E \cap 9Q)$ for each $x \in E \cap 9Q$, and hence for $\sigma(x, E \cap 3Q^+)$ for each $x \in E \cap 3Q^+$. (Here, we use the fact that $E \cap 3Q^+ \subset E \cap 9Q$.)

Fix ϵ_1 small enough so that the preceding arguments hold and so that $\epsilon_1 \leq \epsilon_0/2C_6$. We may assume that $\epsilon \leq \epsilon_1^{D+1}$. Therefore, $\sigma(x, E \cap 3Q^+)$ contains an $(\bar{\mathcal{A}}, x, \epsilon_0, \delta_{Q^+})$ -basis for each $x \in E \cap 3Q^+$. Since $\bar{\mathcal{A}} < \mathcal{A}$, by definition the cube Q^+ is OK, which contradicts the hypothesis that $Q \in CZ^\circ$. This completes the proof of (8), which was the unverified inequality in (2). The proof of Lemma 13.1 is now complete. \square

14. THE JET OF A NEAR OPTIMAL EXTENSION I

14.1. Another Sobolev Inequality. Proposition 2 produces for each Q_ν an associated keystone cube $\mathcal{K}(Q_\nu) = Q_{\kappa(\nu)}^\sharp$ and a finite sequence of CZ cubes $\mathcal{S}_\nu = (Q_{\nu,1}, \dots, Q_{\nu,L_\nu})$ such that

$$(1) \quad \begin{aligned} Q_\nu &= Q_{\nu,1} \leftrightarrow \dots \leftrightarrow Q_{\nu,L_\nu} = Q_{\kappa(\nu)}^\sharp, \text{ with} \\ \delta_{Q_{\nu,k}} &\leq C(1-c)^{k-j} \delta_{Q_{\nu,j}} \text{ for all } 1 \leq j \leq k \leq L_\nu, \text{ where} \\ C > 0 \text{ and } 0 < c < 1 &\text{ are universal constants.} \end{aligned}$$

Lemma 14.1. *Let $F \in L^{m,p}(Q^\circ)$. Then*

$$\sum_{\nu=1}^{\nu_{\max}} \left| J_{x_\nu}(F) - J_{x_{\kappa(\nu)}^\sharp}(F) \right|_\nu^p \lesssim \|F\|_{L^{m,p}(Q^\circ)}^p.$$

Proof. By applying (8.3), we have

$$(2) \quad \begin{aligned} X &:= \sum_{\nu=1}^{\nu_{\max}} \left| J_{x_\nu}(F) - J_{x_{\kappa(\nu)}^\sharp}(F) \right|_\nu^p \\ &\lesssim \sum_{\nu=1}^{\nu_{\max}} \sum_{|\alpha| \leq m-1} \left| \partial^\alpha \left[J_{x_\nu}(F) - J_{x_{\kappa(\nu)}^\sharp}(F) \right] (x_{\kappa(\nu)}^\sharp) \right|_\nu^p \delta_{Q_\nu}^{n-(m-|\alpha|)p}. \end{aligned}$$

Let $x_{\nu,k}$ denote the representative point for $Q_{\nu,k}$. We fix some universal constant $\epsilon' \in (0, 1-n/p)$. Note that $x_{\nu,L_\nu} = x_{\kappa(\nu)}^\sharp$, since $Q_{\nu,L_\nu} = Q_{\kappa(\nu)}^\sharp$. Thus the right-hand

side of (2) is given by

$$\begin{aligned}
& \sum_{\nu=1}^{\nu_{\max}} \sum_{|\alpha| \leq m-1} \delta_{Q_\nu}^{n-(m-|\alpha|)p} \left| \sum_{k=1}^{L_\nu-1} \partial^\alpha [J_{x_{\nu,k}}(F) - J_{x_{\nu,k+1}}(F)] (x_{\kappa(\nu)}^\#) \cdot \delta_{Q_{\nu,k}}^{\frac{n}{p}-(m-|\alpha|)+\epsilon'} \right. \\
& \qquad \qquad \qquad \left. \cdot \delta_{Q_{\nu,k}}^{-\frac{n}{p}+(m-|\alpha|)-\epsilon'} \right|^p \\
& \leq \sum_{\nu=1}^{\nu_{\max}} \sum_{|\alpha| \leq m-1} \delta_{Q_\nu}^{n-(m-|\alpha|)p} \left[\sum_{k=1}^{L_\nu-1} \left| \partial^\alpha [J_{x_{\nu,k}}(F) - J_{x_{\nu,k+1}}(F)] (x_{\kappa(\nu)}^\#) \right|^p \right. \\
& \qquad \qquad \qquad \left. \cdot \delta_{Q_{\nu,k}}^{n-(m-|\alpha|)p+\epsilon'p} \right] \\
& \qquad \qquad \qquad \cdot \left[\sum_{k=1}^{L_\nu-1} \delta_{Q_{\nu,k}}^{-\frac{np'}{p}+(m-|\alpha|)p'-\epsilon'p'} \right]^{p/p'} \\
& \left(\text{by Hölder's inequality; here, } p' \text{ is the dual exponent to } p \text{ so that } \frac{1}{p'} + \frac{1}{p} = 1 \right).
\end{aligned}$$

From (1), we have $\delta_{Q_{\nu,k}} \leq C(1-c)^k \delta_{Q_\nu}$. Thus, using $\epsilon' < 1 - n/p$, we obtain

(3)

$$\begin{aligned}
X & \lesssim \sum_{\nu=1}^{\nu_{\max}} \sum_{|\alpha| \leq m-1} \delta_{Q_\nu}^{n-(m-|\alpha|)p} \cdot \delta_{Q_\nu}^{-n+(m-|\alpha|)p-\epsilon'p} \\
& \quad \cdot \left[\sum_{k=1}^{L_\nu-1} \left| \partial^\alpha [J_{x_{\nu,k}}(F) - J_{x_{\nu,k+1}}(F)] (x_{\kappa(\nu)}^\#) \right|^p \cdot \delta_{Q_{\nu,k}}^{n-(m-|\alpha|)p+\epsilon'p} \right] \\
& = \sum_{\nu=1}^{\nu_{\max}} \delta_{Q_\nu}^{-\epsilon'p} \sum_{k=1}^{L_\nu-1} \sum_{|\alpha| \leq m-1} \left| \partial^\alpha [J_{x_{\nu,k}}(F) - J_{x_{\nu,k+1}}(F)] (x_{\kappa(\nu)}^\#) \right|^p \cdot \delta_{Q_{\nu,k}}^{n-(m-|\alpha|)p+\epsilon'p} \\
& = \sum_{\nu=1}^{\nu_{\max}} \delta_{Q_\nu}^{-\epsilon'p} \sum_{k=1}^{L_\nu-1} \delta_{Q_{\nu,k}}^{\epsilon'p} \left\{ \sum_{|\alpha| \leq m-1} \left| \partial^\alpha [J_{x_{\nu,k}}(F) - J_{x_{\nu,k+1}}(F)] (x_{\kappa(\nu)}^\#) \right|^p \right. \\
& \qquad \qquad \qquad \left. \cdot \delta_{Q_{\nu,k}}^{n-(m-|\alpha|)p} \right\}.
\end{aligned}$$

Thanks to (1) it follows that $|x_{\nu,k} - x_{\kappa(\nu)}^\#| \leq C\delta_{Q_{\nu,k}}$ (recall that $x_{\nu,k} \in Q_{\nu,k}$ and $x_{\kappa(\nu)}^\# \in Q_{\kappa(\nu)}^\#$). Therefore, continuing from (3) and applying (2.2), we have

$$\begin{aligned}
(4) \quad X & \lesssim \sum_{\nu=1}^{\nu_{\max}} \delta_{Q_\nu}^{-\epsilon'p} \sum_{k=1}^{L_\nu-1} \delta_{Q_{\nu,k}}^{\epsilon'p} \left\{ \sum_{|\alpha| \leq m-1} \left| \partial^\alpha [J_{x_{\nu,k}}(F) - J_{x_{\nu,k+1}}(F)] (x_{\nu,k}) \right|^p \delta_{Q_{\nu,k}}^{n-(m-|\alpha|)p} \right\} \\
& = \sum_{\bar{\nu} \leftrightarrow \bar{\nu}'} \sum_{\nu=1}^{\nu_{\max}} \sum_{k=1}^{L_\nu-1} \delta_{Q_\nu}^{-\epsilon'p} \delta_{Q_{\bar{\nu}'}}^{\epsilon'p} |J_{x_{\bar{\nu}}}(F) - J_{x_{\bar{\nu}'}}(F)|_{x_{\bar{\nu}}, \delta_{Q_{\bar{\nu}'}}}^p \mathbb{1}_{Q_{\bar{\nu}}=Q_{\nu,k} \wedge Q_{\bar{\nu}'}=Q_{\nu,k+1}}.
\end{aligned}$$

Thanks to (1), any given $Q_{\bar{\nu}}$ can arise as $Q_{\nu,k}$ (for fixed ν) for at most C distinct k . Thus, from (4) we obtain

$$X \lesssim \sum_{\bar{\nu} \leftrightarrow \bar{\nu}'} \sum_{\nu=1}^{\nu_{\max}} (\delta_{Q_{\bar{\nu}}}/\delta_{Q_{\nu}})^{\epsilon' p} \cdot |J_{x_{\bar{\nu}}}(F) - J_{x_{\bar{\nu}'}}(F)|_{x_{\bar{\nu}}, \delta_{Q_{\bar{\nu}}}}^p \cdot \mathbb{1}_{\exists k \text{ s.t. } Q_{\bar{\nu}}=Q_{\nu,k} \wedge Q_{\bar{\nu}'}=Q_{\nu,k+1}}.$$

If there exists k such that $Q_{\bar{\nu}} = Q_{\nu,k}$, then $Q_{\bar{\nu}} \subset CQ_{\nu}$, as follows from (1). Therefore, we obtain

$$X \lesssim \sum_{\bar{\nu} \leftrightarrow \bar{\nu}'} |J_{x_{\bar{\nu}}}(F) - J_{x_{\bar{\nu}'}}(F)|_{x_{\bar{\nu}}, \delta_{Q_{\bar{\nu}}}}^p \cdot \sum \left\{ \left(\frac{\delta_{Q_{\bar{\nu}}}}{\delta_Q} \right)^{\epsilon' p} : \text{dyadic } Q \text{ such that } Q_{\bar{\nu}} \subset CQ \right\}.$$

The inner sum is at most C , for each fixed $\bar{\nu}$. Hence,

$$\begin{aligned} X &\lesssim \sum_{\bar{\nu} \leftrightarrow \bar{\nu}'} |J_{x_{\bar{\nu}}}(F) - J_{x_{\bar{\nu}'}}(F)|_{x_{\bar{\nu}}, \delta_{Q_{\bar{\nu}}}}^p \lesssim \sum_{\bar{\nu} \leftrightarrow \bar{\nu}'} \|F\|_{L^{m,p}([1.1Q_{\bar{\nu}} \cup 1.1Q_{\bar{\nu}'}] \cap Q^{\circ})}^p \\ &\quad \text{(thanks to (2.3); we also use } x_{\bar{\nu}} \in Q_{\bar{\nu}}, x_{\bar{\nu}'} \in Q_{\bar{\nu}'} \\ &\quad \text{and good geometry of the CZ cubes)} \\ &\lesssim \|F\|_{L^{m,p}(Q^{\circ})}^p \\ &\quad \text{(thanks to the first bullet point in Lemma 5.3} \\ &\quad \text{and good geometry of the CZ cubes).} \end{aligned}$$

This completes the proof of Lemma 14.1. □

14.2. New constraints on extensions. Every Whitney field $\vec{P} \in Wh(E')$ has a natural restriction, denoted $\vec{P}^{\sharp} \in Wh(E^{\sharp})$, to the keystone representative points $E^{\sharp} = \{x_{\mu}^{\sharp} : \mu = 1, \dots, \mu_{\max}\}$. That is,

$$(5) \quad P_{\mu}^{\sharp} = P_{\nu}, \text{ where } \nu \in \{1, \dots, \nu_{\max}\} \text{ is such that } Q_{\nu} = Q_{\mu}^{\sharp}.$$

Definition 13. Let $\vec{P} = (P_{\nu})_{\nu=1}^{\nu_{\max}} \in Wh(E')$ and $P \in \mathcal{P}$. We say that \vec{P} is Coherent with P provided that

$$P_1 = P, \text{ and } \partial^{\alpha} P_{\mu}^{\sharp}(x_{\mu}^{\sharp}) = \partial^{\alpha} P(x_{\mu}^{\sharp}) \text{ for } \mu = 1, \dots, \mu_{\max}, \text{ and } \alpha \in \mathcal{A}.$$

We say that \vec{P} is Constant Along Paths provided that

$$P_{\nu} = P_{\kappa(\nu)}^{\sharp} \text{ for } \nu = 2, \dots, \nu_{\max}.$$

Similarly, we say that a function $F \in L^{m,p}(\mathbb{R}^n)$ is Coherent with P when $J_{E'}(F)$ is Coherent with P , while we say that F is Constant Along Paths when $J_{E'}(F)$ is Constant Along Paths.

The main result of this section states that there exist near-optimal extensions of the data (f, P) that are Coherent with P and Constant Along Paths.

Proposition 3. *Given $(f, P) \in L^{m,p}(E; z)$, there exists $\widehat{F} \in L^{m,p}(\mathbb{R}^n)$ with the following properties:*

- (i) \widehat{F} extends the data (f, P) .
- (ii) $\|\widehat{F}\|_{L^{m,p}(\mathbb{R}^n)} \leq C \cdot \|(f, P)\|_{L^{m,p}(E; z)}$.
- (iii) \widehat{F} is Coherent with P .
- (iv) \widehat{F} is Constant Along Paths.

Proof. By subtracting $P|_E$ from f and subtracting P from the \widehat{F} that we seek, we may assume that $P = 0$. For the remainder of the proof of Proposition 3, $\|(\cdot, \cdot)\|$ denotes the trace seminorm on $L^{m,p}(E; z)$.

First we draw some immediate conclusions from subsection 4.1. Denote the $(C, C\epsilon)$ near-triangular matrix $A^\nu = A^{x_\nu}$ from (4.16) for each $\nu = 1, \dots, \nu_{\max}$. Then (4.7)-(4.9) and (4.16)-(4.17d) imply that the following properties are satisfied by the function

$$(6) \quad \varphi_{\nu, \alpha} := \sum_{\gamma \in \mathcal{A}} A_{\alpha\gamma}^\nu \cdot \varphi_\gamma \quad (\alpha \in \mathcal{A}):$$

$$(7) \quad \varphi_{\nu, \alpha} = 0 \text{ on } E,$$

$$(8) \quad J_{x_\nu}(\varphi_{\nu, \alpha}) = P_\alpha^{x_\nu}, \text{ and}$$

$$(9) \quad \partial^\beta \varphi_{\nu, \alpha}(x_\nu) = \delta_{\alpha\beta} \quad (\beta \in \mathcal{A}).$$

Note that (4.9) states that

$$(10) \quad \|\varphi_\gamma\|_{L^{m,p}(\mathbb{R}^n)} \leq \epsilon \quad (\gamma \in \mathcal{A}).$$

From (8) and Lemma 12.1, we also have

$$(11) \quad |\partial^\beta \varphi_{\nu, \alpha}(x_\nu)| = |\partial^\beta P_\alpha^{x_\nu}(x_\nu)| \leq C \delta_{Q_\nu}^{|\alpha| - |\beta|} \quad (\alpha \in \mathcal{A}, \beta \in \mathcal{M}, \nu = 1, \dots, \nu_{\max}).$$

Since the inverse of a near-triangular matrix is near-triangular with comparable parameters, by (6) we have

$$(12) \quad \varphi_{\nu', \alpha} = \sum_{\beta \in \mathcal{A}} \omega_{\alpha\beta}^{\nu\nu'} \varphi_{\nu, \beta}, \quad \text{where } |\omega_{\alpha\beta}^{\nu\nu'}| \leq C \text{ for } \alpha, \beta \in \mathcal{A} \text{ and } \nu, \nu' \in \{1, \dots, \nu_{\max}\}.$$

Let $F \in L^{m,p}(\mathbb{R}^n)$ satisfy

$$(13) \quad F = f \text{ on } E \text{ and } J_z(F) = 0 \text{ and}$$

$$(14) \quad \|F\|_{L^{m,p}(\mathbb{R}^n)} \leq 2\|(f, 0)\|.$$

We first modify F to construct \widetilde{F} that is Coherent with the zero polynomial. Since $J_z(F) = 0$ and $|z - x_\nu| \leq 1$ for each ν , the Sobolev Inequality (2.3) yields the basic estimate:

$$(15) \quad |\partial^\alpha F(x_\nu)| = |\partial^\alpha (F - J_z(F))(x_\nu)| \lesssim \|F\|_{L^{m,p}(\mathbb{R}^n)} \quad (\alpha \in \mathcal{M}, \nu = 1, \dots, \nu_{\max}).$$

We define

$$(16) \quad F_\nu := F - \sum_{\alpha \in \mathcal{A}} \partial^\alpha F(x_\nu) \varphi_{\nu, \alpha} \quad \text{for each } \nu = 1, \dots, \nu_{\max}.$$

Therefore,

$$(17) \quad F_\nu \stackrel{(7),(13)}{=} f \text{ on } E \text{ and } \partial^\beta F_\nu(x_\nu) \stackrel{(9)}{=} 0 \text{ for all } \beta \in \mathcal{A}.$$

Note that $x_1 = z$. Thanks to (13) and (16), it follows that $F_1 = F$.

Recall the partition of unity $\{\theta_\nu\}_{\nu=1}^{\nu_{\max}}$ that satisfies **(POU1)**-**(POU5)**. We define $\tilde{F} \in L^{m,p}(Q^\circ)$ by

$$\tilde{F}(x) = \sum_{\nu=1}^{\nu_{\max}} F_\nu(x)\theta_\nu(x) \quad (x \in Q^\circ).$$

The following properties hold:

$$(18) \quad \begin{aligned} \tilde{F} &= f \text{ on } E && \text{(thanks to **(POU5)** and (17)),} \\ J_z \tilde{F} &= J_{x_1} \tilde{F} = J_{x_1} F_1 = J_{x_1} F = 0 && \text{(thanks to **(POU4)** and (13)),} \\ \partial^\beta \tilde{F}(x_\nu) &= \partial^\beta F_\nu(x_\nu) = 0 \text{ for } \beta \in \mathcal{A}, \nu = 1, \dots, \nu_{\max} && \text{(thanks to **(POU4)** and (17)).} \end{aligned}$$

Thus, \tilde{F} extends the data $(f, 0)$ and \tilde{F} is Coherent with $0 \in \mathcal{P}$. We next estimate the seminorm $\|\tilde{F}\|_{L^{m,p}(Q^\circ)}$. A straightforward application of Lemma 9.1 shows that

$$(19) \quad \begin{aligned} \|\tilde{F}\|_{L^{m,p}(Q^\circ)}^p &\lesssim \sum_{\nu=1}^{\nu_{\max}} \|F_\nu\|_{L^{m,p}(1.1Q_\nu)}^p + \sum_{\nu \leftrightarrow \nu'} \sum_{\bar{\beta} \in \mathcal{M}} \left| \partial^{\bar{\beta}}(F_\nu - J_{x_{\nu'}} F_{\nu'})(x_\nu) \right|^p \delta_{Q_\nu}^{n+(|\bar{\beta}|-m)p} \\ &\lesssim \sum_{\nu \leftrightarrow \nu'} \left[\|F_\nu\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}^p + \sum_{\bar{\beta} \in \mathcal{M}} \left| \partial^{\bar{\beta}}(F_\nu - F_{\nu'})(x_\nu) \right|^p \delta_{Q_\nu}^{n+(|\bar{\beta}|-m)p} \right], \end{aligned}$$

thanks to the Sobolev Inequality (2.3). (Note that $x_\nu \in Q_\nu$ and $x_{\nu'} \in Q_{\nu'}$, and note that the Sobolev Inequality holds for $U = 1.1Q_\nu \cup 1.1Q_{\nu'}$ whenever $\nu \leftrightarrow \nu'$.)

Let $\nu, \nu' \in \{1, \dots, \nu_{\max}\}$ satisfy $\nu \leftrightarrow \nu'$. To start, we bound the terms from the innermost sum on the right-hand side of (19). Using (12) and (16), we write

$$(20) \quad F_{\nu'} = F - \sum_{\alpha, \beta \in \mathcal{A}} \partial^\alpha F(x_{\nu'}) \omega_{\alpha\beta}^{\nu\nu'} \varphi_{\nu, \beta}.$$

Since \mathcal{A} is monotonic, if $\gamma \in \mathcal{A}$ and $|\gamma'| \leq m - 1 - |\gamma|$, then $\gamma + \gamma' \in \mathcal{A}$, and hence $\partial^{\gamma+\gamma'}(F_{\nu'})(x_{\nu'}) = 0$, thanks to (17). Thus by a Taylor expansion, we have

$$(21) \quad \partial^\gamma J_{x_{\nu'}}(F_{\nu'})(x_\nu) = \sum_{|\gamma'| \leq m-1-|\gamma|} \text{coeff}(\gamma, \gamma') \cdot \partial^{\gamma+\gamma'} F_{\nu'}(x_{\nu'}) \cdot (x_\nu - x_{\nu'})^{\gamma'} = 0 \quad (\gamma \in \mathcal{A}).$$

For each $\bar{\beta} \in \mathcal{M}$, we now bound

$$\begin{aligned}
|\partial^{\bar{\beta}}(F_\nu - F_{\nu'})(x_\nu)| &= \left| \sum_{\beta \in \mathcal{A}} \partial^\beta F(x_\nu) \partial^{\bar{\beta}} \varphi_{\nu, \beta}(x_\nu) - \sum_{\alpha, \beta \in \mathcal{A}} \partial^\alpha F(x_{\nu'}) \omega_{\alpha \beta}^{\nu \nu'} \partial^{\bar{\beta}} \varphi_{\nu, \beta}(x_\nu) \right| \\
&\quad \text{(from (16) and (20))} \\
&\leq \sum_{\beta \in \mathcal{A}} \left| \partial^{\bar{\beta}} \varphi_{\nu, \beta}(x_\nu) \right| \left[\left| \partial^\beta F(x_\nu) - \sum_{\alpha \in \mathcal{A}} \partial^\alpha F(x_{\nu'}) \omega_{\alpha \beta}^{\nu \nu'} \right| \right] \\
&= \sum_{\beta \in \mathcal{A}} \left| \partial^{\bar{\beta}} \varphi_{\nu, \beta}(x_\nu) \right| \left[\left| \partial^\beta F_{\nu'}(x_\nu) \right| \right] \\
&\quad \text{(from (9) and (20))} \\
&= \sum_{\beta \in \mathcal{A}} \left| \partial^{\bar{\beta}} \varphi_{\nu, \beta}(x_\nu) \right| \left[\left| \partial^\beta (F_{\nu'} - J_{x_{\nu'}}(F_{\nu'}))(x_\nu) \right| \right] \\
&\quad \text{(from (21))} \\
&\lesssim \sum_{\beta \in \mathcal{A}} \delta_{Q_\nu}^{|\beta| - |\bar{\beta}|} \delta_{Q_\nu}^{m-n/p - |\beta|} \|F_{\nu'}\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})} \\
&\quad \text{(from (11) and the Sobolev Inequality (2.3))} \\
(22) \quad &\lesssim \delta_{Q_\nu}^{m-n/p - |\bar{\beta}|} \|F_{\nu'}\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}.
\end{aligned}$$

Next, by $|A_{\alpha\gamma}^\nu| \leq C$, (6), (15), (16), the first bullet point in Lemma 5.3 and good geometry of the CZ cubes, we obtain

$$\begin{aligned}
&\sum_{\nu \leftrightarrow \nu'} \|F_\nu\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}^p \\
&\lesssim \sum_{\nu \leftrightarrow \nu'} \left[\|F\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}^p + \sum_{\gamma \in \mathcal{A}} \|\varphi_\gamma\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}^p \|F\|_{L^{m,p}(\mathbb{R}^n)}^p \right] \\
(23) \quad &\lesssim \|F\|_{L^{m,p}(\mathbb{R}^n)}^p + \|F\|_{L^{m,p}(\mathbb{R}^n)}^p \sum_{\gamma \in \mathcal{A}} \|\varphi_\gamma\|_{L^{m,p}(\mathbb{R}^n)}^p \stackrel{(10)}{\lesssim} \|F\|_{L^{m,p}(\mathbb{R}^n)}^p.
\end{aligned}$$

Finally, inserting (22) and (23) into (19), we have

$$(24) \quad \|\tilde{F}\|_{L^{m,p}(Q^\circ)} \lesssim \|F\|_{L^{m,p}(\mathbb{R}^n)}.$$

Next, we modify \tilde{F} to be Constant Along Paths, without ruining (18) or the control on the seminorm. For each $\nu = 2, \dots, \nu_{\max}$, there exists $\bar{\theta}_\nu \in C_0^\infty(Q_\nu)$ that satisfies

$$(25) \quad \bar{\theta}_\nu \equiv 1 \text{ in a neighborhood of } x_\nu;$$

$$(26) \quad \bar{\theta}_\nu \equiv 0 \text{ in a neighborhood of } E \cup \{x_{\nu'} : \nu' \in \{1, 2, \dots, \nu_{\max}\}, \nu' \neq \nu\}; \text{ and}$$

$$(27) \quad |\partial^\alpha \bar{\theta}_\nu| \lesssim \delta_{Q_\nu}^{-|\alpha|} \text{ when } |\alpha| \leq m.$$

(Here, the conditions on E' from Lemma 8.2 are used.) We define $\bar{F} \in L^{m,p}(Q^\circ)$ by

$$\bar{F} = \tilde{F} + \sum_{\nu=2}^{\nu_{\max}} \bar{\theta}_\nu \cdot \left[J_{x_{\kappa(\nu)}}^\#(\tilde{F}) - J_{x_\nu}(\tilde{F}) \right].$$

It is simple to check that

- (a) $\overline{F} = \tilde{F} + 0 = f$ on E ; this follows from (18) and (26).
- (b) \overline{F} is Coherent with $0 \in \mathcal{P}$, and in particular, $J_z \overline{F} = 0$; this follows from (18) and (26). (We also use the fact that $x_{\kappa(\nu)}^\sharp = x_\nu$ whenever Q_ν is keystone; see **(K3)** in Proposition 2.)
- (c) $J_{x_\nu}(\overline{F}) = J_{x_{\kappa(\nu)}^\sharp}(\tilde{F})$ for $\nu = 2, \dots, \nu_{\max}$; this follows from (25) and (26). (Also, see **(K3)** in Proposition 2.)

We now estimate the seminorm

$$\|\overline{F}\|_{L^{m,p}(Q^\circ)}^p \lesssim \|\tilde{F}\|_{L^{m,p}(Q^\circ)}^p + \sum_{\nu=2}^{\nu_{\max}} \sup_{x \in Q_\nu} \left[\sum_{|\alpha+\beta|=m} |\partial^\alpha \bar{\theta}_\nu(x)|^p \cdot \left| \partial^\beta \left(J_{x_\nu}(\tilde{F}) - J_{x_{\kappa(\nu)}^\sharp}(\tilde{F}) \right)(x) \right|^p \right] \delta_{Q_\nu}^n$$

(from $\text{supp}(\bar{\theta}_\nu) \subset Q_\nu$)

$$\lesssim \|\tilde{F}\|_{L^{m,p}(Q^\circ)}^p + \sum_{\nu=2}^{\nu_{\max}} \sup_{x \in Q_\nu} \left[\sum_{|\beta| \leq m} \delta_{Q_\nu}^{n-(m-|\beta|)p} \cdot \left| \partial^\beta \left(J_{x_\nu}(\tilde{F}) - J_{x_{\kappa(\nu)}^\sharp}(\tilde{F}) \right)(x) \right|^p \right]$$

(from (27))

$$\lesssim \|\tilde{F}\|_{L^{m,p}(Q^\circ)}^p + \sum_{\nu=2}^{\nu_{\max}} \left| J_{x_\nu}(\tilde{F}) - J_{x_{\kappa(\nu)}^\sharp}(\tilde{F}) \right|_\nu^p$$

(from (2.2); recall that $|P|_\nu = |P|_{x_\nu, \delta_{Q_\nu}}$ and $|x_\nu - x| \leq C\delta_{Q_\nu}$ for any $x \in Q_\nu$)

$$\lesssim \|\tilde{F}\|_{L^{m,p}(Q^\circ)}^p$$

(from Lemma 14.1).

By Lemma 9.2, there exists $\widehat{F} \in L^{m,p}(\mathbb{R}^n)$ with $\widehat{F} = \overline{F}$ on $0.99Q^\circ$ and $\|\widehat{F}\|_{L^{m,p}(\mathbb{R}^n)} \lesssim \|\overline{F}\|_{L^{m,p}(Q^\circ)}$. Thus,

$$\|\widehat{F}\|_{L^{m,p}(\mathbb{R}^n)} \lesssim \|\overline{F}\|_{L^{m,p}(Q^\circ)} \lesssim \|\tilde{F}\|_{L^{m,p}(Q^\circ)} \stackrel{(24)}{\lesssim} \|F\|_{L^{m,p}(\mathbb{R}^n)} \stackrel{(14)}{\leq} 2\|(f, 0)\|.$$

It follows from $E \cup E' \subset 0.99Q^\circ$ and (a)-(c) that \widehat{F} extends the data $(f, 0)$, \widehat{F} is Coherent with $0 \in \mathcal{P}$ and \widehat{F} is Constant Along Paths. This concludes the proof of Proposition 3. \square

15. A CONSTRAINED PROBLEM

For each $\nu = 1, \dots, \nu_{\max}$, we define $E_\nu = E \cap (1.1)Q_\nu$. For the given function $f : E \rightarrow \mathbb{R}$, we denote $f_\nu = f|_{E_\nu}$ here and throughout.

Proposition 4. *There exist a linear operator $T : L^{m,p}(E; E') \rightarrow L^{m,p}(\mathbb{R}^n)$ and a collection of linear functionals $\Omega' \subset [L^{m,p}(E)]^*$ such that the following hold:*

- T is an extension operator.
- $\|T(f, \vec{P})\|_{L^{m,p}(\mathbb{R}^n)} \leq C\|(f, \vec{P})\|$ for each $(f, \vec{P}) \in L^{m,p}(E; E')$.

- $\sum \{dp(\omega') : \omega' \in \Omega'\} \leq C \cdot \#(E)$.
- T has Ω' -assisted bounded depth.

Moreover,

$$(1) \quad \|(f, \vec{P})\|^p \simeq \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, P_\nu)\|^p + \sum_{\nu \leftrightarrow \nu'} |P_\nu - P_{\nu'}|_\nu^p.$$

Above, $\|(f_\nu, P_\nu)\|$ denotes the trace seminorm $\|(f_\nu, P_\nu)\|_{L^{m,p}(E_\nu; x_\nu)}$ and $\|(f, \vec{P})\|$ denotes the trace seminorm $\|(f, \vec{P})\|_{L^{m,p}(E; E')}$.

Proof. For each $\nu = 1, \dots, \nu_{\max}$, we apply **(L1)**-**(L6)** from section 10 to the subset $\bar{E}_\nu = E_\nu$. Thus there exist linear functionals $\Omega_\nu \subset [L^{m,p}(E_\nu)]^*$ and a linear map $T_\nu : L^{m,p}(E_\nu; x_\nu) \rightarrow L^{m,p}(\mathbb{R}^n)$ (with Ω_ν -assisted bounded depth) such that

$$(2) \quad T_\nu(f_\nu, P_\nu) = f_\nu \text{ on } E_\nu \text{ and } J_{x_\nu}(T_\nu(f_\nu, P_\nu)) = P_\nu \text{ and}$$

$$(3) \quad \|T_\nu(f_\nu, P_\nu)\|_{L^{m,p}(1.1Q_\nu)} \leq \|T_\nu(f_\nu, P_\nu)\|_{L^{m,p}(\mathbb{R}^n)} \simeq \|(f_\nu, P_\nu)\|.$$

Using the partition of unity $\{\theta_\nu\}$ that satisfies **(POU1)**-**(POU5)**, we define

$$F^\circ = T^\circ(f, \vec{P}) = \sum_{\nu=1}^{\nu_{\max}} T_\nu(f_\nu, P_\nu)\theta_\nu.$$

Of course, $T^\circ : L^{m,p}(E; E') \rightarrow L^{m,p}(Q^\circ)$ is a linear map. From **(POU2)** and (2), $T_\nu(f_\nu, P_\nu) = f$ on $\text{supp}(\theta_\nu) \cap E$. Hence, by **(POU5)**, $F^\circ = f$ on E . From **(POU4)** and (2), $J_{E'}F^\circ = \vec{P}$. To summarize, we have shown that

$$(4) \quad F^\circ = f \text{ on } E \text{ and } J_{E'}F^\circ = \vec{P}.$$

From (2), (3) and a straightforward application of Lemma 9.1, we have

$$(5) \quad \|F^\circ\|_{L^{m,p}(Q^\circ)}^p \lesssim \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, P_\nu)\|^p + \sum_{\nu \leftrightarrow \nu'} |P_\nu - P_{\nu'}|_\nu^p.$$

We identify $[L^{m,p}(E_\nu)]^*$ with a subspace of $[L^{m,p}(E)]^*$ through the natural restriction from $L^{m,p}(E)$ to $L^{m,p}(E_\nu)$ and define

$$\Omega' = \bigcup_{\nu=1}^{\nu_{\max}} \Omega_\nu \subset [L^{m,p}(E)]^*.$$

Since Ω_ν satisfies **(L4)** (for the subset $E_\nu = E \cap (1.1)Q_\nu$), we obtain

$$(6) \quad \sum_{\omega \in \Omega'} dp(\omega) \leq \sum_{\nu=1}^{\nu_{\max}} \sum_{\omega \in \Omega_\nu} dp(\omega) \leq \sum_{\nu=1}^{\nu_{\max}} C \cdot \#(E \cap (1.1)Q_\nu),$$

which is bounded by $C' \cdot \#(E)$ due to the first bullet point in Lemma 5.3. This proves the third bullet point of Proposition 4.

Using **(POU2)**, for each fixed $x \in Q^\circ$ we obtain a list of cubes $Q_{\nu_1}, \dots, Q_{\nu_L} \in CZ^\circ$ and linear maps $\psi_{1,x}, \dots, \psi_{L,x} : \mathcal{P} \rightarrow \mathcal{P}$ such that

$$(7) \quad J_x[T^\circ(f, \vec{P})] = \psi_{1,x}(J_x[T_{\nu_1}(f_{\nu_1}, P_{\nu_1})]) + \dots + \psi_{L,x}(J_x[T_{\nu_L}(f_{\nu_L}, P_{\nu_L})]).$$

The list $Q_{\nu_1}, \dots, Q_{\nu_L}$ is composed of the CZ cubes for which $x \in (1.1)Q_{\nu_i}$ (thus, $\theta_{\nu_1}, \dots, \theta_{\nu_L}$ include those cutoff functions which do not vanish in a neighborhood of x), and we define $\psi_{i,x}(P) = J_x(\theta_{\nu_i}P)$ ($i = 1, \dots, L$). The first bullet point in Lemma 5.3 shows that L is controlled by a universal constant. Thus, by (7) and the fact that T_{ν_i} has Ω_{ν_i} -assisted bounded depth, the linear map $J_x T^\circ(\cdot, \cdot) :$

$L^{m,p}(E; E') \rightarrow \mathcal{P}$ has Ω' -assisted bounded depth. Therefore, T° has Ω' -assisted bounded depth.

Next, we apply Lemma 9.2 to the function $F^\circ = T^\circ(f, \vec{P}) \in L^{m,p}(Q^\circ)$. This gives a linear map $T : L^{m,p}(E; E') \rightarrow L^{m,p}(\mathbb{R}^n)$ with Ω' -assisted bounded depth such that $F = T(f, \vec{P})$ satisfies $F = F^\circ$ on $0.99Q^\circ$ and

$$(8) \quad \|F\|_{L^{m,p}(\mathbb{R}^n)}^p \lesssim \|F^\circ\|_{L^{m,p}(Q^\circ)}^p \stackrel{(5)}{\lesssim} \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, P_\nu)\|^p + \sum_{\nu \leftrightarrow \nu'} |P_\nu - P_{\nu'}|_\nu^p.$$

From $E \cup E' \subset 0.99Q^\circ$ and (4), we have

$$(9) \quad F = f \text{ on } E \text{ and } J_{E'}(F) = \vec{P}.$$

Thus the first and fourth bullet points in Proposition 4 hold.

Finally, we must establish (1) and prove that $\|F\|_{L^{m,p}(\mathbb{R}^n)}$ lies within a universal constant factor of the trace seminorm $\|(f, \vec{P})\|_{L^{m,p}(E; E')}$. Let $H \in L^{m,p}(\mathbb{R}^n)$ be an arbitrary function that satisfies

$$(*) \quad H = f \text{ on } E \text{ and } J_{E'}H = \vec{P}.$$

From the Sobolev Inequality (2.3) and $J_{E'}H = \vec{P}$, we have

$$|P_\nu - P_{\nu'}|_\nu^p \lesssim \|H\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}^p.$$

Hence, by good geometry of the CZ cubes,

$$(10) \quad \sum_{\nu \leftrightarrow \nu'} |P_\nu - P_{\nu'}|_\nu^p \lesssim \sum_{\nu \leftrightarrow \nu'} \|H\|_{L^{m,p}(1.1Q_\nu \cup 1.1Q_{\nu'})}^p \lesssim \|H\|_{L^{m,p}(\mathbb{R}^n)}^p.$$

Recall that $E_\nu = (1.1)Q_\nu \cap E \subset (0.9)(1.3)Q_\nu$ and $x_\nu \in Q_\nu \subset (0.9)(1.3)Q_\nu$. Applying Lemma 9.2 with $Q = (1.3)Q_\nu$ and $a = 0.1$ we deduce that $\|(f_\nu, P_\nu)\| \lesssim \|H\|_{L^{m,p}(1.3Q_\nu)}$. Hence, by good geometry of the CZ cubes,

$$(11) \quad \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, P_\nu)\|^p \lesssim \sum_{\nu=1}^{\nu_{\max}} \|H\|_{L^{m,p}(1.3Q_\nu)}^p \lesssim \|H\|_{L^{m,p}(\mathbb{R}^n)}^p.$$

Adding together (10) and (11) shows that the right-hand side of (8) is bounded by $C\|H\|_{L^{m,p}(\mathbb{R}^n)}^p$. Taking the infimum with respect to those H satisfying (*), we obtain

$$\sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, P_\nu)\|^p + \sum_{\nu \leftrightarrow \nu'} |P_\nu - P_{\nu'}|_\nu^p \lesssim \|(f, \vec{P})\|^p.$$

The reverse inequality holds as well, thanks to (8) and (9). Thus, (1) holds. Together, (1) and (8) imply that $\|F\|_{L^{m,p}(\mathbb{R}^n)} \leq C'\|(f, \vec{P})\|_{L^{m,p}(E; E')}$. Thus the second bullet point in Proposition 4 holds. This concludes the proof of Proposition 4. \square

16. THE JET OF A NEAR OPTIMAL EXTENSION II

Recall our notation and assumptions on indexing in the text surrounding (8.2). For each $\mu = \mu_{\min}, \dots, \mu_{\max}$, Lemma 11.2 provides a jet R_μ^\sharp that depends linearly on $(f|_{E \cap 9Q_\mu^\sharp}, P)$ and satisfies

$$(1) \quad \partial^\alpha R_\mu^\sharp(x_\mu^\sharp) = \partial^\alpha P(x_\mu^\sharp) \text{ for every } \alpha \in \mathcal{A}$$

and

$$(2) \quad \|(f_\mu^\sharp, R_\mu^\sharp)\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)} \leq C \inf \left\{ \|(f_\mu^\sharp, R')\|_{L^{m,p}(E \cap 9Q_\mu^\sharp; x_\mu^\sharp)} : R' \in \mathcal{P}, \right. \\ \left. \partial^\alpha R'(x_\mu^\sharp) = \partial^\alpha P(x_\mu^\sharp) \text{ for every } \alpha \in \mathcal{A} \right\}.$$

In the case that Q_1 is keystone, we also define $R_1^\sharp = P$. Finally, we define $R_1 = P$ and $R_\nu = R_{\kappa(\nu)}^\sharp$ for each $\nu = 2, \dots, \nu_{\max}$.

It is immediate that $\vec{R} = (R_\nu)_{\nu=1}^{\nu_{\max}} \in Wh(E')$ is Constant Along Paths, Coherent with P and depends linearly on (f, P) . The main result of this section states that \vec{R} is the jet on E' of a near optimal extension of (f, P) .

Lemma 16.1. *Given $(f, P) \in L^{m,p}(E; z)$, let $\vec{R} \in Wh(E')$ be defined as above. Then,*

$$\|(f, \vec{R})\|_{L^{m,p}(E; E')} \lesssim \|(f, P)\|_{L^{m,p}(E; z)}.$$

Proof. By Proposition 3, it suffices to bound

$$(3) \quad \|(f, \vec{R})\|_{L^{m,p}(E; E')} \lesssim \|(f, \vec{P})\|_{L^{m,p}(E; E')}$$

for each $\vec{P} = (P_\nu)_{\nu=1}^{\nu_{\max}} \in Wh(E')$ that satisfies the following properties:

Constant Along Paths: $P_\nu = P_{\kappa(\nu)}^\sharp$ for each $\nu = 2, \dots, \nu_{\max}$.

Coherent with \mathbf{P} : $P_1 = P$ and $\partial^\alpha P_\mu^\sharp(x_\mu^\sharp) = \partial^\alpha P(x_\mu^\sharp)$

for each $\mu = 1, \dots, \mu_{\max}$, $\alpha \in \mathcal{A}$.

Fix $\vec{P} \in Wh(E')$ as above. Then, we have

$$(4) \quad \begin{aligned} \|(f, \vec{R})\|^p &\simeq \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, R_\nu)\|^p + \sum_{\nu \leftrightarrow \nu'} |R_\nu - R_{\nu'}|_\nu^p \\ &\quad \text{(from (15.1) in Proposition 4)} \\ &\lesssim \sum_{\nu=1}^{\nu_{\max}} [\|(f_\nu, P_\nu)\|^p + \|(0|_{E_\nu}, P_\nu - R_\nu)\|^p] \\ &\quad + \sum_{\nu \leftrightarrow \nu'} [|R_\nu - P_\nu|_\nu^p + |P_\nu - P_{\nu'}|_\nu^p + |P_{\nu'} - R_{\nu'}|_\nu^p] \\ &\quad \text{(from subadditivity of the seminorms)} \\ &\lesssim \sum_{\nu=1}^{\nu_{\max}} [\|(f_\nu, P_\nu)\|^p + \|(0|_{E_\nu}, P_\nu - R_\nu)\|^p + |R_\nu - P_\nu|_\nu^p] + \sum_{\nu \leftrightarrow \nu'} |P_\nu - P_{\nu'}|_\nu^p \\ &\quad \text{(from (8.4) and good geometry of the CZ cubes)} \\ &\simeq \|(f, \vec{P})\|^p + \sum_{\nu=1}^{\nu_{\max}} [\|(0|_{E_\nu}, R_\nu - P_\nu)\|^p + |R_\nu - P_\nu|_\nu^p] \\ &\quad \text{(from (15.1))} \\ &\lesssim \|(f, \vec{P})\|^p + \sum_{\nu=2}^{\nu_{\max}} |R_\nu - P_\nu|_\nu^p \\ &\quad \text{(from } P_1 = P = R_1 \text{ and Lemma 13.1).} \end{aligned}$$

Define X as the second term on the right-hand side of (4). Because \vec{P} and \vec{R} are Constant Along Paths and from (8.3), we have

$$(5) \quad \begin{aligned} X &= \sum_{\nu=2}^{\nu_{\max}} |R_{\nu} - P_{\nu}|_{\nu}^p = \sum_{\nu=2}^{\nu_{\max}} \left| R_{\kappa(\nu)}^{\sharp} - P_{\kappa(\nu)}^{\sharp} \right|_{\nu}^p \\ &\lesssim \sum_{\mu=1}^{\mu_{\max}} \sum_{\kappa(\nu)=\mu} \sum_{|\beta| \leq m-1} |\partial^{\beta}(R_{\mu}^{\sharp} - P_{\mu}^{\sharp})(x_{\mu}^{\sharp})|^p \delta_{Q_{\nu}}^{n+(|\beta|-m)p}. \end{aligned}$$

Next, we estimate the dyadic sum of lengthscales arising above.

If Q_{μ}^{\sharp} is the keystone cube arising from Q_{ν} , then $\delta_{Q_{\nu}} \geq c\delta_{Q_{\mu}^{\sharp}}$ and $Q_{\mu}^{\sharp} \subset CQ_{\nu}$ for a large enough universal constant C and a small enough universal constant c . Hence, for each μ and $\tau > 0$, we have

$$\begin{aligned} \sum_{\kappa(\nu)=\mu} [\delta_{Q_{\nu}}]^{-\tau} &\leq C'(\tau, m, n, p) \\ &\quad \cdot \sum \left\{ [\delta_Q]^{-\tau} : Q \subset \mathbb{R}^n \text{ dyadic with } [\delta_Q \geq c\delta_{Q_{\mu}^{\sharp}} \text{ and } Q_{\mu}^{\sharp} \subset CQ] \right\} \\ &\leq C''(\tau, m, n, p) [\delta_{Q_{\mu}^{\sharp}}]^{-\tau}. \end{aligned}$$

Plugging this inequality into (5), we have

$$(6) \quad \begin{aligned} X &\lesssim \sum_{\mu=1}^{\mu_{\max}} \sum_{|\beta| \leq m-1} |\partial^{\beta}(R_{\mu}^{\sharp} - P_{\mu}^{\sharp})(x_{\mu}^{\sharp})|^p [\delta_{Q_{\mu}^{\sharp}}]^{n+(|\beta|-m)p} = \sum_{\mu=1}^{\mu_{\max}} |R_{\mu}^{\sharp} - P_{\mu}^{\sharp}|_{x_{\mu}^{\sharp}, \delta_{Q_{\mu}^{\sharp}}}^p \\ &= \sum_{\mu=\mu_{\min}}^{\mu_{\max}} |R_{\mu}^{\sharp} - P_{\mu}^{\sharp}|_{x_{\mu}^{\sharp}, \delta_{Q_{\mu}^{\sharp}}}^p, \end{aligned}$$

since when Q_1 is keystone we have $P_1^{\sharp} = P_1 = P = R_1 = R_1^{\sharp}$ by Coherence with P . (Recall the indexing assumption (8.2).)

Note that $\partial^{\alpha}(R_{\mu}^{\sharp})(x_{\mu}^{\sharp}) = \partial^{\alpha}P(x_{\mu}^{\sharp}) = \partial^{\alpha}(P_{\mu}^{\sharp})(x_{\mu}^{\sharp})$ for each $\mu = \mu_{\min}, \dots, \mu_{\max}$ and $\alpha \in \mathcal{A}$. (This follows from Coherence with P .) Next, from Lemma 13.1 and (6), we obtain

$$(7) \quad \begin{aligned} X &\lesssim \sum_{\mu=\mu_{\min}}^{\mu_{\max}} \|(0|_{E \cap 9Q_{\mu}^{\sharp}}, R_{\mu}^{\sharp} - P_{\mu}^{\sharp})\|^p \lesssim \sum_{\mu=\mu_{\min}}^{\mu_{\max}} \left[\|(f|_{E \cap 9Q_{\mu}^{\sharp}}, R_{\mu}^{\sharp})\|^p + \|(f|_{E \cap 9Q_{\mu}^{\sharp}}, P_{\mu}^{\sharp})\|^p \right] \\ &\lesssim \sum_{\mu=\mu_{\min}}^{\mu_{\max}} \|(f|_{E \cap 9Q_{\mu}^{\sharp}}, P_{\mu}^{\sharp})\|^p, \text{ thanks to (2)}. \end{aligned}$$

(The seminorms above are taken in the space $L^{m,p}(E \cap 9Q_{\mu}^{\sharp}, x_{\mu}^{\sharp})$.)

Let $F \in L^{m,p}(\mathbb{R}^n)$ be arbitrary with $F = f$ on E and $J_{E'}(F) = \vec{P}$. Applying Lemma 9.2 with $Q = 10Q_{\mu}^{\sharp}$ and $a = 0.1$, we deduce that $\|(f|_{E \cap 9Q_{\mu}^{\sharp}}, P_{\mu}^{\sharp})\| \lesssim \|F\|_{L^{m,p}(10Q_{\mu}^{\sharp})}$. Thus, (7) yields

$$X \lesssim \sum_{\mu=1}^{\mu_{\max}} \|F\|_{L^{m,p}(10Q_{\mu}^{\sharp})}^p,$$

which is bounded by $C \cdot \|F\|_{L^{m,p}(\mathbb{R}^n)}^p$, thanks to Lemma 7.1. Taking the infimum with respect to $F \in L^{m,p}(\mathbb{R}^n)$ as above, we obtain $X \lesssim \|(f, \vec{P})\|^p$. Thus (3) holds (see (4) and the definition of X that follows), which completes the proof of Lemma 16.1. \square

17. THE EXTENSION OPERATOR

In this section we complete the proof of the Extension Theorem for $(E; z)$. We now fix a sufficiently small universal constant $\epsilon(\mathcal{A}) = \epsilon$ so that the results from previous sections hold.

Proposition 4 provides a bounded linear extension operator $\widehat{T} : L^{m,p}(E; E') \rightarrow L^{m,p}(\mathbb{R}^n)$ and assists $\widehat{\Omega} \subset [L^{m,p}(E)]^*$ so that \widehat{T} has $\widehat{\Omega}$ -assisted bounded depth and such that

$$(1) \quad \sum_{\omega \in \widehat{\Omega}} \text{dp}(\omega) \leq C \cdot \#(E).$$

Let $\vec{R} \in Wh(E')$ be as defined in the previous section. In particular, \vec{R} depends linearly on (f, P) , \vec{R} is Constant Along Paths and \vec{R} is Coherent with P .

We define

$$T(f, P) = \widehat{T}(f, \vec{R}).$$

Since \vec{R} depends linearly on (f, P) and the map \widehat{T} is linear, the map T is clearly linear. Also, since \widehat{T} is an extension operator and $R_1 = P$ (by Coherence of \vec{R} with P), we have

$$T(f, P) = \widehat{T}(f, \vec{R}) = f \text{ on } E \text{ and } J_z(T(f, P)) = J_{x_1}(\widehat{T}(f, \vec{R})) = R_1 = P.$$

Thus, T is an extension operator and we have established **(E1)** in the statement of the Extension Theorem for (E, z) .

We now prove **(E2)**. The first inequality in **(E2)** follows trivially from **(E1)** and the definition of the trace seminorm. Thanks to Lemma 16.1 and the fact that $\widehat{T} : L^{m,p}(E; E') \rightarrow L^{m,p}(\mathbb{R}^n)$ is bounded, we have

$$(2) \quad \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)} = \|\widehat{T}(f, \vec{R})\|_{L^{m,p}(\mathbb{R}^n)} \lesssim \|(f, \vec{R})\|_{L^{m,p}(E; E')} \lesssim \|(f, P)\|_{L^{m,p}(E; z)}.$$

Also, since $\|(f, P)\|_{L^{m,p}(E; z)} \leq \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}$, we may replace every \lesssim with \simeq in the above. This completes the proof of **(E2)**.

Next we estimate the seminorm $\|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}$ and define $M(f, P)$ in order to establish conclusion **(E3)**. Let us define

$$\begin{aligned} \mathcal{I} &:= \{(\nu, \nu') \in \{2, \dots, \nu_{\max}\}^2 : \nu \leftrightarrow \nu' \text{ and } \kappa(\nu) \neq \kappa(\nu')\} \text{ and} \\ \mathcal{I}^\sharp &:= \{(\kappa(\nu), \kappa(\nu')) : (\nu, \nu') \in \mathcal{I}\}. \end{aligned}$$

From **(K2)** in Proposition 2, we learn that

$$(3) \quad \#(\mathcal{I}^\sharp) \leq \#(\mathcal{I}) \leq C \cdot \#(E).$$

Recall that $E_\nu = E \cap (1.1)Q_\nu$ and $f_\nu = f|_{E_\nu}$. From the comment preceding (2) and from Proposition 4,

$$\begin{aligned}
\|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}^p &\simeq \|(f, \vec{R})\|_{L^{m,p}(E; E')}^p \simeq \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, R_\nu)\|^p + \sum_{\nu \leftrightarrow \nu'} |R_\nu - R_{\nu'}|_\nu^p \\
&\simeq \sum_{\nu=1}^{\nu_{\max}} \|(f_\nu, R_\nu)\|^p + \sum_{(\nu, \nu') \in \mathcal{I}} |R_{\kappa(\nu)}^\# - R_{\kappa(\nu')}^\#|_\nu^p \\
(4) \quad &+ \sum_{\substack{2 \leq \nu \leq \nu_{\max} \\ \nu \leftrightarrow 1}} |R_{\kappa(\nu)}^\# - R_1|_\nu^p \\
&\quad \text{(since } \vec{R} \text{ is Constant Along Paths).}
\end{aligned}$$

(Here, $\|(f_\nu, R_\nu)\|$ denotes the trace seminorm on $L^{m,p}(E_\nu; x_\nu)$.) From (8.3), we deduce that

$$\begin{aligned}
\sum_{(\nu, \nu') \in \mathcal{I}} |R_{\kappa(\nu)}^\# - R_{\kappa(\nu')}^\#|_\nu^p &\simeq \sum_{(\nu, \nu') \in \mathcal{I}} \sum_{\alpha \in \mathcal{M}} |\partial^\alpha (R_{\kappa(\nu)}^\# - R_{\kappa(\nu')}^\#)(x_{\kappa(\nu)}^\#)|^p \delta_{Q_\nu}^{n+(|\alpha|-m)p} \\
&= \sum_{(\mu, \mu') \in \mathcal{I}^\#} \sum_{\alpha \in \mathcal{M}} |\partial^\alpha (R_\mu^\# - R_{\mu'}^\#)(x_\mu^\#)|^p \Delta_\alpha(\mu, \mu')^p,
\end{aligned}$$

$$\text{where } \Delta_\alpha(\mu, \mu')^p := \sum \left\{ \delta_{Q_\nu}^{n+(|\alpha|-m)p} : (\nu, \nu') \in \mathcal{I}, \kappa(\nu) = \mu, \kappa(\nu') = \mu' \right\}.$$

Plugging into (4) this formula, the definition of the norm $|\cdot|_\nu$ and the formula $M_\nu(f_\nu, P_\nu) = \left(\sum_{\xi \in \Xi_\nu} |\xi(f_\nu, P_\nu)|^p \right)^{1/p}$ that approximates the trace seminorm $\|(f_\nu, P_\nu)\|$ (see **(L1)**-**(L6)** in section 10), we have

$$\begin{aligned}
\|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}^p &\simeq \sum_{\nu=1}^{\nu_{\max}} \sum_{\xi \in \Xi_\nu} |\xi(f_\nu, R_\nu)|^p \\
(5) \quad &+ \sum_{(\mu, \mu') \in \mathcal{I}^\#} \sum_{\alpha \in \mathcal{M}} |\partial^\alpha (R_\mu^\# - R_{\mu'}^\#)(x_\mu^\#)|^p \Delta_\alpha(\mu, \mu')^p \\
&+ \sum_{\substack{2 \leq \nu \leq \nu_{\max} \\ \nu \leftrightarrow 1}} \sum_{\alpha \in \mathcal{M}} |\partial^\alpha (R_{\kappa(\nu)}^\# - R_1)(x_\nu)|^p \delta_{Q_\nu}^{n-(m-|\alpha|)p}.
\end{aligned}$$

Here we record the fact (also following from **(L1)**-**(L6)**) that there exist exists $\Omega_\nu \subset [L^{m,p}(E_\nu)]^*$ such that

$$\begin{aligned}
&\text{each functional in } \Xi_\nu \text{ has } \Omega_\nu\text{-assisted bounded depth,} \\
(6) \quad &\#(\Xi_\nu) \leq C \cdot \#(E_\nu) \text{ and } \sum_{\omega \in \Omega_\nu} \text{dp}(\omega) \leq C \cdot \#(E_\nu).
\end{aligned}$$

We now simply take

$$(7) \quad M(f, P)^p := \text{the right-hand side of (5).}$$

Thus, $\|T(f, P)\| \simeq M(f, P)$, which establishes **(E3)**.

17.1. Assisted bounded depth. First, we define the collection of linear functionals $\Omega \subset [L^{m,p}(E)]^*$ that satisfies **(E4)**, **(E5)** and **(E6)**. Recall that $R_\mu^\sharp \in \mathcal{P}$ depends linearly on $(f|_{E \cap 9Q_\mu^\sharp}, P)$ for each $\mu = 1, \dots, \mu_{\max}$. Since \vec{R} is Coherent with P and \vec{R} is Constant Along Paths, we have

$$(8) \quad R_1 = P, \text{ and } R_\nu = R_{\kappa(\nu)}^\sharp \text{ for each } \nu = 2, \dots, \nu_{\max}.$$

In this subsection, R_μ^\sharp and R_ν should be considered as \mathcal{P} -valued linear maps on $L^{m,p}(E, z)$. Consider the decomposition

$$(9) \quad R_\mu^\sharp = \widehat{R}_\mu^\sharp + \widetilde{R}_\mu^\sharp, \text{ where } \widehat{R}_\mu^\sharp \in \mathcal{P} \text{ depends only on } f|_{E \cap 9Q_\mu^\sharp} \\ \text{and } \widetilde{R}_\mu^\sharp \in \mathcal{P} \text{ depends only on } P.$$

First we define linear functionals $\omega_{\beta,\mu} : L^{m,p}(E) \rightarrow \mathbb{R}$, for $\mu = 1, \dots, \mu_{\max}$ and $\beta \in \mathcal{M}$, by

$$\omega_{\beta,\mu}(f) = \partial^\beta \left[\widehat{R}_\mu^\sharp \right] (x_\mu^\sharp).$$

Therefore, $\text{dp}(\omega_{\beta,\mu}) \leq \#(E \cap 9Q_\mu^\sharp)$. We define

$$\Omega_{\text{new}} := \{ \omega_{\beta,\mu} : \mu = 1, \dots, \mu_{\max}, \beta \in \mathcal{M} \} \subset [L^{m,p}(E)]^*.$$

Recall the definition of $\widehat{\Omega}$ and Ω_ν in the text surrounding (1) and (6), respectively. Note that $\Omega_\nu \subset [L^{m,p}(E_\nu)]^*$ is included in $[L^{m,p}(E)]^*$ through the natural restriction from $L^{m,p}(E)$ onto $L^{m,p}(E_\nu)$. Let us define

$$(10) \quad \Omega := \widehat{\Omega} \cup \left(\bigcup_{\nu=1}^{\nu_{\max}} \Omega_\nu \right) \cup \Omega_{\text{new}}.$$

Thus, from (1) and (6), we have

$$\begin{aligned} \sum_{\omega \in \Omega} \text{dp}(\omega) &\leq \sum_{\nu} \sum_{\omega \in \Omega_\nu} \text{dp}(\omega) + \sum_{\omega \in \widehat{\Omega}} \text{dp}(\omega) + \sum_{\beta, \mu} \text{dp}(\omega_{\beta,\mu}) \\ &\lesssim \sum_{\nu} \#(E \cap 1.1Q_\nu) + \#(E) + \sum_{\mu} \#(E \cap 9Q_\mu^\sharp), \end{aligned}$$

which is bounded by $C \cdot \#(E)$ thanks to Lemma 5.3 and Lemma 7.1. This proves **(E4)**.

Since \widehat{T} has $\widehat{\Omega}$ -assisted bounded depth, there exists a universal constant $r \in \mathbb{N}$ such that for each point $x \in \mathbb{R}^n$ there exist $\omega_1, \dots, \omega_r \in \widehat{\Omega}$, polynomials $\widehat{P}^1, \dots, \widehat{P}^r \in \mathcal{P}$ and a linear map $\widetilde{\lambda} : L^{m,p}(E; E') \rightarrow \mathcal{P}$ with $\text{dp}(\widetilde{\lambda}) \leq r$ such that

$$(11) \quad J_x [T(f, P)] = J_x \left[\widehat{T}(f, \vec{R}) \right] = \omega_1(f) \widehat{P}^1 + \dots + \omega_r(f) \widehat{P}^r + \widetilde{\lambda}(f, \vec{R}).$$

Thanks to (8), (9), and a change of basis, we obtain

$$J_x [T(f, P)] = \omega_1(f) \widehat{P}^1 + \dots + \omega_r(f) \widehat{P}^r + \bar{\lambda} \left(f, P, \left(\widehat{R}_\mu^\sharp + \widetilde{R}_\mu^\sharp \right)_{\mu=1}^{\mu_{\max}} \right),$$

where $\text{dp}(\bar{\lambda}) \leq C \cdot \text{dp}(\widetilde{\lambda}) \leq C \cdot r$. Thus, by placing aside those terms from $\bar{\lambda}$ that depend on \widehat{R}_μ^\sharp (which depends solely on f), we have

$$J_x [T(f, P)] = \omega_1(f) \widehat{P}^1 + \dots + \omega_r(f) \widehat{P}^r + \omega_{r+1}(f) \widehat{P}^{r+1} + \dots + \omega_{r+s}(f) \widehat{P}^{r+s} + \widehat{\lambda}(f, P),$$

where $\omega_{r+1}, \dots, \omega_{r+s} \in \Omega_{\text{new}}$, $\widehat{P}^{r+1}, \dots, \widehat{P}^{r+s} \in \mathcal{P}$ are independent of (f, P) , and $s \leq C \cdot r$, $\text{dp}(\widehat{\lambda}) \leq \text{dp}(\bar{\lambda}) \leq C \cdot r$. Thus, T has Ω -assisted bounded depth, which establishes **(E5)**.

Finally, we define $\Xi \subset [L^{m,p}(E; z)]^*$ as

$$\Xi = \left\{ \begin{array}{l} \text{linear functionals arising on the right-hand side of (5),} \\ \text{counted with repetition} \end{array} \right\}.$$

By “counted with repetition”, we mean that if some linear functional $\xi : L^{m,p}(E; z) \rightarrow \mathbb{R}$ appears t times on the right-hand side of (5), then we include $t^{1/p}\xi$ in Ξ .

From the definition of M in (7), we find that

$$M(f, P) = \left(\sum_{\xi \in \Xi} |\xi(f, P)|^p \right)^{1/p}.$$

Thus, **(E6)(c)** holds.

From (3), (6) and the good geometry of the CZ cubes, we have

$$\begin{aligned} \#(\Xi) &\leq \sum_{\nu=1}^{\nu_{\max}} \#(\Xi_{\nu}) + C \cdot \#(\mathcal{I}^{\sharp}) + C \cdot \#\{1 \leq \nu \leq \nu_{\max} : \nu \leftrightarrow 1\} \\ &\leq C' \sum_{\nu=1}^{\nu_{\max}} \#(E_{\nu}) + C' \cdot \#(E), \end{aligned}$$

which is controlled by $C'' \cdot \#(E)$, thanks to Lemma 5.3. Thus, **(E6)(b)** holds.

One verifies that each functional $\xi \in \Xi$ (appearing on the right-hand side of (5)) has Ω -assisted bounded depth using (6), (9) and (10). Thus, **(E6)(a)** holds.

We have completed the Induction Step indicated in section 3, thus completing the proof of the Main Lemma for all labels \mathcal{A} . Taking $\mathcal{A} = \emptyset$, we obtain the Extension Theorem for (E, z) whenever $E \subset \mathbb{R}^n$ is finite.

18. PROOFS OF THE MAIN THEOREMS FOR FINITE E

Let $E \subset \mathbb{R}^n$ be a finite subset with cardinality $\#(E) = N$. Pick $z \in \mathbb{R}^n \setminus E$ that satisfies $d(z, E) \leq 1$. The Extension Theorem for (E, z) produces (T, M, Ω, Ξ) that satisfy **(E1)-(E6)**. Thus, T is a bounded extension operator:

(1a) $T(f, P) = f$ on E and $J_z(T(f, P)) = P$ and

(1b) $\|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)} \simeq \|(f, P)\|_{L^{m,p}(E; z)}$ for any data (f, P) .

We also have a formula

$$(2) \quad M(f, P) := \left(\sum_{\xi \in \Xi} |\xi(f, P)|^p \right)^{1/p} \quad \text{with } \#(\Xi) \leq C \cdot \#(E)$$

such that

$$(3) \quad \|(f, P)\|_{L^{m,p}(E; z)} \simeq M(f, P).$$

Moreover, T and the functionals in Ξ have Ω -assisted bounded depth, while the assists Ω satisfy

$$(4) \quad \sum_{\omega \in \Omega} \text{dp}(\omega) \leq C \cdot \#(E).$$

18.1. Theorems 1, 2 and 3 for finite sets. By definition of the trace seminorm, we have

$$(5) \quad \|f\|_{L^{m,p}(E)} = \inf\{\|(f, P)\|_{L^{m,p}(E;z)} : P \in \mathcal{P}\}.$$

We use Lemma 11.1 to choose $R = R(f) \in \mathcal{P}$ depending linearly on f , with

$$M(f, R) \leq C \cdot \inf\{M(f, P) : P \in \mathcal{P}\}.$$

Define $\widehat{M}(f) := M(f, R)$ and $\widehat{T}(f) := T(f, R)$. Since T is an extension operator we have $\widehat{T}(f) = T(f, R) = f$ on E . Therefore, \widehat{T} is an extension operator. From (3) and (5), we have

$$(6) \quad \widehat{M}(f) \simeq \|f\|_{L^{m,p}(E)}.$$

Moreover,

$$\|\widehat{T}(f)\|_{L^{m,p}(\mathbb{R}^n)} = \|T(f, R)\|_{L^{m,p}(\mathbb{R}^n)} \stackrel{(1b),(3)}{\simeq} M(f, R) = \widehat{M}(f) \simeq \|f\|_{L^{m,p}(E)},$$

which completes the proof of Theorem 1.

We now define the collections of linear functionals:

$$\widehat{\Omega} := \Omega \cup \{f \mapsto \partial^\alpha[R(f)](z) : \alpha \in \mathcal{M}\} \text{ and}$$

$$\widehat{\Xi} := \{\text{linear functionals } f \mapsto \xi(f, R(f)) \text{ with } \xi \in \Xi, \text{ counted with repetition}\}.$$

(For the definition of ‘‘counted with repetition’’, see the proof of **(E6)(c)** in the preceding section.)

From (2), we have $\#(\widehat{\Xi}) \leq \#(\Xi) \leq CN$, and also

$$\widehat{M}(f) = M(f, R) = \left(\sum_{\xi \in \widehat{\Xi}} |\xi(f)|^p \right)^{1/p}.$$

Together with (6), this yields the conclusion of Theorem 2.

By standard arguments (e.g., see the previous section), the extension operator \widehat{T} and the linear functionals belonging to $\widehat{\Xi}$ have $\widehat{\Omega}$ -assisted bounded depth. Since $\widehat{\Omega}$ consists of the assists Ω along with $\#(\mathcal{M}) = D$ (possibly) new linear functionals, it follows that

$$\sum_{\omega \in \widehat{\Omega}} \text{dp}(\omega) \leq \sum_{\omega \in \Omega} \text{dp}(\omega) + D \cdot \#(E) \leq C \cdot \#(E),$$

which proves Theorem 3. □

18.2. Inhomogeneous Sobolev spaces. Denote $W^{m,p}(E)$ for the space of real-valued functions on some finite subset $E \subset \mathbb{R}^n$, equipped with the norm

$$\|f\|_{W^{m,p}(E)} := \inf\{\|F\|_{W^{m,p}(\mathbb{R}^n)} : F = f \text{ on } E\}.$$

Here, we consider a variant of the problem solved in the previous section.

$W^{m,p}(\mathbb{R}^n)$ extension problem. Let $f : E \rightarrow \mathbb{R}$ be defined on a finite subset $E \subset \mathbb{R}^n$. Find $G \in W^{m,p}(\mathbb{R}^n)$ that depends linearly on f , with $G|_E = f$ and $\|G\|_{W^{m,p}(\mathbb{R}^n)} \leq C\|f\|_{W^{m,p}(E)}$.

We solve this problem, thereby proving the analogue of Theorem 1 for $W^{m,p}$ and finite E . The obvious analogues of Theorems 2 and 3 for $W^{m,p}$ also hold; we leave their consideration to the reader.

Let Q^1, Q^2, \dots be a tiling of \mathbb{R}^n by unit cubes. If $F^i \in W^{m,p}(1.3Q^i)$ is a near optimal extension of $f|_{E \cap 1.1Q^i}$ for each $i = 1, 2, \dots$, it is easy to verify that

$$G = \sum_{i=1}^{\infty} \theta^i F^i \in W^{m,p}(\mathbb{R}^n) \text{ is a near optimal extension of } f, \text{ where}$$

$\theta^1, \theta^2, \dots$ form a smooth partition of unity on \mathbb{R}^n , with θ^i supported on $1.1Q^i$.

Thus, we may assume that $E \subset (0.9)Q$ for some cube with sidelength $\delta_Q \simeq 1$, and solve for an F that depends linearly on f , such that $F = f$ on E and $\|F\|_{W^{m,p}(Q)}$ is nearly minimal. If we could do this, we could take the F^i to depend linearly on f , and ultimately we could take G to depend linearly on f .

Fix some $z \in 0.9Q$ with $0 < d(z, E) \leq 1$. We use the linear operator $T : L^{m,p}(E; z) \rightarrow L^{m,p}(\mathbb{R}^n)$ that satisfies (1a) and (1b). Let $P \in \mathcal{P}$ be arbitrary. By the Sobolev Inequality and (1a), we have

$$(7) \quad \begin{aligned} \|T(f, P)\|_{W^{m,p}(Q)}^p &\lesssim \|T(f, P)\|_{L^{m,p}(\mathbb{R}^n)}^p + \sum_{|\alpha| \leq m-1} |\partial^\alpha P(z)|^p \\ &\simeq \|(f, P)\|_{L^{m,p}(E; z)}^p + \sum_{|\alpha| \leq m-1} |\partial^\alpha P(z)|^p \quad (\text{thanks to (1b)}). \end{aligned}$$

On the other hand, let $H \in W^{m,p}(Q)$ be arbitrary with $H = f$ on E and $J_z H = P$. Choose $\theta \in C_0^\infty(Q)$ with $\theta \equiv 1$ on a neighborhood of $E \cup \{z\}$, and $|\partial^\alpha \theta| \lesssim 1$ when $|\alpha| \leq m$. Therefore, $\theta H \in W^{m,p}(\mathbb{R}^n)$ satisfies

$$\theta H = f \text{ on } E, \quad J_z(\theta H) = P, \quad \text{and} \quad \|\theta H\|_{W^{m,p}(\mathbb{R}^n)} \lesssim \|H\|_{W^{m,p}(Q)}.$$

Thus, by definition of the trace seminorm and the Sobolev Inequality, we have

$$(8) \quad \|(f, P)\|_{L^{m,p}(E; z)}^p + \sum_{|\alpha| \leq m-1} |\partial^\alpha P(z)|^p \lesssim \|\theta H\|_{W^{m,p}(\mathbb{R}^n)}^p \lesssim \|H\|_{W^{m,p}(Q)}^p.$$

Thus, $T(f, P) \in W^{m,p}(Q)$ is a near optimal extension of (f, P) . Moreover, from (7) and (8) and from (2) and (3), we have

$$\begin{aligned} \|T(f, P)\|_{W^{m,p}(Q)}^p &\simeq \|(f, P)\|_{L^{m,p}(E; z)}^p + \sum_{|\alpha| \leq m-1} |\partial^\alpha P(z)|^p \\ &\simeq \sum_{\xi \in \Xi} |\xi(f, P)|^p + \sum_{|\alpha| \leq m-1} |\partial^\alpha P(z)|^p. \end{aligned}$$

Choose a polynomial $R \in \mathcal{P}$ that depends linearly on f , for which $P = R$ minimizes the right-hand side above to within a universal constant factor. (This is possible thanks to Lemma 11.1.) Thus, $T(f, R) \in W^{m,p}(Q)$ is a near optimal extension of f . This also yields a solution to the $W^{m,p}(\mathbb{R}^n)$ extension problem, as mentioned before.

19. PASSAGE TO INFINITE E

In this section, we deduce Theorem 1 from the known case of finite E . Our plan is as follows.

Let $E \subset \mathbb{R}^n$ be infinite. Pick a countable subset

$$(1) \quad E^\circ = \{x_0, x_1, x_2, \dots\} \subset E, \text{ whose closure contains } E.$$

For each $N \geq 0$, define

$$(2) \quad E_N = \{x_0, x_1, \dots, x_N\}.$$

The known case of Theorem 1 produces a linear map $T_N : L^{m,p}(E_N) \rightarrow L^{m,p}(\mathbb{R}^n)$ such that

$$(3) \quad T_N f = f \text{ on } E_N$$

and

$$(4) \quad \|T_N f\|_{L^{m,p}(\mathbb{R}^n)} \leq C \|f\|_{L^{m,p}(E_N)}$$

for all $f \in L^{m,p}(E_N)$.

We hope to pass from T_N to T by taking a Banach limit as $N \rightarrow \infty$. Recall that a Banach limit is a linear map that carries an arbitrary bounded sequence $(t_N)_{N \geq 0}$ of real numbers to a real number denoted $\text{Blim}_{N \rightarrow \infty} t_N$. The defining properties of a Banach limit are

$$(5) \quad \text{Blim}_{N \rightarrow \infty} t_N = \lim_{N \rightarrow \infty} t_N \text{ whenever } \lim_{N \rightarrow \infty} t_N \text{ exists, and}$$

$$(6) \quad |\text{Blim}_{N \rightarrow \infty} t_N| \leq \limsup_{N \rightarrow \infty} |t_N|.$$

The existence of Banach limits is immediate from the Hahn-Banach theorem. See [6].

Thus, for $f \in L^{m,p}(E)$, we hope to define

$$(7) \quad T f(x) := \text{Blim}_{N \rightarrow \infty} [T_N(f|_{E_N})(x)] \quad \text{for } x \in \mathbb{R}^n$$

and then prove that T is an extension operator as in Theorem 1.

Unfortunately, without further ideas, the above plan is doomed. For instance, suppose $E \subset \{P_0 = 0\}$ for some polynomial $P_0 \in \mathcal{P}$. If T_N is an extension operator for $L^{m,p}(E_N)$, then so is

$$T_N^\# f(x) := T_N f(x) + \mu_N f(x_0) P_0(x) \quad (x \in \mathbb{R}^n),$$

where x_0 is as in (1) and μ_N is any real number. In fact, (3) and (4) hold for the $T_N^\#$, with the same constant as for T_N .

Since the sequence $(\mu_N)_{N \geq 0}$ is arbitrary, there is no way to guarantee that the sequence $(T_N(f|_{E_N})(x))_{N \geq 0}$ will be bounded for fixed x . Consequently, we cannot guarantee that the Banach limit (7) exists. This problem arises because $L^{m,p}(\mathbb{R}^n)$ carries a seminorm, not a norm.

To overcome this difficulty, we normalize the extension operators T_N as follows. Among all finite subsets $S \subset E$, we pick S_0 to minimize the dimension of the vector space $\mathcal{P}(S) = \{P \in \mathcal{P} : P = 0 \text{ on } S\}$. Such an S_0 exists, since any non-empty set of non-negative integers has a minimum.

For any $y \in E$, the subspace $\mathcal{P}(S_0 \cup \{y\}) \subset \mathcal{P}(S_0)$ has dimension no less than that of $\mathcal{P}(S_0)$. Therefore, $\mathcal{P}(S_0 \cup \{y\}) = \mathcal{P}(S_0)$. That is, any $P \in \mathcal{P}$ that vanishes on S_0 must also vanish at y . Thus,

$$(8) \quad S_0 \subset E \text{ is finite}$$

and

$$(9) \quad \text{any polynomial } P \in \mathcal{P} \text{ that vanishes on } S_0 \text{ must also vanish on } E.$$

Let $S_0 = \{y_0, y_1, \dots, y_L\}$. Without loss of generality, we may pick our x_0, x_1, \dots in (1) so that $x_i = y_i$ for $i = 0, \dots, L$. Therefore,

$$(10) \quad S_0 \subset E_N \quad \text{for all } N \geq L.$$

We pick any projection $\pi_0 : \mathcal{P} \rightarrow \mathcal{P}(S_0)$. We will establish the following results.

Lemma 19.1. *For $N \geq L$, the extension operators T_N in (3) and (4) can be picked to satisfy the additional condition*

$$(11) \quad \pi_0 J_{x_0} [T_N f] = 0 \quad \text{for all } f \in L^{m,p}(E_N), \text{ with } x_0 \text{ as in (1)}.$$

Lemma 19.2. *Suppose that the T_N satisfy (11) for $N \geq L$. Then, for any $f \in L^{m,p}(E)$ and for every cube Q , the functions $T_N(f|_{E_N})$ are bounded in $W^{m,p}(Q)$.*

If the T_N satisfy (11), then $[T_N(f|_{E_N})(x)]_{N \geq 0}$ is thus a bounded sequence for each fixed $x \in \mathbb{R}^n$ and $f \in L^{m,p}(E)$. Therefore, the Banach limit in (7) is well defined. Clearly, T is a linear map, taking functions $f \in W^{m,p}(E)$ to functions Tf defined on \mathbb{R}^n . Moreover, for each $k \geq 0$, we have

$$Tf(x_k) = \text{Blim}_{N \rightarrow \infty} T_N(f|_{E_N})(x_k) = f(x_k) \quad \text{with } x_k \text{ as in (1)},$$

simply because $x_k \in E_N$ for $N \geq k$, hence $T_N(f|_{E_N})(x_k) = f(x_k)$ for $N \geq k$; see (3) and (5). Thus,

$$(12) \quad Tf = f \text{ on } E^\circ \quad \text{for any } f \in L^{m,p}(\mathbb{R}^n), \text{ with } E^\circ \subset E \text{ as in (1)}.$$

We do not yet know that $Tf \in L^{m,p}(\mathbb{R}^n)$. Therefore, we prove the following result.

Lemma 19.3. *Let $Q \subset \mathbb{R}^n$ be a cube, and let F_0, F_1, \dots be a bounded sequence in $W^{m,p}(Q)$. Then the function F , defined by*

$$(13) \quad F(x) = \text{Blim}_{N \rightarrow \infty} F_N(x) \quad (x \in Q),$$

belongs to $W^{m,p}(Q)$, and we have

$$\|F\|_{L^{m,p}(Q)} \leq \limsup_{N \rightarrow \infty} \|F_N\|_{L^{m,p}(Q)}.$$

We apply the above lemma to $F_N = T_N(f|_{E_N})$ for $f \in L^{m,p}(E)$. Then $F = Tf$ is given by (13). Since $\|(f|_{E_N})\|_{L^{m,p}(E_N)} \leq \|f\|_{L^{m,p}(E)}$, the estimate (4) and Lemmas 19.2 and 19.3 together imply that Tf belongs to $L^{m,p}(Q)$ for any cube Q , and moreover

$$\|Tf\|_{L^{m,p}(Q)} \leq C \|f\|_{L^{m,p}(E)}.$$

Here we assume the T_N satisfy (11). Since $Q \subset \mathbb{R}^n$ is an arbitrary cube, it follows that $Tf \in L^{m,p}(\mathbb{R}^n)$ and that

$$(14) \quad \|Tf\|_{L^{m,p}(\mathbb{R}^n)} \leq C \|f\|_{L^{m,p}(E)} \quad \text{for all } f \in L^{m,p}(E).$$

Also, since the subset E° in (1) is dense in E , we conclude from (12) that

$$(15) \quad Tf = f \text{ on } E, \text{ for all } f \in L^{m,p}(E).$$

Thus, our extension operator T maps $L^{m,p}(E)$ to $L^{m,p}(\mathbb{R}^n)$ and satisfies (14) and (15). We therefore obtain Theorem 1, once we have Lemmas 19.1, 19.2, and 19.3.

19.1. Proof of Lemma 19.1. To prove Lemma 19.1, let $T_N : L^{m,p}(E_N) \rightarrow L^{m,p}(\mathbb{R}^n)$ satisfy (3) and (4). For $f \in L^{m,p}(E_N)$, define

$$\tilde{T}_N f = T_N f - \pi_0 J_{x_0}(T_N f) \in L^{m,p}(\mathbb{R}^n) \quad \text{since } \pi_0 J_{x_0}(T_N f) \in \mathcal{P}.$$

Note that $\pi_0 J_{x_0}(\tilde{T}_N f) = \pi_0 [J_{x_0}(T_N f) - \pi_0 J_{x_0}(T_N f)] = 0$, where we have used the fact that $J_{x_0}(T_N f) \in \mathcal{P}$ (and the fact that $\pi_0^2 = \pi_0$). Also, since $\pi_0 J_{x_0}(T_N f) \in \mathcal{P}$, we have $\|\pi_0 J_{x_0}(T_N f)\|_{L^{m,p}(\mathbb{R}^n)} = 0$; hence

$$\|\tilde{T}_N f\|_{L^{m,p}(\mathbb{R}^n)} = \|T_N f\|_{L^{m,p}(\mathbb{R}^n)} \leq C \|f\|_{L^{m,p}(E_N)} \quad \text{by (4)}.$$

Finally, since $\pi_0 J_{x_0}(T_N f) \in \mathcal{P}(S_0)$, we have $\pi_0 J_{x_0}(T_N f) = 0$ on S_0 , hence also on E , by (9). Therefore,

$$\tilde{T}_N f = f \text{ on } E, \text{ by (3).}$$

Thus, $\tilde{T}_N : L^{m,p}(E_N) \rightarrow L^{m,p}(\mathbb{R}^n)$ is a linear map that satisfies (3), (4) and (11). The proof of Lemma 19.1 is complete.

19.2. Proof of Lemma 19.2. To establish Lemma 19.2, we use the following result.

Lemma 19.4. *For any cube Q containing x_0 there exist constants $A_1(Q), A_2(Q) > 0$ such that for all $F \in W^{m,p}(Q)$ with $\pi_0 J_{x_0}(F) = 0$, we have the estimate*

$$(16) \quad \sum_{|\alpha| \leq m-1} |\partial^\alpha F(x_0)| \leq A_1(Q) \cdot \left[\|F\|_{L^{m,p}(Q)} + \max_{y \in S_0} |F(y)| \right],$$

and therefore

$$(17) \quad \|F\|_{W^{m,p}(Q)} \leq A_2(Q) \cdot \left[\|F\|_{L^{m,p}(Q)} + \max_{y \in S_0} |F(y)| \right].$$

Proof. Fix a cube Q containing x_0 , and suppose (16) fails. Then there exists a sequence of functions $F_N \in W^{m,p}(Q)$ ($N \geq 0$) such that

$$(18) \quad \sum_{|\alpha| \leq m-1} |\partial^\alpha F_N(x_0)| = 1$$

and

$$(19) \quad \pi_0 J_{x_0}(F_N) = 0,$$

but

$$(20) \quad \|F_N\|_{L^{m,p}(Q)} \rightarrow 0 \text{ as } N \rightarrow \infty \text{ and}$$

$$(21) \quad \max_{y \in S_0} |F_N(y)| \rightarrow 0 \text{ as } N \rightarrow \infty.$$

By (18) and (20), the functions F_N are bounded in $W^{m,p}(Q)$, and hence in $C^{m-1,s}(Q)$ with $s = 1 - n/p \in (0,1)$. By Ascoli's theorem, a subsequence of the (F_N) converges in $C^{m-1}(Q)$ to a function $F \in C^{m-1,s}(Q)$. From (18)-(21), we

obtain the following properties of F :

(22)

$$\sum_{|\alpha| \leq m-1} |\partial^\alpha F(x_0)| = 1;$$

(23)

$$\pi_0 J_{x_0}(F) = 0;$$

(24)

$$\|F\|_{\dot{C}^{m-1,s}(Q)} \leq \limsup_{N \rightarrow \infty} \|F_N\|_{\dot{C}^{m-1,s}(Q)} \leq C \limsup_{N \rightarrow \infty} \|F_N\|_{L^{m,p}(Q)} = 0; \text{ and}$$

(25)

$$F = 0 \text{ on } S_0.$$

From (24) we see that F is a polynomial, $F \in \mathcal{P}$. By (25), we have $F \in \mathcal{P}(S_0)$, hence $\pi_0 J_{x_0}(F) = \pi_0 F = F$. Therefore, $F = 0$ by (23). This contradicts (22).

The above contradiction shows that (16) cannot fail. Conclusion (17) follows at once from (16) and the Sobolev Inequality. The proof of Lemma 19.4 is complete. \square

To prove Lemma 19.2, we fix a cube $Q \subset \mathbb{R}^n$. Without loss of generality, we may suppose that Q contains x_0 .

For fixed $f \in L^{m,p}(E)$ and for any $N \geq L$, we apply Lemma 19.4 to $F_N = T_N(f|_{E_N})$. Note that $F_N \in W^{m,p}(Q)$, and $\pi_0 J_{x_0}(F_N) = 0$ by (11). Hence, Lemma 19.4 applies, and we learn that

$$(26) \quad \|F_N\|_{W^{m,p}(Q)} \leq A_2(Q) \cdot \left[\|F_N\|_{L^{m,p}(Q)} + \max_{y \in S_0} |F_N(y)| \right].$$

On the other hand (4) yields the estimates

$$(27) \quad \|F_N\|_{L^{m,p}(Q)} \leq \|F_N\|_{L^{m,p}(\mathbb{R}^n)} \leq C \|(f|_{E_N})\|_{L^{m,p}(E_N)} \leq C \|f\|_{L^{m,p}(E)}.$$

Also, since $S_0 \subset E_N$, (3) yields $F_N = f|_{E_N}$ on S_0 , i.e.,

$$(28) \quad F_N = f \text{ on } S_0.$$

Putting (27) and (28) into (26), we find that

$$\|F_N\|_{W^{m,p}(Q)} \leq CA_2(Q) \left[\|f\|_{L^{m,p}(E)} + \max_{y \in S_0} |f(y)| \right] \text{ for } N \geq L.$$

Therefore, the functions F_0, F_1, F_2, \dots form a bounded subset of $W^{m,p}(Q)$, completing the proof of Lemma 19.2.

19.3. Proof of Lemma 19.3. The proof of Lemma 19.3 uses the following simple observation. As before, we take $s = 1 - n/p \in (0, 1)$.

Lemma 19.5. *Let $A > 0$ be a constant, and let $Q \subset \mathbb{R}^n$ be a cube.*

For each multi-index α of order $|\alpha| \leq m-1$, let $f^{(\alpha)}$ be a function on Q . Assume the following estimates:

$$(29) \quad |f^{(\alpha)}(x+h) - f^{(\alpha)}(x) - \sum_{j=1}^n f^{(\alpha+\mathbb{1}_j)}(x)h_j| \leq A|h|^{1+s}$$

$$\text{for } |\alpha| \leq m-2, x \in Q, h = (h_1, \dots, h_n), x+h \in Q,$$

$$(30) \quad |f^{(\alpha)}(x) - f^{(\alpha)}(y)| \leq A|x-y|^s \text{ for } |\alpha| \leq m-1, x, y \in Q.$$

(Here, $\mathbb{1}_j$ denotes the j^{th} unit multi-index so that $\partial^{\mathbb{1}_j} = \frac{\partial}{\partial x_j}$.)

Then $f^{(0)} \in C^{m-1,s}(Q)$, and $f^{(\alpha)} = \partial^\alpha f^{(0)}$ on Q for each α of order $|\alpha| \leq m-1$.

Proof. Hypotheses (29) and (30) show that, for $|\alpha| \leq m-2$, the function $f^{(\alpha)}$ belongs to $C^1(Q)$, and $\partial_{x_j} f^{(\alpha)} = f^{(\alpha+\mathbb{1}_j)}$. Thus, $f^{(0)} \in C^{m-1}(Q)$ and $\partial^\alpha f^{(0)} = f^{(\alpha)}$ for $|\alpha| \leq m-1$. Hypothesis (30) now shows that the derivatives of $f^{(0)}$ up to order $m-1$ are Lipschitz- s . Hence $f^{(0)} \in C^{m-1,s}(Q)$, completing the proof of Lemma 19.5. \square

Proof of Lemma 19.3. Let Q, F_0, F_1, \dots, F be as in the hypotheses of Lemma 19.3. For $|\alpha| \leq m-1$, $N \geq 0$, let

$$F_N^{(\alpha)} = \partial^\alpha F_N \in C^{m-1-|\alpha|,s}(Q).$$

Since the F_N are bounded in $W^{m,p}(Q)$, the following estimates hold for some constant A :

$$(31) \quad |F_N^{(\alpha)}(x)| \leq A \quad \text{for } x \in Q,$$

$$(32) \quad |F_N^{(\alpha)}(x+h) - F_N^{(\alpha)}(x) - \sum_{j=1}^n F_N^{(\alpha+\mathbb{1}_j)}(x)h_j| \leq A|h|^{1+s}$$

for $|\alpha| \leq m-2$, $x \in Q$, $h = (h_1, \dots, h_n)$, $x+h \in Q$,

$$(33) \quad |F_N^{(\alpha)}(x) - F_N^{(\alpha)}(y)| \leq A|x-y|^s \quad \text{for } |\alpha| \leq m-1, x, y \in Q.$$

Thanks to (31), the Banach limit

$$(34) \quad F^{(\alpha)}(x) = \text{Blim}_{N \rightarrow \infty} F_N^{(\alpha)}(x) \quad (x \in Q)$$

is well defined. Note that

$$(35) \quad F^{(0)}(x) = \text{Blim}_{N \rightarrow \infty} F_N(x) = F(x), \quad \text{with } F \text{ as in the statement of Lemma 19.3.}$$

Applying (6), we may pass to the Banach limit and deduce from (31), (32), and (33) for $F_N^{(\alpha)}$ the corresponding estimates for the $F^{(\alpha)}$. Lemma 19.5 now shows that $F = F^{(0)} \in C^{m-1,s}(Q)$ and that $\partial^\alpha F = F^{(\alpha)}$ for each $|\alpha| \leq m-1$.

Hence, to prove that $F \in W^{m,p}(Q)$ and that

$$\|F\|_{L^{m,p}(Q)} \leq \limsup_{N \rightarrow \infty} \|F_N\|_{L^{m,p}(Q)},$$

it is enough to prove that

$$(36) \quad \left| \int_Q \partial_{x_j} \varphi \cdot F^{(\alpha)} dx \right| \leq \limsup_{N \rightarrow \infty} \|F_N\|_{L^{m,p}(Q)} \cdot \|\varphi\|_{L^{p'}(Q)}$$

for any test function $\varphi \in C_0^\infty(Q)$ and any $|\alpha| = m-1$, $j \in \{1, \dots, n\}$. Here, p' is the dual exponent to p , and we use the definition of the $L^{m,p}$ -seminorm:

$$\|F\|_{L^{m,p}(Q)}^p = \max_{|\alpha|=m} \int_Q |\partial^\alpha F(x)|^p dx.$$

Thus, Lemma 19.3 reduces to the task of proving (36) for any given $\varphi \in C_0^\infty(Q)$, $|\alpha| = m-1$, $j \in \{1, \dots, n\}$. Fix such φ, α, j , and let M be any real number greater than $\limsup_{N \rightarrow \infty} \|F_N\|_{L^{m,p}(Q)}$.

For N large enough we have $\|F_N\|_{L^{m,p}(Q)} \leq M$, hence

$$(37) \quad \left| \int_Q \partial_{x_j} \varphi \cdot F_N^{(\alpha)} dx \right| \leq M \|\varphi\|_{L^{p'}(Q)}$$

since $F_N^{(\alpha)} = \partial^\alpha F_N$.

We will derive (36) by passing to the Banach limit in (37). To do so, we simply approximate the integrals in (36) and (37) by Riemann sums.

We know that

$$(38) \quad |F_N^{(\alpha)}(x) - F_N^{(\alpha)}(y)| \leq A|x - y|^s \quad \text{and} \quad |F^{(\alpha)}(x) - F^{(\alpha)}(y)| \leq A|x - y|^s$$

for $x, y \in Q$, with A independent of N .

Let $\delta > 0$ be a small number (later, we will take $\delta \rightarrow 0^+$), and let $\{Q_\nu\}$ be a partition of Q into subcubes with center $(Q_\nu) = z_\nu$ and with sidelength $\delta_{Q_\nu} < \delta$. Then (31) and (38) together imply the estimates

$$(39) \quad \left| \int_Q \partial_{x_j} \varphi \cdot F_N^{(\alpha)} dx - \sum_{\nu=1}^{\nu_{\max}} \partial_{x_j} \varphi(z_\nu) \cdot F_N^{(\alpha)}(z_\nu) \cdot \delta_{Q_\nu}^n \right| \leq CA\delta^s \cdot \delta_{Q_\nu}^n$$

and

$$(40) \quad \left| \int_Q \partial_{x_j} \varphi \cdot F^{(\alpha)} dx - \sum_{\nu=1}^{\nu_{\max}} \partial_{x_j} \varphi(z_\nu) \cdot F^{(\alpha)}(z_\nu) \cdot \delta_{Q_\nu}^n \right| \leq CA\delta^s \cdot \delta_{Q_\nu}^n,$$

with C independent of N .

In particular, (39) shows that the sequence $\left(\int_Q \partial_{x_j} \varphi \cdot F_N^{(\alpha)} dx \right)_{N \geq 0}$ is bounded.

Since $F^{(\alpha)}(z_\nu) = \text{Blim}_{N \rightarrow \infty} F_N^{(\alpha)}(z_\nu)$ for each ν , property (6) of the Banach limit, together with (39), implies that

$$(41) \quad \left| \left[\text{Blim}_{N \rightarrow \infty} \int_Q \partial_{x_j} \varphi \cdot F_N^{(\alpha)} dx \right] - \sum_{\nu=1}^{\nu_{\max}} \partial_{x_j} \varphi(z_\nu) \cdot F^{(\alpha)}(z_\nu) \cdot \delta_{Q_\nu}^n \right| \leq CA\delta^s \cdot \delta_{Q_\nu}^n.$$

(In (41), the Banach limit in square brackets is well defined, thanks to (37).)

Comparing (40) and (41), we see that

$$\left| \int_Q \partial_{x_j} \varphi \cdot F^{(\alpha)} dx - \left[\text{Blim}_{N \rightarrow \infty} \int_Q \partial_{x_j} \varphi \cdot F_N^{(\alpha)} dx \right] \right| \leq 2CA\delta^s \cdot \delta_{Q_\nu}^n.$$

Since $\delta > 0$ is arbitrarily small, we conclude that

$$(42) \quad \int_Q \partial_{x_j} \varphi \cdot F^{(\alpha)} dx = \text{Blim}_{N \rightarrow \infty} \int_Q \partial_{x_j} \varphi \cdot F_N^{(\alpha)} dx.$$

From (37), (42) and property (6) of the Banach limit, we obtain the estimate

$$(43) \quad \left| \int_Q \partial_{x_j} \varphi \cdot F^{(\alpha)} dx \right| \leq M \|\varphi\|_{L^{p'}(Q)}.$$

This implies the desired estimate (36), since M in (43) is an arbitrary real number greater than $\limsup_{N \rightarrow \infty} \|F_N\|_{L^{m,p}(Q)}$.

We reduced the proof of Lemma 19.3 to (36), and we have now proven (36). This completes the proof of Lemma 19.3 and, with it, the proof of Theorem 1.

19.4. **Epilogue.** Given $E \subset \mathbb{R}^n$ (possibly infinite), we can also “construct” a linear extension operator $T : W^{m,p}(E) \rightarrow W^{m,p}(\mathbb{R}^n)$ by applying the case of finite E and passing to a Banach limit. This exercise is a much easier version of the argument we just explained for $L^{m,p}$. Since $\|\cdot\|_{W^{m,p}(\mathbb{R}^n)}$ is a norm (rather than a seminorm), the pitfalls that we worked to avoid will no longer arise. We leave the details to the reader.

By using a Banach limit, we have sacrificed all knowledge of the structure of our linear extension operator T . It would be very interesting to gain some understanding of that structure. Such an understanding is achieved for the C^m case in [11] and for the deeper case of $C^{m,s}$ and related spaces in [18].

ACKNOWLEDGMENTS

The authors are grateful to B. Klartag and A. Naor for introducing them to the idea of sparsification and for several lively discussions. They are also grateful to the NSF, the ONR, and the American Institute of Mathematics (AIM) for generous support. Key ideas arose during fruitful workshops held at AIM and at the College of William and Mary. The authors are grateful to P. Shvartsman and Nahum Zobin for lively discussions.

REFERENCES

- [1] Joshua D. Batson, Daniel A. Spielman, and Nikhil Srivastava, *Twice-Ramanujan sparsifiers*, STOC’09—Proceedings of the 2009 ACM International Symposium on Theory of Computing, ACM, New York, 2009, pp. 255–262. MR2780071 (2012f:05169)
- [2] Edward Bierstone, Pierre D. Milman, and Wiesław Pawłucki, *Differentiable functions defined in closed sets. A problem of Whitney*, Invent. Math. **151** (2003), no. 2, 329–352, DOI 10.1007/s00222-002-0255-6. MR1953261 (2004h:58009)
- [3] Yuri Brudnyi and Pavel Shvartsman, *Generalizations of Whitney’s extension theorem*, Internat. Math. Res. Notices **3** (1994), 129 ff., approx. 11 pp. (electronic), DOI 10.1155/S1073792894000140. MR1266108 (95c:58018)
- [4] Yuri Brudnyi and Pavel Shvartsman, *The Whitney problem of existence of a linear extension operator*, J. Geom. Anal. **7** (1997), no. 4, 515–574, DOI 10.1007/BF02921632. MR1669235 (2000a:46051)
- [5] Paul B. Callahan and S. Rao Kosaraju, *A decomposition of multidimensional point sets with applications to k -nearest-neighbors and n -body potential fields*, J. Assoc. Comput. Mach. **42** (1995), no. 1, 67–90, DOI 10.1145/200836.200853. MR1370371 (96i:68033)
- [6] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part I*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988. General theory; With the assistance of William G. Bade and Robert G. Bartle; Reprint of the 1958 original; A Wiley-Interscience Publication. MR1009162 (90g:47001a)
- [7] Lawrence C. Evans, *Partial differential equations*, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, RI, 1998. MR1625845 (99e:35001)
- [8] Charles L. Fefferman, *A sharp form of Whitney’s extension theorem*, Ann. of Math. (2) **161** (2005), no. 1, 509–577, DOI 10.4007/annals.2005.161.509. MR2150391 (2006h:58008)
- [9] Charles Fefferman, *Whitney’s extension problem for C^m* , Ann. of Math. (2) **164** (2006), no. 1, 313–359, DOI 10.4007/annals.2006.164.313. MR2233850 (2007g:58013)
- [10] Charles Fefferman, *C^m extension by linear operators*, Ann. of Math. (2) **166** (2007), no. 3, 779–835, DOI 10.4007/annals.2007.166.779. MR2373373 (2009b:46069)
- [11] Charles Fefferman, *The structure of linear extension operators for C^m* , Rev. Mat. Iberoam. **23** (2007), no. 1, 269–280, DOI 10.4171/RMI/495. MR2351135 (2009c:41003)
- [12] Charles Fefferman, *Fitting a C^m -smooth function to data. III*, Ann. of Math. (2) **170** (2009), no. 1, 427–441, DOI 10.4007/annals.2009.170.427. MR2521121 (2011e:58014)
- [13] Charles Fefferman and Arie Israel, *The jet of an interpolant on a finite set*, Rev. Mat. Iberoam. **27** (2011), no. 1, 355–360, DOI 10.4171/RMI/639. MR2815741 (2012f:41022)

- [14] Charles Fefferman and Bo'az Klartag, *Fitting a C^m -smooth function to data. I*, Ann. of Math. (2) **169** (2009), no. 1, 315–346, DOI 10.4007/annals.2009.169.315. MR2480607 (2011g:58011)
- [15] Charles Fefferman and Bo'az Klartag, *Fitting a C^m -smooth function to data. II*, Rev. Mat. Iberoam. **25** (2009), no. 1, 49–273, DOI 10.4171/RMI/569. MR2514338 (2011f:58020)
- [16] Georges Glaeser, *Étude de quelques algèbres tayloriennes*, J. Analyse Math. **6** (1958), 1–124; erratum, insert to 6 (1958), no. 2 (French). MR0101294 (21 #107)
- [17] A. Israel, *A bounded linear extension operator for $L^{2,p}(\mathbb{R}^2)$* , Annals of Math. (to appear).
- [18] Garving K. Luli, *$C^{m,\omega}$ extension by bounded-depth linear operators*, Adv. Math. **224** (2010), no. 5, 1927–2021, DOI 10.1016/j.aim.2010.01.027. MR2646115 (2011g:58012)
- [19] G.K. Luli, *Sobolev extension in one-dimension* (2008), notes available at <http://www.math.princeton.edu/~gluli/TH/notes.pdf>.
- [20] P. Shvartsman, *Sobolev W_p^1 -spaces on closed subsets of \mathbf{R}^n* , Adv. Math. **220** (2009), no. 6, 1842–1922, DOI 10.1016/j.aim.2008.09.020. MR2493183 (2010b:46077)
- [21] P. Shvartsman, *Lipschitz spaces generated by the Sobolev-Poincaré inequality and extensions of Sobolev functions* (preprint) (2011).
- [22] P. Shvartsman, *On the sum of a Sobolev space and a weighted L_p -space* (preprint) (2011).
- [23] Daniel A. Spielman and Nikhil Srivastava, *Graph sparsification by effective resistances*, STOC'08, ACM, New York, 2008, pp. 563–568, DOI 10.1145/1374376.1374456. MR2582687 (2011c:05340)
- [24] Daniel A. Spielman and Shang-Hua Teng, *Nearly-linear time algorithms for graph partitioning, graph sparsification, and solving linear systems*, Proceedings of the 36th Annual ACM Symposium on Theory of Computing, ACM, New York, 2004, pp. 81–90 (electronic), DOI 10.1145/1007352.1007372. MR2121588 (2005i:05188)
- [25] Hassler Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc. **36** (1934), no. 1, 63–89, DOI 10.2307/1989708. MR1501735
- [26] Hassler Whitney, *Differentiable functions defined in closed sets. I*, Trans. Amer. Math. Soc. **36** (1934), no. 2, 369–387, DOI 10.2307/1989844. MR1501749
- [27] Hassler Whitney, *Functions differentiable on the boundaries of regions*, Ann. of Math. (2) **35** (1934), no. 3, 482–485, DOI 10.2307/1968745. MR1503174

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, FINE HALL, WASHINGTON ROAD, PRINCETON, NJ 08544

DEPARTMENT OF MATHEMATICS, NEW YORK UNIVERSITY-COURANT INSTITUTE, WARREN WEAVER HALL, 251 MERCER STREET, NEW YORK, NY 10012-1185

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06520