

1. P. APPELL, "Sur une classe de polynômes à deux variables et le calcul approché des intégrales doubles," *Annales de la Faculté des Sciences de Toulouse*, v. 4, 1890, p. H.1-H.20.
2. C. BIRINDELLI, "Su nuove formule interpolatorie del Picone per funzioni in piu variabili e loro contributo al calcolo numerico degli integrali multipli," *Compositio Mathematica*, v. 10, 1952, p. 117-167.
3. W. BURNSIDE, "An approximate quadrature formula," *Messenger of Math.*, v. 37, 1908, p. 166-167.
4. C. F. GAUSS, "Methodus nova integralium valores per approximationem inveniendi," *Werke*, Göttingen, v. 3, 1866, p. 163-196.
5. P. C. HAMMER, O. J. MARLOWE, & A. H. STROUD, "Numerical integration over simplexes and cones," *MTAC*, v. 10, 1956, p. 130-137.
6. P. C. HAMMER & ARTHUR H. STROUD, "Numerical integration over simplexes," *MTAC*, v. 10, 1956, p. 137-139.
7. PRESTON C. HAMMER & A. WAYNE WYMORE, "Numerical evaluation of multiple integrals I," *MTAC*, v. 11, 1957, p. 59-67.
8. J. C. MAXWELL, "On approximate multiple integration between limits by summation," *Camb. Phil. Soc., Proc.*, v. 3, 1877, p. 39-47.
9. M. PICONE, "Vedute generali sull' interpolazione e qualche loro conseguenza," *Annali della Scuola Normale Superiore di Pisa*, Ser. III, v. 5, 1951, p. 193-244.
10. MICHAEL SADOWSKY, "A formula for approximate computation of a triple integral," *Amer. Math. Mon.*, v. 47, 1940, p. 539-543.
11. G. SZEGÖ, *Orthogonal Polynomials*, Amer. Math. Soc. Colloquium Publication, v. 23, New York, 1939.
12. G. W. TYLER, "Numerical integration of functions of several variables," *Can. J. Math.*, v. 5, 1953, p. 393-412.
13. RICHARD VON MISES, "Numerische Berechnung mehrdimensionaler Integrale," *Z. Angew. Math. Mech.*, v. 34, 1954, p. 201-210.

## On a Generalization of the Prime Pair Problem

**1. Introduction.** One of the many unsolved problems in the theory of prime numbers concerns the celebrated conjecture by A. de Polignac [1] about primes which differ by two. Prime pairs (or twin primes) are generally believed to constitute an infinite set but to date no one has succeeded in proving or disproving this conjecture.

It was established by Brun [2] that the sum of the reciprocals of all prime pairs is bounded but whether this boundedness is due to the finiteness of the set or to the "thinness" of an infinite set has not been determined.

**2. Prime  $n$ -tuples.** In generalizing from prime pairs to prime  $n$ -tuples we must first define what is meant by a prime  $n$ -tuple.

*Definition:* a prime  $n$ -tuple is a set of  $n$  odd primes  $p_1, p_2, \dots, p_n$  such that the difference  $\Delta n = p_n - p_1$  is a non-trivial minimum.

The meaning of "non-trivial" will now be explained. For  $n = 3$  we can exhibit a set of three consecutive odd primes such that  $\Delta 3 = 4$  (i.e., 3, 5, 7). We consider this a trivial minimum, however, because (with one exception) in every set of three consecutive odd integers one of the integers in the set is composite since it is divisible by 3. The single exception to this rule is the case where the number divisible by 3 is the number 3 itself. Generally, we consider a set of  $n$  consecutive primes a trivial prime  $n$ -tuple if one of the primes of the set is *always* a factor of one of the terms of any sequence of consecutive odd integers having the same value for  $\Delta n$ .



determined above we shall have proved that  $\Delta 8 = 26$ . The following primes complete the proof.

11, 13, 17, 19, 23, 29, 31, 37.

When conditions (A) and (B) have been established, the procedure described here gives *all* the essentially different possible types of  $n$ -tuples for a given  $n$  and by solving one or more linear indeterminate equations the exact form of each type of  $n$ -tuple can be obtained. Thus, for  $n = 8$ , there are, at most, three different types of 8-tuples, viz.,

- (a)  $210x + 11, 13, 17, 19, 23, 29, 31, 37$
- (b)  $210x + 173, 179, 181, 187, 191, 193, 197, 199$
- (c)  $30x + 17, 19, 23, 29, 31, 37, 41, 43$ .

**5. Some further questions.** The preceding discussion raises some new questions:

(3) Are there values of  $n$  such that  $\Delta n > U + 2$ ?

The writer has verified that  $\Delta n = U + 2$  for all  $n \leq 10$ . On the other hand an exhaustive search of the first two billion numbers failed to disclose a single 19-tuple of the types prescribed by the above procedure.

(4) Are there some possible types of  $n$ -tuples for which no  $n$ -tuple exists?

For example, there are, at most, two types of 10-tuples; a 10-tuple for one type has been exhibited but not for the other; one possibility is that none exists for the other type.

**6. Numerical results.** The writer has computed  $\Delta n$  for  $n \leq 26$  and determined all the possible types of  $n$ -tuples for each  $n$  in this range; these are tabulated in Table I. It is convenient to describe a given type of  $n$ -tuple by specifying the smallest prime in the sequence with respect to an appropriate modulus followed by the succession of differences between the constituent primes. It is observed that a kind of "duality" exists between the different types of  $n$ -tuples for a given  $n$  such that if the set of differences for a given  $n$ -tuple is known, its dual will have these same differences in reverse order. Some  $n$ -tuples are self dual and do not yield distinct types, such as  $n = 6$ . An examination of Table I shows that for  $n = 2, 4$ , and 6 there is only one type of  $n$ -tuple; for  $n = 8$ , there are 3 types; for  $n = 3, 9, 15$ , and 22 there are 4 types; for  $n = 13$  there are 6 types; for all others in the table two types are given, each the dual of the other. In a recent attempt to discover another  $n$ -tuple that is self dual the writer ascertained that for  $n = 41$  there are 8 distinct types of possible 41-tuples.

The numerical results obtained in this investigation were performed on an IBM 701 electronic calculator. Four separate programs were written. Program I was designed to search for 4-tuples; an exhaustive search of all numbers  $\leq 5,073,379$  was made and 549 4-tuples were discovered.

Program II was designed to search for 7-tuples of the types

- (A)  $210x + 11, 13, 17, 19, 23, 29, 31$
- (B)  $210x + 179, 181, 187, 191, 193, 197, 199$ .

Seventeen 7-tuples of type (A) and twenty-four of type (B) were discovered in an exhaustive search of all numbers  $\leq 157,131,419$ .

TABLE I

$n$	$\Delta n$	$m$	$p \pmod{m}$	differences
2	2	6	5	2
3	6	30	11	2,4
3	6	30	17	2,4
3	6	30	7	4,2
3	6	30	13	4,2
4	8	30	11	2,4,2
5	12	30	11	2,4,2,4
5	12	30	7	4,2,4,2
6	16	210	97	4,2,4,2,4
7	20	210	11	2,4,2,4,6,2
7	20	210	179	2,6,4,2,4,2
8	26	30	17	2,4,6,2,6,4,2
8	26	210	11	2,4,2,4,6,2,6
8	26	210	173	6,2,6,4,2,4,2
9	30	30	13	4,2,4,6,2,6,4,2
9	30	30	17	2,4,6,2,6,4,2,4
9	30	210	11	2,4,2,4,6,2,6,4
9	30	210	169	4,6,2,6,4,2,4,2
10	32	210	11	2,4,2,4,6,2,6,4,2
10	32	210	167	2,4,6,2,6,4,2,4,2
11	36	2310	1271	2,4,2,4,6,2,6,4,2,4
11	36	2310	1003	4,2,4,6,2,6,4,2,4,2
12	42	2310	1271	2,4,2,4,6,2,6,4,2,4,6
12	42	2310	997	6,4,2,4,6,2,6,4,2,4,2
13	48	2310	1271	2,4,2,4,6,2,6,4,2,4,6,6
13	48	2310	991	6,6,4,2,4,6,2,6,4,2,4,2
13	48	2310	1259	2,10,2,4,2,4,6,2,6,4,2,4
13	48	2310	1003	4,2,4,6,2,6,4,2,4,2,10,2
13	48	2730	1199	2,6,6,4,2,4,6,2,6,4,2,4
13	48	2730	1483	4,2,4,6,2,6,4,2,4,6,2,6,4,2
14	50	30030	15131	2,4,2,4,6,2,6,4,2,4,6,6,2
14	50	30030	14849	2,6,6,4,2,4,6,2,6,4,2,4,2
15	56	30030	15131	2,4,2,4,6,2,6,4,2,4,6,6,2,6
15	56	30030	14843	6,2,6,6,4,2,4,6,2,6,4,2,4,2
15	56	210	17	2,4,6,2,6,4,2,4,6,6,2,6,4,2
15	56	210	137	2,4,6,2,6,6,4,2,4,6,2,6,4,2
16	60	30030	6943	4,2,4,6,2,6,4,2,4,6,6,2,6,4,2
16	60	30030	23027	2,4,6,2,6,6,4,2,4,6,2,6,4,2,4
17	66	30030	6943	4,2,4,6,2,6,4,2,4,6,6,2,6,4,2,6
17	66	30030	23021	6,2,4,6,2,6,6,4,2,4,6,2,6,4,2,4
18	70	30030	6943	4,2,4,6,2,6,4,2,4,6,6,2,6,4,2,6,4
18	70	30030	23017	4,6,2,4,6,2,6,6,4,2,4,6,2,6,4,2,4
19	76	5 10510	2 17153	4,2,4,6,2,6,4,2,4,6,6,2,6,4,2,6,4,6
19	76	5 10510	2 93281	6,4,6,2,4,6,2,6,6,4,2,4,6,2,6,4,2,4
20	80	30030	29	2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,2,4,2
20	80	30030	29921	2,4,2,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2
21	84	30030	29	2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,2,4,2,4
21	84	30030	29917	4,2,4,2,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2
22	90	5 10510	19	4,6,2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,2,4,2
22	90	5 10510	5 10401	2,4,2,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2,6,4
22	90	5 10510	23	6,2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,2,4,2,4
22	90	5 10510	5 10397	4,2,4,2,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2,6
23	94	5 10510	19	4,6,2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,2,4,2,4
23	94	5 10510	5 10397	4,2,4,2,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2,6,4
24	100	5 10510	2 17153	4,2,4,6,2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,6,2,4
24	100	5 10510	2 93257	4,2,6,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2,6,4,2,4
25	110	30030	29	2,6,4,2,4,6,6,2,6,4,2,6,4,6,8,4,2,4,2,4,14,4,6,2
25	110	30030	29891	2,6,4,14,4,2,4,2,4,8,6,4,6,2,4,6,2,6,6,4,2,4,6,2
26	114	96 99690	34 64999	2,12,4,2,4,6,2,6,4,2,4,6,8,6,4,2,6,4,6,8,4,2,4,2,4
26	114	96 99690	62 34577	4,2,4,2,4,8,6,4,6,2,4,6,8,6,4,2,4,6,2,6,4,2,4,12,2

TABLE II

<i>1st term of n-tuple</i>	<i>n</i>	<i>Type</i>
11	10	A
1277	9	C
5639	7	B
88789	9	B
1 13143	9	C
1 13147	9	C
1 65701	7	A
2 84723	8	B
6 26609	7	B
8 55709	9	B
10 68701	7	A
11 46773	8	B
25 80647	8	C
65 60993	8	B
75 40439	7	B
85 73429	7	B
119 00501	7	A
157 60091	8	A
178 43459	7	B
185 04371	7	A
190 89599	7	B
207 37877	8	C
210 36131	7	A
240 01709	7	B
256 58441	8	A
394 31921	7	A
429 81929	7	B
435 34019	7	B
450 02591	7	A
678 16361	7	A
691 56533	8	B
733 73537	8	C
742 66249	9	B
761 70527	8	C
792 08399	7	B
804 27029	7	B
841 04549	7	B
868 18211	7	A
879 88709	7	B
936 25991	8	A
1006 58627	8	C
1240 66079	7	B
1247 16071	7	A
1284 69149	7	B
1347 64997	8	C
1362 61241	7	A
1379 43347	8	C
1401 17051	7	A
1442 14319	7	B
1546 35191	7	A
1571 31419	7	B

Program III was designed to search for 8-tuples of the type

$$(C) \quad 30x + 17, 19, 23, 29, 31, 37, 41, 43.$$

Ten 8-tuples of this type were obtained in a search of all numbers  $\leq 137,943,347$ .

Program IV was the unsuccessful attempt to find a 19-tuple. As a check on the above calculations program II was recently re-run on an IBM 650 magnetic drum calculator; there was 100 percent agreement between the results of the two machines.

An examination of Table I shows that all  $n$ -tuples in a given range of numbers for  $7 \leq n \leq 15$  can be determined by examining only the  $n$ -tuples obtained from the three types given above. This subsequent calculation yielded three 8-tuples and one 10-tuple for type (A), four 8-tuples and three 9-tuples for type (B) and three 9-tuples for type (C). These results are tabulated in Table II. Since a 9-tuple may also be an 8-tuple, and a 7-tuple, etc., the highest ranking designation is given in each case.

**7. Conclusion.** The prime pair problem has been shown to be the simplest case of a more general problem. Essentially, the problem poses two questions:

- (5) How "clustered" can a given number of primes be?
- (6) How many such "clusters" are there?

The first question is considered equivalent to the question, what is the smallest range of natural numbers that can contain  $n$  primes? The scheme proposed in this paper gives a necessary range; it is suggested but not proved generally that this range is sufficient. It is proved to be both necessary and sufficient for all  $n$ -tuples for  $n \leq 10$ .

G. H. Hardy [3] and W. W. Ball [4] have made conjectures concerning the number of prime pairs and higher  $n$ -tuples under a given magnitude but the results of this investigation are too meager to expect any degree of accuracy from the formulas that have been proposed.

The programs for this investigation were not the most efficient since the writer had a limited amount of time to write, debug and run them. With a more elegant approach to the programming and by using the more sophisticated logic of the IBM 704 these calculations could be accelerated by a factor of 200 or more.

The writer wishes to express his thanks to the referee for his comments and suggestions.

HERSCHEL F. SMITH

Applied Science Division  
International Business Machines  
Chicago, Illinois

1. A. DE POLIGNAC, *Nouvelles Annales de Math.*, v. 8, 1849, p. 428.
2. EDMUND LANDAU, *Vorlesungen über Zahlentheorie*, Chelsea Pub. Co., New York, v. 1, 1947, p. 71.
3. G. H. HARDY & J. E. LITTLEWOOD, "Some problems of 'Partitio Numerorum'; III: On the expression of a number as a sum of primes," *Acta Math.*, v. 44, 1923, p. 1-70.
4. W. W. BALL, *Mathematical Recreations and Essays*, 11th ed., p. 64.