

# Two Theorem Tables of Matrix Algebra

By Gilbert C. Best

**Abstract.** It is shown in this paper how it is possible, by a slight modification of the mode of presentation, to state many of the theorems of matrix algebra in table form. Due to the large number of theorems presented the tables give only the statement of each theorem. It is not considered feasible to also attempt to indicate sources and proofs.

**1. Introduction.** Many theorems are of such a nature that they can be stated in table form. Take, for example, the theorem which reads, "If  $A$  is a non-singular symmetric matrix then so is  $A^{-1}$ ." As a first step we restate this in "operator" form. We define the "inverse-forming" operator  $O^{-1}$  to be such that  $O^{-1}A = A^{-1}$ , where  $A$  is assumed non-singular. Then the theorem reads, "The property of symmetry is invariant for the operator  $O^{-1}$ ." Finally, defining " $v$ " to indicate invariance, the theorem may be presented in an "operator vs. property" table by a  $v$  in row  $O^{-1}$  and column "symmetry." This theorem appears in Table I in row 3 column 6. This method of presentation has the obvious advantage of considerable compactness.

Again, consider the theorem

$$(1) \quad (A^T)^{-1} = (A^{-1})^T.$$

Defining a "transposing" operator  $O^T$  such that  $O^T A = A^T$  this theorem can be rewritten in "operator" form as :

$$(2) \quad O^T O^{-1} A = O^{-1} O^T A.$$

The theorem can then be stated as: "The operators  $O^T$  and  $O^{-1}$  are commutative." In this last form the theorem in question is readily presented in an "operator vs. operator" table by defining  $k$  to indicate commutativeness. Then (1) is given as a  $k$  in row  $O^T$  and column  $O^{-1}$ , or, equally well, by a  $k$  in row  $O^{-1}$  and column  $O^T$ .

**2. Operators.** In general, any transformation of a matrix may be considered to define an operator. Also, for implied given fixed coefficients  $\alpha_1, \dots, \alpha_m$  and an ordered set  $(A_1, A_2, \dots, A_m)$  of matrices, the linear combination  $\sum_{i=1}^m \alpha_i A_i$  may be considered an operator. Similarly, any function of a matrix or matrices may be considered an operator. For example, the forming of the determinant, the extraction of the characteristic roots or characteristic vectors, or the product of two matrices, may all be considered operators. A numbered list of the operators used in this paper is shown below. The operators in the rows of Table 1 and the rows and columns of Table 2 preserve the same numbering.

**3. Properties.** The idea of a property also may be generalized beyond the well-known and clearly defined properties such as symmetry, skew-symmetry, singularity, positive definiteness, etc. For example, the fact that a given matrix  $A$  has specific roots, e.g., 2, -5, 7 say, is a property, and this property is unaffected by a

---

Received April 22, 1960.

LIST OF OPERATORS

#	Symbol	Explanation
0	$I$	Identity Operator
1	$O^T$	Transposer
2	$\bar{O}$	Conjugator, i.e., sends $A = (a_{ij})$ into $\bar{A} = (\bar{a}_{ij})$
3	$O^{-1}$	Inverse Former (if inverse exists)
4	$O^A$	Adjoint Former (or adjugate, see [1])
5	$O^{(k)}$	$k$ th Compound Former, see [1]
6	$PQ$	Equivalent Matrix Former*
7	$PP^T$	Congruent Matrix Former*
8	$PP^T$	Hermitely Congruent Matrix Former*
9	$PP^{-1}$	Similar Matrix Former*
10	$PP^{\perp}$	Orthogonally Similar Matrix Former*
11	$UU^{\perp}$	Unitarily Similar Matrix Former*
12	$\Sigma$	Sum Former
13	$O^{\alpha}$	Former of Product with Scalar $\alpha \neq 0$
14	$LNCO$	Former of Linear Combination (coefficients assumed fixed)*
15	$\Pi$	Former of Product of Matrices
16	$O^P$	Former of Matrix Raised to Power $P$
17	$POL$	Former of a Polynomial from a Matrix (coefficients assumed fixed)*
18	$FUNC$	Function Taker*
19	$DET$	Determinant Taker
20	$tr$	Trace Taker
21	$R$	Characteristic Root Taker
22	$V$	Characteristic Vector Taker

\* Operator matrices or coefficients are over the same field as the matrix  $A$  to which they are applied unless otherwise indicated.

LIST OF PROPERTIES

#	Property
1	Diagonal
2	Block Diagonal (also called "direct sum")
3	Triangular. 2 types* : upper and lower
4	Block Triangular. 2 types* : upper and lower
5	Non-Singular
6	Symmetric
7	Hermitian
8	Skew Symmetric
9	Skew Hermitian
10	Orthogonal
11	Unitary
12	Normal
13	Diagonable, i.e., similar to a diagonal matrix
14	Symmetric Positive Definite (real)
15	Hermitian Positive Definite
16	Definite. 4 types* positive definite, positive semi-definite, negative semi-definite, negative definite
17	Specific Rank
18	Specific Index
19	Specific Invariant Factors
20	Specific Characteristic Equation or Characteristic Roots
21	Specific Characteristic Vectors
22	Specific Determinant

\*  $v$  in Table 1 implies that the matrix remains the same type.

similarity transformation. If a matrix has a specific determinant, rank, or index, that is also a property. A list of properties used is shown below, and the columns of Table 1 preserve the same numbering.

**4. Rules of Application.** If an operator is applied to more than one matrix at a time the following rules hold:

(1) If the operator  $O$  is applicable to a single matrix, i.e., is a matrix-to-matrix operator, and  $O$  is applied to the ordered set  $(A_1, A_2, \dots, A_m)$ , then one obtains  $(OA_1, OA_2, \dots, OA_m)$ .

(2) If the operator  $O$  is applicable to two matrices and yields one matrix and  $O$  is applied to the ordered set  $(A_1, A_2, \dots, A_m)$ , then  $O$  is first applied to  $A_1, A_2$  yielding  $A_2'$ , then  $O$  is applied to  $A_2'A_3$  yielding  $A_3'$ , etc.

If a set of matrices  $(A_1, A_2, \dots, A_m)$  is said to have a property  $P$ , then each matrix of the set has property  $P$ .

TABLE 1  
*Operators vs. Properties*

Op.	Diagonal	Block Diagonal	Triangular	Block Triangular	Non-singular	Symmetric	Hermitian	Skew Symmetric	Skew Hermitian	Orthogonal	Unitary	Normal	Diagonable	Sym. Pos. Def.	Herm. Pos. Def.	Definite	Rank (Has Specific)	Index	Invariant Factors	Ch. Roots	Ch. Vectors	Determinant	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
1 $O^T$	v	v	t	t	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v		v	1
2 $\bar{O}$	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v			2
3 $O^{-1}$	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v			3
4 $O^A$	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v		v		4
5 $O^{(k)}$	v		v		v	v	v				v						a						5
6 $PQ$					v	v											v		v				6
7 $PP^T$					v	v	v		v					v			v		v	v			7
8 $P\bar{P}^T$					v	v		v						v		v	v		v	v			8
9 $PP^{-1}$					v	v			v						v	v	v		v	v			9
10 $PP^\perp$					v	v		v		v			v		v	v	v		v	v		v	10
11 $UU^\perp$					v		v		v		v	v			v	b	v		v	v		v	11
12 $\Sigma$	v	v	v	v	v	v	r	v	r			v	v		v	b	v		v			v	12
13 $O^\alpha$	v	v	v	v	v	v	r	v	r			v	v	+	+	+	v	+	v			v	13
14 $LNCO$	F	F	F	F	F	F	v	v	v				v				+						14
15 $\Pi$	v	v	v	v	v	k	k			v	v			k	k								15
16 $O^P$	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v	v				v		16
17 $POL$	F	F	F	F	F	F	v	v	v	v	F	F		r	r	r					v		17
18 $FUNC$	F	F	F	F	F	F	v				F	F		r	r	r					v		18

Symbol	Explanation
a	maintains type for even $n$ but for $n$ odd: neg $\rightarrow$ pos
b	maintains type, though a change from semi into non-semi is possible
F	invariant for $LNCO, POL, FUNC$ over a field $F$ for matrices over $F$
k	invariant if and only if the matrices commute
r	invariant for $\alpha, LNCO, POL, FUNC$ over the real field
t	changes type
v	invariant
+	for $\alpha > 0$

TABLE 2  
Operators vs. Operators

Op.	$O^r$	$O$	$O^{-1}$	$O^A$	$O^{(k)}$	$PQ$	$PPT$	$PPT$	$PP^{-1}$	$PP_{\perp}$	$UU_{\perp}$	$\Sigma$	$O^*$	$LNCO$	$\Pi$	$O^P$	$POL$	$FUNC$	$DET$	$tr$	$R$	$V$
1	$O^r$	$I$	$k$	$k$	$k$		$k$			$k$		$k$	$k$	$k$	$o$	$k$	$k$	$k$	$k$	$k$	$k$	$k$
2	$O$	$k$	$I$	$k$	$k$							$k$	$k$	$k$	$r$	$k$	$k$	$r$	$k$	$k$	$k$	$k$
3	$O^{-1}$	$k$	$k$	$I$	$k$				$k$			$k$		$o$	$o$	$k$	$k$					
4	$O^A$	$k$	$k$	$k$	$k$				$k$	$k$				$o$	$o$	$k$	$k$					
5	$O^{(k)}$	$k$	$k$	$k$	$k$						$k$			$o$	$o$	$k$	$k$					
6	$PQ$											$k$	$k$	$k$	$k$	$k$	$k$					
7	$PPT$	$k$										$k$	$k$	$k$	$k$	$k$	$k$					
8	$PPT$											$k$	$k$	$k$	$k$	$k$	$k$					
9	$PP^{-1}$			$k$	$k$							$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$
10	$PP_{\perp}$	$k$		$k$	$k$							$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$
11	$UU_{\perp}$			$k$	$k$							$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$
12	$\Sigma$	$k$	$k$			$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$									$k$
13	$O^*$	$k$	$k$			$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$									
14	$LNCO$	$k$	$k$	$r$		$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$	$k$							$k$	$k$
15	$\Pi$	$o$	$k$	$k$	$o$	$k$			$k$	$k$	$k$	$k$	$k$						$k$			
16	$O^P$	$k$	$k$	$k$	$k$	$k$			$k$	$k$	$k$	$k$	$k$								$k$	$k$
17	$POL$	$k$	$k$	$k$	$k$	$k$			$k$	$k$	$k$	$k$	$k$								$k$	$k$
18	$FUNC$	$k$	$k$	$r$	$k$	$k$			$k$	$k$	$k$	$k$	$k$			$k$					$k$	$k$
19	$DET$	$k$	$k$	$k$	$k$	$k$			$k$	$k$	$k$	$k$	$k$			$k$					$k$	$k$
20	$tr$	$k$	$k$	$k$	$k$	$k$			$k$	$k$	$k$	$k$	$k$		$k$	$k$					$k$	$k$
21	$R$	$k$	$k$	$k$	$k$	$k$			$k$	$k$	$k$	$k$	$k$			$k$	$k$				$k$	$k$
22	$V$	$k$	$k$	$k$	$k$	$k$			$k$	$k$	$k$	$k$	$k$			$k$	$k$				$k$	$k$

Symbol Explanation  
 $I$  two applications of operator equivalent to the identity operator  
 $o$  order of product reverses, e.g. if  $P$  = operator for row in table  $P\Pi(AB) = \Pi P(BA) = P(B)*P(A)$   
 $k$  operators are commutative  
 $r$  operators are commutative for  $LNCO, POL, FUNC$  over the real field

3703 Tulsa Way  
 Fort Worth-7, Texas

1. A. C. AITKEN, *Determinants and Matrices*, Oliver & Boyd, London, 1951, p. 51, p. 90.