

## TABLE ERRATA

**397.**—PAUL F. BYRD & MORRIS D. FRIEDMAN, *Handbook of Elliptic Integrals for Engineers and Physicists*, Springer-Verlag, Berlin, 1954.

The following corrections should be made in the table entitled Values of the Function  $KZ(\beta, k)$ , on pp. 336–343.

| $\sin^{-1} k$ | $\beta$ | <i>for</i> | <i>read</i> |
|---------------|---------|------------|-------------|
| 15°           | 44°     | .027204    | .027203     |
| 40°           | 57°     | .196336    | .196349     |
|               | 64°     | .171978    | .171980     |
|               | 73°     | .124059    | .124061     |
| 85°           | 63°     | 1.982530   | 1.982526    |
|               | 87°     | .548499    | .558435     |
| 87°           | 22°     | 1.229612   | 1.229589    |
|               | 44°     | 2.154030   | 2.153771    |
|               | 79°     | 1.931185   | 1.930751    |
| 88°           | 73°     | 2.635400   | 2.635330    |
| 89°           | 8°      | .616197    | .616207     |
|               | 71°     | 3.351047   | 3.350992    |
|               | 86°     | 2.081462   | 2.081437    |

HENRY E. FETTIS  
JAMES C. CASLIN

Applied Mathematics Laboratory  
Aerospace Research Laboratories  
Wright-Patterson Air Force Base, Ohio

EDITORIAL NOTE: An additional serious error in this table was noted by D. Caligo (*MTAC*, v. 13, 1959, p. 141, MTE **269**). For further notices of errata in this book, see *Math. Comp.*, v. 18, 1964, p. 532, MTE **352**, and p. 687, MTE **359**.

**398.**—HENRY E. FETTIS & JAMES C. CASLIN, *Tables of Elliptic Integrals of the First, Second and Third Kind*, Report ARL 64-232, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio, December, 1964.

In Table III (pp. 44–93), corresponding to  $k^2 = 1.00$ , the following *additive* corrections should be made, in units of the last decimal place.

| $\alpha^2$ | $\phi$ |       |       |       |       |       |       |
|------------|--------|-------|-------|-------|-------|-------|-------|
|            | 65.0°  | 70.0° | 75.0° | 80.0° | 82.5° | 85.0° | 87.5° |
| –1.0       |        |       |       |       |       | 1     | 6     |
| –.9        |        |       |       |       | 1     | 2     | 7     |
| –.8        |        |       |       |       | 1     | 2     | 7     |
| –.7        |        |       |       | 1     | 1     | 2     | 7     |
| –.6        |        |       |       |       | 1     | 2     | 9     |
| –.5        |        |       |       |       | 1     | 2     | 9     |
| –.4        |        |       |       |       | 1     | 2     | 10    |

|     |   |   |   |   |    |    |     |
|-----|---|---|---|---|----|----|-----|
| -.3 |   |   | 1 | 1 | 1  | 3  | 10  |
| -.2 |   |   |   |   | 1  | 2  | 10  |
| -.1 |   |   |   |   | 2  | 3  | 11  |
| +.1 |   |   | 1 | 1 | 1  | 3  | 14  |
| +.2 |   |   |   |   | 1  | 4  | 16  |
| +.3 |   |   |   |   | 1  | 4  | 18  |
| +.4 |   | 1 |   |   | 2  | 5  | 22  |
| +.5 |   | 1 | 1 | 1 | 3  | 7  | 26  |
| +.6 |   |   |   |   | 2  | 8  | 32  |
| +.7 |   | 1 | 1 | 2 | 4  | 10 | 42  |
| +.8 |   |   |   | 1 | 3  | 14 | 63  |
| +.9 | 1 | 1 | 1 | 4 | 10 | 27 | 122 |
| 1.0 |   |   |   |   |    | 1  | 3   |

These errors in the table of 10D values of the elliptic integral of the third kind are attributable to a programming error, which resulted in the value of  $k^2$  being set equal to  $1 - 10^{-16}$  instead of 1.

HENRY E. FETTIS

Applied Mathematics Research Laboratory  
Wright-Patterson Air Force Base, Ohio

EDITORIAL NOTE: For a review of these tables see *Math. Comp.*, v. 19, 1965, p. 509, R MT 81

399.—MILTON ABRAMOWITZ & IRENE A. STEGUN, Editors, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series, No. 55, U. S. Government Printing Office, Washington, D. C., third printing, March 1965.

On p. 333, formula 8.2.7 should read

$$P_{-\mu-1/2}^{-\nu-1/2} \left[ \frac{z}{(z^2-1)^{1/2}} \right] = \frac{(z^2-1)^{1/4} e^{-i\mu\pi} Q_{\nu}^{\mu}(z)}{(\frac{1}{2}\pi)^{1/2} \Gamma(\nu + \mu + 1)}$$

and the left side of formula 8.2.8 should read

$$Q_{-\mu-1/2}^{-\nu-1/2} \left[ \frac{z}{(z^2-1)^{1/2}} \right].$$

On p. 334, the left side of formula 8.6.11 should read  $-Q_{\nu}^{-1/2}(z)$ .

On p. 335, in formula 8.8.2 the factor  $(z^2-1)^{-\mu/2}$  on the right side should be replaced by  $(z^2-1)^{\mu/2}$ .

HENRY E. FETTIS

Applied Mathematics Research Laboratory  
Wright-Patterson Air Force Base, Ohio

On p. 783, in formula 22.9.8 the third column should read  $(1 - \ln R^2)/2$ , and in formula 22.9.11 the third column should read  $R^{-1}(1 - xz + R)^{-1/2}$ .

VAN E. WOOD

Battelle Memorial Institute  
Columbus, Ohio

Recalculation of the coefficients in the Maclaurin series for  $1/\Gamma(z)$  to more than 25D has revealed the following corrections to be required in the 16D table in 6.1.34 on p. 256. The final decimal digits in  $c_k$  corresponding to  $k = 3, 8, 10, 12, 16$ , and 17 should each be increased by a unit; the final digits in  $c_{11}$  and  $c_{24}$  should each be decreased by a unit, while the value  $c_{25}$  should be decreased by two final units. Also, the sign of  $c_{26}$  should be changed to minus.

This supplements and emends the corrections made by Isaacson and Salzer (*MTAC*, v. 1, 1943, p. 124, MTE 19) in the corresponding original table of Bourguet (*Acta Math.*, v. 2, 1883, pp. 261–295).

J. W. W.

EDITORIAL NOTE: An independent calculation of  $c_{23}$  shows that the value, 206, given in the NBS Handbook is correct—contrary to the assertion made in MTE 393. In fact  $c_{23} = -0.013\ 20\ 58326\ 05356\ 479\ \dots$

400.—A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER & F. G. TRICOMI, *Higher Transcendental Functions*, Volume 2, McGraw-Hill Book Co., New York, 1953.

On p. 187, the right side of equation (34) should read

$$T_{n+m}(x) + T_{n-m}(x).$$

VAN E. WOOD

401.—A. ERDÉLYI, W. MAGNUS, F. OBERHETTINGER & F. G. TRICOMI, *Tables of Integral Transforms*, McGraw-Hill Book Co., New York, 1954.

In Volume I, p. 218, in transform 4.23(18), for  $\frac{1}{2}\sigma$ ,  $\frac{1}{2}\sigma + \frac{1}{2}$ , read  $\sigma$ ,  $\sigma + \frac{1}{2}$ . Also, the second convergence condition on the right should read  $\operatorname{Re} p > 2 |\operatorname{Re} \lambda|$  if  $m = n - 1$ .

In Volume II, pp. 128–129, in transform 10.2(9), the denominator parameters in the first  ${}_1F_2$  should be  $1 - \mu - (\rho + \nu)/2$ ,  $1 - \mu - (\rho - \nu)/2$ , while the numerator parameter in the second  ${}_1F_2$  should be  $(\rho + \nu)/2$ .

In Volume II, p. 153, in transform 10.3(88), for  $-\lambda x^2$ , read  $\lambda x^2$ . Also change the convergence conditions on the right to read

$$\begin{aligned} \operatorname{Re} y > 0 & \quad \text{if } p < q - 1; \\ \operatorname{Re} y > 2 |\operatorname{Re} \lambda| & \quad \text{if } p = q - 1. \end{aligned}$$

VAN E. WOOD

402.—G. E. ROBERTS & H. KAUFMAN, *Table of Laplace Transforms*, W. B. Saunders, Philadelphia, Pennsylvania, 1966.

On p. 116, in transform 33.2.1(18), for  $c/2$ ,  $(c + 1)/2$ , read  $c$ ,  $c + \frac{1}{2}$ . Also, the last convergence condition should read  $\operatorname{Re} s > 2 |\operatorname{Re} k|$  if  $p = q - 1$ .

On p. 112, transform 32.1(3) is a special case of the preceding, and the convergence conditions should accordingly be

$$\begin{aligned} \operatorname{Re} s > 2 |\operatorname{Re} c|, & \quad q = p + 1; \\ \operatorname{Re} s > 0, & \quad q > p + 1. \end{aligned}$$

VAN E. WOOD

403.—D. H. LEHMER, *Guide to Tables in the Theory of Numbers*, National Research Council, National Academy of Sciences, Washington, D. C., 1941, reprinted 1961.

On p. 162, in section 2, [f<sub>1</sub>], it is erroneously stated that  $10^8 + 2271$ ,  $10^8 + 4291$ , and  $10^8 + 4909$  should be deleted from the list of primes given on pp. 97–98 of *Tavole di Numeri Primi entro Limiti Diversi e Tavole Affini*, by L. Poletti, Milan, 1920. In fact, these numbers are prime.

There exists an additional error in Poletti's table; namely,  $10^8 + 9513$  is not prime, since it is divisible by 1531.

M. F. JONES

M. LAL

Memorial University of Newfoundland  
St. John's, Newfoundland  
Canada

EDITORIAL NOTE: The primality of the first three numbers cited can be verified by consulting C. L. Baker & F. J. Gruenberger, *The First Six Million Prime Numbers*, The Microcard Foundation, Madison, Wisconsin, 1959. (See *Math. Comp.*, v. 15, 1961, p. 82, RMT 4.)

404.—D. N. LEHMER, *List of Prime Numbers from 1 to 10,006,721*, Publication No. 165, Carnegie Institution of Washington, Washington, D. C., 1914; reprinted by Hafner Publishing Co., New York, 1956.

A table of the Riemann function  $P(x)$  is given on pp. xiii–xvi. The entries therein should each be decreased by a unit for the following 11 values of  $x$ :

|           |           |           |           |
|-----------|-----------|-----------|-----------|
| 750,000   | 1,000,000 | 2,400,000 | 3,450,000 |
| 5,050,000 | 6,350,000 | 9,250,000 | 9,650,000 |
| 9,750,000 | 9,850,000 | 9,950,000 |           |

and the entry corresponding to  $x = 4,700,000$  should be increased by a unit.

In the same table the columns headed "Tchebycheff" do not constitute, as the author erroneously states (p. ix), a tabulation of

$$\int_2^x dy/\ln y,$$

but of

$$Li(x) = \lim_{\epsilon \rightarrow 0} \int_0^{1-\epsilon} dy/\ln y + \int_{1-\epsilon}^x dy/\ln y.$$

(The same error occurs in D. C. Mapes, "Fast method for computing the number of primes less than a given limit," *Math. Comp.*, v. 17, 1963, pp. 179–185.) These tabular values of  $Li(x)$  should be decreased by a unit for the following 11 values of  $x$ :

|           |           |           |           |
|-----------|-----------|-----------|-----------|
| 650,000   | 1,200,000 | 2,150,000 | 4,400,000 |
| 4,550,000 | 5,350,000 | 5,550,000 | 8,200,000 |
| 8,350,000 | 8,450,000 | 8,800,000 |           |

and the entry for  $x = 9,950,000$  should be increased by a unit.

M. F. JONES

M. LAL