

Stability of Bounded Solutions of Linear Functional Equations

By Joel N. Franklin

Abstract. The weak sequential compactness of reflexive Banach spaces is used to explain the fact that certain ill-posed, linear problems become well-posed if the solutions are required to satisfy a prescribed bound. Applications are made to the computability of solutions of ill-posed problems associated with elliptic and parabolic partial differential equations.

1. Introduction. In defining the concept of a *well-posed* problem, J. Hadamard [1] recognized that the *existence* and the *uniqueness* of the solution are not enough to ensure the computability of the solution. Numerical computation requires that the solution have *stability*, or continuous dependence on the data.

F. John [2], [3], [4] and other authors have shown that, for certain problems whose solutions have existence and uniqueness but not stability, if the solutions considered are required to satisfy a prescribed bound, stability results. An extensive bibliography appears in the paper [5] by L. E. Payne. All these results depend on precise definitions of "continuous dependence on data" which are suitable for the particular problems discussed. Much of the work preceding the present paper contains not merely qualitative statements of stability, but quantitative inequalities satisfied in particular problems.

The present paper contains only the qualitative observation that an elementary result, Theorem 1, on the weak topology of reflexive Banach spaces yields the stability of uniformly bounded solutions of a large class of linear functional equations. Several new applications are made, including a theorem on the stability of solutions of the final-value problem corresponding to the general initial-value problem discussed in the book [6] by R. D. Richtmyer.

2. Notation and Review. We shall use the letter T to stand for a bounded linear operator mapping a Banach space B_1 into a Banach space B_2 . We suppose that the range TB_1 is dense in B_2 , and that $Tz = 0$ only if $z = 0$.

If B is a Banach space, the notation B^* represents the conjugate Banach space of bounded linear functionals f mapping B into the real numbers. If $x \in B$ and $f \in B^*$, the notation (x, f) represents the number which results when f is applied to x . If $B^{**} = B$, the space B is called *reflexive* [7]. Hilbert spaces are reflexive Banach spaces. If $1 < p < \infty$, the space L^p of measurable functions $x(t)$ for which

$$\int_a^b |x(t)|^p dt < \infty$$

Received August 26, 1970.

AMS 1970 subject classifications. Primary 65M30, 65N99, 65R05; Secondary 65J05.

Key words and phrases. Well-posed, ill-posed, continuous dependence on data, stability, backwards heat-equation, final-value problem, elliptic continuation.

Copyright © 1971, American Mathematical Society

is reflexive. In fact, $(L^p)^* = L^q$ if $1/p + 1/q = 1$. However, the Banach space C of continuous functions under the maximum-norm is not reflexive.

If $\{x_n\}$ is a sequence of B , we say that it converges, or converges *strongly*, to x if $\|x_n - x\| \rightarrow 0$ as $n \rightarrow \infty$; we then write $x_n \rightarrow x$. We say that x_n converges *weakly* to x if, for each f in B^* , (x_n, f) tends to the limit (x, f) as $n \rightarrow \infty$; we then write $x_n \rightharpoonup x$. A convergent sequence is weakly convergent, but the converse is false.

If g lies in the range of T , the equation

$$(1) \quad Tu = g$$

has a unique solution, $u \in B_1$, since we have supposed $Tz = 0$ only if $z = 0$. Now the question of stability is this: *If $Tu_n = g_n$, and if the g_n tend in some sense to g , do the u_n tend to u ?*

The trivial case is that in which T has a bounded inverse, T^{-1} . Then $u_n \rightarrow u$ if and only if $g_n \rightarrow g$, and $u_n \rightharpoonup u$ if and only if $g_n \rightharpoonup g$. The first assertion is obvious, and the second follows at once from the existence of bounded adjoint operators T^* and $(T^{-1})^*$. We note that if B_1 is finite-dimensional, then T does have a bounded inverse, since we assume that T is one-to-one, and that TB_1 is dense in B_2 .

If B_1 is infinite-dimensional, T may lack a bounded inverse and still satisfy our requirements. A simple example is given by the mapping of the space, C , of continuous functions $x(\tau)$ on $0 \leq \tau \leq 1$ into $L^2[0, 1]$ by the transformation

$$Tx(\tau) = \int_0^\tau x(\sigma) d\sigma.$$

Many examples are of the following form: Let B_1 and B_2 be identical Hilbert spaces. Let T be a completely continuous, selfadjoint operator whose eigenvalues λ_n are nonzero, and whose eigenvectors v_n form a basis for B_1 . Then a bounded inverse T^{-1} cannot exist; for if $\|v_n\| = 1$, then $v_n \rightarrow 0$ and hence $\lambda_n v_n = Tv_n \rightarrow 0$.

3. Stability of Bounded Solutions. After proving the following general theorem, we will explain the need for reflexivity, for boundedness, and for the use of weak, instead of strong, convergence.

THEOREM 1. *Let T be a bounded operator mapping a reflexive Banach space, B_1 , into a dense subset of a Banach space, B_2 . Let $Tz = 0$ only if $z = 0$. Let*

$$(1) \quad Tu_n = g_n \quad (n = 1, 2, \dots).$$

Assume that the u_n are bounded: $\|u_n\| \leq \beta$. Then $g_n \rightarrow g$ in B_2 if and only if there is a point u in B_1 for which $u_n \rightarrow u$, with $Tu = g$.

Proof. We assert that $T^*B_2^*$ is dense in B_1^* . Otherwise, there would be an element f_0 in B_1^* but not in the closure of $T^*B_2^*$. Using the Hahn-Banach theorem, we now construct a functional $\Phi(f)$ for all $f \in B_1^*$, with

$$(2) \quad \Phi(f_0) \neq 0, \quad \text{but} \quad \Phi(f) = 0 \quad \text{for } f \in T^*B_2^*.$$

Since $\Phi \in B_1^{**} = B_1$, there is a point x_0 in B_1 such that

$$(3) \quad \Phi(f) = (x_0, f) \quad \text{for all } f \in B_1^*.$$

If $f = T^*h \in T^*B_2^*$, then (2) and (3) imply

$$0 = \Phi(f) = (x_0, f) = (x_0, T^*h) = (Tx_0, h).$$

Thus, $0 = (Tx_0, h)$ for all $h \in B_2^*$, which implies $Tx_0 = 0$, and hence $x_0 = 0$. Now (3) implies $\Phi(f) \equiv 0$ for $f \in B_1^*$, which contradicts the first half of (2).

If $u_n \rightarrow u \in B_1$, then $Tu_n = g_n \rightarrow Tu = g$ because, for all $h \in B_2^*$,

$$(4) \quad (g_n - Tu, h) = (T(u_n - u), h) = (u_n - u, T^*h) \rightarrow 0.$$

Conversely, assuming Tu_n converges weakly, we can show that u_n converges weakly. Let $Tu_n \rightarrow g$. Let f be given in B_1^* . Since $T^*B_2^*$ has been shown to be dense in B_1^* , given $\epsilon > 0$ we can find a point $h \in B_2^*$ for which $\|f - T^*h\| < \epsilon$. Since

$$(5) \quad (u_n - u_m, f) = (T(u_n - u_m), h) + (u_n - u_m, f - T^*h)$$

we deduce, for all n and m ,

$$(6) \quad |(u_n - u_m, f)| \leq |(Tu_n - Tu_m, h)| + 2\beta\epsilon.$$

Since $\{Tu_n\}$ converges weakly, and since ϵ is arbitrarily small, we conclude that $(u_n - u_m, f) \rightarrow 0$ as n and $m \rightarrow \infty$. Therefore, for every $f \in B_1^*$, there exists a limit

$$(7) \quad L(f) = \lim_{n \rightarrow \infty} (u_n, f).$$

Now $L(f)$ is a bounded linear functional; indeed, $\|L\| \leq \beta$, since all $\|u_n\| \leq \beta$. Hence, $L \in B_1^{**} = B_1$; and so L has a representation

$$(8) \quad L(f) \equiv (u, f) \quad \text{for all } f \in B_1^*$$

where u is some point in B_1 , independent of f . From (7) and (8) we conclude that $u_n \rightarrow u$. Finally, $Tu = g$ because, for all $h \in B_2^*$,

$$(Tu - g, h) = \lim_{n \rightarrow \infty} (Tu_n - g, h) = 0.$$

Having proved the theorem, we will explain the need for its hypotheses.

Univalence. If $Tz = 0$ for some $z \neq 0$, the theorem is false. For if $x_n = (-1)^n z$, then x_n does not tend to zero weakly (or strongly), while $g_n = Tx_n \equiv 0$.

Reflexiveness. If we do not require B_1 to be reflexive, the theorem is false. For example, let B_1 be the space, C , of real-valued continuous functions $x = \psi(t)$ defined for $0 \leq t \leq 1$. Let $B_2 = L^2[0, 1]$. Let

$$Tx = \int_0^t \psi(\tau) d\tau \quad (0 \leq t \leq 1).$$

Note that $Tz = 0$ only if $z = 0$. If $u_n = \psi_n(t) = \cos nt$, then

$$g_n = Tu_n = n^{-1} \sin nt \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Moreover, the u_n are uniformly bounded:

$$\|u_n\| = \max_{0 \leq t \leq 1} |\cos nt| = 1.$$

But, if Φ is the functional which evaluates a function $\psi(t)$ for $t = \pi/4$, then

$$\Phi(u_n) = \cos(n\pi/4)$$

which diverges as $n \rightarrow \infty$. Hence, the sequence $\{u_n\}$ is not weakly convergent even though g_n is strongly convergent. Incidentally, this example and Theorem 1 provide an independent proof that the Banach space C is not reflexive; cf. [7, p. 214].

Weak convergence. If we do not use the notion of weak convergence, but use only strong convergence, the theorem reduces to the trivial case, in which T has a bounded inverse:

Assertion. Let T be a bounded operator mapping a Banach space B_1 into a dense subset of a Banach space B_2 , with $Tz = 0$ only if $z = 0$. (Here we do not need the assumption that B_1 is reflexive.) Let T have the property that a sequence $\{x_n\}$ must converge strongly if it is bounded and if $\{Tx_n\}$ converges strongly. Then $TB_1 = B_2$, and T has a bounded inverse, T^{-1} , mapping B_2 into B_1 .

Proof. To show $TB_1 = B_2$, let y be any point in B_2 . Since TB_1 is assumed to be dense in B_2 , there is a sequence $\{x_n\}$ in B_1 such that $Tx_n \rightarrow y$. If $\{x_n\}$ has a bounded subsequence, $\{x'_n\}$, the hypothesis implies that x'_n has a strong limit, x , and hence $Tx = y \in TB_1$. If $\{x_n\}$ has no bounded subsequence, then $\|x_n\| \rightarrow \infty$. If $x_n \neq 0$ for $n \geq N$, define the points

$$u_{2n-1} = x_n/\|x_n\|, \quad u_{2n} = 0 \quad \text{for } n \geq N.$$

Since $Tx_n \rightarrow y$ while $\|x_n\| \rightarrow \infty$, we have $Tu_n \rightarrow 0$. Since $\|u_n\| \leq 1$, the hypothesis implies the convergence of $\{u_n\}$. But $\{u_n\}$ diverges because $\|u_{2n-1} - u_{2n}\| = 1$. Therefore, $\{x_n\}$ does have a bounded subsequence, and $TB_1 = B_2$.

Hence, T has an inverse, T^{-1} . If T^{-1} were not bounded, there would be a sequence of unit vectors, v_n , for which $Tv_n \rightarrow 0$. Then $Tx_n \rightarrow 0$ if $x_{2n-1} = v_n$ and $x_{2n} = 0$. The hypothesis now implies that $\{x_n\}$ converges, which is impossible because $\|x_{2n-1} - x_{2n}\| = 1$. This completes the proof of the assertion.

Boundedness. If we omit the assumption that $\{u_n\}$ is bounded, the theorem is false. For example, let T be any bounded linear operator mapping a Hilbert space into itself, with unit eigenvectors v_1, v_2, \dots and associated nonzero eigenvalues $\lambda_1, \lambda_2, \dots$ tending to zero as $n \rightarrow \infty$. Let $u_n = |\lambda_n|^{-1/2}v_n$. Then $Tu_n \rightarrow 0$ while $\|u_n\| \rightarrow \infty$. Since a weakly convergent sequence must be bounded, $\{u_n\}$ is not weakly convergent.

4. An Example for the Heat Equation. To show how the theorem can be used, we will consider the stability of the flux of heat depending upon a temperature. Consider the conduction of heat in a semi-infinite, homogeneous rod $0 \leq x < \infty$ in an interval of time $0 \leq t \leq t_1$. If $\varphi(x, t)$ is the temperature, assume that it obeys the equation

$$(1) \quad \frac{\partial \varphi}{\partial t} = \frac{\partial^2 \varphi}{\partial x^2} \quad (0 \leq x < \infty, 0 \leq t \leq t_1).$$

Let the temperature at $x = 0$ be given:

$$(2) \quad \varphi(0, t) = g(t) \quad (0 \leq t \leq t_1)$$

and let the unknown be the flux of heat into the end at $x = 0$:

$$(3) \quad -\frac{\partial}{\partial x} \varphi(0, t) = u(t) \quad (0 \leq t \leq t_1).$$

We assume the initial and boundary conditions

$$(4) \quad \varphi(x, 0) \equiv 0 \quad \text{and} \quad \varphi(+\infty, t) \equiv 0.$$

Define the Laplace transform

$$\Phi(x, s) = \int_0^\infty e^{-st} \varphi(x, t) dt \quad (\text{Re } s > 0)$$

and let the boundary temperature, $g(t)$, and flux, $u(t)$, have the transforms $G(s)$ and $U(s)$. Then the preceding equations yield

$$s\Phi(x, s) = \frac{\partial^2}{\partial x^2} \Phi(x, s)$$

and

$$(5) \quad \Phi(x, s) = G(s) \exp(-x\sqrt{s}), \quad U(s) = -\frac{\partial}{\partial x} \Phi(0, s) = \sqrt{s} G(s).$$

Since $s^{-1/2}$ is the Laplace transform of $(t/\pi)^{1/2}$, the convolution theorem yields

$$(6) \quad \frac{1}{\sqrt{\pi}} \int_0^t (t - \tau)^{-1/2} u(\tau) d\tau = g(t) \quad (0 \leq t \leq t_1).$$

This is an equation, $Tu = g$, for which we will discuss the stability of the solution, u . We will not use the explicit solution of the Abel integral equation (6), which is

$$(7) \quad u(t) = \frac{d}{dt} \frac{1}{\sqrt{\pi}} \int_0^t (t - \tau)^{-1/2} g(\tau) d\tau.$$

For the sake of definiteness, consider the example

$$(8) \quad g_0(t) \equiv 1, \quad u_0(t) = (\pi t)^{-1/2} \quad (0 \leq t \leq t_1).$$

The function $u(t)$ is not square integrable, but it does lie in the reflexive Banach space $B_1 = L^p[0, t_1]$ if $1 < p < 2$. We will now show that T is a bounded linear operator mapping $L^p[0, t_1]$ into $L^2[0, t_1]$.

Given the equation (6), with $u \in L^2$, we must show that

$$(9) \quad \int_0^{t_1} g^2(t) dt < \infty.$$

From (6) we find

$$(10) \quad \int_0^{t_1} g^2(t) dt = \int_0^{t_1} \int_0^{t_1} H(\sigma, \tau) u(\sigma) u(\tau) d\sigma d\tau$$

where

$$(11) \quad H(\sigma, \tau) = \frac{1}{\pi} \int_{\max(\sigma, \tau)}^{t_1} [(t - \tau)(t - \sigma)]^{-1/2} dt$$

or

$$(12) \quad H(\sigma, \tau) = \frac{1}{\pi} \log \frac{2t_1 - \sigma - \tau + 2[(t_1 - \sigma)(t_1 - \tau)]^{1/2}}{|\tau - \sigma|}.$$

Since

$$0 \leq H(\sigma, \tau) \leq \frac{1}{\pi} \log \frac{4t_1}{|\tau - \sigma|},$$

we have, for all $q > 0$,

$$(13) \quad \eta_q \equiv \int_0^{t_1} \int_0^{t_1} \{H(\sigma, \tau)\}^q d\sigma d\tau < \infty.$$

If $q = (1 - p^{-1})^{-1}$, Hölder's inequality applied to (10) yields

$$(14) \quad \int_0^{t_1} g^2(t) dt \leq (\eta_q)^{1/q} \left(\int_0^{t_1} |u(t)|^p dt \right)^{2/p} < \infty.$$

This completes the proof that T is bounded.

If $g(t) = 0$, then the boundary-value problem (1), (2), (4) has the unique solution $\varphi(x, t) \equiv 0$. Then $u(t) = -\varphi_x(0, t) = 0$. Therefore, $Tu = 0$ only if $u = 0$.

Theorem 1 now yields the following result: Let $u_n(t)$ satisfy

$$(15) \quad \int_0^{t_1} |u_n(t)|^p dt \leq \beta^p \quad (n = 1, 2, \dots)$$

where β is a finite bound independent of n . Let $\{g_n(t)\}$ converge weakly in L^2 to $g_0(t)$, i.e., let

$$(16) \quad \int_0^{t_1} [g_n(t) - g_0(t)]w(t) dt \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

for all $w(t) \in L^2$. Then $\{u_n(t)\}$ converges weakly in L^p to $u_0(t)$, i.e.,

$$(17) \quad \int_0^{t_1} [u_n(t) - u_0(t)]\psi(t) dt \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

for all $\psi(t) \in L^q$, where $q = (1 - p^{-1})^{-1}$.

It is noteworthy that T does not have a bounded inverse, T^{-1} , mapping L^2 into L^p . To see this, let $u_n(t) = \cos nt$. Then

$$(18) \quad g_n(t) = Tu_n(t) = \frac{1}{\sqrt{\pi}} \int_0^t (t - \tau)^{-1/2} \cos n\tau d\tau$$

which is $O(n^{-1/2})$ as $n \rightarrow \infty$ uniformly in t for $0 \leq t \leq t_1$. Therefore, $\|g_n\| \rightarrow 0$ as $n \rightarrow \infty$. But, since $p < 2$,

$$\|u_n\|^p = \int_0^{t_1} |\cos nt|^p dt \geq \int_0^{t_1} (\cos nt)^2 dt \rightarrow \frac{1}{2}t_1 > 0 \quad \text{as } n \rightarrow \infty.$$

Thus, $\|g_n\| \rightarrow 0$ while $\|Tg_n\|$ is bounded away from zero.

The practical use of the weak convergence (17) is illustrated as follows. Usually one is not so much interested in the instantaneous flux, $u(t)$, as in the total flux

$$(19) \quad F = \int_{t_0}^{t_0 + \Delta t} u(t) dt$$

during some nonzero time interval, $t_0 \leq t \leq t_0 + \Delta t$. Let $\psi(t) \equiv 1$ in this interval and $\psi(t) \equiv 0$ elsewhere. Then

$$(20) \quad F_n = \int_{t_0}^{t_0 + \Delta t} u_n(t) dt = \int_0^{t_1} u_n(t)\psi(t) dt.$$

Since $\psi(t) \in L^q$, the weak convergence (17) implies that $F_n \rightarrow F_0$ as the data $g_n(t)$

tends to $g_0(t)$ provided that the instantaneous fluxes $u_n(t)$ are uniformly bounded in L^2 , which is the assumption (15).

5. Application to Elasticity. Consider the biharmonic equation

$$(1) \quad \Delta^2 \varphi = 0$$

in two dimensions. If φ and $\partial\varphi/\partial n$ are prescribed on a simple closed curve, C_1 , and if P is a point in the domain, D_1 , enclosed by C_1 , then there is a formula (cf. Garabedian [8, p. 266])

$$(2) \quad \varphi(P) = \int_{C_1} \left[A(P, Q)\varphi(Q) + B(P, Q) \frac{\partial\varphi}{\partial n}(Q) \right] ds$$

where $A(P, Q)$ and $B(P, Q)$ are analytic functions of the Cartesian coordinates of P in the open set D_1 .

Let C_2 be a simple closed curve in D_1 , as in the figure.




FIGURE 1

Suppose that φ and $\partial\varphi/\partial n$ are prescribed on the inner curve, C_2 . We now inquire about the stability of the solution, φ , at a point, P , outside C_2 . We will consider only functions φ which solve the biharmonic equation in the larger domain, D_1 , and which satisfy on the outer curve, C_1 , an inequality

$$(3) \quad \int_{C_1} \{ [\varphi(Q)]^2 + [\partial\varphi(Q)/\partial n]^2 \} ds \leq \beta^2$$

where β is a finite number independent of φ

Let g be defined as the pair of real-valued functions $\varphi(P_2)$, $\partial\varphi(P_2)/\partial n$ for P_2 on C_2 . Let

$$(4) \quad \|g\|^2 = \int_{C_2} \{ [\varphi(P_2)]^2 + [\partial\varphi(P_2)/\partial n]^2 \} ds.$$

Similarly, let u be the pair of functions $\varphi(Q)$, $\partial\varphi(Q)/\partial n$; and define $\|u\|^2$ by the left-

hand side of (3). By using Green's formula (2) and its first derivatives, with P replaced by P_2 on C_2 , one obtains an operator, T , which relates u to g :

$$(5) \quad Tu = g.$$

Let B_1 be the Hilbert space of pairs of functions $\varphi(Q), \psi(Q)$ which are square-integrable on C_1 . Similarly, define B_2 on C_2 . Then Green's formula shows that T is a bounded linear operator mapping B_1 into B_2 . The application of T to polynomials in two variables shows that TB_1 is dense in B_2 . Moreover, $Tz = 0$ only if $z = 0$; for if φ and $\partial\varphi/\partial n$ vanish on C_2 , then $\varphi \equiv 0$ in D_2 , and the analyticity of φ implies that $\varphi \equiv 0$ in the larger domain, D_1 , so that φ and $\partial\varphi/\partial n$ also vanish on C_1 .

Let g_1, g_2, \dots be a sequence of data converging weakly in B_2 . This will be true if the corresponding functions $\varphi_1(P_2), \varphi_2(P_2), \dots$ and their normal derivatives on C_2 converge uniformly for P_2 on C_2 . Less is required; since the $\|g_k\|$ have a uniform bound, it suffices to assume that the limit

$$(6) \quad \lim_{k \rightarrow \infty} \int_{C_2} \left[a(P_2)\varphi_k(P_2) + b(P_2) \frac{\partial\varphi_k(P_2)}{\partial n} \right] ds$$

exists for every pair of functions, $a(P_2)$ and $b(P_2)$, chosen from a dense subset of $L^2(C_2)$.

Theorem 1 now implies the weak convergence of u_k in B_1 , i.e., the weak convergence of the pair of functions $\varphi, \partial\varphi/\partial n$ on the Cartesian product $L^2(C_1) \times L^2(C_1)$.

If, as in the figure, P is a fixed point between C_2 and C_1 , Green's formula (2) represents the number $\varphi(P)$ as an inner product in B_1 ; and differentiation of Green's formula represents all derivatives of φ as inner products. Hence, the weak convergence of u_k implies that φ_k and all of its derivatives converge pointwise as $k \rightarrow \infty$, i.e., as the data $\varphi, \partial\varphi/\partial n$ converge on the *inner* curve, C_2 . Of course, if P had been chosen inside C_2 , the convergence would have been an immediate consequence of Green's formula for the inner region, D_2 .

6. Stability of Harmonic Continuation. If, in the preceding section, we had considered Laplace's equation, $\Delta\varphi = 0$, instead of the biharmonic equation, using boundary values only for φ and not for $\partial\varphi/\partial n$, we would have obtained a classical result: If $\Delta\varphi_k = 0$ in the larger domain, D_1 , and if the φ_k are uniformly bounded in D_1 , and if the φ_k converge on the *inner* curve, C_2 , then φ_k and all its derivatives converge at points P between C_2 and C_1 , as well as at points inside C_2 .

If E is an elliptic operator of order $2k$, these results generalize immediately for the boundary-value problem with prescribed values for φ and its normal derivatives of orders less than k .

7. The Final-Value Problem. Richtmyer's book [6] discusses the initial-value problem

$$(1) \quad dv(t)/dt = Av(t) \quad (0 \leq t \leq t_1), \quad v(0) = u,$$

where, for each t , $v(t)$ is an element in a Banach space, B . The operator A is a linear, bounded or unbounded operator whose domain is dense in B , and whose range lies in B . There is a solution-operator, $E(t)$, which is supposed to have a bounded extension to all of B . Thus, if $v(t_1) = g$ and $E(t_1) = T$, the initial-value problem, to determine

g from u , is well-posed; it has the solution

$$(2) \quad Tu = g.$$

The form (1) has been used to discuss wave motion, heat transfer, neutron transport, and elastic vibration.

By the *final-value* problem, we mean that of determining $u = v(0)$ from $g = v(t_1)$ in a well-posed initial-value problem (1). The backward heat equation is of this type. It is an ill-posed problem, since the solutions are unstable. However, the backward wave equation is well-posed. Under certain general conditions, we will show that *uniformly bounded* solutions of the final-value problem are stable.

We will give a commonly-met condition on A which ensures that $Tz = 0$ only if $z = 0$. Let there be a set of points f_1, f_2, \dots in B^* such that, for all w in the domain of A

$$(3) \quad (Aw, f_n) = \alpha_n(w, f_n) \quad (n = 1, 2, \dots)$$

where $\alpha_1, \alpha_2, \dots$ are certain scalars. Thus, the points f_n are eigenvectors in the domain of A^* . Assume that the $\{f_n\}$ are *complete* in B^* , in the sense that

$$(4) \quad (z, f_n) = 0 \quad \text{for all } n \text{ only if } z = 0.$$

For example, if $B = L^2[0, 1]$, and if $A\psi(x) = \psi''(x)$, the second derivative of $w = \psi(x)$, and if the domain of A consists of the functions $\psi(x)$ with two continuous derivatives such that $\psi(0) = \psi(\pi) = 0$, then $B = B^*$, $A = A^*$, and we may choose

$$f_n = \sin n\pi x, \quad \alpha_n = -n^2\pi^2.$$

Then (4) states that the set $\{\sin n\pi x\}$ is complete in $L^2[0, 1]$.

If A satisfies the preceding condition, then $Tz = 0$ only if $z = 0$. For let $v(t)$ satisfy the initial-value problem (1) with $v(0) = z$. Now, for each f_n ,

$$(5) \quad \frac{d}{dt}(v(t), f_n) = (Av(t), f_n) = \alpha_n(v(t), f_n) \quad (0 \leq t \leq t_1)$$

by the assumption (3). Therefore, for $0 \leq t \leq t_1$ and for $n = 1, 2, \dots$,

$$(6) \quad (v(t), f_n) = (z, f_n) \exp(\alpha_n t).$$

When $t = t_1$, we have $v(t_1) = Tz = 0$, and hence (6) implies

$$(7) \quad 0 = (z, f_n) \quad (n = 1, 2, \dots).$$

Now (4) implies $z = 0$.

Applied to the preceding example, this argument implies the well-known uniqueness of solutions to the final-value problem for the heat equation $\varphi_t = \varphi_{xx}$ with boundary condition $\varphi = 0$ for $x = 0, \pi$. If, more generally,

$$A\psi(x) = (p(x)\psi'(x))' + q(x)\psi(x)$$

with $p(x) > 0$, and with $p(x)$, $p'(x)$, and $q(x)$ continuous, the completeness of eigenfunctions for Sturm-Liouville operators implies the uniqueness of solutions to the final-value problem for

$$(8) \quad \frac{\partial \varphi}{\partial t}(x, t) = \frac{\partial}{\partial x} \left(p(x) \frac{\partial \varphi}{\partial x} \right) + q(x)\varphi.$$

To ensure that TB is dense in B , it suffices to assume that eigenvectors w_1, w_2, \dots whose finite linear combinations are dense in B . For then, if $Aw_r = \beta_r w_r$, we have

$$(9) \quad T \sum_{r=1}^n \gamma_r w_r = \sum_{r=1}^n \gamma_r \exp(\beta_r t_1) w_r$$

and the linear combinations (9) are dense in B .

In summary, we have proved the following result:

LEMMA. *If A has eigenvectors w_n whose finite linear combinations are dense in B , and if A^* has eigenvectors f , which are complete in B^* , then the well-posed initial-value problem (1) has a solution-operator, T , which maps B into a dense subset of B , with $Tz = 0$ only if $z = 0$.*

Theorem 1 now yields the following criterion for the stability of solutions of the terminal-value problem:

THEOREM 2. *Let the well-posed initial-value problem (1) be defined for a reflexive Banach space, B . Let $Tu_n = g_n$ ($n = 1, 2, \dots$) where the initial values u_n are uniformly bounded: $\|u_n\| \leq \beta$. Let A satisfy the conditions of the lemma. Then the weak convergence of the final values, g_n , implies the weak convergence of the initial values, u_n .*

The weak convergence of the initial values ensures the *strong* convergence of the vectors $v_r(t)$ for $0 < t < t_1$ if the solution operator $E(t)$ is completely continuous for $t > 0$. This is true, for instance, for the heat equation (8) involving a general Sturm-Liouville operator, A . If a complete set of eigenvectors is the set of orthonormal functions $w_n = \varphi_n(x)$, then the eigenvalues β_n tend to $+\infty$ like a constant multiple of n^2 , and

$$(10) \quad E(t) \sum_1^\infty \gamma_n \varphi_n(x) = \sum_1^\infty \gamma_n \exp(-\beta_n t) \varphi_n(x).$$

Here we assume $\sum \gamma_n^2 < \infty$, and the convergence of the infinite series $\sum \gamma_n \varphi_n(x)$ is understood in the mean-square sense. If the initial values u_1, u_2, \dots have series representations

$$(11) \quad u_k = \sum_{n=1}^\infty c_{kn} \varphi_n(x) \quad (k = 1, 2, \dots)$$

and if

$$u = \sum_{n=1}^\infty c_n \varphi_n(x)$$

in L^2 , then weak convergence $u_k \rightarrow u$ is equivalent to the conditions

$$(12) \quad \sum_{n=1}^\infty c_{kn}^2 \leq c^2 \quad \text{independent of } k,$$

$$(13) \quad \lim_{k \rightarrow \infty} c_{kn} = c_n \quad \text{for each fixed } n.$$

Applying (10), we find, for $t > 0$,

$$(14) \quad v_k(t) = E(t)u_k = \sum_{n=1}^\infty \gamma_{kn} \exp(-\beta_n t) \varphi_n(x).$$

Since $\beta_n \rightarrow +\infty$ as $n \rightarrow \infty$, the conditions of weak convergence, (12) and (13), imply the strong convergence

$$(15) \quad \lim_{k \rightarrow \infty} \|E(t)u_k - E(t)u\| = 0 \quad \text{if } t > 0.$$

This follows at once from the inequality

$$(16) \quad \left\| \sum_{n=1}^{\infty} (\gamma_{kn} - \gamma_n) \exp(-\beta_n t) \varphi_n(x) \right\|^2 \\ \leq \sum_{n=1}^{N-1} (\gamma_{kn} - \gamma_n)^2 \exp(-2\beta_n t) + 2 \exp(-2\beta_N t) (c^2 + \sum \gamma_n^2)$$

which holds for each $N \geq 1$.

In this example, we are even able to deduce the *pointwise* convergence

$$(17) \quad \varphi_k(x_0, t) \rightarrow \varphi(x_0, t) \quad \text{as } k \rightarrow \infty \text{ if } t > 0.$$

For fixed x_0 and $t > 0$, the real number $\varphi_k(x_0, t)$ is the inner product of the vector $u_k = \varphi_k(x, 0)$ with the vector in L^2 whose Fourier series is

$$(18) \quad s = \sum_{n=1}^{\infty} [\varphi_n(x_0) \exp(-\beta_n t)] \varphi_n(x)$$

where, since $t_0 > 0$,

$$(19) \quad \|s\|^2 = \sum_{n=1}^{\infty} [\varphi_n(x_0) \exp(-\beta_n t)]^2 < \infty.$$

(Here we have used the result $\sum \beta_n^{-1} \varphi_n^2(x_0) < \infty$ from Sturm-Liouville theory.) Now the weak convergence $u_k \rightharpoonup u$ yields

$$(20) \quad (u_k, s) \rightarrow (u, s) \quad \text{as } k \rightarrow \infty$$

which is the asserted pointwise convergence (17).

Acknowledgment. I wish to thank Professors David Boyd and Stanley A. Weiss for their helpful comments.

Applied Mathematics Department
California Institute of Technology
Pasadena, California 91109

1. J. HADAMARD, *Lectures on Cauchy's Problem in Linear Partial Differential Equations*, Yale Univ. Press, New Haven, Conn., 1923.

2. F. JOHN, "Numerical solution of the equation of heat conduction for preceding times," *Ann. Mat. Pura Appl.* (4), v. 40, 1955, pp. 129-142. MR 19, 323.

3. F. JOHN, *Numerical Solution of Problems which are not Well Posed in the Sense of Hadamard*, Proc. Sympos. Numerical Treatment of Partial Differential Equations with Real Characteristics (Rome, 1959), Libreria Eredi Virgilio Veschi, Rome, 1959, pp. 103-116. MR 21 #6704.

4. F. JOHN, "Continuous dependence on data for solutions of partial differential equations with a prescribed bound," *Comm. Pure Appl. Math.*, v. 13, 1960, pp. 551-585. MR 24 #A317.

5. L. E. PAYNE, *On Some Nonwell Posed Problems for Partial Differential Equations*, Proc. Adv. Sympos. Numerical Solutions of Nonlinear Differential Equations (Madison, Wis., 1966), Wiley, New York, 1966, pp. 239–263. MR 35 #4606.
6. R. D. RICHTMYER, *Difference Methods for Initial-Value Problems*, Interscience Tracts in Pure and Appl. Math., Tract 4, Interscience, New York, 1957. MR 20 #438.
7. F. RIESZ & B. SZ.-NAGY, *Lecons d'Analyse Fonctionnelle*, Akad. Kiadó, Budapest, 1953; English transl., Ungar, New York, 1955. MR 15, 132; MR 17, 175.
8. P. R. GARABEDIAN, *Partial Differential Equations*, Wiley, New York, 1964. MR 28 #5247.
9. J. R. GANNON, *Some Numerical Results for the Solution of the Heat Equation Backwards in Time*, Proc. Adv. Sympos. Numerical Solutions of Nonlinear Differential Equations (Madison, Wis., 1966), Wiley, New York, 1966, pp. 21–54. MR 34 #7037.
10. J. DOUGLAS, JR., *Approximate Continuation of Harmonic and Parabolic Functions*, Proc. Sympos. Numerical Solution of Partial Differential Equations (Univ. of Maryland, 1965), Academic Press, New York, 1966, pp. 353–364. MR 34 #2206.
11. R. SAYLOR, "Numerical elliptic continuation," *SIAM J. Numer. Anal.*, v. 4, 1967, pp. 575–581. MR 36 #5513.