

An Equation of Mordell

By Andrew Bremner

Abstract. All integer solutions of the Diophantine equation $6y^2 = (x + 1)(x^2 - x + 6)$ are found.

1. Mordell [1] asks if all the integer solutions of the Diophantine equation $6y^2 = (x + 1)(x^2 - x + 6)$ are given by $x = -1, 0, 2, 7, 15$ and 74 . It is shown that there are precisely seven integer solutions, the seventh with $x = 767$.

Consideration of factorization gives, for some integers a, b ,

$$\begin{aligned} \begin{cases} x^2 - x + 6 = 6b^2 \\ x + 1 = a^2 \end{cases} & \text{ or } \begin{cases} x^2 - x + 6 = 3b^2 \\ x + 1 = 2a^2 \end{cases} & \text{ or} \\ \begin{cases} x^2 - x + 6 = 2b^2 \\ x + 1 = 3a^2 \end{cases} & \text{ or } \begin{cases} x^2 - x + 6 = b^2 \\ x + 1 = 6a^2 \end{cases} \end{aligned}$$

and the latter case is impossible modulo 3. We thus obtain, on eliminating x , the three quartic equations,

- (i) $a^4 - 3a^2 + 8 = 6b^2$,
- (ii) $4a^4 - 6a^2 + 8 = 3b^2$,
- (iii) $9a^4 - 9a^2 + 8 = 2b^2$.

The standard technique in dealing with equations of this type is to factorize in the appropriate quadratic extension of the integers, which here is $\mathbf{Z}[(1 + \sqrt{-23})/2]$, to obtain a finite set of equations of the form,

$$a^2 = f(v, w), \quad 1 = g(v, w),$$

where f, g are homogeneous quadratic forms.

We need to know some details of the quadratic field $\mathbf{Q}(\sqrt{-23})$. The class-number of the ring of integers is 3; and we denote the ideal factorizations of 2 and 3 by $(2) = \mathfrak{p}_2 \bar{\mathfrak{p}}_2$, $(3) = \mathfrak{p}_3 \bar{\mathfrak{p}}_3$ where a bar denotes conjugacy, and $\mathfrak{p}_2 \mathfrak{p}_3 = ((1 + \sqrt{-23})/2)$.

Thus in Eq. (i), $(2a^2 - 3)^2 + 23 = 24b^2$ implies the ideal equation

$$\left(\frac{2a^2 - 3 \pm \sqrt{-23}}{2} \right) = \mathfrak{q} \bar{\mathfrak{q}} \mathfrak{b}^2 \quad \text{where } \mathfrak{q} \bar{\mathfrak{q}} = (6) \text{ and } \mathfrak{b} \text{ is some integral ideal.}$$

There are essentially two possibilities, $\mathfrak{q} = \mathfrak{p}_2 \mathfrak{p}_3$ and $\mathfrak{q} = \bar{\mathfrak{p}}_2 \bar{\mathfrak{p}}_3$. In the former instance, \mathfrak{b} is principal, and in the latter, $\mathfrak{b} \bar{\mathfrak{p}}_2$ is principal.

Since $\mathfrak{p}_3 \bar{\mathfrak{p}}_2^{-1} = ((1 + \sqrt{-23})/4)$ we have, respectively,

Received October 29, 1974.

AMS (MOS) subject classifications (1970). Primary 10B10.

Key words and phrases. Diophantine equation.

Copyright © 1975, American Mathematical Society

$$\pm \left(\frac{2a^2 - 3 \pm \sqrt{-23}}{2} \right) = \left(\frac{1 + \sqrt{-23}}{2} \right) \left(\frac{u + v\sqrt{-23}}{2} \right)^2$$

and

$$\pm \left(\frac{2a^2 - 3 \pm \sqrt{-23}}{2} \right) = \left(\frac{1 + \sqrt{-23}}{4} \right) \left(\frac{u + v\sqrt{-23}}{2} \right)^2$$

for some integers u, v satisfying $u \equiv v \pmod{2}$. Thus we have, respectively,

$$\left\{ \begin{array}{l} -(2a^2 - 3) = \frac{u^2 - 46uv - 23v^2}{4} \\ 1 = \frac{u^2 + 2uv - 23v^2}{4} \end{array} \right.$$

and

$$\left\{ \begin{array}{l} (2a^2 - 3) = \frac{u^2 - 46uv - 23v^2}{8} \\ -1 = \frac{u^2 + 2uv - 23v^2}{8} \end{array} \right.$$

where the signs in each equation have been determined by a congruence modulo 3.

In the former case, putting $u + v = 2w$, we obtain

$$\text{I: } \begin{cases} a^2 = w^2 + 12wv - 12v^2 \\ 1 = w^2 - 6v^2. \end{cases}$$

In the latter case, $u^2 + 2uv + v^2 \equiv 0 \pmod{8}$, so $u + v = 4w$ say; then

$$\text{II: } \begin{cases} a^2 = 6v^2 - 12vw - 2w^2 \\ 1 = 3v^2 - 2w^2. \end{cases}$$

In similar manner (ii) gives rise to

$$\text{III: } \begin{cases} a^2 = -2w^2 + 12wv - 6v^2 \\ 1 = 4w^2 - 3v^2 \end{cases}$$

and (iii) to the three pairs

$$\text{IV: } \begin{cases} a^2 = v(9v + 16w) \\ 1 = 9v^2 - 8w^2 \end{cases} \quad \text{V: } \begin{cases} a^2 = v(v + 8w) \\ 1 = v^2 - 18w^2 \end{cases}$$

and

$$\text{VI: } \begin{cases} a^2 = 32w(v - 9w) \\ 1 = v^2 - 288w^2. \end{cases}$$

Of these six pairs of equations, V and VI may be treated by simple descent arguments. For instance, in VI, we have that $w = m^2$ or $2m^2$ after change of sign if necessary: so it suffices to determine all integer solutions of the equations $1 = v^2 - 18m^4$ and $1 = v^2 - 72m^4$, respectively. This is readily achieved by means of a classical descent argument; but we can quote Ljunggren [2] to say that the only integer solutions of the former are $(\pm r, \pm m) = (1, 0)$ and $(17, 1)$, and of the latter $(\pm r, \pm m) = (1, 0)$. These give the solutions $a = 0$ and $a = 16$ of Eq. (iii) whence solutions $x = -1, 767$ of the original equation.

Each of the four remaining pairs of equations represents the intersection of two quadrics in three-dimensional space; the method of solution, as exploited by Cassels [3], is to consider the singular elements in the pencil of the quadrics. Such singular quadrics are given by $f - \lambda g$, where $\det(f - \lambda g) = 0$: that is, a linear combination of f and g which is a perfect square. In general, of course, λ is a quadratic irrational. We can thus rewrite each pair of equations in the form $a^2 - \mu L(v, w)^2 = \lambda$ for some $\mu \in \mathbf{Q}(\lambda)$, where $L(v, w)$ is a homogeneous linear form with coefficients in $\mathbf{Q}(\lambda)$. We now work over $\mathbf{Q}(\delta)$ where $\delta^2 = \mu$ and equate $(a + L\delta)$ and $(a - L\delta)$ as ideals, to two ideal factors of λ in $\mathbf{Q}(\delta)$, noting that the two factors must be conjugate over $\mathbf{Q}(\mu)$. All the ideals are principal, so using the appropriate arithmetical details of the field $\mathbf{Q}(\delta)$, we can equate coefficients of elements of an integer base; in particular, it is clear that the coefficient of δ^2 in $a + L\delta$ is zero, and the resulting equation is completely solved by congruence considerations.

As an illustration, consider Eq. II. The singular quadrics in this pencil are obtained by taking a linear combination which is a perfect square: so let $3(2 + \lambda)v^2 - 12vw - 2(1 + \lambda)w^2$ be a perfect square. Then $36 = -6(1 + \lambda)(2 + \lambda)$ or $\lambda^2 + 3\lambda + 8 = 0$. Taking $\lambda = (-3 - \sqrt{-23})/2$ we obtain

$$a^2 - (1 + \sqrt{-23}) \left[w - \frac{1 - \sqrt{-23}}{4} v \right]^2 = \frac{3 + \sqrt{-23}}{2},$$

and accordingly work in $\mathbf{Q}(\delta)$ where $\delta^2 = 1 + \sqrt{-23}$. We need some arithmetical details of this field; certainly $\tau = (\delta^3 - 2\delta^2 + 2\delta + 4)/8$ is an algebraic integer, since $\tau^2 - \tau((1 - \sqrt{-23})/2) + 1 = 0$. The discriminant of $R = \mathbf{Z}[1, \delta, \delta^2/2, \tau]$ is $2^3 \cdot 3 \cdot 23^2$, whence R is indeed the ring of integers of the field (for 23 certainly ramifies, so 23^2 divides the discriminant, and Stickelberger's criterion says that the discriminant is congruent to 0 or 1 modulo 4). It is also readily calculated by standard techniques that τ is a fundamental unit for the field, and that we have the factorization, $(2) = \mathfrak{q}_2(\mathfrak{q}'_2)^2$, where $\mathfrak{q}_2 = \mathfrak{p}_2$, $(\mathfrak{q}'_2)^2 = \bar{\mathfrak{p}}_2$, with $\mathfrak{p}_2 = (2, (1 + \sqrt{-23})/2)$, $\bar{\mathfrak{p}}_2 = (2, (1 - \sqrt{-23})/2)$.

The equation now becomes in terms of ideals,

$$\left(a + \delta \left(w - \frac{v}{2} \right) + \frac{\delta^3}{4} v \right) \left(a - \delta \left(w - \frac{v}{2} \right) - \frac{\delta^3}{4} v \right) = (\mathfrak{q}'_2)^6;$$

and since the two ideals on the left are conjugate over $\mathbf{Z}[(1 + \sqrt{-23})/2]$ we must have

$$\left(a + \delta \left(w - \frac{v}{2} \right) + \frac{\delta^3}{4} v \right) = (\mathfrak{q}'_2)^3 = \left(\frac{\delta^3 - 6\delta + 16}{4} \right).$$

Because there are no nontrivial roots of unity in $\mathbf{Q}(\delta)$ we now obtain $a + \delta(w - v/2) + (\delta^3/4)v = \pm((\delta^3 - 6\delta + 16)/4)\tau^n$ for some integer n . This exponential equation is solved by first comparing coefficients of δ^2 , using the fact that $\tau^5 \equiv -1 \pmod{7}$; a congruence modulo a suitable power of 7 then shows that the only solutions are given by $n = 0$ or -3 . These give $a = 4$ as solution of (i), and $x = 15$ as a solution of the original equation.

The complete details of the proof are to appear in my Ph.D. Thesis. I gratefully thank Professors Swinnerton-Dyer and Cassels for their advice and encouragement.

Emmanuel College
Cambridge, England

1. L. J. MORDELL, *Diophantine Equations*, Pure and Appl. Math., vol. 30, Academic Press, London and New York, 1969, p. 259. MR 40 #2600.
2. W. LJUNGGREN, "Einige Eigenschaften der Einheiten reeller quadratischer und rein-biquadratischer Zahlkörper," *Oslo Vid.-Akad. Skrifter*, v. 1, 1936, no. 12.
3. J. W. S. CASSELS, "Integral points on certain elliptic curves," *Proc. London Math. Soc.* (3), v. 14A, 1965, pp. 55–57. MR 31 #2200.