

A Curiosity of Low-Order Explicit Runge-Kutta Methods

By J. Oliver

Abstract. By introducing an additional parameter into the first stage of the explicit Runge-Kutta process, new formulae of second and third order are derived, offering improved error bounds in the second-order case.

1. Introduction. The first stage in an explicit Runge-Kutta process for the solution of an initial-value problem for a system of ordinary differential equations, $y' = f(x, y)$, at any point (x_n, y_n) , has hitherto been presented as being necessarily the evaluation of $k_1 = hf(x_n, y_n)$. While there is no alternative to using the current known value y_n if the method is indeed to be explicit in the sense of Butcher [4], there is no corresponding computational reason why x_n should not be replaced by $x_n + \alpha_1 h$, for some α_1 chosen in the same way as the other parameters in the Runge-Kutta process. It might seem at first sight that this additional parameter would introduce an extra degree of freedom into the algebraic equations governing the parameters, and we shall show that this is indeed the case for the two-stage process. Furthermore, we obtain a new two-stage second-order method for which the truncation error bound is smaller than the previous minimum error bound found by Ralston [1]. In the three-stage case, we find that the number of degrees of freedom is in fact reduced by choosing α_1 to be nonzero, but nevertheless one such family of methods exists, while for four stages the parameters are over-determined unless $\alpha_1 = 0$.

The fact that α_1 must necessarily vanish for fourth- and higher-order methods is one possible explanation of why earlier authors have apparently overlooked this exception to the general rule in the second- and third-order cases. Another reason may be the practice [3], [4] of considering the differential system $y' = f(y)$ in which x is treated as a dependent variable whose derivative is unity. Although it is usually true, as Butcher [3] claims, that no loss of generality results from taking f to be independent of x , what follows here can be viewed as a counterexample to this assertion.

2. The General Equations. Following Ralston [1], [2] we write the general explicit Runge-Kutta method with m stages as

$$(2.1) \quad y_{n+1} - y_n = \sum_{i=1}^m w_i k_i$$

where the w_i 's are constants and

$$(2.2) \quad k_i = hf \left(x_n + \alpha_i h, y_n + \sum_{j=1}^{i-1} \beta_{ij} k_j \right)$$

Received August 9, 1974.

AMS (MOS) subject classifications (1970). Primary 65L05.

Key words and phrases. Runge-Kutta methods.

Copyright © 1975, American Mathematical Society

with $h = x_{n+1} - x_n$ and the α_i 's and β_{ij} 's also constants. Defining the operators

$$(2.3) \quad D = \partial/\partial x + (\partial/\partial y)f_n \quad \text{and} \quad D_i = \alpha_i(\partial/\partial x) + \left(\sum_{j=1}^{i-1} \beta_{ij} \right) (\partial/\partial y)f_n,$$

where $f_n = f(x_n, y_n)$, and matching both the powers of h up to h^4 and the operators in the expansion of the right-hand side of (2.1) about (x_n, y_n) with the corresponding terms in the Taylor series expansion of $y_{n+1} - y_n$, we obtain eight equations in the case $m = 4$ (from which the cases $m = 2, 3$ can readily be obtained). We give these equations in full, since in earlier presentations such as [2] the terms involving D_1 were absent.

$$\begin{aligned} h: & (w_1 + w_2 + w_3 + w_4)f = f, \\ h^2: & (w_1 D_1 f + w_2 D_2 f + w_3 D_3 f + w_4 D_4 f) = \frac{1}{2} Df, \\ h^3: & \frac{1}{2}(w_1 D_1^2 f + w_2 D_2^2 f + w_3 D_3^2 f + w_4 D_4^2 f) = \frac{1}{6} D^2 f, \\ & f_y[(w_2 \beta_{21} + w_3 \beta_{31} + w_4 \beta_{41})D_1 f \\ & \quad + (w_3 \beta_{32} + w_4 \beta_{42})D_2 f + w_4 \beta_{43} D_3 f] = \frac{1}{6} f_y Df, \\ (2.4) \quad h^4: & (w_1 D_1^3 f + w_2 D_2^3 f + w_3 D_3^3 f + w_4 D_4^3 f) = \frac{1}{24} D^3 f, \\ & \frac{1}{2} f_y[(w_2 \beta_{21} + w_3 \beta_{31} + w_4 \beta_{41})D_1^2 f \\ & \quad + (w_3 \beta_{32} + w_4 \beta_{42})D_2^2 f + w_4 \beta_{43} D_3^2 f] = \frac{1}{24} f_y D^2 f, \\ & [w_2 D_2 f_y \beta_{21} D_1 f + w_3 D_3 f_y (\beta_{31} D_1 f + \beta_{32} D_2 f) \\ & \quad + w_4 D_4 f_y (\beta_{41} D_1 f + \beta_{42} D_2 f + \beta_{43} D_3 f)] = \frac{1}{8} Df_y Df, \\ & f_y^2[(w_3 \beta_{32} \beta_{21} + w_4 \beta_{42} \beta_{21} + w_4 \beta_{43} \beta_{31})D_1 f + w_4 \beta_{43} \beta_{32} D_2 f] = \frac{1}{24} f_y^2 Df. \end{aligned}$$

It is then customary to argue that these equations can only be satisfied independently of $f(x, y)$ if the ratios

$$(2.5) \quad D_j f / Df \quad (j = 1, 2, 3, 4) \quad \text{and} \quad D_j f_y / Df_y \quad (j = 2, 3, 4)$$

are constant, which implies that

$$(2.6) \quad \alpha_i = \sum_{j=1}^{i-1} \beta_{ij} \quad (i = 1, 2, 3, 4)$$

and in particular that $\alpha_1 = 0$. We now show, however, that this argument is invalid in the second-order case.

3. Two-Stage Methods. Matching powers of h up to h^2 , and putting $w_3 = w_4 = 0$ in (2.4) yields the following three equations corresponding to the terms indicated.

$$(3.1) \quad \begin{aligned} hf: & \quad w_1 + w_2 = 1, \\ h^2 f_x: & \quad \alpha_1 w_1 + \alpha_2 w_2 = \frac{1}{2}, \\ h^2 f_y f: & \quad w_2 \beta_{21} = \frac{1}{2}. \end{aligned}$$

We thus have a two-parameter family when $\alpha_1 \neq 0$,

$$(3.2) \quad \alpha_2 = \frac{1}{2w_2} + \alpha_1 \left(1 - \frac{1}{w_2}\right), \quad \beta_{21} = \frac{1}{2w_2}, \quad w_1 = 1 - w_2,$$

including as a special case the usual one-parameter family obtained when $\alpha_1 = 0$.

Writing the truncation error T_m in an m th-order method, applied to a single differential equation for ease of analysis, as

$$(3.3) \quad T_m = c_m h^{m+1} + O(h^{m+2}),$$

the coefficient c_2 of h^3 is here given by

$$(3.4) \quad c_2 = \frac{1}{6}D^2f - \frac{1}{2}(w_1D_1^2f + w_2D_2^2f) + \frac{1}{6}f_yDf - w_2\beta_{21}f_yD_1f;$$

and, as expected, no choice of the free parameters can make this vanish independently of $f(x, y)$.

Ralston [1], [2] obtained an upper bound on $|c_2|$ by assuming that in a suitable region about (x_n, y_n) ,

$$(3.5) \quad |f(x, y)| < M \quad \text{and} \quad |\partial^{i+j}f/\partial x^i\partial y^j| < L^{i+j}/M^{j-1},$$

where M and L are constants and $i + j \leq 2$, and he showed that for $\alpha_1 = 0$ the minimum value of this bound was $ML^2/3$, obtained by setting $\alpha_2 = 2/3$, and thus

$$k_1 = hf(x_n, y_n),$$

$$(3.6) \quad k_2 = hf(x_n + 2h/3, y_n + 2k_1/3),$$

$$y_{n+1} = y_n + k_1/4 + 3k_2/4.$$

A discussion of the relevance of this type of bound to practical computation was also given by Ralston [1], [2].

Under this same assumption (3.5), we obtain from (3.4) using (2.3) and (3.2) a corresponding bound when $\alpha_1 \neq 0$ of

$$(3.7) \quad |c_2| < \left[\left| \frac{1}{2} \left(\frac{1}{3} - \alpha_1 + \alpha_1^2 \right) - \frac{(\frac{1}{2} - \alpha_1)^2}{2w_2} \right| + \left| \frac{1}{3} - \frac{1}{2}\alpha_1 - \frac{(\frac{1}{2} - \alpha_1)}{2w_2} \right| + \frac{1}{2} \left| \frac{1}{3} - \frac{1}{4w_2} \right| + \frac{1}{2} \left| \frac{1}{3} - \alpha_1 \right| + \frac{1}{6} \right] ML^2;$$

and investigation shows that this attains a smaller minimum value, namely $7ML^2/27$, for the method

$$k_1 = hf(x_n + h/3, y_n),$$

$$(3.8) \quad k_2 = hf(x_n + 5h/9, y_n + 2k_1/3),$$

$$y_{n+1} = y_n + k_1/4 + 3k_2/4.$$

On the basis of Ralston's truncation error criterion, therefore, (3.8) is preferable to all previously published second-order methods.

4. Three-Stage Methods. When $m = 3$, matching of the powers of h up to h^3 requires that, from (2.4),

$$\begin{aligned}
 &w_1 + w_2 + w_3 = 1, \\
 &w_1 D_1 f + w_2 D_2 f + w_3 D_3 f = \frac{1}{2} Df, \\
 (4.1) \quad &w_1 D_1^2 f + w_2 D_2^2 f + w_3 D_3^2 f = \frac{1}{3} D^2 f, \\
 &(w_2 \beta_{21} + w_3 \beta_{31}) f_y D_1 f + w_3 \beta_{32} f_y D_2 f = \frac{1}{6} f_y Df.
 \end{aligned}$$

In order to satisfy these equations independently of f , we must have

$$(4.2) \quad \beta_{21} = \alpha_2 \quad \text{and} \quad \beta_{31} + \beta_{32} = \alpha_3$$

so that $D_j f = \alpha_j Df$ ($j = 2, 3$). If α_1 is not to vanish, then since $D_1 f/Df$ will depend upon f , Eqs. (4.1) can only be satisfied by requiring $w_1 = 0$, in which case they reduce to

$$\begin{aligned}
 (4.3) \quad &w_2 + w_3 = 1, \quad \alpha_2 w_2 + \alpha_3 w_3 = \frac{1}{2}, \quad \alpha_2^2 w_2 + \alpha_3^2 w_3 = \frac{1}{3}, \\
 &\alpha_1 (w_2 \alpha_2 + w_3 \beta_{31}) = 0, \quad w_3 \beta_{32} \alpha_2 = \frac{1}{6}.
 \end{aligned}$$

Note that the usual choice of $\alpha_1 = 0$ effectively removes the penultimate equation of (4.3), and since w_1 can then also be nonzero, the system obtained from (4.1) has two degrees of freedom. By contrast, we have here seven equations (4.2) and (4.3) in eight parameters, possessing the one-parameter solution

$$(4.4) \quad \alpha_2 = \beta_{21} = \frac{1}{3}, \quad \alpha_3 = 1, \quad \beta_{31} = -1, \quad \beta_{32} = 2, \quad w_2 = \frac{3}{4}, \quad w_3 = \frac{1}{4}$$

with α_1 arbitrary, and thus specifying the third-order method

$$\begin{aligned}
 (4.5) \quad &k_1 = hf(x_n + \alpha_1 h, y_n), \quad k_2 = hf(x_n + h/3, y_n + k_1/3), \\
 &k_3 = hf(x_n + h, y_n - k_1 + 2k_2), \quad y_{n+1} = y_n + 3k_2/4 + k_3/4.
 \end{aligned}$$

The coefficient c_3 of h^4 in the truncation error when (4.5) is applied to a single differential equation is

$$\begin{aligned}
 (4.6) \quad &c_3 = -\frac{1}{216} D^3 f + \frac{1}{72} f_y D^2 f - \frac{1}{24} [(1 - 4\alpha_1) f_x + ff_y] Df_y \\
 &+ \frac{1}{24} [(1 - 4\alpha_1) f_x + ff_y] f_y^2
 \end{aligned}$$

and under similar assumptions (3.5) on f and its derivatives for $i + j \leq 3$ the resulting bound on $|c_3|$ attains its minimum value of $(47/216)ML^3$ when $\alpha_1 = 1/4$. Unfortunately, this bound is nearly double the value of $ML^3/9$ pertaining to the optimum third-order method with $\alpha_1 = 0$ advocated by Ralston [1].

5. Four-Stage Methods. In an attempt to find parameters which satisfy all the Eqs. (2.4) with $\alpha_1 = 0$, thus giving a four-stage method of fourth order, we follow, for the same reasons, the approach which proved successful in the three-stage case and put

$$(5. \quad \beta_{21} = \alpha_2, \quad \beta_{31} + \beta_{32} = \alpha_3, \quad \beta_{41} + \beta_{42} + \beta_{43} = \alpha_4, \quad w_1 = 0.$$

Using (5.1) to substitute for $\beta_{21}, \beta_{31}, \beta_{41}$ and w_1 in (2.4) and rearranging, we find that the following eleven equations must be satisfied by the ten parameters involved.

$$\begin{aligned}
 w_2 + w_3 + w_4 &= 1, & \alpha_2 w_2 + \alpha_3 w_3 + \alpha_4 w_4 &= \frac{1}{2}, & \alpha_2^2 w_2 + \alpha_3^2 w_3 + \alpha_4^2 w_4 &= \frac{1}{3}, \\
 \alpha_2^3 w_2 + \alpha_3^3 w_3 + \alpha_4^3 w_4 &= \frac{1}{4}, & \alpha_1 \left(w_3 \beta_{32} + w_4 \beta_{42} + w_4 \beta_{43} - \frac{1}{2} \right) &= 0, \\
 \alpha_2 w_3 \beta_{32} + \alpha_2 w_4 \beta_{42} + \alpha_3 w_4 \beta_{43} &= \frac{1}{6}, & \alpha_2^2 w_3 \beta_{32} + \alpha_2^2 w_4 \beta_{42} + \alpha_3^2 w_4 \beta_{43} &= \frac{1}{12}, \\
 \alpha_1 \left(\alpha_3 w_3 \beta_{32} + \alpha_4 w_4 \beta_{42} + \alpha_4 w_4 \beta_{43} - \frac{1}{3} \right) &= 0, \\
 \alpha_2 \alpha_3 w_3 \beta_{32} + \alpha_2 \alpha_4 w_4 \beta_{42} + \alpha_3 \alpha_4 w_4 \beta_{43} &= \frac{1}{8}, \\
 \alpha_1 \left(w_4 \beta_{43} \beta_{32} - \frac{1}{6} \right) &= 0, & \alpha_2 w_4 \beta_{43} \beta_{32} &= \frac{1}{24}.
 \end{aligned}
 \tag{5.2}$$

It is clear that the usual choice of $\alpha_1 = 0$ effectively removes three of these constraints, leaving only eight equations in nine parameters, and noting that w_1 need not then be set to zero, there are two degrees of freedom. On the other hand if we insist that $\alpha_1 \neq 0$, then it is possible though tedious to verify that Eqs. (5.2) are indeed independent, and so no four-stage explicit method with $\alpha_1 \neq 0$ can be of fourth order. A similar situation applies if more stages are introduced in order to achieve correspondingly higher order.

6. Conclusions. We have shown that if the parameter α_1 is not arbitrarily set to zero, then new families of second and third (but not fourth) order are obtained, and that on the basis of Ralston's truncation error criterion a particular one of these new methods (3.8) should be employed when a second-order Runge-Kutta method is to be used for starting the solution of an initial-value problem or changing the interval.

Computing Centre
University of Essex
Colchester, England

1. A. RALSTON, "Runge-Kutta methods with minimum error bounds," *Math. Comp.*, v. 16, 1962, pp. 431-437. MR 27 #940.
2. A. RALSTON, *A First Course in Numerical Analysis*, McGraw-Hill, New York, 1965. MR 32 #8479.
3. J. C. BUTCHER, "Coefficients for the study of Runge-Kutta integration processes," *J. Austral. Math. Soc.*, v. 3, 1963, pp. 185-201. MR 27 #2109.
4. J. C. BUTCHER, "Implicit Runge-Kutta processes," *Math. Comp.*, v. 18, 1964, pp. 50-64. MR 28 #2641.