

Application of Method of Collocation on Lines for Solving Nonlinear Hyperbolic Problems

By E. N. Houstis*

Abstract. A collocation on lines procedure based on piecewise polynomials is applied to initial/boundary value problems for nonlinear hyperbolic partial differential equations. Optimal order a priori estimates are obtained for the error of approximation. The Crank-Nicholson discretization in time is studied and convergence rates of the collocation-Crank-Nicholson procedure are established. Finally, the superconvergence is verified at particular points for linear hyperbolic problems.

Introduction. We consider the nonlinear hyperbolic problem

$$p(x, t, u)D_t^2u - q(x, t, u)D_x^2u = f(x, t, u, D_xu), \quad (x, t) \in (0, 1) \times (0, T],$$

subject to the initial conditions

$$u(x, 0) = u_0, \quad D_tu(x, 0) = u_1, \quad x \in (0, 1),$$

and to Dirichlet boundary conditions for $t > 0$. We examine the convergence of the collocation on lines procedure using piecewise polynomials with continuous first derivatives as the approximating functions.

In Section 4 we obtain optimal-order asymptotic estimates for the error of the approximation in the L_∞ -norm. In Section 5, the Crank-Nicholson discretization of the resulting system of ordinary differential equations is studied and convergence rates of the collocation on lines-Crank-Nicholson procedure are established. Finally, in Section 6 the superconvergence phenomenon is established locally for a linear hyperbolic problem.

The method of collocation on lines was proposed first by Kantorovich [7]. The convergence of this method for a problem of mathematical physics was investigated by E. B. Karpilovskaya [8]. Yartsev [11], [10] proved convergence for linear elliptic and biharmonic type problems using trigonometric polynomials as basis functions. Douglas and Dupont [3], have studied the same method using piecewise cubic Hermite polynomials for a nonlinear parabolic problem and in [4] verified the superconvergence locally for the heat equation. Finally, Douglas and Dupont [5] generalized and extended their results in [3], [4]. The results in this paper are from the author's thesis [6].

1. Preliminary Results. Let $\Delta_x = (x_i)_0^N$ be a partition of $[0, 1]$, $I = [0, 1]$, $h_j \equiv |x_{j+1} - x_j|$, $I_j \equiv [x_j, x_{j+1}]$ and $h \equiv \max_j |x_{j+1} - x_j|$. Throughout this paper

Received June 10, 1975; revised June 7, 1976.

AMS (MOS) subject classifications (1970). Primary 65N35.

Key words and phrases. Collocation on lines method, nonlinear hyperbolic problems.

*This research was partially supported by NSF Grant GP-32940X.

Copyright © 1977, American Mathematical Society

we denote by \mathbf{P}_r the set of polynomials of degree less than r and \mathbf{P}_{r,Δ_x} the set of functions that are polynomials of degree $r - 1$ in each subinterval $[x_i, x_{i+1}]$. We take $-1 < \rho_1 < \rho_2 < \dots < \rho_k < 1$ and $w_j > 0, j = 1, \dots, k$, to be Gaussian points and weights, respectively, so that

$$\int_{-1}^{+1} p(x)dx = \sum_{i=1}^k p(\rho_i)w_i, \quad p \in \mathbf{P}_{2k}([-1, 1]).$$

The Gaussian points and weights in the subinterval $[x_j, x_{j+1}]$ are

$$\xi_{kj+i} \equiv (x_j + x_{j+1})/2 + \rho_i h_j/2, \quad w_i^* = h_j w_i/2, \quad i = 1, \dots, k.$$

We introduce two pseudo-inner products corresponding to Gaussian quadrature and composite Gaussian quadrature:

$$(f, g)_{h_j} \equiv \frac{h_j}{2} \sum_{i=1}^k w_i f(\xi_{kj+i}) \cdot g(\xi_{kj+i}),$$

and

$$(f, g)_h \equiv \sum_{j=0}^{N-1} (f, g)_{h_j},$$

with

$$|f|_h \equiv \sum_{j=0}^{N-1} (f, f)_{h_j}.$$

For later use, we state without proof the lemmas:

LEMMA 1.1. *The seminorm $|f|_h$ is positive definite for all $f \in \mathbf{P}_{k+2,\Delta_x} \cap C^1[0, 1]$ with $f(0) = f(1) = 0$.*

LEMMA 1.2. *If $f, g \in \mathbf{P}_{k+2,\Delta_x} \cap C^1[0, 1]$, then*

$$\begin{aligned} & - (D_x^2 f, g)_h = (D_x f, D_x g) - D_x f \cdot g|_0^1 \\ (1.1) \quad & + \frac{(k+1)k}{(2k)!} \sum_j \frac{D_x^{k+1} f_j}{(k+1)!} \cdot \frac{D_x^{k+1} g_j}{(k+1)!} \int_{x_j}^{x_{j+1}} \prod_{i=1}^k (x - \xi_{kj+i})^2 dx. \end{aligned}$$

LEMMA 1.3. *If $f \in \{v \in \mathbf{P}_{k+2,\Delta_x} \cap C^1, v(0) = v(1) = 0\}$, then*

$$(1.2) \quad (D_x f, D_x f) \leq - (D_x^2 f, f)_h \leq 2(D_x f, D_x f)$$

and

$$(1.3) \quad |D_x f|_h^2 \leq (D_x f, D_x f).$$

LEMMA 1.4. *If $f \in \mathbf{P}_{k+2,\Delta_x} \cap C^1[0, 1]$, then*

$$(1.4) \quad |f|_h \leq \lambda \|f\|_{L^2(I)},$$

where λ is the maximum eigenvalue of the matrix $A_{k+1} \equiv [\sum_{i=1}^k w_i L_i(\rho_i) L_j(\rho_i)]$ and L_i denotes the i th degree Legendre polynomials in $[-1, 1]$.

Let H^k be the Sobolev space of functions having L^2 -derivatives of order k on I and $H_0^k \equiv \{u \in H^k | u(0) = u(1) = 0\}$.

LEMMA 1.5. For $f \in H^1$ we have

$$(1.5) \quad (D_x f, D_x f) + |f|_h^2 \geq \frac{1}{4} \|f\|_{H^1(I)}^2.$$

The above lemmas are established in [6], proofs also appear in [5]. Lemmas 1.2, 1.3 and 1.5 have been first proved for the case of cubic Hermite polynomials by Douglas and Dupont [3].

2. Approximation Theory. In [6] we show that $R_k(x) \equiv D_x^k(1-x^2)^{k+2}$, $k = 0, 1, \dots$, on $(-1, 1)$ are orthogonal polynomials. By Rodrigues' formula we see that $D_x^2 R_k(x) = D_x^{k+2}(1-x^2)^{k+2}$ is a multiple of the Legendre polynomial on the interval $(-1, 1)$. We now establish some properties of these polynomials.

LEMMA 2.1. If $k \geq 3$,

$$(2.1) \quad (D_x^\mu R_{k-2}, x^\nu)_h = 0, \quad \mu = 0, 1, 2, \nu \leq \mu.$$

PROOF. Since $D_x^\mu R_{k-2} x^\nu$ is a polynomial of degree $k+2-\mu+\nu$, we have for $k \geq 3$,

$$(D_x^\mu R_{k-2}, x^\nu)_h = \int_{-1}^1 D_x^\mu R_{k-2} x^\nu dx.$$

Lemma 2.1 now follows by using integration by parts and the fact that $D_x^\mu R_{k-2}$ vanishes at $x = \pm 1$ and $D_x^2 R_{k-2}$ vanishes at the Gaussian points. Note that for $k \geq 2$,

$$(D_x R_{k-2}, 1)_h = (D_x^2 R_{k-2}, x^\nu)_h = 0.$$

We define an interpolation operator

$$T_h: C^1(I) \rightarrow P_{k+2, \Delta_x} \cap C^1(I)$$

such that

$$\begin{aligned} (T_h v)(x_l) &= v(x_l), \\ (D_x T_h v)(x_l) &= (D_x v)(x_l), \quad l = 0, 1, \dots, N, \\ (T_h v)(\tau_{i,j}) &= v(\tau_{i,j}), \quad i = 1, \dots, k, j = 1, \dots, N, \end{aligned}$$

where $\tau_{i,j} \equiv x_j + \sigma_i(x_{j+1} - x_j)$ and the σ_i 's are the roots in the interval $(0, 1)$ of the orthogonal polynomials $R_{k-2}(x)$.

LEMMA 2.2. Assume that $u \in H^{k+4}(I)$ and let $e \equiv u - T_h u$. Then there is a constant K independent of h so that

$$|D_x^l e|_h \leq Kh^{k-l+2} \|u\|_{H^{k+2}(I)}, \quad l = 0, 1,$$

$$|D_x^2 e|_h \leq Kh^{k-l+1} \|u\|_{H^{k+3}(I)},$$

$$|(D_x e, 1)_h| \leq Kh^{2k+5/2} \|u\|_{H^{k+3}(I)},$$

$$|(D_x^2 e, 1)_h| \leq Kh^{2k+5/2} \|u\|_{H^{k+4}(I)}.$$

PROOF. It follows easily from Lemma 2.1 and Peano's Kernel Theorem [9].

3. Collocation on Lines. In this section we consider the problem of approximating the solution of the nonlinear hyperbolic equation

$$(3.1) \quad p(x, t, u)D_t^2 u - q(x, t, u)D_x^2 u = f(x, t, u, D_x u), \quad (x, t) \in (0, 1) \times (0, T],$$

subject to the initial conditions

$$(3.2) \quad u(x, 0) = \alpha_1(x), \quad D_t u(x, 0) = \alpha_2(x), \quad 0 < x < 1,$$

and the boundary conditions

$$(3.3) \quad u(0, t) = 0, \quad u(1, t) = 0, \quad 0 < t \leq T.$$

Assume that the coefficients satisfy

$$(3.4) \quad 0 < c_1 \leq p(x, t, u) \leq C_1, \quad c_2 \leq q(x, t, u) \leq C_2,$$

for $0 \leq x \leq 1$, $0 \leq t \leq T$ and $-\infty < u < +\infty$. Also, we assume that p, q, f are continuously differentiable functions of their arguments and uniformly bounded.

Throughout, we assume that this problem has a solution, u .

Let $S_{\Delta_x} \equiv \mathbf{P}_{k+2, \Delta_x} \cap C^1$ and $S_{\Delta_x} \cap H_0^1$ be spanned by the basis functions $\{B_i\}_1^{kN}$. We seek an approximation $u_h(x, t)$ to u of the form

$$u_h(x, t) = \sum_{i=1}^{kN} \beta_i(t) B_i(x).$$

The coefficients $\{\beta_i(t)\}_{i=1}^{kN}$ as functions of time are the solutions of the nonlinear ordinary differential equations

$$(3.5) \quad \{p(u_h)D_t^2 u_h - q(u_h)D_x^2 u_h - f(u_h, D_x u_h)\}(\xi_i, t) = 0, \\ 0 < t \leq T, i = 1, \dots, kN,$$

and

$$(3.6) \quad u_h(\xi_i, 0) = \hat{\alpha}_1(\xi_i), \quad D_t u_h(\xi_i, 0) = \hat{\alpha}_2(\xi_i), \quad k = 1, \dots, kN,$$

where $\hat{\alpha}_1, \hat{\alpha}_2$ are the S_{Δ_x} -interpolants of $\alpha_1(x), \alpha_2(x)$ respectively.

Although these are the equations which one solves in practice, the analysis is more conveniently made if one considers the equivalent problem of finding $u_h \in S_{\Delta_x} \cap H_0^1$ such that

$$(3.7) \quad (p(u_h)D_t^2 u_h - q(u_h)D_x^2 u_h - f(u_h, D_x u_h), B_i)_h = 0, \\ 0 < t \leq T, i = 1, \dots, kN,$$

and

$$(3.8) \quad u_h(\xi_i, 0) = \hat{\alpha}_1(\xi_i), \quad D_t u_h(\xi_i, 0) = \hat{\alpha}_2(\xi_i), \quad i = 1, \dots, kN.$$

LEMMA 3.1. *The collocation method (3.5), (3.6) and the discrete Galerkin method (3.7), (3.8) each possess a unique solution for $0 < t \leq T$. Moreover, these solutions are identical if the processes are started from the same initial values.*

PROOF. It follows from Lemma 4.1 in [5].

4. Error Analysis. In this section, we find a priori error bounds for the collocation on lines procedure. We consider the problem of finding $u_h \in S_{\Delta_x} \cap H_0^1$ such that

$$(4.1) \quad (p(u_h)D_t^2 u_h - D_x^2 u_h - f(u_h, D_x u_h), v)_h = 0, \quad 0 < t \leq T,$$

for all $v \in S_{\Delta_x} \cap H_0^1$.

In order to find estimates for the error $u - u_h$ in the L_∞ -norm, we assume that $u(\cdot, t) \in C^1(I)$ and define $w(\cdot, t) \equiv T_h u$ which is in S_{Δ_x} . Then we find a priori bounds for the difference $w - u_h \in S_{\Delta_x}$; and applying known approximation results to the difference $u - w$, we obtain bounds for the error of the collocation on lines procedure.

If X is a normed space and $\psi: [0, T] \rightarrow X$, define

$$\|\psi\|_{L^2(0,T;X)} = \int_0^T \|\psi(t)\|_X^2 dt, \quad \|\psi\|_{L^\infty(0,T;X)} = \sup_{0 \leq t \leq T} \|\psi(t)\|_X.$$

THEOREM 4.1. *If*

(i) *the coefficients in (3.1) have bounded third derivatives and satisfy conditions (3.4),*

(ii) $u \in L^\infty(0, T; H^{k+4})$, $D_t u \in L^2(0, T; H^{k+4})$ and $D_t^2 u \in L^2(0, T; H^{k+4})$,

(iii) $u_h(x, 0)$, $D_t u_h(x, 0)$ are the S_{Δ_x} interpolants of $u(x, 0)$ and $D_t u(x, 0)$, respectively, then for the error of approximation we have

$$\|u - u_h\|_{L^\infty(0,T;L^\infty)} \leq K [\|u\|_{L^\infty(0,T;H^{k+4}(I))} + \|D_t u\|_{L^2(0,T;H^{k+4}(I))} + \|D_t^2 u\|_{L^2(0,T;H^{k+4}(I))}] h^{k+2},$$

where K is a constant independent of h and u .

PROOF. Let $\eta \equiv u - w$ and $\zeta \equiv w - u_h$. Then (3.1), (4.1) imply that

$$(4.2) \quad \begin{aligned} (p(u_h)D_t^2 \zeta - D_x^2 \zeta, v)_h &= (-p_u^1 \zeta D_t^2 w - p(w)D_t^2 \eta - p_u^2 \eta D_x^2 u, v)_h \\ &+ (D_x^2 \eta, v)_h + ([f(w, D_x u) - f(w, D_x w)], v)_h \\ &+ (f_u^1 \eta + f_u^2 \zeta + f_{D_x u}^3 u D_x \zeta, v)_h. \end{aligned}$$

In (4.2) we choose $v = D_t \zeta$ and in [6] we show that

$$(4.3) \quad \begin{aligned} &\frac{1}{2} [\sqrt{p(u_h)} D_t \zeta|_h^2 + |\zeta|_h^2 - (D_x^2 \zeta, \zeta)_h] \\ &\leq K \int_0^t \{|\zeta|_h^2 + |D_x \zeta|_h^2\} d\tau + \int_0^t \{|\eta|_h^2 + |D_t^2 \eta|_h^2\} d\tau + K \int_0^t |D_t \zeta|_h^2 \\ &+ K \{|\zeta|_h^2(0) - (D_x^2 \zeta, \zeta)_h(0) + \sqrt{p(u_h)} D_t \zeta|_h^2(0)\} \\ &+ \int_0^t (D_x^2 \eta, D_t \zeta)_h d\tau + \int_0^t (f(w, D_x u) - f(w, D_x w), D_t \zeta)_h d\tau. \end{aligned}$$

Integration by parts gives

$$\int_0^t (D_x^2 \eta, D_t \zeta)_h d\tau = (D_x^2 \eta, \zeta)_h|_0^t - \int_0^t (D_t D_x^2 \eta, \zeta)_h d\tau,$$

and

$$\int_0^t (f(w, D_x u) - f(w, D_x w), D_t \zeta)_h d\tau = (f(w, D_x u) - f(w, D_x w), \zeta)_h \Big|_0^t - \int_0^t (D_t \{f(w, D_x u) - f(w, D_x w)\}, \zeta)_h d\tau.$$

Using Poincaré’s inequality, the elementary inequality $|cd| \leq (\frac{1}{4}p)c^2 + pd^2$ and Lemma 2.2 in [6] we have obtained

$$(4.4) \quad \left| \int_0^t (D_t D_x^2 \eta, \zeta)_h d\tau \right| \leq \frac{1}{16} \int_0^t [-(D_x^2 \zeta, \zeta)_h + |\zeta|_h^2] d\tau + K \sum_{i=1}^N h_j^{2k+4} \int_0^t \|D_t u(\cdot, \tau)\|_{H^{k+4}(I_j)}^2 d\tau.$$

Using Taylor’s theorem, we can easily show that

$$(f(w, D_x u) - f(w, D_x w), w - u_h)_h = \sum_{j=1}^N (f_{D_x u}(w, D_x w)|_{x=\xi_{k(j-1)+1}} D_x \eta + \omega h_j D_x \eta, \zeta)_{h_j},$$

where ω is bounded independent of h_j . It follows from Lemma 2.2

$$|(f_{D_x u}|_{x=\xi_{k(j-1)+1}} D_x \eta, \zeta)_{h_j}| \leq K h_j^{k+2} [\|u(\cdot, t)\|_{H^{k+3}(I_j)} |\zeta|_{h_j} + \|u(\cdot, t)\|_{H^{k+3}(I_j)} \|D_x \zeta(\cdot, t)\|_{L^2(I_j)}],$$

and

$$|(\omega h_j D_x \eta, \zeta)_{h_j}| \leq K h_j^{k+2} \|u(\cdot, t)\|_{H^{k+2}(I_j)} |\zeta|_{h_j}.$$

Moreover, we obtain

$$(f(w, D_x u) - f(w, D_x w), \zeta)_h \leq \frac{1}{16} [-(D_x^2 \zeta, \zeta)_h + |\zeta|_h^2] + K \sum_{j=1}^N h_j^{2k+4} \|u(\cdot, t)\|_{H^{k+3}(I_j)}^2.$$

Following similar arguments as above, we show that

$$(4.5) \quad \int_0^t (D_t \{f(w, D_x u) - f(w, D_x w)\}, \zeta)_h d\tau \leq \frac{1}{16} \int_0^t [-(D_x^2 \zeta, \zeta)_h + |\zeta|_h^2] d\tau + K \sum_{j=1}^N h_j^{2k+4} \int_0^t [\|u(\cdot, \tau)\|_{H^{k+3}(I_j)}^2 + \|D_t u(\cdot, \tau)\|_{H^{k+3}(I_j)}^2] d\tau.$$

It follows from (4.3)–(4.5), (1.3) and Gronwall’s Lemma [6] that

$$\begin{aligned}
& \|w - u_h\|_{L^\infty(0,T;L^\infty)}^2 \\
& \leq K [\|(w - u_h)(\cdot, 0)\|_{H^1(I)}^2 + \|D_t(w - u_h)(\cdot, 0)\|_{L^2(I)}^2] \\
(4.6) \quad & + \sum_{j=1}^N h_j^{2k+4} \{ \|u\|_{L^\infty(0,T;H^{k+4}(I_j))}^2 \\
& \quad + \|D_t u\|_{L^2(0,T;H^{k+4}(I_j))}^2 + \|D_t^2 u\|_{L^2(0,T;H^{k+4}(I_j))}^2 \}.
\end{aligned}$$

It is an elementary consequence of Peano's Kernel Theorem that

$$(4.7) \quad \|u - w\|_{L^\infty(0,T;L^\infty)}^2 \leq K \sum_{j=1}^N h_j^{2k+4} \|u\|_{L^\infty(0,T;H^{k+4}(I_j))}^2.$$

Finally, from (4.6), (4.7) and assumption (iii) it follows that

$$\begin{aligned}
& \|u - u_h\|_{L^\infty(L^\infty(I))} \\
& \leq Kh^{k+2} [\|u\|_{L^\infty(H^{k+4}(I))} + \|D_t u\|_{L^2(H^{k+4}(I))} + \|D_t^2 u\|_{L^2(H^{k+4}(I))}].
\end{aligned}$$

This concludes the proof of Theorem 4.1.

5. Computational Considerations. In this section, we discuss the question of actually solving the system of ordinary differential equations (3.5), (3.6).

Let

$$\begin{aligned}
& u_h^j \equiv u_h^j(x) = u_h^j(x, t^j), \quad t^j \equiv j \Delta t, \quad \Delta t = T/N, \\
(5.1) \quad & v^{j+\frac{1}{2}} \equiv (v^{j+1} + v^j)/2, \quad v^{j,\frac{3}{4}} \equiv \frac{1}{4} v^{j+1} + \frac{1}{2} v^j + \frac{1}{4} v^{j-1},
\end{aligned}$$

$$\partial_t v^{j+\frac{1}{2}} \equiv (v^{j+1} - v^j)/\Delta t, \quad \partial_t^2 v^j \equiv (v^{j+1} - 2v^j + v^{j-1})/(\Delta t)^2.$$

Then the Crank-Nicholson-Collocation approximation $\{u_h^j\}_0^N$ is defined such that

$$\begin{aligned}
(5.2) \quad (i) \quad & \{p(t^j, u_h^{j,\frac{3}{4}}) \partial_t^2 u_h^j - q(t^j, u_h^{j,\frac{3}{4}}) D_x^2 u_h^{j,\frac{3}{4}} - f(t^j, u_h^{j,\frac{3}{4}}, D_x u_h^{j,\frac{3}{4}})\}(\xi_i) = 0, \\
& i = 1, \dots, kN, j = 0, \dots, N-1,
\end{aligned}$$

$$(ii) \quad u_h^j(0) = u_h^j(1) = 0, \quad j = 0, \dots, N.$$

At the end of this section we discuss the choice of u_h^0, u_h^1 . In order to analyze the convergence of the solution of (5.2) we consider the equivalent to (5.2) normalized problem

$$(5.3) \quad (p(t^j, u_h^{j,\frac{3}{4}}) \partial_t^2 u_h^j, v)_h - (D_x^2 u_h^{j,\frac{3}{4}}, v)_h = (f(t^j, u_h^{j,\frac{3}{4}}, D_x u_h^{j,\frac{3}{4}}), v)_h,$$

$$v \in S_{\Delta x} \cap H_0^1, \quad 0 \leq j < N.$$

Also, we introduce the notation

$$\|u\|_{L_{\Delta t}^2(0,T;X)}^2 \equiv \sum_{0 < t^j < T} \|u^j\|_X^2 \Delta t,$$

$$\|u\|_{L_{\Delta t}^\infty(0,T;X)}^2 \equiv \max_{0 < t^j < T} \|u^j\|_X^2,$$

$$\|u\|_{L^2_{\Delta t}(0,T;X)}^2 \equiv \sum_{0 < t^j < T} \|u^j\|_X^2 \Delta t.$$

THEOREM 5.1. *Assume the hypotheses (i), (ii) of Theorem 4.1 hold. Further, assume $D_t^3 u, D_t^4 u$ are in $L^\infty(0, T; L^2(I))$ and*

$$\|(u_h - w)^{1/2}\|_{H^1(I)} + \|\partial_t(u_h - w)^{1/2}\|_{L^2(I)} = O(h^{k+2}).$$

For Δt sufficiently small there exists a unique solution of the Crank-Nicholson-Collocation equations (5.2) and for the error of approximation we have

$$\|u - u_h\|_{L^\infty_{\Delta t}(0,T;L^\infty)} \leq C(h^{k+2} + (\Delta t)^2),$$

where C depends on u and is independent of $h, \Delta t$.

PROOF. It is easily seen that a unique solution of (5.2) exists under assumption (i) and (3.3) for Δt sufficiently small. Throughout this proof we use the notation $w \equiv T_h u, \eta \equiv u - w$ and $\xi \equiv u_h - w$. First, we observe that u satisfies

$$(5.4) \quad (p(u^{j,1/4}) \partial_t^2 u^j)_h - (D_x^2 u^{j,1/4}, v)_h = (f(u^{j,1/4}, D_x u^{j,1/4}), v)_h + (e^j, v)_h$$

for $v \in S_{\Delta x} \cap H_0^1$, where $\|e^j\|_{L^2(I)} = O(\Delta t^2) \|D_t^4 u\|_{L^2(I)}$.

After straightforward calculations and the application of the Mean Value Theorem, we obtain

$$\begin{aligned} & (p(u_h^{j,1/4}) \partial_t^2 \xi^j, v)_h - (D_x^2 \xi^j, v) \\ &= (p^* \xi^{j,1/4} \partial_t^2 w^j, v)_h + (p(w^{j,1/4}) \partial_t^2 \eta^j, v)_h \\ (5.5) \quad &+ (p^{**} \eta^{j,1/4} \partial_t^2 u^j, v)_h + (e^j, v)_h - (D_x^2 \eta^{j,1/4}, v)_h \\ &+ (f_u^* \eta^{j,1/4} + f_u^{**} \xi^{j,1/4} + f_{D_x u}^* D_x \xi^{j,1/4}, v)_h \\ &+ (f(w^{j,1/4}, D_x w^{j,1/4}) - f(w^{j,1/4}, D_x u^{j,1/4}), v)_h. \end{aligned}$$

In (5.5), we choose as test function $v = (\xi^{j+1} - \xi^{j-1})/2t$ and then we obtain

$$\begin{aligned} & \frac{1}{2\Delta t} \{ [|\sqrt{p(u_h^{j,1/4})} \partial_t \xi^{j+1/2}|_h^2 + |\xi^{j+1/2}|_h^2 - (D_x^2 \xi^{j+1/2}, \xi^{j+1/2})_h] \\ & \quad - [|\sqrt{p(u_h^{j,1/4})} \partial_t \xi^{j-1/2}|_h^2 + |\xi^{j-1/2}|_h^2 - (D_x^2 \xi^{j-1/2}, \xi^{j-1/2})_h] \} \\ & \leq C [|\xi^{j+1/2}|_h^2 + |\xi^{j-1/2}|_h^2 + |\partial_t^2 \eta^j|_h^2 + |\eta^{j,1/4}|_h^2 + |\partial_t \xi^{j+1/2}|_h^2 \\ (5.6) \quad & \quad + |\partial_t \xi^{j-1/2}|_h^2 + |D_x \xi^{j,1/4}|_h^2 + |e^j|_h^2] \\ & \quad + \left| \left(D_x^2 \eta^{j,1/4}, \frac{\xi^{j+1} - \xi^{j-1}}{2\Delta t} \right)_h \right| \\ & \quad + \left| \left(f(w^{j,1/4}, D_x w^{j,1/4}) - f(w^{j,1/4}, D_x u^{j,1/4}), \frac{\xi^{j+1} - \xi^{j-1}}{2\Delta t} \right)_h \right|, \end{aligned}$$

where C is a generic constant.

$$\begin{aligned}
 & |\partial_t \xi^{n-1/2}|_h^2 + |\xi^{n-1/2}|_h^2 - (D_x^2 \xi^{n-1/2}, \xi^{n-1/2})_h \\
 & \leq C \{ - (D_x^2 \xi^{1/2}, \xi^{1/2})_h + |\xi^{1/2}|_h^2 |\partial_t \xi^{1/2}|_h^2 \} \\
 & + K \Delta t \sum_{j=1}^{n-1} \sum_{i=0}^{N-1} h_i^{2k+4} [\|u^j\|_{H^{k+3}(I_i)}^2 + \|D_t^2 u^j\|_{H^{k+4}(I_i)}^2] \\
 (5.9) \quad & + C \Delta t \sum_{j=1}^{n-1} \{ |\partial_t^2 \eta^j|_h^2 + |\eta^{j-1/2}|_h^2 \} + \Delta t \sum_{j=1}^{n-1} [\|e_2^j\|_{L^2(I)}^2 + \|e_3^j\|_{L^2(I)}^2] \\
 & + K \left[\max_{0 \leq t^j \leq T} \sum_{i=0}^{N-1} h_i^{2k+4} \|u^j\|_{H^{k+4}(I_i)}^2 + \max_{0 \leq t^j \leq T} \|e_1^j\|_{L^2(I)}^2 \right].
 \end{aligned}$$

Finally from Lemma 1.4, 2.2 and inequality (5.9), we conclude that

$$\begin{aligned}
 (5.10) \quad \|\xi\|_{L^\infty_{\Delta t}(0,T;L^\infty)} & \leq C [\|\xi^{1/2}\|_{H^1(I)} + \|\partial_t \xi^{1/2}\|_{L^2(I)}] \\
 & + Kh^{k+2} [\|u\|_{\tilde{L}^2_{\Delta t}(0,T;H^{k+3}(I))} + \|D_t^2 u\|_{\tilde{L}^2(0,T;H^{k+4}(I))}] \\
 & + \|u\|_{L^\infty_{\Delta t}(0,T;H^{k+4}(I))}] \\
 & + c(u)\Delta t^2,
 \end{aligned}$$

where C and K are generic constants independent of u , h , Δt and $c(u)$ independent of h , Δt . From the results of Section 2 we easily see that

$$(5.11) \quad \|\eta\|_{L^\infty_{\Delta t}(0,T;L^\infty)} \leq Ch^{k+2} \|u\|_{L^\infty_{\Delta t}(0,T;H^{k+2})}.$$

Therefore, the inequalities (5.10) and (5.11) imply

$$\|u - u_h\|_{L^\infty_{\Delta t}(0,T;L^\infty)} \leq c(u) (h^{k+2} + (\Delta t)^2),$$

provided

$$\|\xi^{1/2}\|_{H^1(I)} + \|\partial_t \xi^{1/2}\|_{L^2(I)} \leq ch^{k+2},$$

where $c(u)$ is independent of h and Δt . This concludes the proof of Theorem 5.1.

It remains to discuss the choice of u_h^0 and u_h^1 . We choose $u_h^0 \equiv T_h u(x, 0)$ and $u_h^1 \equiv T_h \tilde{u}$ where

$$\tilde{u} \equiv u(x, 0) + \Delta t D_t u(x, 0) + \frac{(\Delta t)^2}{2} D_t^2 u(x, 0) + \frac{(\Delta t)^3}{6} D_t^3 u(x, 0);$$

the derivatives $D_t^2 u$ and $D_t^3 u$ are evaluated using the differential equation.

6. The Superconvergence Phenomenon. Consider the linear hyperbolic problem

$$(6.1) \quad p(x, t) D_t^2 u - D_x^2 u = f(x, t), \quad (x, t) \in (0, 1) \times (0, T),$$

subject to initial conditions

$$(6.2) \quad u(x, 0) = \varphi_1(x), \quad D_t u(x, 0) = \varphi_2(x), \quad 0 \leq x \leq 1,$$

and boundary conditions

$$(6.3) \quad u(0, t) = 0, \quad u(1, t) = 0, \quad 0 < t \leq T.$$

Also, we assume for all $(x, t) \in [0, 1] \times [0, T]$,

$$(6.4) \quad 0 < m < p(x, t) \leq M, \quad 0 < m \leq q(x, t) \leq M.$$

Let u_h denote the collocation on lines approximation defined from (3.5) and (3.6) where p, q and f are independent of u . Throughout we denote by $L \equiv pD_t^2 - D_x^2, \|u\|_{j,i} \equiv \sup \{D_x^\alpha D_t^\beta u(x, t) | x \in I, \alpha \leq j, \beta \leq i\}$ and $x_{i-1/2} \equiv (i - 1/2)h_j$. By Peano's Kernel Theorem [9] we obtain

$$\begin{aligned} L(u - T_h u)(\xi_{kj+i}, t) &= \sum_{i=1}^{s-2} \{D_x^{k+i+1} D_t^2 u(x_{j-1/2}) \psi_i(\rho_i) - D_x^{k+i+3} u(x_{j-1/2}) \psi_{i+2}''(\rho_i)\} h_j^{k+i+1} \\ &\quad - D_x^{k+3} u(x_{j-1/2}) \psi_2''(\rho_i) h_j^{k+1} + O(h_j^{k+s} [\|u\|_{k+s+2,0} + \|u\|_{k+s+2,2}]), \end{aligned}$$

where

$$\psi_i(x) = \frac{1}{(k+i+1)!} A_i(x) R_{k-2}(x)$$

with A_i a polynomial of degree $i - 1$. In order to cancel the term of h_j^{k+1} accuracy we make a correction to $T_h u$ defined locally by the following relations, $\delta_0(\cdot, t) \in P_{k+2, \Delta_x} \cap C^1$ with

$$\begin{aligned} h^{s-1} D_x^2 \delta_0(\xi_{kj+i}, t) &= D_x^{k+3} u(x_{j-1/2}) \psi_2''(\rho_i), \quad i = 1, \dots, k, j = 0, \dots, N-1, \\ \delta_0(x_j, t) = D_x \delta(x_j, t) &= 0, \quad j = 0, 1, \dots, N. \end{aligned}$$

Now, in order to cancel the h_j^{k+1} order terms we define a new correction in the following way: first we introduce the function

$$v(y) = \begin{cases} 0, & y \leq 0, \\ 3y^2 - 2y^3, & 0 \leq y \leq 1, \\ 1, & 1 \leq y, \end{cases}$$

which obviously belongs to C^1 and define for $x \in I_j$

$$E_j(x, t) \equiv \lambda_{1,i} D_x^{k+i+1} D_t^2 u(x_{j-1/2}) v\left(\frac{x-x_j}{h_j}\right) - \lambda_{2,i} D_x^{k+i+3} u(x_{j-1/2}) v\left(\frac{x-x_j}{h_j}\right),$$

where $\lambda_{1,i} \equiv -\psi_i(\rho_i)/v''(\rho_i), \lambda_{2,i} \equiv -\psi_{i+2}''(\rho_i)/v''(\rho_i)$. Also, we define

$$\begin{aligned} \delta_l(x, t) &\equiv \sum_{j=0}^{N-1} h_j^{l+3-s} \{E_j(x, t) - xE_j(1, t)\} \\ &= \sum_{j=0}^{N-1} \{\lambda_{1,l} D_x^{k+l+1} D_t^2 u(x_{j-\frac{1}{2}}) - \lambda_{2,l} D_x^{k+l+3} u(x_{j-\frac{1}{2}})\} \left(v \left(\frac{x - x_j}{h_j} \right) - x \right). \end{aligned}$$

In [6] we show that the $\lambda_{\alpha,l}$ for $\alpha = 1, 2$ are well defined and easily obtain

$$L(u - \bar{u})(\xi_{kj+i}, t) = O(h_j^{k+s} [\|u\|_{k+s+2,0} + \|u\|_{k+s+2,2}]),$$

where

$$\bar{u} = T_h u + h_j^{k+s} \sum_{l=0}^{s-2} \delta_l.$$

THEOREM 6.1. *Let u denote the solution of the problem (6.1) to (6.4) such that $u \in L^\infty(0, T; H^{k+s+4})$, $s \leq k$ and u_h is the collocation on lines approximation of u defined by (3.5), (3.6). Then the error of approximation at the nodes satisfies*

$$\begin{aligned} \max_j \|(u - u_h)(x_j, \cdot)\|_{L^\infty(0,T)} &\leq Ch^{k+s} [\|u\|_{k+s+2,0} + \|u\|_{k+s+2,2}] \\ &\quad + C[\|D_t(u_h - \bar{u})\|_{L^2(I)}(0) + \|u_h - \bar{u}\|_{H^1(I)}(0)], \end{aligned}$$

where C is a constant independent of u and h and $s \leq k$.

PROOF. We define

$$\rho(\xi_{kj+i}, t) \equiv L(u_h - \bar{u})(\xi_{kj+i}, t),$$

where

$$|\rho(\xi_{kj+i}, t)| \leq Ch_j^{k+s} [\|u\|_{k+s+2,0} + \|u\|_{k+s+2,2}];$$

and we form the relation

$$(\rho, D_t(u_h - \bar{u}))_h = (D_t^2(u_h - \bar{u}), D_t(u_h - \bar{u}))_h - (D_x^2(u_h - \bar{u}), D_t(u_h - \bar{u}))_h.$$

We apply the elementary inequality $\frac{1}{2}a^2 + \frac{1}{2}b^2 \geq ab$ to obtain

$$|\rho|_h^2 + |D_t(u_h - \bar{u})|_h^2 \geq D_t |D_t(u_h - \bar{u})|_h^2 - D_t (D_x^2(u_h - \bar{u}), u_h - \bar{u})_h.$$

In the above inequality we add the inequality

$$\frac{1}{2} D_t |u_h - \bar{u}|_h^2 \leq \frac{1}{2} |D_t(u_h - \bar{u})|_h^2 + \frac{1}{2} |u_h - \bar{u}|_h^2$$

to obtain

$$\begin{aligned} |\rho|_h^2 + |u_h - \bar{u}|_h^2 + 2|D_t(u_h - \bar{u})|_h^2 \\ \geq D_t \{ |u_h - \bar{u}|_h^2 + |D_t(u_h - \bar{u})|_h^2 \} - D_t (D_x^2(u_h - \bar{u}), u_h - \bar{u})_h. \end{aligned}$$

We integrate from 0 to t and apply Gronwall's Lemma to get

$$\begin{aligned} & C \int_0^T |\rho_h^2(\tau)| d\tau + |u_h - \bar{u}|_h^2(0) + |D_t(u_h - \bar{u})|_h^2(0) - (D_x^2(u_h - \bar{u}), u_h - \bar{u})_h(0) \\ & \geq |u_h - \bar{u}|_h^2 + |D_t(u_h - \bar{u})|_h^2 - (D_x^2(u_h - \bar{u}), u_h - \bar{u})_h. \end{aligned}$$

It follows from Lemmas 1.3, 1.4, 1.5 that

$$\begin{aligned} & C \left\{ \max_t |\rho|_h + \|D_t(u_h - \bar{u})\|_{L^2(I)}(0) + \|u_h - \bar{u}\|_{H^1(I)}(0) \right\} \\ & \geq \|u_h - \bar{u}\|_{H^1(I)} + |D_t(u_h - \bar{u})|_h^2. \end{aligned}$$

In particular, we have

$$C \left\{ \max_t |\rho|_h + \|D_t(u_h - \bar{u})\|_{L^2(I)}(0) + \|u_h - \bar{u}\|_{H^1(I)}(0) \right\} \geq \|u_h - \bar{u}\|_{L^\infty(I)}.$$

It is easy to see that

$$|(u - \bar{u})(x_j, t)| \leq Ch^{k+s} [\|u\|_{k+s-1,2} + \|u\|_{k+s+1,0}],$$

where $h = \max_j h_j$. Consequently, we have

$$\begin{aligned} \max_t \max_{0 \leq j \leq N} |(u - u_h)(x_j, t)| & \leq Ch^{k+s} [\|u\|_{k+s+2,0} + \|u\|_{k+s+2,2}] \\ & + C \{ \|D_t(u_h - \bar{u})\|_{L^2(0)} + \|u_h - \bar{u}\|_{H^1(I)}(0) \}. \end{aligned}$$

This completes the proof of Theorem 6.1.

Now, we consider the problem of choosing initial values, in order to obtain maximum accuracy. It is clear that the following

$$u_h(x, 0) = T_h \varphi_1 + h^{k+s} \delta_0(x, 0), \quad D_t u_h(x, 0) = T_h \varphi_2 + h^{k+s} \delta_0(x, 0)$$

yields $(u_h - \bar{u})(0) = O(h^{k+s})$ in H^1 norm, and $D_t(u_h - \bar{u})(0) = O(h^{k+s})$ in L^2 norm.

Department of Computer Sciences
Purdue University
West Lafayette, Indiana 47904

1. I. S. BEREZIN & N. P. ŽIDKOV, *Computing Methods*, Vols. I, II, Fizmatgiz, Moscow, 1962; English transl., Addison-Wesley, Reading, Mass.; Pergamon Press, New York, 1965. MR 22 #12685; 30 #4372.
2. E. A. CODDINGTON, *An Introduction to Ordinary Differential Equations*, Prentice-Hall, Englewood Cliffs, N.J., 1961. MR 23 #A3869.
3. JIM DOUGLAS, JR. & TODD DUPONT, "A finite element collocation method for quasi-linear parabolic equations," *Math. Comp.*, v. 27, 1973, pp. 17-28. MR 49 #4266.
4. JIM DOUGLAS, JR. & TODD DUPONT, "A super convergence result for the approximate solution of the heat equation by a collocation method," *Mathematical Foundations of Finite Element Method with Applications to Partial Differential Equations* (A. K. Aziz, Editor), Academic Press, New York, 1972.
5. JIM DOUGLAS, JR. & TODD DUPONT, *Collocation Methods for Parabolic Equations in a Single Space Variable (Based on C^1 -Piecewise-Polynomial Spaces)*, Springer Lecture Notes in Math., Vol. 385, Springer-Verlag, Berlin and New York, 1974.

6. E. N. HOUSTIS, *Finite Element Methods for Solving Initial/Boundary Value Problems*, Doctoral thesis, Purdue University, 1974.
7. L. V. KANTOROVIČ, "Sur une méthode de résolution approchée d'équations différentielles aux dérivées partielles," *C. R. Acad. (Dokl.) Sci. URSS*, v. 2, 1934, pp. 532–536. (Russian)
8. È. B. KARPILOVSKAJA, "Convergence of a collocation method for certain boundary-value problems of mathematical physics," *Sibirsk. Mat. Ž.*, v. 4, 1963, pp. 632–640. (Russian) MR 27 #6402.
9. M. H. SCHULTZ, *Spline Analysis*, Prentice-Hall, Englewood Cliffs, N.J., 1973. MR 50 #15270.
10. Yu. P. YARTSEV, "Convergence of the collocation method on lines," *Differencial'nye Uravnenija*, v. 3, 1967, pp. 1606–1613 = *Differential Equations*, v. 3, 1967, pp. 838–842.
11. Yu. P. YARTSEV, "The method of line collocation," *Differencial'nye Uravnenija*, v. 4, 1968, pp. 925–932 = *Differential Equations*, v. 4, 1968, pp. 481–485.