

# Evaluation of the Integral $\int_0^\infty \frac{t^{2\alpha-1} J_\nu(x\sqrt{1+t^2})}{(1+t^2)^{\alpha+\beta-1}} dt$

By Paul W. Schmidt\*

**Abstract.** Methods are developed for evaluating the integral

$$I_\nu^{\alpha\beta}(x) = \int_0^\infty \frac{t^{2\alpha-1} J_\nu(x\sqrt{1+t^2})}{(1+t^2)^{\alpha+\beta-1}} dt,$$

where  $J_\nu(t)$  is the Bessel function of the first kind and order  $\nu$ ,  $\alpha > 0$ ,  $\beta > 1/4$ , and  $\nu$  is real. Only  $I_\nu^{1/2,1}(x)$  and  $I_\nu^{\alpha,\nu/2+1-\alpha}(x)$  are included in previously published tables of integrals of Bessel functions. The integrals  $I_1^{1/2,1/2}(x)$  and  $I_2^{1/2,1}(x)$  are used in a technique developed by I. S. Fedorova for calculating the diameter distribution of long circular cylinders from small-angle x-ray, light, or neutron scattering data.

The  $I_\nu^{\alpha\beta}(x)$  are shown to be proportional to a  $G$  function. From this result, power series expansions and recurrence relations are developed for use in evaluating the  $I_\nu^{\alpha\beta}(x)$ . A convenient expression is obtained for the quantity required in Fedorova's method for computing diameter distributions.

**I. Introduction.** In the method which Fedorova has recently developed [1], [2] for using light, small-angle x-ray, or small-angle neutron scattering data to calculate the diameter distribution of assemblies of independently-scattering long circular cylinders, the quantity

$$\Phi(a) = aI_1^{1/2,1/2}(a) - I_2^{1/2,1}(a)$$

must be evaluated, where

$$(1) \quad I_\nu^{\alpha\beta}(x) = \int_0^\infty \frac{t^{2\alpha-1} J_\nu(x\sqrt{1+t^2})}{(1+t^2)^{\alpha+\beta-1}} dt,$$

$x > 0$ , and  $J_\nu(x)$  is the Bessel function of the first kind and order  $\nu$ . The conditions

$$(2) \quad \alpha > 0, \beta > 1/4,$$

ensure that the integral  $I_\nu^{\alpha\beta}(x)$  exists for all real  $\nu$ .

As Fedorova has pointed out [2],  $I_\nu^{1/2,1}(x)$  can be expressed [3] in terms of Bessel functions. Only  $I_\nu^{1/2,1}(x)$  and  $I_\nu^{\alpha,\nu/2+1-\alpha}(x)$  have been listed in tables of integrals of Bessel functions.

In Section II,  $I_\nu^{\alpha\beta}(x)$  is shown to be expressible in terms of a  $G$  function [4]. A series expansion and some recurrence relations are developed which are useful for evaluating  $I_\nu^{\alpha\beta}(x)$ . By use of some of the recurrence relations, in Section III the quantity  $\Phi(a)$  used in Fedorova's diameter distribution method is expressed in terms of Bessel functions.

Received May 24, 1976; revised September, 1976, December 14, 1976 and May 12, 1977.

AMS (MOS) subject classifications (1970). Primary 33A40; Secondary 44A20.

\*Work supported by the U. S. National Science Foundation.

## II. Some Techniques for Calculating the $I_\nu^{\alpha\beta}(x)$ .

(a) *Power Series.* The Bessel functions  $J_\nu(x)$  are special cases of the  $G$  function [4] and can be expressed [5]

$$J_\nu(x) = G_{0,2}^{1,0} \left( \frac{x^2}{4} \middle| \frac{\nu}{2}, -\frac{\nu}{2} \right).$$

With this result, the integral  $I_\nu^{\alpha\beta}(x)$  defined in (1) can be written [6], [7]

$$\begin{aligned} (3) \quad I_\nu^{\alpha\beta}(x) &= \frac{1}{2} \Gamma(\alpha) G_{1,3}^{2,0} \left( \frac{x^2}{4} \middle| \alpha + \beta - 1, \frac{\nu}{2}, -\frac{\nu}{2} \right) \\ &= \frac{1}{2} \Gamma(\alpha) \left\{ \frac{\Gamma\left(\frac{\nu}{2} - \beta + 1\right)}{\Gamma\left(\frac{\nu}{2} + \beta\right) \Gamma(\alpha)} \left(\frac{x^2}{4}\right)^{\beta-1} {}_1F_2 \left( \begin{matrix} 1 - \alpha \\ \beta - \frac{\nu}{2}, \beta + \frac{\nu}{2} \end{matrix} \middle| -\frac{x^2}{4} \right) \right. \\ &\quad \left. + \frac{\Gamma\left(\beta - 1 - \frac{\nu}{2}\right)}{\Gamma(1 + \nu) \Gamma\left(\alpha + \beta - 1 - \frac{\nu}{2}\right)} {}_1F_2 \left( \begin{matrix} 2 + \frac{\nu}{2} - \alpha - \beta \\ 2 + \frac{\nu}{2} - \beta, 1 + \nu \end{matrix} \middle| -\frac{x^2}{4} \right) \right\}, \end{aligned}$$

where the  ${}_1F_2$  functions are hypergeometric functions.

When this  $G$  function is expanded in a power series [7],  $I_\nu^{\alpha\beta}(x)$  can be expressed

$$(4) \quad I_\nu^{\alpha\beta}(x) = \frac{\frac{\pi}{2} \Gamma(\alpha)}{\sin\left(\frac{2\beta - \nu}{2} \pi\right)} [S_\nu^{\alpha\beta}(x) - T_\nu^{\alpha\beta}(x)],$$

where

$$\begin{aligned} S_\nu^{\alpha\beta}(x) &= \sum_{k=0}^{\infty} \frac{(x/2)^{2k+2\beta-2}}{k! \Gamma\left(\beta + \frac{\nu}{2} + k\right) \Gamma\left(\beta - \frac{\nu}{2} + k\right) \Gamma(\alpha - k)}, \\ T_\nu^{\alpha\beta}(x) &= \sum_{k=0}^{\infty} \frac{(x/2)^{2k+\nu}}{k! \Gamma\left(\alpha + \beta - \frac{\nu}{2} - 1 - k\right) \Gamma\left(2 - \beta + \frac{\nu}{2} + k\right) \Gamma(1 + \nu + k)}. \end{aligned}$$

In  $S_\nu^{\alpha\beta}(x)$  and  $T_\nu^{\alpha\beta}(x)$  and in all other series introduced below, terms are defined to be zero if their numerator is finite and in their denominator they contain a gamma function with argument equal to zero or a negative integer.

The expression for  $I_\nu^{\alpha\beta}(x)$  can also be obtained by the residue theorem from the Mellin-Barnes representation for the function  $G_{1,3}^{2,0}$  (see [4, p. 146]).

Equation (4) does not apply when  $2\beta - \nu$  is an even integer. The series expansion for  $\beta = j + 1 + \nu/2$ , where  $j$  is an integer, can be obtained from (4) by finding the limit of this equation as  $\beta$  approaches  $j + 1 + \nu/2$ . This series can be expressed

$$(5) \quad I_\nu^{\alpha, j+1+\nu/2}(x) = \frac{1}{2} \Gamma(\alpha) [U_\nu^{\alpha j}(x) + V_\nu^{\alpha j}(x)],$$

where

$$U_\nu^{\alpha j}(x) = (-1)^j \sum_{k=0}^{\infty} \frac{[-2\gamma - 2 \log_e(x/2) + P_k^{\alpha j}](x/2)^{2k+2j_++\nu}}{k!(k + |j|)!\Gamma(\alpha + j_- - k)\Gamma(1 + \nu + j_+ + k)},$$

$$V_\nu^{\alpha 0}(x) = 0,$$

$$V_\nu^{\alpha j}(x) = \sum_{k=0}^{|j|-1} \frac{(-1)^k (|j| - 1 - k)!(x/2)^{2k+2j_-+\nu}}{k!\Gamma(\alpha + j_+ - k)\Gamma(1 + \nu + k + j_-)}, \quad j \neq 0,$$

$$j_+ = \frac{j + |j|}{2}, \quad j_- = \frac{j - |j|}{2}.$$

$$P_k^{\alpha j} = Q(1 + k) + Q(1 + |j| + k) + Q(1 + j_+ + \nu + k) - Q(\alpha + j_- - k)$$

$$(6) \quad Q(z) = \psi(z) + \gamma = \sum_{i=0}^{\infty} \left[ \frac{1}{1+i} - \frac{1}{z+i} \right],$$

$$\psi(z) = \frac{d}{dz} [\log_e \Gamma(z)]$$

and  $\gamma$  is Euler's constant.

The relation [8]

$$(7) \quad \frac{Q(z)}{\Gamma(z)} = \frac{Q(1-z)}{\Gamma(z)} - \Gamma(1-z)\cos(z\pi)$$

was employed in obtaining  $V_\nu^{\alpha j}(x)$  and also is useful in evaluating  $U_\nu^{\alpha j}(x)$  when any of the  $Q(z)$  has an argument equal to zero or a negative integer.

The expression for  $Q(z)$  can often be simplified [8].

The power series (4) and (5) can often be used for calculating the  $I_\nu^{\alpha\beta}(x)$ . Other techniques, such as rational approximations and Chebyshev expansions, may also be convenient.

(b) *Recurrence Relations.* When the  $G$  function in (3) is expressed as a contour integral [9], the  $I_\nu^{\alpha\beta}(x)$  can be written

$$(8) \quad I_\nu^{\alpha\beta}(x) = \frac{\Gamma(\alpha)}{4\pi i} \int_L \frac{\Gamma(\beta - 1 - s)\Gamma\left(\frac{\nu}{2} - s\right)}{\Gamma(\alpha + \beta - 1 - s)\Gamma\left(1 + \frac{\nu}{2} + s\right)} \left(\frac{x}{2}\right)^{2s} ds.$$

The integration contour  $L$  is the first path described in [9], which also lists the conditions under which (8) is valid.

From (8),

$$(9) \quad \frac{x}{2\nu} [I_{\nu-1}^{\alpha\beta}(x) + I_{\nu+1}^{\alpha\beta}(x)] = \frac{\Gamma(\alpha)}{4\pi i} \int_L \frac{\Gamma(\beta - 1 - s)\Gamma\left(\frac{\nu-1}{2} - s\right)}{\Gamma(\alpha + \beta - 1 - s)\Gamma\left(\frac{\nu+3}{2} + s\right)} \left(\frac{x}{2}\right)^{2s+1} ds.$$

The recurrence relation

$$(10) \quad \frac{x}{2\nu} [I_{\nu-1}^{\alpha\beta}(x) + I_{\nu+1}^{\alpha\beta}(x)] = I_{\nu}^{\alpha,\beta+1/2}(x)$$

then is obtained from (9) by the change of variable  $r = s + \frac{1}{2}$ .

Similar calculations show that

$$(11) \quad (2\beta + 2\alpha - \nu - 4)I_{\nu}^{\alpha\beta}(x) = x[I_{\nu+1}^{\alpha,\beta-3/2}(x) - I_{\nu+1}^{\alpha,\beta-1/2}(x)] + (2\beta - \nu - 4)I_{\nu}^{\alpha,\beta-1}(x),$$

$$(12) \quad I_{\nu}^{\alpha+1,\beta-1}(x) + I_{\nu}^{\alpha,\beta}(x) = I_{\nu}^{\alpha,\beta-1}(x),$$

and that

$$(13) \quad x^{-\nu} \frac{d}{dx} [x^{\nu} I_{\nu}^{\alpha\beta}(x)] = I_{\nu-1}^{\alpha,\beta-1/2}(x),$$

$$(14) \quad x^{\nu} \frac{d}{dx} [x^{-\nu} I_{\nu}^{\alpha\beta}(x)] = -I_{\nu+1}^{\alpha,\beta-1/2}(x).$$

Integral relations corresponding to (13) and (14) are

$$(15) \quad I_{\nu}^{\alpha\beta}(x) = x^{-\nu} \int_0^x y^{\nu} I_{\nu-1}^{\alpha,\beta-1/2}(y) dy,$$

$$(16) \quad I_{\nu}^{\alpha\beta}(x) = -x^{\nu} \int_x^{\infty} y^{-\nu} I_{\nu+1}^{\alpha,\beta-1/2}(y) dy.$$

After the  $I_{\nu}^{\alpha\beta}(x)$  have been expressed in terms of known functions or evaluated by the power series or by some other technique for  $\alpha_0 \leq \alpha < \alpha_0 + 1$  and  $\beta_0 \leq \beta < \beta_0 + \frac{1}{2}$  for the necessary values of  $\nu$ , the recurrence relations (10), (11), and (12) can be used to find the  $I_{\nu}^{\alpha\beta}(x)$  for larger values of  $\alpha$  and  $\beta$ . Relations (13)–(16) are convenient when the  $I_{\nu}^{\alpha\beta}(x)$  are expressed in terms of functions which can be easily integrated or differentiated.

When  $\nu = 0$ , (10) cannot be used. Equation (11) then can be employed.

When the recurrence relations are used successively many times, the possibility of round-off errors must be considered. This effect can be especially noticeable if (10) is used when  $x/(2\nu)$  is large with respect to 1.

The special cases  $I_{\nu}^{1/2,1}(x)$  and  $I_{\nu}^{\alpha,\nu/2+1-\alpha}(x)$ , which, as was mentioned in Section I, are often found in tables of integrals of Bessel functions, also follow from (3).

Thus, with the  $G$ -function representation [10] of the product  $J_{\nu}(x)Y_{\nu}(x)$ , the expression

$$(17) \quad I_{\nu}^{1/2,1}(x) = -\frac{\pi}{2} J_{\nu/2}(x/2)Y_{\nu/2}(x/2)$$

can be obtained. Also, substitution of (17) in (13) gives

$$(18) \quad I_{\nu}^{1/2,1/2}(x) = -\frac{\pi}{4} \left[ J_{(\nu-1)/2}\left(\frac{x}{2}\right)Y_{(\nu+1)/2}\left(\frac{x}{2}\right) + J_{(\nu+1)/2}\left(\frac{x}{2}\right)Y_{(\nu-1)/2}\left(\frac{x}{2}\right) \right].$$

Another useful equation

$$I_0^{1/2,3/2}(x) = \int_x^{\infty} \frac{\sin y}{y} dy$$

can be obtained either from (16) and (17) or by letting  $\alpha = 1/2$ ,  $\beta = 3/2$ , and  $\nu = 0$  in the equation expressing  $I_{\nu}^{\alpha\beta}(x)$  in terms of the hypergeometric functions  ${}_1F_2$  [4, p. 223, Eq. 25].

From (8) and the  $G$ -function representation [9], [5] of  $J_{\nu}(x)$ ,

$$I_{\nu}^{\alpha, \nu/2 + 1 - \alpha}(x) = \frac{1}{2} \Gamma(\alpha) (x/2)^{-\alpha} J_{\nu - \alpha}(x).$$

This result is one of Sonine's formulas [11].

**III. The Expression for  $\Phi(a)$ .** With (17), (18), and the relation [12]

$$Y_{\nu-1}(x)J_{\nu}(x) - Y_{\nu}(x)J_{\nu-1}(x) = 2/(\pi x),$$

the quantity  $\Phi(a)$  employed in Fedorova's method of computing the diameter distribution for assemblies of cylinders [1], [2] can be written

$$\begin{aligned} \Phi(a) &= a I_1^{1/2, 1/2}(a) - I_2^{1/2, 1}(a) \\ (19) \quad &= \frac{\pi}{2} Y_1(a/2) [J_1(a/2) - a J_0(a/2)] - 1. \end{aligned}$$

Equation (19) is convenient for calculating diameter distributions by Fedorova's method.

Substitution of the asymptotic expansions for the Bessel functions in (19) gives the asymptotic approximation previously obtained [2] by Fedorova by a different method.

**Acknowledgements.** I would like to thank I. S. Fedorova and B. A. Fedorov for several interesting and stimulating discussions about calculating diameter distributions from scattering data and about the integrals which occur in  $\Phi(a)$ . I also wish to express my appreciation to James Casteel and Brian DeFacio for their help in the preparation of the manuscript.

Physics Department  
University of Missouri  
Columbia, Missouri 65201

1. I. S. FEDOROVA, *Dokl. Akad. Nauk SSSR*, v. 223, 1975, p. 1007.
2. I. S. FEDOROVA & J. COLLOID, *Interface Sci.*, v. 59, 1977, pp. 100-101.
3. W. MAGNUS & F. OBERHETTINGER, *Formulas and Theorems for the Functions of Mathematical Physics*, Chelsea, New York, 1954, p. 32.
4. Y. L. Luke, *The Special Functions and Their Approximations*, Vol. I, Sect. 5.2, Academic Press, New York and London, 1969.
5. Ref. 4, p. 226, Eq. (7).
6. Ref. 4, p. 170, Eq. (6).
7. Ref. 4, p. 145, Eq. (7).
8. Ref. 4, pp. 11-13.
9. Ref. 4, pp. 143-144.
10. Ref. 4, p. 229, Eq. (30).
11. G. N. WATSON, *A Treatise on the Theory of Bessel Functions*, 2nd ed., Cambridge Univ. Press, Cambridge, London & New York, 1966, p. 417, Eq. (5).
12. Ref. 11, p. 77, Eq. (12).