

A Collocation-Galerkin Method for Poisson's Equation on Rectangular Regions

By Julio César Díaz

Abstract. A collocation-Galerkin method is defined for Poisson's equation on the unit square, using tensor products of continuous piecewise polynomials. Optimal L^2 and H_0^1 orders of convergence are established. This procedure requires fewer quadratures than the corresponding Galerkin procedure.

Introduction. The collocation-Galerkin method was first introduced by Díaz [2], [3] for the two-point boundary value problems and optimal L^2 -rates of convergence were established for a particular choice of the collocation points; namely, the affine images of the roots of a Jacobi polynomial. In [6] Wheeler derived optimal L^p -estimates and applied this method to a one space dimensional parabolic problem. In [4] Dunn and Wheeler analyzed some collocation- H^{-1} -Galerkin methods and established optimal L^p -estimates for any choice of the collocation points. Archer and Díaz [1] have applied similar ideas to a one-dimensional first order hyperbolic problem and derived optimal L^2 -estimates. Here, a collocation-Galerkin method is defined for Poisson's equation on the unit square, using tensor products of continuous piecewise polynomials and the collocation points are based on the roots of a Jacobi polynomial. Optimal L^2 - and H_0^1 -estimates are established. On the basis of computational complexity the collocation-Galerkin method is intermediate between the Galerkin method and the collocation method. It has an advantage over the Galerkin procedure for the same space in that the integrals involve the product of the approximate solution and a piecewise linear function, thus the integrals are simpler and, of course, there are fewer of them. Also, the continuity conditions on the approximate solution are weaker than those required of the collocation approximation defined by Prenter and Russell [5].

In the following section, the collocation-Galerkin method is defined and existence and uniqueness are shown using some semidiscrete bilinear forms. In the last section, the error analysis is presented. The analysis consists of reducing the problem to some one-dimensional problems for which the results of [3] can be applied.

The Collocation-Galerkin Method. Consider the boundary value problem

$$(1) \quad \begin{aligned} -\Delta u &= f, & \text{on } \Omega, \\ u &= 0, & \text{on } \partial\Omega, \end{aligned}$$

where $\Omega = (0, 1) \times (0, 1)$. We shall assume that there exists a unique u and that it is sufficiently smooth.

Received March 8, 1976; revised March 24, 1978.

AMS (MOS) subject classifications (1970). Primary 65N35, 65N30.

© 1979 American Mathematical Society
0025-5718/79/0000-0005/\$03.00

For a partition $\delta = \{x_i\}_{i=1}^{N+1}$ of the unit interval $I = [0, 1]$ satisfying,

$$0 = x_0 < x_1 < \cdots < x_{N+1} = 1,$$

let

$$y_l = x_l, \quad l = 0, \dots, N+1,$$

$$I_i = [x_i, x_{i+1}], \quad h_i = x_{i+1} - x_i, \quad i = 0, \dots, N,$$

$$h = \max_{0 \leq i \leq N} h_i.$$

If $r \geq 1$ is an integer, let $P_r(E)$ denote the class of polynomials of degree at most r on the set E , and define

$$M'_0 \equiv M'_0(\delta) = \{V \in C(I) \mid V \in P_r(I_i), i = 0, \dots, N, V(0) = V(1) = 0\}.$$

Let $\varphi_1, \dots, \varphi_{r-1}$ and positive $\omega_1, \dots, \omega_{r-1}$ be the unique choices such that

$$(2) \quad \int_0^1 x(1-x)p(x) dx = \sum_{j=1}^{r-1} \omega_j p(\varphi_j), \quad \forall p \in P_{2r-3}(0, 1).$$

That is, $\varphi_1, \dots, \varphi_{r-1}$ are the roots of the Jacobi polynomial J_{r-1} of degree $r-1$ on $[0, 1]$ with respect to the weight function $x(1-x)$. The collocation points are tensor products of affine transformations of the roots of J_{r-1} onto each subinterval. More precisely, let $x_{ij} = x_i + h_i \varphi_j$, $y_{lk} = y_l + h_l \varphi_k$, $i, l = 0, \dots, N$; $j, k = 1, \dots, r-1$.

The collocation-Galerkin method for the approximate solution of (1) consists in finding $U \in M = M'_0 \otimes M'_0$ satisfying

$$(3.i) \quad \Delta U(x_{ij}, x_{lk}) + f(x_{ij}, x_{lk}) = 0, \quad \begin{array}{l} i, l = 0, \dots, N, \\ j, k = 1, \dots, r-1, \end{array}$$

$$(3.ii) \quad \int_I \frac{\partial^2}{\partial x^2} (U(x_{ij}, \xi)) V(\xi) d\xi - \int_I \frac{\partial}{\partial y} (U(x_{ij}, \xi)) V'(\xi) d\xi \\ + \int_I f(x_{ij}, \xi) V(\xi) d\xi = 0, \quad \begin{array}{l} i = 0, \dots, N, \\ j = 1, \dots, r-1, \end{array} \quad \forall V \in M^1_0,$$

$$(3.iii) \quad - \int_I \frac{\partial}{\partial x} (U(\xi, y_{lk})) V'(\xi) d\xi + \int_I \frac{\partial^2}{\partial y^2} (U(\xi, y_{lk})) V(\xi) d\xi \\ + \int_I f(\xi, y_{lk}) V(\xi) d\xi = 0, \quad \begin{array}{l} l = 0, \dots, N, \\ k = 1, \dots, r-1, \end{array} \quad \forall V \in M^1_0,$$

$$(3.iv) \quad (\nabla U, \nabla V) = (f, V), \quad \forall V \in M^1_0 \otimes M^1_0,$$

where (\cdot, \cdot) denotes the L^2 -inner product over Ω .

In order to demonstrate the existence and uniqueness of the collocation-Galerkin approximation U , bilinear forms $D(\cdot, \cdot)$, \mathcal{D} and \mathcal{I} are introduced. Let

$$Z^r_0 = \{V \in M^r_0 \mid V(x_i) = 0, i = 1, \dots, N\}.$$

Then

$$(4) \quad M^r_0 = M^1_0 \oplus Z^r_0.$$

For $v \in Z_0^r$ and a function g defined on I , let

$$D_i(g, V) = \sum_{j=1}^{r-1} h_i \omega_j \frac{g(x_{ij})V(x_{ij})}{\varphi_j(1-\varphi_j)},$$

and set

$$D(g, V) = \sum_{i=0}^N D_i(g, V).$$

Two semidiscrete bilinear forms $\mathcal{D}(\cdot, \cdot)$ and $\mathcal{I}(\cdot, \cdot)$ are defined as follows; for $V \in M_0^r$, there are unique $V_1 \in M_0^1$ and $V_2 \in Z_0^r$ such that $V = V_1 + V_2$, and for g defined on each I_i and $g \in L^2(I)$, let

$$\mathcal{I}(g, V) = D(g, V_2) + \int_0^1 g(x)V_1(x) dx.$$

For $g \in H^1(I)$ such that g'' is defined on each I_i ,

$$\mathcal{D}(g, V) = -D(g'', V_2) + \int_0^1 g'(x)V_1'(x) dx.$$

If $g, V \in M_0^r$, relationship (2) and integration by parts imply that

$$(5) \quad \mathcal{D}(g, V) = \int_0^1 g'(x)V'(x) dx.$$

Notice that if $p \in \mathbf{P}_{2r-2}(0, 1)$, then $p(x) = p_{2r-2} \cdot J_{r-1}^2(x) + q(x)$, where p_{2r-2} is the coefficient of x^{2r-2} in $p(x)$, and $q \in \mathbf{P}_{2r-3}(0, 1)$. Then it is straightforward to show that

$$\begin{aligned} \sum_{j=1}^{r-1} \omega_j p(\varphi_j) &= \int_0^1 x(1-x)p(x) dx \\ &+ \frac{1}{(2r)!} \frac{d^{2r}}{dx^{2r}} [x(1-x)p(x)] \int_0^1 x(1-x)J_{r-1}^2(x) dx. \end{aligned}$$

From this relationship it follows that if $g \in M_0^r$, then

$$(6) \quad \mathcal{I}(g, g) \geq \int_0^1 g^2(x) dx.$$

When considering functions of more than one variable a subscript, x or y , will be used to denote the variable to which the bilinear form is being applied. Bilinear forms on the two variables are formed by taking tensor products of \mathcal{D} and \mathcal{I} . From (4), it follows that

$$M = (Z_0^r \otimes Z_0^r) \oplus (Z_0^r \otimes M_0^1) \oplus (M_0^1 \otimes Z_0^r) \oplus (M_0^1 \otimes M_0^1) \equiv N_1 \oplus N_2 \oplus N_3 \oplus N_4.$$

Thus, for $V \in M$ there are unique $V_m \in N_m$, $m = 1, \dots, 4$, such that

$$V = V_1 + V_2 + V_3 + V_4.$$

And for g an L^2 -function defined on all of Ω , let

$$\begin{aligned} I_x \otimes I_y(g, V) &= \sum_{i,l=0}^N \left\{ \sum_{k=1}^{r-1} \sum_{j=1}^{r-1} h_i h_l \omega_k \omega_j \frac{g(x_{ij}, y_{lk})V_1(x_{ij}, y_{lk})}{\varphi_k(1-\varphi_k)\varphi_j(1-\varphi_j)} \right\} \\ &+ \int_I D(g(\cdot, y), V_2(\cdot, y)) dy + \int_I D(g(x, \cdot), V_3(x, \cdot)) dx + (g, V_4). \end{aligned}$$

The other bilinear forms $\mathcal{D}_x \otimes I_y$ and $I_x \otimes \mathcal{D}_y$ are similarly defined. Using these bilinear forms, a related variational approximation is defined. Let $W \in M$ be the solution to

$$(7) \quad \mathcal{D}_x \otimes I_y(W, V) + I_x \otimes \mathcal{D}_y(W, V) = I_x \otimes I_y(f, V), \quad \forall V \in M.$$

It follows from the definition of the bilinear forms that $U \in M$, the solution to (3), satisfies equations (7). Thus, in order to show existence and uniqueness of $U \in M$ satisfying (3) it is enough to show that if $f = 0$ then $W = 0$. Relationship (5) implies that W satisfies

$$\int_0^1 I_y(W_x, V_x) dx + \int_0^1 I_x(W_y, V_y) dy = I_x \otimes I_y(f, V), \quad \forall V \in M.$$

Choosing $V = W$ and using (6), it follows that

$$\int_0^1 \int_0^1 W_x^2 dx dy + \int_0^1 \int_0^1 W_y^2 dx dy \leq I_x \otimes I_y(f, W),$$

thus

$$(\nabla W, \nabla W) \leq I_x \otimes I_y(f, W);$$

hence if $f = 0$, $W = 0$.

Error Analysis. As before let u denote the solution to the boundary value problem (1) and $U \in M$ the collocation-Galerkin solution to (7). In this section, estimates for $u - U$ are derived. Those estimates are given in the following theorem.

THEOREM. *Let u be the solution to (1) and $U \in M$ the collocation-Galerkin approximation to u defined by (3). Then, if u is sufficiently smooth, there exists a constant C independent of h and u such that*

$$\|u - U\|_{L^2} + h \|u - U\|_{H_0^1} \leq Ch^s \|u\|_{H^s}, \quad 1 \leq s \leq r + 1.$$

Before proving the theorem, some basic estimates are derived. For $e = u - U$, let $\psi \in H^2(\Omega)$ satisfy

$$\begin{aligned} -\Delta \psi &= e, & \text{on } \Omega, \\ \psi &= 0, & \text{on } \partial\Omega, \end{aligned}$$

then

$$\|e\|_{L^2}^2 = (e, e) = -(e, \Delta \psi) = (\nabla e, \nabla \psi) = (\nabla e, \nabla(\psi - \chi)), \quad \chi \in M_0^1 \otimes M_0^1,$$

using (3.iv). Thus, by Cauchy-Schwarz inequality

$$\|e\|_{L^2}^2 \leq \|e\|_{H_0^1} \|\psi - \chi\|_{H_0^1}, \quad \chi \in M_0^1 \otimes M_0^1,$$

the approximation properties of the space $M_0^1 \otimes M_0^1$ imply that

$$\inf_{\chi \in M_0^1 \otimes M_0^1} \|\psi - \chi\|_{H_0^1} \leq Ch \|\psi\|_{H^2},$$

also by elliptic regularity

$$\|\psi\|_{H^2} \leq C \|e\|_{L^2},$$

thus it follows from these inequalities that

$$(9) \quad \|e\|_{L^2} \leq Ch \|e\|_{H_0^1}.$$

Therefore, it is sufficient to derive an H_0^1 -norm estimate of the error. In the derivation of this estimate the following lemma, proved in [3], will play an important role.

LEMMA 1. $\Phi \in H^{r-1}(0, 1)$, let $g \in H^2(0, 1) \cap H_0^1(0, 1)$ satisfy $g'' = \Phi$. Also, let $G \in M_0^r$ be the collocation-Galerkin approximation to g satisfying

$$G''(x_{ij}) = \Phi(x_{ij}), \quad i = 0, \dots, N; j = 1, \dots, r-1,$$

and

$$\int_0^1 G'V' dx + \int_0^1 \Phi V dx = 0, \quad \forall V \in M_0^1;$$

then, there exists a constant C independent of h such that

$$\int_0^1 |\epsilon|^2 dx + h^2 \int_0^1 |\epsilon'|^2 dx \leq Ch^{2s} \int_0^1 \left| \frac{d^s}{dx^s} g \right|^2 dx, \quad 1 \leq s \leq r+1,$$

where $\epsilon = g - G$.

Also, a space M_x and an element $q \in M_x$, a projection of u into M_x , are introduced. Then error estimates for $u - q$ and $q - U$ will be obtained in Lemmas 2 and 3, respectively. The desired estimates will follow from the triangle inequality.

The space M_x is defined by

$$M_x = \left\{ \sum_s \alpha_s(y) V_s(x), \{V_s\} \text{ basis for } M_0^r \text{ and } \alpha_s \in H_0^1(I) \cap H^2(I) \right\}.$$

The function $q \in M_x$ satisfies the following equations

$$(10.i) \quad (u - q)_{xx}(x_{ij}, y) + (u - q)_{yy}(x_{ij}, y) = 0, \\ i = 0, \dots, N; j = 1, \dots, r-1, y \in (0, 1),$$

$$(10.ii) \quad - \int_0^1 (u - q)_{xx}(\xi, y) V'(\xi) d\xi + \int_0^1 (u - q)_{yy}(\xi, y) V(\xi) d\xi = 0,$$

$$V \in M_0^1, y \in (0, 1).$$

This system of equations leads to a system of second order two-point boundary value problems, the collocation-Galerkin solution of which corresponds to the solution U of (3). Moreover, U satisfies

$$(11) \quad \mathcal{D}_x \otimes I_y(U - q, V) + I_x \otimes \mathcal{D}_y(U - q, V) = 0, \quad \forall V \in M.$$

This fact will play an important role later. Using the bilinear forms \mathcal{D} and I , a weak formulation of (10) can be written as

$$(12) \quad \int_0^1 \mathcal{D}_x((u - q), v) dy + \int_0^1 I_x((u - q)_y, v_y) dy = 0, \quad \forall v \in M_x.$$

The choice $v = q$ and the use of relationships (5) and (6) in (12) give

$$\int_0^1 \int_0^1 q_x^2 dx dy + \int_0^1 \int_0^1 q_y^2 dx dy \leq \int_0^1 \mathcal{D}_x(u, q) dy + \int_0^1 I_x(u_y, q_y) dy,$$

from which existence and uniqueness of q follow. Notice that if for any $\alpha \geq 0$, $\partial^\alpha u / \partial y^\alpha \in H = H_0^1(\Omega) \cap H^2(\Omega)$, then $\partial^\alpha q / \partial y^\alpha \in M_x$. The estimates of $u - q$ are given in the following lemma.

LEMMA 2. Let u be the solution to (1) and $q \in M_x$ the solution to (12). Then, if u is sufficiently smooth, there exists a constant C independent of h such that, for $\alpha \geq 0$

$$(13) \quad \left\| \frac{\partial^\alpha}{\partial y^\alpha} (u - q) \right\|_{L^2} + h \left\| \frac{\partial^{\alpha+1}}{\partial x \partial y^\alpha} (u - q) \right\|_{L^2} \\ \leq Ch^s \left\{ \left\| \frac{\partial^{s+\alpha} u}{\partial x^s \partial y^\alpha} \right\|_{L^2} + \left\| \frac{\partial^{s+\alpha} u}{\partial x^{s-1} \partial y^{\alpha+1}} \right\|_{L^2} \right\}, \quad 1 \leq s \leq r + 1.$$

Proof. Let $\eta = u - q$. It is straightforward to show that equations (10.ii) imply that

$$(14) \quad \|\eta\|_{L^2} \leq Ch \|\nabla \eta_x\|_{L^2}.$$

Thus, it suffices to estimate $\|\eta_x\|_{L^2}$. For this a map $w \in M_x$, for which the results of Lemma 1 can be applied directly, is introduced. Estimates of the difference of w and q will be derived and estimates for $u - q$ will follow from the triangle inequality. Define the map w by

$$(u - w)_{xx}(x_{ij}, y) = 0, \quad i = 0, \dots, N; j = 1, \dots, r - 1; y \in (0, 1),$$

and

$$\int_0^1 (u - w)_x(\xi, y) V_x(\xi) d\xi = 0, \quad V \in M_0^1, y \in (0, 1).$$

Equivalently, w is the solution of

$$(15) \quad \int_0^1 \mathcal{D}_x(u - w, v) dy = 0, \quad v \in M_x.$$

Notice that, as for q , if for any $\alpha \geq 0$, $\partial^\alpha u / \partial y^\alpha \in H$, then $\partial^\alpha w / \partial y^\alpha \in M_x$. From Lemma 1 it follows that there exists a constant C independent of h satisfying for $\alpha \geq 0$,

$$(16) \quad \left\| \frac{\partial^\alpha (u - w)}{\partial y^\alpha} \right\|_{L^2} + h \left\| \frac{\partial^{\alpha+1}}{\partial x \partial y^\alpha} (u - w) \right\|_{L^2} \leq Ch^s \left\| \frac{\partial^{s+\alpha}}{\partial x^s \partial y^\alpha} u \right\|_{L^2}, \quad 1 \leq s \leq r + 1.$$

Estimates of the difference between w and q are now derived. Let $\epsilon = q - w$. Equations (12) and (15) imply

$$\int_0^1 \mathcal{D}_x(\epsilon, v) dy + \int_0^1 I_x((\epsilon)_y, v_y) dy = \int_0^1 I_x((u - w)_y, v_y) dy, \quad \forall v \in M_x.$$

Since ϵ and $v \in M_x$, then by (5)

$$\int_0^1 \mathcal{D}_x(\epsilon, v) dy = \int_0^1 \int_0^1 \epsilon_x v_x dx dy = (\epsilon_x, V_x),$$

and for $v = \epsilon$ by (6)

$$\int_0^1 I_x(\epsilon_y, \epsilon_y) dy \geq \int_0^1 \int_0^1 \epsilon_y^2 dx dy = (\epsilon_y, \epsilon_y).$$

In [3], it was shown that if $g \in H^s(I)$, $1 \leq s \leq r + 1$ and $V \in M_0^r$, then

$$(17) \quad \left| I(g, V) - \int_0^1 g V dx \right| \leq Ch^s \left\{ \int_0^1 \left(\frac{d^s g}{dx^s} \right)^2 dx \right\}^{1/2} \left\{ \int_0^1 V^2 dx \right\}^{1/2}.$$

Thus

$$\begin{aligned} \|\epsilon_x\|_{L^2}^2 + \|\epsilon_y\|_{L^2}^2 &\leq ((u-w)_y, \epsilon_y) + \int_0^1 \left\{ I_x((u-w)_y, \epsilon_y) - \int_0^1 (u-w)_y \epsilon_y dx \right\} dy \\ &\leq ((u-w)_y, \epsilon_y) + \int_0^1 \left\{ Ch \left(\int_0^1 (u-w)_{xy}^2 dx \right)^{\frac{1}{2}} \left(\int_0^1 \epsilon_y^2 dx \right)^{\frac{1}{2}} \right\} dy \\ &\leq \frac{1}{4\epsilon} \|(u-w)_y\|_{L^2}^2 + 2\xi \|\epsilon_y\|_{L^2}^2 + \frac{C^2 h^2}{4\xi} \|(u-w)_{xy}\|_{L^2}^2, \end{aligned}$$

where the inequality $2ab \leq a^2 + b^2$ has been used. Take $\xi = \frac{1}{2}$, then

$$(18) \quad \|(w-q)_x\|_{L^2}^2 \leq C \{ \|(u-w)_y\|_{L^2}^2 + h^2 \|(u-w)_{xy}\|_{L^2}^2 \}.$$

Estimates (14), (16) and (18) and the triangle inequality complete the proof of the lemma.

It remains to obtain estimates of the difference between U and q . In order to obtain these estimates, an auxiliary function $W \in M$ is introduced which satisfies

$$(19) \quad (W-q)_{yy}(x, y_{lk}) = 0, \quad l = 0, \dots, N; k = 1, \dots, r-1, x \in (0, 1),$$

$$- \int_0^1 (W-q)_y(x, \xi) V_y(\xi) d\xi = 0, \quad V \in M_0^1, x \in (0, 1),$$

or equivalently

$$\mathcal{D}_y(W-q, V) = 0, \quad \forall V \in M_0^r, x \in (0, 1).$$

In particular

$$(20) \quad I_x \otimes \mathcal{D}_y(W-q, V) = 0, \quad \forall V \in M.$$

Equations (19) and Lemma 1 imply that for $\alpha = 0, 1$

$$\left\| \frac{\partial^\alpha}{\partial x^\alpha} (q-W) \right\|_{L^2} + h \left\| \frac{\partial^{\alpha+1}}{\partial x^\alpha \partial y} (q-W) \right\|_{L^2} \leq Ch^s \left\| \frac{\partial^{s+\alpha}}{\partial x^\alpha \partial y^s} q \right\|_{L^2}.$$

Setting $q = q - u + u$ and using estimate (13), it follows that

$$(21) \quad \|q-W\|_{L^2} + h \{ \|(q-W)_x\|_{L^2} + \|(q-W)_y\|_{L^2} \} + h^2 \|(q-W)_{xy}\|_{L^2} \leq Ch^s \|u\|_{H^s}.$$

In the following lemma, estimates of the difference between U and W are given.

LEMMA 3. Let $U \in M$ be the solution to (3), $q \in M_x$ the solution to (12) and $W \in M$ the solution to (19). Then there exists a constant C independent of h such that

$$(22) \quad \|\nabla(U-W)\|_{L^2}^2 \leq C \{ \|(q-W)_x\|_{L^2}^2 + h^2 \|(q-W)_{xy}\|_{L^2}^2 \}.$$

Proof. Let $E = U - W$. Using (11) and (20); it follows that

$$\mathcal{D}_x \otimes I_y(E, V) + I_x \otimes \mathcal{D}_y(E, V) = \mathcal{D}_x \otimes I_y(q-W, V), \quad \forall V \in M,$$

or equivalently using (5)

$$\int_0^1 I_y(E_x, V_x) dx + \int_0^1 I_x(E_y, V_y) dy = \int_0^1 I_y((q-W)_x, V_x) dx, \quad \forall V \in M,$$

with $V = E$ and using (6), it follows that

$$\|E_x\|_{L^2}^2 \leq \int_0^1 I_y(E_x, E_x) dx,$$

and

$$\|E_y\|_{L^2}^2 \leq \int_0^1 I_x(E_y, E_y) dy.$$

Hence

$$\begin{aligned} \|E_x\|_{L^2}^2 + \|E_y\|_{L^2}^2 &\leq \int_0^1 I_y((q - W)_x, E_x) dy \\ &\leq ((q - W)_x, E_x) + \int_0^1 \left\{ I_y((q - W)_x, E_x) - \int_0^1 (q - W)_x E_x dx \right\} dy \\ &\leq ((q - W)_x, E_x) + \int_0^1 \left\{ Ch \left(\int_0^1 (q - W)_{xy}^2 dy \right)^{1/2} \left(\int_0^1 E_x^2 dy \right)^{1/2} \right\} dx \\ &\leq \frac{1}{4\epsilon} \|(q - W)_x\|_{L^2}^2 + 2\epsilon \|E_x\|_{L^2}^2 + \frac{C^2 h^2}{4\epsilon} \|(q - W)_{xy}\|_{L^2}^2, \end{aligned}$$

where the inequality $2ab \leq a^2 + b^2$ has been used. With $\epsilon = 1/4$ and $\epsilon = 1/2$, estimate (22) follows completing the proof of the lemma.

The Theorem follows from estimates (9), (13), (21) and (22) and the triangle inequality.

Remarks. 1. Although the same partition has been used for both sides, the argument holds for different partitions.

2. The Galerkin procedure for the same space requires $\{(N + 1)(r - 1) + N\}^2$ two-dimensional quadratures while the procedure described by (3) needs only N^2 two-dimensional quadratures plus $2(N + 1)(r - 1)$ one-dimensional quadratures.

3. In general the solution can only be asserted to lie in $H^{3-\epsilon}(\Omega)$, for any $\epsilon > 0$.

Department of Mathematics
University of Kentucky
Lexington, Kentucky 40506

1. D. ARCHER & J. C. DÍAZ, "A collocation-Galerkin method for a first order hyperbolic equation." (To appear.)

2. JULIO CÉSAR DÍAZ VELASCO, *A Hybrid Collocation-Galerkin Method for the Two Point Boundary Value Problem Using Continuous Piecewise Polynomials Spaces*, Ph.D. Thesis, Rice Univ., Houston, Texas, 1974.

3. J. C. DÍAZ, "A collocation-Galerkin method for the two point boundary value problem using continuous piecewise polynomial spaces," *SIAM J. Numer. Anal.*, v. 14, 1977, pp. 844–858.

4. R. J. DUNN & M. F. WHEELER, "Some collocation-Galerkin methods for two-point boundary-value problems," *SIAM J. Numer. Anal.*, v. 13, 1976, pp. 720–733.

5. P. M. PRENTER & R. D. RUSSELL, "Orthogonal collocation for elliptic partial differential equations," *SIAM J. Numer. Anal.*, v. 13, 1976, pp. 923–939.

6. M. F. WHEELER, "A C^0 collocation-Galerkin method for two-point boundary-value problems and one space dimensional parabolic problems," *SIAM J. Numer. Anal.*, v. 14, 1977, pp. 71–90.