

A Lower Bound for the Norm of the Theta Operator*

By L. Alayne Parson

Abstract. The Poincaré theta operator maps the space of holomorphic functions with period one onto the space of cusp forms for a finitely generated Fuchsian group. It is easy to show that the norm of the operator does not exceed one. In the case of the classical modular group and weight six, it is now shown that the norm is bounded below by .927.

1. Definition of the Theta Operator. Let Γ be a finitely generated Fuchsian group acting on the upper half plane \mathfrak{H} and containing the translation $Sz = z + 1$. Let $A_q(\Gamma)$ be the space of cusp forms of weight q . $A_q(\Gamma)$ is then a Banach space with norm

$$\|f\|_{A_q(\Gamma)} = \iint_{\mathfrak{H} = \mathfrak{H}/\Gamma} |f(z)| y^{q-2} dx dy.$$

When $\Gamma = \Gamma_\infty = \langle S \rangle$, denote $A_q(\Gamma_\infty)$ by A_q . The Poincaré θ -operator, θ_q , which is defined by

$$\theta_q(f) = \sum_{A \in \Gamma/\Gamma_\infty} f(Az) A'^q(z),$$

is a surjective continuous linear operator from A_q to $A_q(\Gamma)$. That $\|\theta_q\| \leq 1$ is trivial. (See, for instance, Kra [1, Chapter 3].) Although many people suspect that $\|\theta_q\| < 1$ for various choices of q and Γ , this is not known, one way or another, in even a single case. Also, the standard lower bound on the norm is $\|\theta_q\| \geq (q-1)/(2q-1)$ (see Kra [1, p. 91]).

2. The Modular Group and Weight Six. The purpose of this note is to improve the lower bound on the theta operator norm for the modular group $\Gamma(1)$ and weight $q = 6$. In this case it is shown that $\|\theta_6\| \geq .927$, which is a marked improvement on the standard lower bound of $5/11$. $A_6(\Gamma(1))$ is the one-dimensional space of modular cusp forms which is spanned by the famous discriminant function

$$\Delta(z) = e^{2\pi iz} \prod_{n=1}^{\infty} (1 - e^{2\pi inz})^{24}.$$

The Fourier expansion at ∞ is $\Delta(z) = \sum_{n=1}^{\infty} \tau(n) e^{2\pi inz}$, where $\tau(n)$ is the celebrated Ramanujan tau function.

To improve the lower bound on $\|\theta_6\|$ it is necessary to evaluate $\|\theta_6(f)\|_{A_6(\Gamma(1))} / \|f\|_{A_6}$ for various choices of f for which both $\|\theta_6(f)\|_{A_6(\Gamma(1))}$ and $\|f\|_{A_6}$ are easy to evaluate accurately. Specifically, use is made of the following result.

Received July 14, 1982.

1980 *Mathematics Subject Classification*. Primary 10D12, 30F35.

*Research partially supported by NSF grant MCS-8202025.

©1983 American Mathematical Society
 0025-5718/83 \$1.00 + \$.25 per page

THEOREM. Set $p_k(z) = \sum_{n=0}^k a_n e^{2\pi i n z}$ and

$$f_k(z) = e^{2\pi i z} p_k^2(z) = \sum_{n=1}^{2k+1} b_n e^{2\pi i n z}.$$

Then

$$\|f_k\|_{A_6} = \frac{4!}{(2\pi)^5} \sum_{n=0}^k \frac{|a_n|^2}{(1+2n)^5}$$

and

$$\|\theta_6(f_k)\|_{A_6(\Gamma(1))} = \frac{10! \|\Delta\|_{A_6(\Gamma(1))}}{(4\pi)^{11} (\Delta, \Delta)} \left| \sum_{n=1}^{2k+1} \frac{b_n \tau(n)}{n^{11}} \right|,$$

where $(\Delta, \Delta) = \int_{\mathfrak{H}/\Gamma(1)} |\Delta(z)|^2 y^{10} dx dy$ is the Petersson scalar product of $\Delta(z)$ with itself.

Proof. Since

$$\|f_k\|_{A_6} = \int_0^\infty \int_0^1 |f_k(z)| y^4 dx dy = \int_0^\infty \int_0^1 e^{-2\pi y} |p_k(z)|^2 y^4 dx dy,$$

an application of Parseval's formula gives that

$$\|f_k\|_{A_6} = \int_0^\infty e^{-2\pi y} y^4 \sum_{n=0}^k |a_n|^2 e^{-4\pi n y} dy = \frac{4!}{(2\pi)^5} \sum_{n=0}^k \frac{|a_n|^2}{(2n+1)^5}.$$

Next note that $\theta_6(f_k) = \sum_{n=1}^{2k+1} b_n \theta_6(e^{2\pi i n z})$. However, since $A_6(\Gamma(1))$ has dimension one, $\theta_6(e^{2\pi i n z}) = c_n \Delta$, where c_n can be determined easily using properties of the scalar product. $\theta_6(e^{2\pi i n z})$ is generally denoted by G_n , the n th Poincaré series. Then $(G_n, \Delta) = (c_n \Delta, \Delta)$ and $c_n = (G_n, \Delta) / (\Delta, \Delta)$. However, by the scalar product formula (Rankin [5, p. 149]),

$$(G_n, \Delta) = \frac{10! \tau(n)}{(4\pi n)^{11}}.$$

As a result

$$\|\theta_6(f_k)\|_{A_6(\Gamma(1))} = \frac{10! \|\Delta\|_{A_6(\Gamma(1))}}{(4\pi)^{11} (\Delta, \Delta)} \left| \sum_{n=1}^{2k+1} \frac{b_n \tau(n)}{n^{11}} \right|.$$

In order to use the above results it is necessary to know the values of (Δ, Δ) and $\|\Delta\|_{A_6(\Gamma(1))}$. Lehmer [2] has shown that

$$(\Delta, \Delta) = 10^{-6} \times 1.03529048.$$

$\|\Delta\|_{A_6(\Gamma(1))}$ was evaluated at Ohio State on the Amdahl 470 using a nested Simpson's rule in double-precision Fortran giving

$$\|\Delta\|_{A_6(\Gamma(1))} = .00070225689.$$

3. The Lower Bound. Now consider polynomials $p_k(z)$ with $a_0 \neq 0$. For $k = 1$ to 30, the approximate maximum of $\|\theta_6(f_k)\|_{A_6(\Gamma(1))} / \|f_k\|_{A_6}$ was calculated, where the coefficients a_1, \dots, a_k of $p_k(z)$ were taken to be real. The results are listed in Table I and show that $\|\theta_6\| \geq .927$. All calculations were carried out in basic on Apple II⁺.

The approximate maximum was found using a Hooke-Jeeves pattern search method as implemented by Nash in [3] with tolerance .001. These calculations were verified for $k = 5, 10, 15, 20$ using the Nelder-Meade simplex method as coded by O'Neill in [4] and were found to be in excellent agreement.

TABLE I

k	approximate maximum of $\ \theta_6(f_k)\ _{A_6(\Gamma(1))} / \ f_k\ _{A_6}$
1	.852608215
2	.881141356
3	.893379912
4	.893629489
5	.903739297
6	.909550921
7	.910936172
8	.913052751
9	.913054225
10	.913458802
11	.916965021
12	.916981747
13	.919648617
14	.920731796
15	.921503011
16	.922274684
17	.922296736
18	.922403535
19	.922519335
20	.924591551
21	.926187880
22	.926225627
23	.926568015
24	.926636356
25	.926636555
26	.926643190
27	.926922205
28	.927056661
29	.927062756
30	.927068371

Department of Mathematics
The Ohio State University
Columbus, Ohio 43210

1. I. KRA, *Automorphic Forms and Kleinian Groups*, Benjamin, Reading, Mass., 1972.
2. D. H. LEHMER, "Ramanujan's function $\tau(n)$," *Duke Math J.*, v. 10, 1943, pp. 483-492.
3. J. C. NASH, "Function minimizations," *Interface Age*, v. 7, 1982, pp. 34-42.
4. R. O'NEILL, "Function minimization using a simplex procedure," *Applied Statistics*, v. 20, 1971, pp. 338-345.
5. R. RANKIN, *Modular Forms and Functions*, Cambridge Univ. Press, Cambridge, 1977.