

## CORRIGENDA

M. LAX & G. P. AGRAWAL, "Evaluation of Fourier integrals using *B*-splines," *Math. Comp.*, v. 39, 1982, p. 547. Replace Eq. (A18) with

$$\Phi_{2,3} = h \left( \frac{2}{3} + \frac{\lambda}{2} + \frac{7\lambda^2}{30} + \frac{\lambda^3}{12} + \frac{31\lambda^4}{1260} + \dots \right).$$

This correction has no effect on any of our results, since the code used the correct equation.

M. LAX AND G. P. AGRAWAL

Physics Department  
City College of the City University of New York  
New York, New York, 10031

Bell Laboratories  
Murray Hill, New Jersey 07974

P. KEAST, G. FAIRWEATHER & J. C. DIAZ, "A computational study of finite element methods for second order linear two-point boundary value problems," *Math. Comp.*, v. 40, 1983, pp. 499–518.

On page 502, Lemma 1 should read:

LEMMA 1. Let  $\{\phi_j\}_{j=1}^K$  and  $\{\psi_j\}_{j=1}^K$  denote the basis functions spanning the *B*-splines of order *K* on the subintervals  $I_1$  and  $I_{\text{NINT}}$ , respectively. Then

$$\begin{aligned} \phi_{j+1}^{(i)}(x) \Big|_{x=Z_0^+} &= l_{ij}, & i, j &= 0, 1, \dots, K-1, \\ \psi_{j+1}^{(i)}(x) \Big|_{x=Z_{\text{NINT}}^-} &= u_{ij}, \end{aligned}$$

where

$$\begin{aligned} l_{ij} &= 0 & \text{if } i < j, \\ u_{ij} &= 0 & \text{if } K - i > j + 1. \end{aligned}$$

On page 503, line 1 after Figure 1, for  $|\lambda_1| + |\lambda_0| \neq 0$  read  $|\lambda_1| \cdot |\lambda_0| \neq 0$ .

On page 505, line 1 after Figure 4, for  $|\lambda_0| + |\lambda_1| > 0$  read  $|\lambda_0| \cdot |\lambda_1| > 0$ .

On page 507, line 12 from foot, in the differential equation, for  $1 + (x - \bar{x})$  read  $1 + \alpha(x - \bar{x})$ .

On page 507, line 10 from foot, for  $\alpha^{-1}\alpha$  read  $\alpha^{-1} + \alpha$ .

G. FAIRWEATHER

Department of Computer Science  
University of Toronto  
Toronto, Ontario M5S 1A7, Canada