

## Odd Triperfect Numbers Are Divisible By Eleven Distinct Prime Factors

By Masao Kishore

**Abstract.** We prove that an odd triperfect number has at least eleven distinct prime factors.

**1. Introduction.** A positive number  $N$  is called a triperfect number if  $\sigma(N) = 3N$  where  $\sigma(N)$  is the sum of the positive divisors of  $N$ . Six even triperfect numbers are known:

$$\begin{aligned} &2^{14} \cdot 5 \cdot 7 \cdot 19 \cdot 31 \cdot 151, \\ &2^{13} \cdot 3 \cdot 11 \cdot 43 \cdot 127, \\ &2^9 \cdot 3 \cdot 11 \cdot 31, \\ &2^8 \cdot 5 \cdot 7 \cdot 19 \cdot 37 \cdot 73, \\ &2^5 \cdot 3 \cdot 7, \\ &2^3 \cdot 3 \cdot 5. \end{aligned}$$

However, the existence of an odd triperfect (OT) number is an open question. McDaniel [4] and Cohen [2] proved that an OT number has at least nine distinct prime factors; the author proved that it has at least ten prime factors [3], and Beck and Najjar [1] showed that it exceeds  $10^{50}$ .

In this paper we prove

**THEOREM.** *If  $N$  is OT,  $N$  has at least eleven distinct prime factors.*

**2. Proof of Theorem.** Throughout this paper we let

$$N = \prod_{i=1}^{10} p_i^{a_i},$$

where  $p_i$ 's are odd primes,  $p_1 < \cdots < p_{10}$  and  $a_i$ 's are positive integers. We call  $p_i^{a_i}$  a component of  $N$  and write  $p_i^{a_i} \parallel N$ .

The following lemmas are easy to prove:

**LEMMA 1.** *If  $N$  is OT,  $a_i$ 's are even for  $1 \leq i \leq 10$ .*

**LEMMA 2.** *If  $N$  is OT and  $q$  is a prime factor of  $\sigma(p_i^{a_i})$  for some  $i$ , then  $q = 3$  or  $q = p_j$  for some  $j$ ,  $1 \leq j \leq 10$ .*

The following lemmas are stated in [5].

---

Received April 4, 1983; revised June 5, 1984.  
1980 *Mathematics Subject Classification.* Primary 10A20.

©1985 American Mathematical Society  
0025-5718/85 \$1.00 + \$.25 per page

**LEMMA 3.** *Suppose  $q$  is a prime,  $q \geq 2$  and  $a \geq 1$ . Then  $\sigma(q^a)$  has a prime factor  $p$  such that  $a + 1$  is the order of  $q$  modulo  $p$  except for  $q = 2$  and  $a = 5$  and for  $q = a$  Mersenne prime and  $a = 1$ . In particular  $a + 1 \mid p - 1$ .*

**LEMMA 4.** *Suppose  $p$  is a Fermat prime (3, 5, 17, etc.),  $q$  is an odd prime and  $a$  is even. If  $p^b \mid \sigma(q^a)$ , then  $q \equiv 1 \pmod{p}$ ,  $p^b \mid a + 1$ , and  $\sigma(q^a)$  has  $b$  distinct prime factors congruent to 1 modulo  $p$ .*

**LEMMA 5.** *If  $N$  is OT,  $17 \nmid N$ .*

*Proof.* Suppose  $N$  is OT. Since the three smallest primes  $\equiv 1 \pmod{17}$  are 103, 137, and 239 and

$$\frac{3}{2} \frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{19}{18} \frac{103}{102} \frac{137}{136} \frac{239}{238} < 3,$$

$N$  has at most two primes  $\equiv 1 \pmod{17}$ . Suppose  $p^a$  and  $q^b$  are components of  $N$  and  $p \equiv q \equiv 1 \pmod{17}$ . If  $17^c \mid N$  and  $c \geq 4$ , then  $17^2 \mid \sigma(p^a)$  or  $17^2 \mid \sigma(q^b)$ , and, by Lemma 4,  $N$  would have two more primes  $\equiv 1 \pmod{17}$ , a contradiction. Hence  $17^4 \nmid N$ . Suppose  $17^2 \mid \mid N$ . Then  $N$  has a component  $307^d$  because  $\sigma(17^2) = 307$ . Then  $17 \nmid \sigma(307^d)$  because  $16661 \cdot 36857 \mid \sigma(307^{16})$ ,  $\sigma(307^{16}) \mid \sigma(307^d)$  and

$$\frac{3}{2} \frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{17}{16} \frac{19}{18} \frac{307}{306} \frac{16661}{16660} \frac{36857}{36856} < 3.$$

Hence  $N$  has another component  $p^b$  such that  $17^2 \mid \sigma(p^b)$ . Then we get a contradiction again. Hence  $17 \nmid N$ . Q.E.D.

The proof of the following lemma is easy.

**LEMMA 6.** *If  $N$  is OT,  $p_9 \leq 283$ .*

**LEMMA 7.** *If  $N$  is OT and  $5^a \mid \mid N$ , then  $a = 2$ ,  $5^2 \mid \sigma(P_{10}^{a_{10}})$  and  $p_{10} \geq 311$ .*

*Proof.* Suppose  $N$  is OT,  $p^b$  is a component of  $N$  and  $5 \mid \sigma(p^b)$ . By Lemma 4,  $p \equiv 1 \pmod{5}$ ,  $5 \mid b + 1$  and  $\sigma(p^4) \mid \sigma(p^b)$ . If  $61 \leq p \leq 281$ , then  $\sigma(p^4)$  has a prime factor  $q$  such that

$$\frac{3}{2} \frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{19}{18} \frac{23}{22} \frac{29}{28} \frac{p}{p-1} \frac{q}{q-1} < 3, \quad \text{or}$$

$\sigma(p^4)$  has prime factors  $q$  and  $r$  such that

$$\frac{3}{2} \frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{19}{18} \frac{23}{22} \frac{p}{p-1} \frac{q}{q-1} \frac{r}{r-1} < 3.$$

Hence  $p = 11, 31$  or  $41$  or  $p \geq 311$ .

Suppose  $p = 11, 31$  or  $41$ . If  $5^2 \mid \sigma(p^b)$ ,  $5^2 \mid b + 1$  by Lemma 4. Then  $\sigma(p^{24}) \mid \sigma(p^b)$  and  $\sigma(p^{24})$  has two distinct prime factors  $> 283$ , contradicting Lemma 6. Hence  $5^2 \nmid \sigma(p^b)$ . Since  $3221 \mid \sigma(11^4)$ ,  $17351 \mid \sigma(31^4)$  and  $579281 \mid \sigma(41^4)$ ,  $5^2 \nmid \sigma(\prod_{i=1}^9 p_i^{a_i})$  and  $p_{10} = 3221, 17351$  or  $579281$  and  $5 \mid \sigma(p_{10}^{a_{10}})$ . However,  $\sigma(p_{10}^4)$  has a prime factor  $> 283$ , contradicting Lemma 6. Hence  $p \geq 311$  and  $5^a \mid \sigma(p^b)$ .

If  $a \geq 4$ , then by Lemma 4,  $N$  would have four more primes  $\equiv 1 \pmod{5}$ , which is a contradiction because

$$\frac{3}{2} \frac{5}{4} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{19}{18} \frac{31}{30} \frac{41}{40} \frac{61}{60} \frac{311}{310} < 3.$$

Hence  $a = 2$ . Q.E.D.

LEMMA 8. If  $N$  is OT,  $p_9 \leq 71$ .

*Proof.* By Lemma 6,  $31 = \sigma(5^2) \mid N$ . Since

$$\frac{3}{2} \frac{\sigma(5^2)}{5^2} \frac{7}{6} \frac{11}{10} \frac{13}{12} \frac{19}{18} \frac{23}{22} \frac{31}{30} \frac{73}{72} \frac{311}{310} < 3,$$

$p_9 \leq 71$ . Q.E.D.

*Proof of Theorem.* If  $N$  is OT, then by Lemmas 4 and 7,  $5^2 \mid \sigma(p_{10}^{a_{10}})$ ,  $5^2 \mid a_{10} + 1$  and  $\sigma(p_{10}^{24}) \mid (p_{10}^{a_{10}})$ . By Lemma 3,  $\sigma(p_{10}^{24})$  has a prime factor  $q$  such that  $25 \mid q - 1$ . Hence  $q = 25b + 1$  for some  $b$ . Since  $q$  is a prime,  $b \neq 1$  or  $2$ . Then  $q > 71$  and  $q \neq p_{10}$ , contradicting Lemma 8. Q.E.D.

Department of Mathematics  
East Carolina University  
Greenville, North Carolina 27834

1. W. E. BECK & R. M. NAJAR, "A lower bound for odd triperfects," *Math. Comp.*, v. 38, 1982, pp. 249–251.
2. G. L. COHEN, "On odd perfect numbers II, Multiperfect numbers and quasiperfect numbers," *J. Austral. Math. Soc.*, v. 29, 1980, pp. 369–384.
3. M. KISHORE, "Odd triperfect numbers," *Math. Comp.*, v. 42, 1984, pp. 231–233.
4. W. MCDANIEL, "On odd multiply perfect numbers," *Boll. Un. Mat. Ital.* (4), v. 3, 1970, pp. 185–190.
5. C. POMERANCE, "Odd perfect numbers are divisible by at least seven distinct primes," *Acta Arith.*, v. 25, 1973/1974, pp. 265–300.