

## CORRIGENDA

FRANÇOIS MORAIN, *On the lcm of the differences of eight primes*, Math. Comp. **52** (1989), 225–229.

On p. 225 it was stated that if

$$r(Q) = \text{lcm}(q_j - q_i)_{1 \leq i < j \leq 8},$$

where  $Q = \{q_1, \dots, q_8\}$  is a set of eight odd primes with  $q_1 < \dots < q_8$ , then

- Erdős has conjectured that  $5040 \mid r(Q)$  for any  $Q$ ;
- **Theorem 1.** *For every  $Q$ ,  $5040 \mid r(Q)$ .*

Both assertions are wrong. It should have been:

- Erdős has conjectured that  $5040 \leq r(Q)$  for any  $Q$ ;
- **Theorem 1.** *For every  $Q$ ,  $5040 \leq r(Q)$ .*

Actually, this is what is proved in the paper. Indeed, it is possible to find examples of sets  $Q$  for which 5040 does not divide  $r(Q)$ . J. Leech has proposed  $r(\{210n + 199, n = 1(1)8\}) = 2^3 3^2 5^2 7^2$  and R. A. Morris  $r(\{11, 17, 19, 23, 29, 41, 47, 53\}) = 2^3 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 17$ . As a matter of fact, the smallest  $\rho$  for which there exists a set  $Q$  such that  $r(Q) = \rho$  and  $2^3 \parallel \rho$  is  $\rho = 2^3 3^2 \cdot 5 \cdot 7 \cdot 11$  with  $Q = (\{17, 19, 23, 29, 37, 41, 47, 59\})$  for instance.

F. MORAIN \*

Institut National de Recherche en Informatique et en Automatique (INRIA)  
Domaine de Voluceau, B. P. 105  
78153 Le Chesnay Cedex (France)  
& Département de Mathématiques  
Université Claude Bernard  
69622 Villeurbanne Cedex (France)  
E-mail: morain@inria.inria.fr

---

\* On leave from the French Department of Defense, Délégation Générale pour l'Armement.