

COMPUTATION OF THE TRIVARIATE NORMAL INTEGRAL

ZVI DREZNER

ABSTRACT. We propose a simple and efficient way to calculate trivariate normal probabilities. The algorithm is based on a formula for the partial derivative of the trivariate probability with respect to a correlation coefficient.

1. INTRODUCTION

The trivariate normal distribution is defined by a correlation matrix R ,

$$R = \begin{pmatrix} 1 & \rho_{12} & \rho_{13} \\ \rho_{12} & 1 & \rho_{23} \\ \rho_{13} & \rho_{23} & 1 \end{pmatrix}.$$

The trivariate probability $L_3(h_1, h_2, h_3; \rho_{12}, \rho_{13}, \rho_{23})$ (or $L_3(h; R)$) is [6]:

$$(1) \quad \begin{aligned} &L_3(h_1, h_2, h_3; \rho_{12}, \rho_{13}, \rho_{23}) \\ &= \frac{1}{\sqrt{8\pi^3|R|}} \int_{h_1}^{\infty} \int_{h_2}^{\infty} \int_{h_3}^{\infty} e^{-x^T R^{-1} x/2} dx_3 dx_2 dx_1, \end{aligned}$$

where $x^T = (x_1, x_2, x_3)$. The L function is by definition

$$(2) \quad L_3(h_1, h_2, h_3; R) = \Pr[(X_1 > h_1) \cap (X_2 > h_2) \cap (X_3 > h_3)].$$

The bivariate and univariate normal distributions are defined by [3, 6]:

$$(3) \quad L_2(h_1, h_2; \rho) = \Pr[(X_1 > h_1) \cap (X_2 > h_2)],$$

$$(4) \quad L_1(h) = \Pr(X > h).$$

An efficient way, by using a five-point Gaussian quadrature, to calculate $L_1(h)$ and $L_2(h; \rho)$ to an accuracy of $2 \cdot 10^{-7}$ for $-1 \leq \rho \leq 1$ is given in [5]. Applying the same method, with Gaussian quadrature based on more than five points, yields any desired accuracy. Our goal here is to calculate L_3 to the same accuracy for most values of the ρ_{ij} 's. Therefore, it is assumed that if values of L_1 and L_2 are required in our calculations, then they are easily available.

The calculation of the integral (1) was a topic of research for a long time. See [6] for a survey of various computation methods and approximations. More

Received by the editor April 30, 1992.

1991 *Mathematics Subject Classification.* Primary 65U05, 65D20, 62H10.

Key words and phrases. Trivariate normal, multivariate normal, computation.

recent works on the multivariate normal integral that can be used to calculate the trivariate integral are [4, 9, 10]. In this paper we present a very efficient and accurate integration method specific to the trivariate normal integral.

2. ON THE DETERMINANT OF R

The determinant of R is

$$(5) \quad |R| = 1 - \rho_{12}^2 - \rho_{13}^2 - \rho_{23}^2 + 2\rho_{12}\rho_{13}\rho_{23}.$$

A correlation matrix must be positive semidefinite, which translates into $-1 \leq \rho_{ij} \leq 1$, and $|R| \geq 0$. It can be easily shown that $|R| \leq 1$. One cannot have a correlation matrix for which $|R| < 0$, and therefore we do not need to worry about calculations involving "illegal" matrices for which $|R| < 0$.

Another issue concerning the determinant of R is whether it is possible that certain values of the ρ 's yield a positive determinant but smaller values yield a negative $|R|$. This issue is important because if we integrate from one value of ρ to another we must make sure that the path of integration passes through points for which $|R| > 0$. Intuitively, it seems that if we have a "legal" correlation matrix, then a correlation matrix with smaller ρ 's must also be legal. This is not necessarily true. Consider the following example. The three ρ 's are $33/65$, $60/65$, $52/65$. By direct calculations $|R| = 0$. If the first ρ is changed to $34/65$, $|R| > 0$, and if it is changed to $32/65$, then $|R| < 0$. However, if one ρ is held constant and the other two are changed proportionally, then if the original $|R|$ is positive, so is the whole path. To prove this, define

$$|R(t)| = 1 - t^2\rho_{12}^2 - t^2\rho_{13}^2 - \rho_{23}^2 + 2t^2\rho_{12}\rho_{13}\rho_{23}.$$

Theorem 1. *If $|R(t)|$ is positive for $t = 1$, then it is positive for $0 \leq t \leq 1$.*

Proof. We need to show that $\rho_{12}^2 + \rho_{13}^2 - 2\rho_{12}\rho_{13}\rho_{23} \geq 0$. If $\rho_{12}\rho_{13} \geq 0$, then $\rho_{12}^2 + \rho_{13}^2 - 2\rho_{12}\rho_{13}\rho_{23} = (\rho_{12} - \rho_{13})^2 + 2\rho_{12}\rho_{13}(1 - \rho_{23}) \geq 0$. If $\rho_{12}\rho_{13} < 0$, then $\rho_{12}^2 + \rho_{13}^2 - 2\rho_{12}\rho_{13}\rho_{23} = (\rho_{12} + \rho_{13})^2 - 2\rho_{12}\rho_{13}(1 + \rho_{23}) \geq 0$. \square

A similar result can also be shown: if all the three ρ 's are reduced proportionally, then $|R|$ increases. Simply, a reduction by a factor of \sqrt{t} can be applied three times, once for each possible pair of ρ 's, and the result is that each ρ is reduced by a factor of t .

3. CALCULATION OF THE TRIVARIATE PROBABILITY

There are a few specific cases where L_3 can be expressed in formulas using L_1 and L_2 . They are summarized in Table 1.

TABLE 1. Some specific cases

| ρ_{12} | ρ_{13} | ρ_{23} | $L_3(h_1, h_2, h_3; R)$ |
|-------------|-------------|-------------|------------------------------------|
| 0 | 0 | 0 | $L_1(h_1)L_2(h_2)L_1(h_3)$ |
| 0 | 0 | ρ | $L_1(h_1)L_2(h_2, h_3; \rho)$ |
| ρ | ρ | 1 | $L_2(h_1, \max\{h_2, h_3\}; \rho)$ |

The trivariate case for the general formula developed by Plackett [6, 8] is

$$(6) \quad \frac{\partial L_3}{\partial \rho_{12}} = Z_2(h_1, h_2; \rho_{12})L_1(h'_3),$$

where Z_2 is the density function of L_2 :

$$(7) \quad Z_2(x, y; \rho) = \frac{\exp(-\frac{x^2+y^2-2\rho xy}{2(1-\rho^2)})}{2\pi\sqrt{1-\rho^2}}$$

and h'_3 is

$$(8) \quad h'_3 = \frac{h_3(1 - \rho_{12}^2) - (\rho_{13} - \rho_{23}\rho_{12})h_1 - (\rho_{23} - \rho_{13}\rho_{12})h_2}{\sqrt{(1 - \rho_{12}^2)|R|}}$$

We suggest two ways to apply (6). The first one is

$$(9) \quad L_3(h; R) = L_3(h; 0) + \int_0^1 \sum_{i < j} \rho_{ij} \frac{\partial L_3(h; t\rho_{12}, t\rho_{13}, t\rho_{23})}{\partial(t\rho_{ij})} dt.$$

Another approach is based on the second row in Table 1:

$$(10) \quad L_3(h; R) = L_3(h; 0, 0, \rho_{23}) + \int_0^1 \sum_{i=2}^3 \rho_{1i} \frac{\partial L_3(h; t\rho_{12}, t\rho_{13}, \rho_{23})}{\partial(t\rho_{1i})} dt,$$

where $L_3(h; 0, 0, \rho_{23}) = L_1(h_1)L_2(h_2, h_3; \rho_{23})$.

By Theorem 1, $|R(t)| > 0$ on the whole integration interval as long as $|R| > 0$ for the original correlation matrix (at $t = 1$).

Since (9) and (10) are smoother for small ρ 's, we suggest resorting the variables such that ρ_{23} is the largest (in absolute value), and using (10) for integrating on the other two ρ 's. It is interesting to note that integrating one ρ at a time, as suggested in [8], may lead to a negative $|R|$ as explained before Theorem 1. The integration can be done by Simpson's rule [1], or preferably by the Gauss quadrature formula based on Legendre polynomials [1]. Expression (10) is definitely superior to (9).

4. ORTHANT PROBABILITIES

As a result of (6)-(9), the orthant probabilities can be calculated explicitly. If $h = 0$, then $h'_3 = 0$, and $L_1(h'_3) = 1/2$. Substituting into (6), we get

$$\frac{\partial L_3}{\partial \rho_{12}} = \frac{Z_2(0; \rho_{12})}{2}$$

Substituting into (9), we get

$$L_3(0; R) = \frac{1}{8} + \frac{1}{2} \sum_{i < j} [L_2(0; \rho_{ij}) - L_2(0; 0)] = \frac{1}{8} + \frac{1}{4\pi} \sum_{i < j} \sin^{-1} \rho_{ij},$$

which is the result given in [2, 7].

5. COMPUTATIONAL EXPERIENCE

A FORTRAN program was written on an XT-compatible with math co-processor. We used the Gaussian quadrature based on Legendre polynomials with only five points [1]. This quadrature method was proven to be very effective in calculating bivariate probabilities and requires very little computational effort [5]. Using a K -point quadrature formula requires the calculation of one bivariate integral, K exponents and $K + 1$ normal (univariate) probabilities. For $K = 5$, the program requires about 0.03 seconds on the XT-compatible.

Problems with h_i for $i = 1, 2, 3$ in $[-3, 3]$ and the ρ_{ij} 's in $[-\rho_{\max}, \rho_{\max}]$ were randomly generated using uniform distributions. If the determinant $|R|$ is negative, the problem is discarded and a new one generated. The errors in the five-point quadrature formula were obtained by comparing with results calculated with a larger K and double-precision arithmetic. We also compared with the results obtained by the program in [4]. We encountered a very interesting phenomenon. For $\rho_{\max} \leq 0.5$, the error never exceeded 10^{-7} . For larger ρ_{\max} (even for $\rho_{\max} = 0.9$) the error was limited by 10^{-7} in almost all cases but the error has reached 10^{-5} in very few cases. We believe that the error is more dependent on $|R|$ than it is on large ρ 's. So, we experimented with $\rho_{\max} = 1$ (i.e., generating any possible ρ) but allowed only problems where the determinant $|R|$ is greater than a given cutoff point between zero and one. For each cutoff experiment, 1000 feasible problems were generated. It was found that when the cutoff was 0.3 or higher, then the error was less than 10^{-7} for all 1000 problems. (Calculations were done in single-precision arithmetic.) For smaller cutoffs the errors were reasonable and are summarized in Table 2.

TABLE 2. Experimentation with trivariate calculations

| $ R $ Cutoff | Max Error |
|--------------|-----------|
| 0.200 | 0.0000001 |
| 0.175 | 0.0000001 |
| 0.150 | 0.0000001 |
| 0.125 | 0.0000002 |
| 0.100 | 0.0000007 |
| 0.075 | 0.0000014 |
| 0.050 | 0.0000026 |
| 0.025 | 0.0000044 |
| 0.000 | 0.0000117 |

In conclusion, we presented a very efficient way to calculate trivariate probabilities. If the determinant $|R|$ is greater than 0.15, the error in the calculation is less than 10^{-7} . For smaller determinants (which means that the variables are close to being linearly dependent) the five-point Gaussian quadrature formula may give a higher error. If a better accuracy is sought, the quadrature can be done with more points, or Simpson's formula can be applied.

APPENDIX. A FORTRAN program

```

FUNCTION TV(H,R,IER)
C   H contains the the values of h1, h2, h3.
C   R contains the values of r12, r13, r23 in that order.
C   IER=1 Negative determinant.
C       =0 Calculation completed.
C   The functions UV (Univariate) and BV (Bivariate) can be taken from [5].
DIMENSION X(5),W(5),H(3),R(3)
DATA X/.04691008,.23076534,.5,.76923466,.95308992/
DATA W/.018854042,.038088059,.0452707394,.038088059,.018854042/
IER=1
H1=H(1)
H2=H(2)
H3=H(3)
R12=R(1)
R13=R(2)
R23=R(3)
DO 1 K=1,3
IF(ABS(R23).GE.ABS(R12).AND.ABS(R23).GE.ABS(R13))GO TO 2
HH=H1
H1=H2
H2=H3
H3=HH
RR=R12
R12=R23
R23=R13
R13=RR
1 CONTINUE
2 CONTINUE
H12=H1*H2
H13=H1*H3
H122=(H1*H1+H2*H2)*0.5
H132=(H1*H1+H3*H3)*0.5
TV=0.
DO 3 I=1,5
RR12=R12*X(I)
RR13=R13*X(I)
DEL=1.-RR12*RR12-RR13*RR13-R23*R23+2*RR12*RR13*R23
IF(DEL.LE.0.)RETURN
FAC=SQRT(DEL)
RR122=1.-RR12*RR12
RR133=1.-RR13*RR13
F1=RR13-R23*RR12
F2=R23-RR12*RR13
F3=RR12-R23*RR13
HP1=(H3*RR122-H1*F1-H2*F2)/FAC/SQRT(RR122)
HP2=(H2*RR133-H1*F3-H3*F2)/FAC/SQRT(RR133)
TV=TV+W(I)*EXP((RR12*H12-H122)/RR122)/SQRT(RR122)*UV(HP1)*R12
3 TV=TV+W(I)*EXP((RR13*H13-H132)/RR133)/SQRT(RR133)*UV(HP2)*R13
TV=UV(H1)*BV(H2,H3,R23)+TVE1
IER=0
RETURN
END

```

BIBLIOGRAPHY

1. M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions*, National Bureau of Standards, Washington, D.C., 7th printing, 1968.
2. F. N. David, *A note on the evaluation of the multivariate normal integral*, *Biometrika* **40** (1953), 458-459.

3. Z. Drezner, *Computation of the bivariate normal integral*, *Math. Comp.* **32** (1978), 277–279.
4. —, *Computation of the multivariate normal integral*, *ACM Trans. Math. Software* **18** (1992), 470–480.
5. Z. Drezner and G. O. Wesolowsky, *On the computation of the bivariate normal integral*, *J. Statist. Comput. Simulation* **35** (1990), 101–107.
6. N. L. Johnson and S. Kotz, *Distributions in statistics: Continuous multivariate distributions*, Wiley, New York, 1972.
7. D. B. Owen, *Orthant probabilities*, *Encyclopedia of Statistical Sciences*, vol. 6, 1985, pp. 521–523.
8. R. L. Plackett, *A reduction formula for normal multivariate integrals*, *Biometrika* **41** (1954), 351–360.
9. M. J. Schervish, *Multivariate normal probabilities with error bound*, *Appl. Statist.* **33** (1984), 81–87.
10. —, *Correction to algorithm AS 195: multivariate normal probabilities with error bound*, *Appl. Statist.* **34** (1985), 103–104.

DEPARTMENT OF MANAGEMENT SCIENCE, SCHOOL OF BUSINESS ADMINISTRATION AND ECONOMICS, CALIFORNIA STATE UNIVERSITY AT FULLERTON, FULLERTON, CALIFORNIA 92634
E-mail address: drezner@fullerton.edu