

BOUNDARY ELEMENT MONOTONE ITERATION SCHEME FOR SEMILINEAR ELLIPTIC PARTIAL DIFFERENTIAL EQUATIONS

YUANHUA DENG, GOONG CHEN, WEI-MING NI, AND JIANXIN ZHOU

ABSTRACT. The monotone iteration scheme is a constructive method for solving a wide class of semilinear elliptic boundary value problems. With the availability of a supersolution and a subsolution, the iterates converge monotonically to one or two solutions of the nonlinear PDE. However, the rates of such monotone convergence cannot be determined in general. In addition, when the monotone iteration scheme is implemented numerically through the boundary element method, error estimates have not been analyzed in earlier studies. In this paper, we formulate a working assumption to obtain an exponentially fast rate of convergence. This allows a margin δ for the numerical implementation of boundary elements within the range of monotone convergence. We then interrelate several approximate solutions, and use the Aubin-Nitsche lemma and the triangle inequalities to derive error estimates for the Galerkin boundary-element iterates with respect to the $H^r(\Omega)$, $0 \leq r \leq 2$, Sobolev space norms. Such estimates are of optimal order. Furthermore, as a peculiarity, we show that for the nonlinearities that are of separable type, “higher than optimal order” error estimates can be obtained with respect to the mesh parameter h . Several examples of semilinear elliptic partial differential equations featuring different situations of existence/nonexistence, uniqueness/multiplicity and stability are discussed, computed, and the graphics of their numerical solutions are illustrated.

1. INTRODUCTION

Numerical solutions of nonlinear partial differential equations (PDEs) are important in applications. Historically, such work is done primarily by the finite difference methods (FDM) and finite element methods (FEM). While boundary element methods (BEM) have steadily gained popularity among engineers and scientists in their work of computing solutions of PDEs, owing to the very nature of their formulation, BEM are still regarded by many people as mainly applicable to

Received by the editor May 4, 1994 and, in revised form, March 1, 1995.

1991 *Mathematics Subject Classification.* Primary 31B20, 35J65, 65N38.

Key words and phrases. Numerical PDE, boundary elements, potential theory, nonlinear PDE, elliptic type.

The first, second, and fourth authors were supported in part by AFOSR Grant 91-0097 and NSF Grant DMS 9404380.

The third author was supported in part by NSF Grants DMS 9101446 and 9401333.

The second author was on sabbatical leave at the Institute of Applied Mathematics, National Tsing Hua University, Hsinchu, Taiwan, ROC. Supported in part by a grant from the National Science Council of the Republic of China.

linear PDEs. Nevertheless, progress has been made; several researchers have applied BEM to encompass nonlinearities in elliptic *boundary value problems* (BVPs) with some success. For example,

- (i) when the governing equation itself is linear, with nonlinearities occurring only in the boundary conditions, then the applicability of BEM is quite ready. A nonlinear integral equation can be formulated on the boundary; an essential feature and an advantage of BEM are retained. Error analysis for such problems has been done by Ruotsalainen and Wendland [15], e.g.;
- (ii) when the governing equation is nonlinear, with the nonlinearities occurring in the lowest-order terms (leading to a semilinear PDE), Brebbia and Walker [5], and Sakakihara [16] have considered iteration schemes for such problems. Accurate numerical results for certain examples have been obtained. But no error analysis and convergence for the discretized scheme were established in [5] and [16].

The semilinear elliptic PDEs considered in [5] and [16] are of the form

$$(1.1) \quad \begin{cases} \Delta u(x) = F(x, u(x)) & \text{on } \Omega, \\ Bu(x) = g(x) & \text{on } \partial\Omega, \end{cases}$$

where Ω is a simply-connected bounded domain in \mathbb{R}^N with C^∞ smooth boundary $\partial\Omega$, $F: \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ is smooth such that $F(x, u)$ is nonlinear in u , B is a linear boundary operator of the form

$$(1.2) \quad Bu = u \quad \text{or} \quad Bu = \frac{\partial u}{\partial n} + \alpha(\cdot)u, \quad \text{with } \alpha(x) \geq 0 \quad \forall x \in \partial\Omega,$$

and n is the unit outward normal on $\partial\Omega$. A *straightforward* iteration scheme suggested by Brebbia and Walker in [5] goes as follows:

$$(1.3) \quad \begin{cases} \text{Initialize: Take an initial guess } u_0(x); \\ \text{Iterate: Solve, for } n = 0, 1, 2, \dots \\ \quad \begin{cases} \Delta u_{n+1}(x) = F(x, u_n(x)), & x \in \Omega, \\ (Bu_{n+1})(x) = g(x), & x \in \partial\Omega, \end{cases} \end{cases}$$

where at each iteration $u_{n+1}(\cdot)$ is solved by a boundary element method. Although this scheme appears quite natural and sometimes even produces nice numerical results, unfortunately, for “strong” nonlinearities, it fails to converge in general, as will be shown in §2.

On the other hand, the iteration scheme used by Sakakihara in [16] is the *monotone iteration scheme*:

$$(1.4) \quad \begin{cases} (1) \text{ Choose a } \lambda > 0 \text{ “sufficiently large”}; \quad (\text{cf. (2.9)}) \\ (2) \text{ Initialize: Take an initial guess } u_0(x); \\ (3) \text{ Iterate: Solve, for } n = 0, 1, 2, \dots \\ \quad \begin{cases} \Delta u_{n+1}(x) - \lambda u_{n+1}(x) = -\lambda u_n(x) + F(x, u_n(x)), & x \in \Omega, \\ (Bu_{n+1})(x) = g(x), & x \in \partial\Omega. \end{cases} \\ \text{(Here } u_{n+1} \text{ is also solved by a boundary element method, but} \\ \text{other general numerical methods may also be used.)} \end{cases}$$

The monotone iteration scheme is a well-known *constructive* method for solving semilinear elliptic PDEs of the type (1.1). When some supersolutions and subsolutions (also known as *barriers*) are available, monotone convergence of the iterates $u_{n+1}(x)$ (of the undiscretized scheme) can be established [2, 12, 17]. In [16], Sakakihara used the Green's formula, i.e., the commonly called *direct approach*, to formulate boundary integral equations (BIEs) to solve each $u_{n+1}(x)$. The convergence of boundary layer densities in those BIEs can be readily established, yielding the convergence of $u_n(x)$ to some $u(x)$ satisfying (1.1), i.e., a solution of (1.1). Nevertheless, in practical BEM computations, a BIE formulation of (1.4) must be *discretized*. Analysis of convergence and error for the discretized problem requires the knowledge of regularities of several singular integral operators involved in the BIEs. Such work is missing in [16].

In boundary element computations of PDEs, *indirect formulations* using only simple- or double-layer potentials to represent the solution are known to have several important advantages, one among which is a significant reduction of the amount of quadrature involved. Here, in our opinion, using the simple-layer potential representation (plus a volume potential to account for the right-hand side $-\lambda u_n(x) + F(x, u_n(x))$ in (1.4), see the second integral in (3.8)) is preferable to the direct formulation adopted in [16]. This will be the basic approach taken in this paper. We will undertake the task of establishing convergence and error estimates for the discretized Galerkin boundary-element scheme corresponding to this approach. In our attempt to establish convergence, we are faced with some fundamental issues in the study of nonlinear PDEs: nonexistence of solutions, or the existence of multiple solutions, some of which are known to be *unstable* [12, 17]. (Unstable solutions may be briefly explained as a saddle-point type critical point on the solution manifold, which must be obtained through a mountain-pass lemma [3] kind of argument and are thus unobtainable as solutions of a maximization/minimization variational problem. Or, unstable solutions may be regarded as an unstable equilibrium of a corresponding dynamic problem.) Only after these questions are properly understood can one talk about which solution the numerical iterates are converging to. Indeed, this may help explain why earlier workers had not been totally successful in deriving any error estimates or convergence for the boundary element monotone iteration scheme. Not surprisingly, a certain additional working assumption (see [H] in §4) is needed in order for the Galerkin boundary element scheme to converge. This assumption, although looking somewhat restrictive at first glance, is actually natural because of the needs to be able to estimate errors, especially after seeing several types of concrete examples of semilinear PDEs analyzed and computed in §7.

The organization of the paper is as follows:

- (i) In §2, we provide some further background material and discussion.
- (ii) In §3, we prove the *antimonotone* convergence property of simple-layer densities of the (undiscretized) boundary integral equations according to our formulation.
- (iii) In §4, we establish convergence and error estimates for a Galerkin boundary element scheme, subject to the Dirichlet boundary condition. By comparing various intermediate approximate solutions, error estimates $\|u_h^n - u_\infty\|_{H^r(\Omega)}$ are obtained which are sharp for the $H^r(\Omega)$ Sobolev space norms, $0 \leq r \leq 2$.
- (iv) A *distinctive feature* of BEM for semilinear PDEs is that we are able to obtain

“higher than optimal order” error estimates with respect to h , provided that the nonlinearity is *separable* (cf. (5.1)). This is presented in §5.

- (v) Error estimates for the semilinear PDE subject to the Neumann or Robin boundary conditions are discussed in §6.
- (vi) In §7, some typical cases of semilinear elliptic PDEs are discussed and computed, with numerical solutions illustrated.

The *collocation scheme* is computationally more convenient than the Galerkin one. Our numerical results in §7 are obtained via the former instead of the latter. At this moment we are still trying to analyze errors for the collocation scheme. This, along with several other related problems, has been partially studied [8] and will appear elsewhere.

2. A STRAIGHTFORWARD ITERATION SCHEME AND THE MONOTONE ITERATION SCHEME

We first review the straightforward iteration scheme (1.3). We show through an ODE example that (1.3) does not converge in general.

Example 2.1. Consider

$$(2.1) \quad \begin{cases} \frac{d^2}{dx^2}u(x) - u^3(x) = 0, & 0 < x < 1, \\ u(0) = u(1) = 0. \end{cases}$$

The differential operator $\frac{d^2}{dx^2}$ represents the simplest elliptic operator in one dimension. Multiplying the equation in (2.1) by u and integrating by parts, we obtain

$$\int_0^1 [|u'(x)|^2 + u^4(x)] dx = 0.$$

Therefore, (2.1) admits a unique solution that is trivial. (This trivial solution is, of course, stable.)

Let us find iterative solutions of (2.1) by (1.3):

$$(2.2) \quad \begin{cases} \text{Set } u_0(x) = C > 0; \\ \text{Solve, for } n = 0, 1, 2, \dots \\ \quad \begin{cases} u''_{n+1}(x) = u_n^3(x), & x \in [0, 1], \\ u_{n+1}(0) = u_{n+1}(1) = 0. \end{cases} \end{cases}$$

For $n = 0$, for example, we get

$$\begin{aligned} u''_1(x) &= C^3, \\ u_1(x) &= \frac{C^3}{2}x^2 + \alpha_1x + \beta_1, \\ u_1(0) = u_1(1) = 0 &\Rightarrow \beta_1 = 0, \alpha_1 = -\frac{C^3}{2}, \end{aligned}$$

hence

$$(2.3) \quad u_1(x) = \frac{C^3}{2}(x^2 - x).$$

In general, we have

$$(2.4) \quad u_n(x) = \alpha_n x^{m_n} + \dots,$$

where the leading coefficient α_n and the power m_n can be uniquely determined recursively:

$$(2.5) \quad \begin{cases} \alpha_0 = C, & m_0 = 0, \\ \alpha_{n+1} = \frac{\alpha_n^3}{(3m_n + 2)(3m_n + 1)}, & m_{n+1} = 3m_n + 2, \quad n = 0, 1, 2, \dots \end{cases}$$

From (2.5), it is not difficult to show that if we choose $C > \frac{2}{3}k_0$ for any $k_0 \geq 6$, then

$$\alpha_n > k_0 \left(m_n + \frac{2}{3} \right) \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Consequently, the iterates $u_n(x)$ in (2.4) cannot converge to the unique solution $u(x) \equiv 0$ of (2.1). \square

Thus, the direct iteration scheme (1.3) does not work for Example 2.1 if the initial state $u_0(x) \equiv C$ is not small. In contrast, the monotone iteration scheme (1.4) works for Example 2.1 for any $C \in \mathbb{R}$ if $u_0(x) \equiv C_i, i = 1, 2$, are properly chosen, because

(i) if $C_1 > 0$, then $\bar{u}(x) \equiv C_1$ satisfies

$$\begin{cases} \bar{u}''(x) - \bar{u}(x)^3 = -C_1^3 < 0 & \text{on } (0,1), \\ \bar{u}(x) = C_1 > 0 & \text{at } x = 0, 1. \end{cases}$$

Therefore, $\bar{u}(x)$ is a supersolution, cf. (2.7).

(ii) If $C_2 < 0$, then $\bar{v}(x) \equiv C_2$ satisfies

$$\begin{cases} \bar{v}''(x) - \bar{v}(x)^3 = -C_2^3 > 0 & \text{on } (0,1), \\ \bar{v}(x) = C_2 < 0 & \text{at } x = 0, 1. \end{cases}$$

Therefore $\bar{v}(x)$ is a subsolution, cf. (2.8).

If we choose $\lambda > 0$ and $C \in \mathbb{R}$ such that

$$\lambda > \max(3C_1^2, 3C_2^2), \quad C \in (C_2, C_1),$$

then for the nonlinearity $F(x, u) = u^3$ in Example 2.1, we have

$$\lambda - \frac{\partial F}{\partial u} = \lambda - 3u^2 > 0 \quad \forall u \in (C_2, C_1).$$

The monotone iteration scheme (1.4) will work for Example 2.1, by Theorem 2.2 below.

To make this paper sufficiently self-contained, in the following we state a version of the monotone iteration theorem.

Theorem 2.2 ([2, 12, 17]). *Let $F(x, u)$ be C^1 with respect to $(x, u) \in \bar{\Omega} \times \mathbb{R}$. Consider the boundary value problem*

$$(2.6) \quad \begin{cases} \Delta u(x) - F(x, u(x)) = 0 & \text{on } \Omega, \\ Bu(x) = g(x) & \text{on } \partial\Omega, \end{cases}$$

where B is given by (1.2) with $\alpha \in C^\infty(\partial\Omega)$, $\alpha(x) \geq 0$, and $\alpha(x) \neq 0$ if $Bu \neq u$ on $\partial\Omega$, and $g \in C^2(\partial\Omega)$. Let $\bar{u}, \bar{v} \in C^2(\bar{\Omega})$ satisfy $\bar{u} \geq \bar{v}$ as well as

$$(2.7) \quad \begin{cases} \Delta \bar{u}(x) - F(x, \bar{u}(x)) \leq 0, & x \in \Omega, \\ B\bar{u}(x) \geq g(x), & x \in \partial\Omega, \end{cases}$$

$$(2.8) \quad \begin{cases} \Delta \bar{v}(x) - F(x, \bar{v}(x)) \geq 0, & x \in \Omega, \\ B\bar{v}(x) \leq g(x), & x \in \partial\Omega. \end{cases}$$

(We call \bar{u} and \bar{v} , respectively, a supersolution and a subsolution for satisfying (2.7) and (2.8).) Choose a number $\lambda > 0$ such that

$$(2.9) \quad \lambda - \frac{\partial F(x, u)}{\partial u} > 0 \quad \forall (x, u) \in \bar{\Omega} \times [a, b], \quad a = \inf \bar{v}, \quad b = \sup \bar{u},$$

and such that the operator $(\Delta - \lambda, B|_{\partial\Omega} = 0)$ has its spectrum strictly contained in the open left-half complex plane. Then the mapping

$$(2.10) \quad T: \phi \mapsto w, \quad w = T\phi,$$

$$(2.11) \quad \phi \in C^2(\bar{\Omega}), \quad \phi(x) \in [a, b] \quad \forall x \in \bar{\Omega},$$

where $w(x)$ is the unique solution of the BVP

$$\begin{cases} \Delta w(x) - \lambda w(x) = -\lambda \phi(x) + F(x, \phi(x)), & x \in \Omega, \\ Bw(x) = g(x), & x \in \partial\Omega, \end{cases}$$

is monotone, i.e., for any ϕ_1, ϕ_2 satisfying (2.11) and $\phi_1 \leq \phi_2$, we have

$$(2.12) \quad T\phi_1, T\phi_2 \text{ satisfy (2.11) and } T\phi_1 \leq T\phi_2 \text{ on } \Omega.$$

Consequently, by letting $F_\lambda(x, u) \equiv -\lambda u + F(x, u)$, the iterations

$$(2.13) \quad \begin{cases} u_0(x) = \bar{u}(x), \\ (\Delta - \lambda)u_{n+1}(x) = F_\lambda(x, u_n(x)), & x \in \Omega, \quad n = 0, 1, 2, \dots, \\ Bu_{n+1} = g & \text{on } \partial\Omega, \end{cases}$$

and

$$(2.14) \quad \begin{cases} v_0(x) = \bar{v}(x), \\ (\Delta - \lambda)v_{n+1}(x) = F_\lambda(x, v_n(x)), & x \in \Omega, \quad n = 0, 1, 2, \dots, \\ Bv_{n+1} = g & \text{on } \partial\Omega, \end{cases}$$

yield iterates u_n and v_n satisfying

$$(2.15) \quad \bar{v} = v_0 \leq v_1 \leq \dots \leq v_n \leq \dots \leq u_n \leq \dots \leq u_1 \leq u_0 = \bar{u},$$

so that the limits

$$(2.16) \quad u_\infty(x) \equiv \lim_{n \rightarrow \infty} u_n(x)$$

and

$$(2.17) \quad v_\infty(x) \equiv \lim_{n \rightarrow \infty} v_n(x)$$

exist in $C^2(\bar{\Omega})$. We have

- (i) $v_\infty \leq u_\infty$ on $\bar{\Omega}$;
- (ii) v_∞ and u_∞ are both stable solutions of (2.6);
- (iii) If $v_\infty \neq u_\infty$, then there exists an unstable solution $\phi \in C^2(\bar{\Omega})$ of (2.6) such that

$$v_\infty \leq \phi \leq u_\infty. \quad \square$$

Remark 2.3. The rate of convergence in (2.16) and (2.17) is not available in general. To make the numerical analysis possible in our subsequent development, obviously some additional working assumption is needed. Otherwise, numerical discretizations cause errors and make the computed solutions deviate outside the range of monotone iterations (2.9). This will be formulated in the important [H] condition in §4. □

Remark 2.4. (a) We note that there is a useful further formulation of the above theorem in the weak (i.e., distributional) sense.

(b) For simplicity of presentation, from now on, we will assume that F, g, \bar{u}, \bar{v} and $\partial\Omega$ (consequently, also u_∞ and v_∞) are all $C^\infty(\bar{\Omega})$, considering that this is not the main issue here. □

3. FORMULATION OF BOUNDARY INTEGRAL EQUATIONS BASED ON THE SIMPLE-LAYER REPRESENTATION

Let $E(x)$ be the fundamental solution satisfying

$$(3.1) \quad (\Delta - \lambda)E(x; \lambda) = -\delta(x), \quad x \in \mathbb{R}^N,$$

where $\delta(x)$ is the Dirac delta function. It is known ([6, p. 341], e.g.) that

$$(3.2) \quad E(x; \lambda) = \frac{1}{4} (i)^{\frac{N}{2}} \left(\frac{\sqrt{\lambda}}{2\pi|x|} \right)^{\frac{N-2}{2}} H_{\frac{N}{2}-1}^{(1)}(i\sqrt{\lambda}|x|), \quad N \geq 2,$$

where $H_{\frac{N}{2}-1}^{(1)}(\cdot)$ is the Hankel function of order $\frac{N}{2} - 1$ of the first kind. In particular, we have

$$(3.3) \quad E(x; \lambda) = \frac{1}{2\pi} K_0(\sqrt{\lambda}|x|), \quad N = 2,$$

where K_0 is the Macdonald function of order 0, and

$$(3.4) \quad E(x; \lambda) = \frac{e^{-\sqrt{\lambda}|x|}}{4\pi|x|}, \quad N = 3.$$

Let the assumptions of Theorem 2.2 be satisfied. We first consider the case when the boundary condition is Dirichlet:

$$(3.5) \quad Bu(x) \equiv u(x) = g(x), \quad \text{on } \partial\Omega,$$

and defer the discussion of the Neumann and Robin boundary conditions to §6. For the boundary integral formulation of the monotone iteration

$$(3.6) \quad \begin{cases} (\Delta - \lambda)u_{n+1}(x) = F_\lambda(x, u_n(x)), & x \in \Omega, \\ u_{n+1}(x) = g(x), & x \in \partial\Omega, \end{cases}$$

the boundary integral equation by the standard *direct formulation* for the unknown $\partial u_{n+1}/\partial n$ is

$$(3.7) \quad \begin{aligned} \int_{\partial\Omega} E(x-y; \lambda) \frac{\partial u_{n+1}(y)}{\partial n} d\sigma_y &= \frac{1}{2}g(x) + \int_{\partial\Omega} \frac{\partial E(x-y; \lambda)}{\partial n_y} g(y) d\sigma_y \\ &\quad - \int_{\Omega} E(x-y; \lambda) F_\lambda(y, u_n(y)) dy, \quad x \in \partial\Omega, \end{aligned}$$

as given in Sakakihara [16, (3.3)]. Once $\partial u_{n+1}/\partial n$ is solved in (3.7), u_{n+1} on the entire Ω can be obtained by quadratures on Ω and $\partial\Omega$.

The *indirect formulation* is based upon the following ansatz ([6, 9, 10])

$$(3.8) \quad \begin{aligned} u_{n+1}(x) &= (\tilde{S}_\lambda \eta_{n+1})(x) - \tilde{V}_\lambda(u_n)(x) \\ &\equiv \int_{\partial\Omega} E(x-y; \lambda) \eta_{n+1}(y) d\sigma_y - \int_{\Omega} E(x-y; \lambda) F_\lambda(y, u_n(y)) dy, \quad x \in \bar{\Omega}, \end{aligned}$$

where $\eta_{n+1}(\cdot)$ is an unknown boundary layer density defined on $\partial\Omega$ to be solved by the BIE

$$(3.9) \quad \int_{\partial\Omega} E(x-y; \lambda) \eta_{n+1}(y) d\sigma_y = g(x) + \int_{\Omega} E(x-y; \lambda) F_\lambda(y, u_n(y)) dy, \quad x \in \partial\Omega.$$

Comparing (3.9) with (3.7), we see that a large amount of numerical quadrature (involved in the evaluation of $\int_{\partial\Omega} \frac{\partial E}{\partial n_y} g d\sigma_y$) is bypassed. This approach is more advantageous.

Let $a, b \in \mathbb{R}$, $\lambda > 0$ be chosen such that for any $\phi \in C^\infty(\bar{\Omega})$,

$$(3.10) \quad \frac{\partial F(x, \phi(x))}{\partial u} - \lambda < 0 \quad \forall \phi: a \leq \phi(x) \leq b, \quad x \in \bar{\Omega}.$$

For any $\phi \in C^\infty(\bar{\Omega})$, the volume potential in (3.8) defines a continuous mapping

$$\tilde{V}_\lambda: C^\infty(\bar{\Omega}) \rightarrow C^\infty(\bar{\Omega}), \quad \tilde{V}_\lambda(\phi)(x) \equiv \int_{\Omega} E(x-y; \lambda) F_\lambda(y, \phi(y)) dy, \quad x \in \bar{\Omega},$$

cf. [6, p. 216], e.g. By the trace theorem, we have

$$(3.11) \quad \tilde{V}_{\lambda, \Gamma}(\phi) \equiv \tilde{V}_\lambda(\phi)|_{\partial\Omega} \in C^\infty(\partial\Omega).$$

The trace of the simple-layer potential \tilde{S}_λ in (3.8) is denoted by S_λ :

$$(3.12) \quad S_\lambda \eta \equiv \tilde{S}_\lambda \eta|_{\partial\Omega}.$$

It is well known that S_λ has the following regularity:

$$(3.13) \quad \begin{cases} S_\lambda: H^r(\partial\Omega) \rightarrow H^{r+1}(\partial\Omega), \text{ isomorphically, } \forall r \in \mathbb{R}, \\ S_\lambda \text{ is a strongly elliptic pseudodifferential operator of order } -1 \text{ on } \partial\Omega. \end{cases}$$

Now define a mapping \mathcal{T} on $C_{a,b}^\infty(\bar{\Omega}) \equiv \{\phi \in C^\infty(\bar{\Omega}) \mid a \leq \phi(x) \leq b \text{ on } \bar{\Omega}\}$:

$$(3.14) \quad \eta = \mathcal{T}\phi,$$

where η is a function defined on $\partial\Omega$ satisfying the BIE

$$(3.15) \quad \int_{\partial\Omega} E(x-y; \lambda) \eta(y) d\sigma_y = \int_{\Omega} E(x-y; \lambda) F_\lambda(y, \phi(y)) dy, \quad x \in \partial\Omega.$$

By (3.11), and (3.13)–(3.15), we have $\eta \in C^\infty(\partial\Omega)$, and that

$$(3.16) \quad \mathcal{T}: C_{a,b}^\infty(\bar{\Omega}) \rightarrow C^\infty(\partial\Omega)$$

is continuous.

We now establish the main theorem in this section.

Theorem 3.1. *The mapping \mathcal{T} in (3.14) is antimonotone, i.e.,*

$$\phi_1, \phi_2 \in C_{a,b}^\infty(\bar{\Omega}), \quad \phi_1(x) \leq \phi_2(x) \quad \forall x \in \bar{\Omega}$$

implies

$$(3.17) \quad \mathcal{T}\phi_1 \geq \mathcal{T}\phi_2 \quad \text{on } \partial\Omega.$$

Proof. Let

$$(3.18) \quad \eta_i = \mathcal{T}\phi_i, \quad i = 1, 2, \quad \text{on } \partial\Omega.$$

Extend ϕ_1 and ϕ_2 to \mathbb{R}^N by

$$(3.19) \quad \tilde{\phi}_i(x) = \begin{cases} 0, & x \in \Omega^c \equiv \mathbb{R}^N \setminus \overline{\Omega}, \\ \phi_i(x), & x \in \overline{\Omega}, \end{cases} \quad i = 1, 2.$$

Then through a localization procedure ([6, Chap. 6]) it is not difficult to show that

$$(3.20) \quad \tilde{\phi}_i \in H^{\frac{1}{2}-\varepsilon}(\mathbb{R}^N), \quad \text{for any } \varepsilon > 0, i = 1, 2.$$

From the definition of $\tilde{S}_\lambda \eta$ in (3.8), by abuse of notation, extend $\tilde{S}_\lambda \eta$ to all \mathbb{R}^N , i.e.,

$$(\tilde{S}_\lambda \eta)(x) = \int_{\partial\Omega} E(x-y; \lambda) \eta(y) d\sigma_y, \quad x \in \mathbb{R}^N.$$

We also extend \tilde{V}_λ to \mathbb{R}^N by defining

$$(3.21) \quad (\tilde{V}_\lambda \tilde{\phi})(x) = \int_{\Omega} E(x-y; \lambda) \tilde{F}_\lambda(y, \phi(y)) dy, \quad x \in \mathbb{R}^N,$$

where

$$(3.22) \quad \tilde{F}_\lambda(x, \phi(x)) \equiv \begin{cases} 0, & \text{if } x \in \Omega^c, \\ F_\lambda(x, \phi(x)), & \text{if } x \in \Omega. \end{cases}$$

From (3.16), (3.19), (3.20), (3.22) and [6, Chap. 6], we have the regularities

$$(3.23) \quad \eta_i \in C^\infty(\partial\Omega), \quad \tilde{V}_\lambda \tilde{\phi}_i \in H_{\text{loc}}^{\frac{5}{2}-\varepsilon}(\mathbb{R}^N), \quad \text{for any } \varepsilon > 0.$$

Now define

$$(3.24) \quad w_i^\pm(x) = (\tilde{S}_\lambda \eta_i)(x) - (\tilde{V}_\lambda \phi_i)(x), \quad i = 1, 2, \\ \text{“+” if } x \in \Omega^c, \quad \text{“−” if } x \in \Omega.$$

By (3.14) and (3.15), we have $w_i^-|_{\partial\Omega} \equiv 0, i = 1, 2$. Also, on Ω , we have

$$(\Delta - \lambda)w_i^-(x) = F_\lambda(x, \phi_i(x)) = -\lambda\phi_i(x) + F(x, \phi_i(x)), \quad i = 1, 2.$$

Therefore, w_i^- is the (unique) solution of the BVP

$$(3.25) \quad \left. \begin{cases} (\Delta - \lambda)w_i^-(x) = -\lambda\phi_i(x) + F(x, \phi_i(x)) & \text{on } \Omega, \\ w_i^-(x) = 0 & \text{on } \partial\Omega, \end{cases} \right\} \quad i = 1, 2.$$

Similarly, from (3.15) and (3.22)–(3.24), w_i^+ satisfies the exterior BVP

$$(3.26) \quad \left. \begin{cases} (\Delta - \lambda)w_i^+(x) = 0 & \text{on } \Omega^c, \\ w_i^+(x) = 0 & \text{on } \partial\Omega, \end{cases} \right\} \quad i = 1, 2.$$

We now show that w_i^+ , $i = 1, 2$, satisfying (3.24) and (3.26), are trivial. For $x \in \Omega^c$, $|x|$ large, and $y \in \Omega$, we have

$$(3.27) \quad \begin{aligned} |x - y| &= \sqrt{|x|^2 - 2\langle x, y \rangle + |y|^2} \\ &= |x| \left[1 - \frac{1}{2} \left(\frac{2\langle x, y \rangle - |y|^2}{|x|^2} \right) + \frac{1}{2} \left(-\frac{1}{2} \right) \left(\frac{-2\langle x, y \rangle + |y|^2}{|x|^2} \right)^2 \pm \dots \right]. \end{aligned}$$

The fundamental solution $E(x; \lambda)$ in (3.2) is known to have fast decay for $|x|$ large when $\lambda > 0$. For example, when $N = 3$, this is obvious from (3.4). For $N = 2$, we have [1]

$$(3.28) \quad \begin{aligned} E(x, \lambda) &= \frac{1}{2\pi} K_0(\sqrt{\lambda}|x|) \\ &\sim \frac{1}{2} \sqrt{\frac{1}{2\sqrt{\lambda}|x|}} e^{-\sqrt{\lambda}|x|} \left\{ 1 - \frac{1}{8\sqrt{\lambda}|x|} \pm \dots \right\}. \end{aligned}$$

Using (3.27) and (3.28) in (3.24), we obtain

$$(3.29) \quad w_i^+(x) = \begin{cases} \frac{1}{2} \lambda^{-\frac{1}{4}} \sqrt{\frac{1}{2|x|}} [1 + \mathcal{O}(|x|^{-\frac{1}{2}})] \left\{ \int_{\partial\Omega} e^{-\sqrt{\lambda}|x-y|} \eta(y) d\sigma_y \right. \\ \qquad \qquad \qquad \left. - \int_{\Omega} e^{-\sqrt{\lambda}|x-y|} F_{\lambda}(y, \phi_i(y)) dy \right\}, & N = 2; \\ \frac{1}{4\pi} \frac{1}{|x|} [1 + \mathcal{O}(|x|^{-1})] \left\{ \int_{\partial\Omega} e^{-\sqrt{\lambda}|x-y|} \eta(y) d\sigma_y \right. \\ \qquad \qquad \qquad \left. - \int_{\Omega} e^{-\sqrt{\lambda}|x-y|} F_{\lambda}(y, \phi_i(y)) dy \right\}, & N = 3. \end{cases}$$

For space dimension $N > 3$, similar expressions can be obtained. Note that each integral inside the parentheses above decay with an exponential rate $e^{-\sqrt{\lambda}|x|}$ for $|x|$ large.

Let

$$\Omega_R^c = \Omega^c \cap \{x \in \mathbb{R}^N \mid |x| < R\}, \quad \text{for large } R > 0.$$

From (3.26), we obtain

$$\begin{aligned} 0 &= \int_{\Omega^c} [(\Delta - \lambda)w_i^+(x)]w_i^+(x)dx \\ &= \lim_{R \rightarrow \infty} \left[\int_{\partial\Omega_R^c} \frac{\partial w_i^+(x)}{\partial n} w_i^+(x) d\sigma - \int_{\Omega_R^c} (|\nabla w_i^+(x)|^2 + \lambda|w_i^+(x)|^2) dx \right] \\ &= - \int_{\Omega^c} (|\nabla w_i^+(x)|^2 + \lambda|w_i^+(x)|^2) dx, \end{aligned}$$

where the limit of the boundary integral over $\partial\Omega_R^c$ tends to 0 as $R \rightarrow \infty$ because

$$\begin{aligned} \int_{\partial\Omega_R^c} \frac{\partial w_i^+(x)}{\partial n} w_i^+(x) d\sigma &= \int_{\partial\Omega^c} \frac{\partial w_i^+(x)}{\partial n} w_i^+(x) d\sigma + \int_{|x|=R} \frac{\partial w_i^+(x)}{\partial r} w_i^+(x) d\sigma \\ &= \int_{|x|=R} \frac{\partial w_i^+(x)}{\partial r} w_i^+(x) d\sigma \quad (r = |x|), \end{aligned}$$

and the above integral tends to zero as $R \rightarrow \infty$ by (3.29). Hence $w_i^+(x) \equiv 0$ on Ω^c , $i = 1, 2$. Define

$$(3.30) \quad w(x) = \begin{cases} w_2^+(x) - w_1^+(x), & x \in \Omega^c, \\ w_2^-(x) - w_1^-(x), & x \in \Omega. \end{cases}$$

Then

$$(3.31) \quad w(x) \equiv 0 \quad \text{for } x \in \Omega^c.$$

On Ω , the function w satisfies

$$(3.32) \quad \begin{aligned} (\Delta - \lambda)w(x) &= -\lambda[\phi_2(x) - \phi_1(x)] + [F(x, \phi_2(x)) - F(x, \phi_1(x))] \\ &= [\phi_2(x) - \phi_1(x)] \cdot \left[-\lambda + \frac{F(x, \phi_2(x)) - F(x, \phi_1(x))}{\phi_2(x) - \phi_1(x)} \right] \\ &\leq 0 \quad \text{by (3.12),} \end{aligned}$$

and on $\partial\Omega$, it satisfies $w \equiv 0$. Thus,

$$(3.33) \quad \begin{cases} (\Delta - \lambda)w \leq 0 & \text{on } \Omega, \\ w|_{\partial\Omega} = 0 & \text{on } \partial\Omega. \end{cases}$$

If $w(x) < 0$ for some $x \in \Omega$, then

$$(3.34) \quad \min_{x \in \bar{\Omega}} w(x) < 0.$$

Assume that

$$w(x_0) = \min_{x \in \bar{\Omega}} w(x).$$

Then $x_0 \notin \Omega$ by Hopf's boundary point lemma [13]. Thus $x_0 \in \partial\Omega$. But

$$(3.35) \quad 0 > w(x_0) = 0, \quad \text{by the boundary condition in (3.33), a contradiction.}$$

Therefore,

$$(3.36) \quad w(x) \geq 0 \quad \text{on } \Omega.$$

Let $\tilde{V}_\lambda \tilde{\phi}_i$ be the volume potential in (3.23). Since $\tilde{F}_\lambda(\cdot, \phi_i(\cdot)) \in H^{\frac{1}{2}-\varepsilon}(\mathbb{R}^N)$ (just as in (3.20)), by elliptic regularity we have

$$(3.37) \quad \tilde{V}_\lambda \tilde{\phi}_i \in H^{\frac{5}{2}-\varepsilon}(\mathbb{R}^N), \quad i = 1, 2, \text{ for any } \varepsilon > 0.$$

Further, let

$$\begin{aligned} (\tilde{V}_\lambda \tilde{\phi}_i)^+ &= \tilde{V}_\lambda \tilde{\phi}_i(x) & \text{if } x \in \Omega^c, \\ (\tilde{V}_\lambda \tilde{\phi}_i)^-(x) &= \tilde{V}_\lambda \tilde{\phi}_i(x) & \text{if } x \in \Omega. \end{aligned}$$

Then the trace theorem gives

$$\frac{\partial}{\partial n} (\tilde{V}_\lambda \tilde{\phi}_i)^\pm \in H^{1-\varepsilon}(\partial\Omega), \quad \text{for any } 0 < \varepsilon < 1.$$

Choose any $\varepsilon: 0 < \varepsilon < 1$. Then by the continuity of the normal trace operation we have

$$(3.38) \quad \frac{\partial}{\partial n}(\tilde{V}_\lambda \tilde{\phi}_i)^+ - \frac{\partial}{\partial n}(\tilde{V}_\lambda \tilde{\phi}_i)^- = \frac{\partial}{\partial n} \tilde{V}_\lambda \tilde{\phi}_i - \frac{\partial}{\partial n} \tilde{V}_\lambda \tilde{\phi}_i = 0 \quad \text{on } \partial\Omega, i = 1, 2.$$

Also, let

$$(3.39) \quad (\tilde{S}_\lambda \eta_i)^\pm(x) = \int_{\partial\Omega} E(x - \xi; \lambda) \eta_i(\xi) d\sigma_\xi, \quad i = 1, 2,$$

“+” if $x \in \Omega^c$, “-” if $x \in \Omega$.

Then, on $\partial\Omega$,

$$(3.40) \quad \begin{aligned} \frac{\partial}{\partial n}[w^+(x) - w^-(x)] &= - \frac{\partial}{\partial n} w^-(x) \quad (\text{by (3.31)}) \\ &= \lim_{h \downarrow 0} \frac{w^-(x - nh) - w^-(x)}{h} \\ &= \lim_{h \downarrow 0} \frac{w^-(x - nh)}{h} \quad (x \in \partial\Omega) \\ &\geq 0 \quad \text{by (3.36)}. \end{aligned}$$

But from (3.38), we also have

$$\begin{aligned} &\frac{\partial}{\partial n}[w^+(x) - w^-(x)] \\ &= \frac{\partial}{\partial n} \{ [(\tilde{S}_\lambda \eta_2)^+(x) - ((\tilde{V}_\lambda \tilde{\phi}_2)^+(x)) - ((\tilde{S}_\lambda \eta_1)^+(x) - (\tilde{V}_\lambda \tilde{\phi}_1)^+(x))] \\ &\quad - [((\tilde{S}_\lambda \eta_2)^-(x) - (\tilde{V}_\lambda \tilde{\phi}_2)^-(x)) - ((\tilde{S}_\lambda \eta_1)^-(x) - (\tilde{V}_\lambda \tilde{\phi}_1)^-(x))] \} \\ &= \frac{\partial}{\partial n} \{ [(\tilde{S}_\lambda \eta_2)^+(x) - (\tilde{S}_\lambda \eta_1)^+(x)] - [(\tilde{S}_\lambda \eta_2)^-(x) - (\tilde{S}_\lambda \eta_1)^-(x)] \} \\ &= -\eta_2(x) - (-\eta_1(x)) \quad (\text{by [6, Corollary 6.5.1]}) \\ &= \eta_1(x) - \eta_2(x) \geq 0 \quad \text{by (3.40)}. \quad \square \end{aligned}$$

Hence the proof is complete.

Some numerical evidence of Theorem 3.1 is provided in Fig. 7.2.

Corollary 3.2. *Let $g \in C^\infty(\partial\Omega)$ be given. Let (3.10) be satisfied. Define*

$$(3.41) \quad \mathcal{T}_g: C_{a,b}^\infty(\bar{\Omega}) \rightarrow C^\infty(\partial\Omega), \quad \mathcal{T}_g \phi = \eta,$$

where η is the solution of the BIE

$$(3.42) \quad \int_{\partial\Omega} E(x - y; \lambda) \eta_{n+1}(y) d\sigma_y = g(x) + \int_{\Omega} E(x - y; \lambda) F_\lambda(y, \phi(y)) dy, \quad x \in \partial\Omega.$$

Then \mathcal{T}_g is also antimonotone. □

Corollary 3.3. *Let the assumptions of Theorem 2.1 be satisfied. Let the monotone iterations*

$$(3.43) \quad \begin{cases} u_0(x) = \bar{u}(x) \in C^\infty(\bar{\Omega}), \\ (\Delta - \lambda)u_{n+1}(x) = F_\lambda(x, u_n(x)), & x \in \Omega, \\ u_{n+1}(x) = g(x), & x \in \partial\Omega, \end{cases}$$

and

$$(3.44) \quad \begin{cases} v_0(x) = \bar{v}(x) \in C^\infty(\bar{\Omega}), \\ (\Delta - \lambda)v_{n+1}(x) = F_\lambda(x, v_n(x)), & x \in \Omega, \\ v_{n+1}(x) = g(x), & x \in \partial\Omega, \end{cases}$$

have iterative solutions u_{n+1} and v_{n+1} represented, for $n = 0, 1, 2, \dots$, as

$$(3.45) \quad u_{n+1}(x) = \int_{\partial\Omega} E(x-y; \lambda)\mu_{n+1}(y)d\sigma_y - \int_{\Omega} E(x-y; \lambda)F_\lambda(y, u_n(y))dy, \quad x \in \bar{\Omega},$$

$$(3.46) \quad v_{n+1}(x) = \int_{\partial\Omega} E(x-y; \lambda)\nu_{n+1}(y)d\sigma_y - \int_{\Omega} E(x-y; \lambda)F_\lambda(y, v_n(y))dy, \quad x \in \bar{\Omega},$$

where the simple-layer densities $\mu_{n+1}(\cdot)$ and $\nu_{n+1}(\cdot)$ are determined from

$$(3.47) \quad \mu_{n+1} = \mathcal{T}_g u_n, \quad \nu_{n+1} = \mathcal{T}_g v_n, \quad n = 0, 1, 2, \dots$$

Then

$$(3.48) \quad \mu_\infty = \lim_{n \rightarrow \infty} \mu_n, \quad \nu_\infty = \lim_{n \rightarrow \infty} \nu_n$$

exist in $C^\infty(\partial\Omega)$ such that

$$(3.49) \quad \mu_1 \leq \mu_2 \leq \dots \leq \mu_n \leq \dots \rightarrow \mu_\infty \leq \nu_\infty \leftarrow \nu_n \leq \dots \leq \nu_2 \leq \nu_1. \quad \square$$

As in Remark 2.4, we again note that the rate of convergence of (3.48) is not available in general.

4. ERROR ESTIMATES FOR THE GALERKIN BOUNDARY ELEMENT SCHEME

We first recall some basic regularity properties of the operators corresponding to the simple-layer and volume potentials. For $\phi \in C^\infty(\bar{\Omega})$, let $\mathcal{V}_\lambda\phi$ denote the volume potential defined by

$$(4.1) \quad (\mathcal{V}_\lambda\phi)(x) = \int_{\Omega} E(x-y; \lambda)\phi(y)dy, \quad x \in \bar{\Omega}.$$

Then it is known (cf. [6, Theorem 6.3.1], e.g.) that \mathcal{V}_λ can be extended to be an operator such that

$$(4.2) \quad \begin{cases} \mathcal{V}_\lambda: H^r(\Omega) \rightarrow H^{r+2}(\Omega) \text{ continuous, } \forall r \geq -1. \\ \|\mathcal{V}_\lambda\phi\|_{H^{r+2}(\Omega)} \leq \bar{C}_r \|\phi\|_{H^r(\Omega)}, \quad \forall \phi \in H^r(\Omega), \quad r \geq -1, \end{cases}$$

for a constant $\overline{C}_r > 0$ independent of ϕ . The trace of (4.1) induces an operator $\mathcal{V}_{\lambda,\Gamma}$ defined by

$$(4.3) \quad (\mathcal{V}_{\lambda,\Gamma}\phi)(x) \equiv (\mathcal{V}_\lambda\phi)(x), \quad x \in \partial\Omega;$$

it satisfies

$$(4.4) \quad \begin{cases} \mathcal{V}_{\lambda,\Gamma}: H^r(\Omega) \rightarrow H^{r+\frac{3}{2}}(\partial\Omega) \text{ continuous, } \forall r \geq -1, \\ \text{there exists } \overline{C}_r > 0 \text{ such that} \\ \|\mathcal{V}_{\lambda,\Gamma}\phi\|_{H^{r+\frac{3}{2}}(\partial\Omega)} \leq \overline{C}_r \|\phi\|_{H^r(\Omega)}, \quad \forall \phi \in H^r(\Omega), r \geq -1. \end{cases}$$

Note the relationship

$$(4.5) \quad \tilde{V}_\lambda\phi = \mathcal{V}_\lambda F(\cdot, \phi(\cdot))$$

from (3.8).

Let the simple-layer potential \tilde{S}_λ be as defined in (3.8), and let S_λ be the corresponding trace as given in (3.12). It is known ([6, Chap. 6]) that

$$(4.6) \quad \begin{cases} \tilde{S}_\lambda: H^r(\partial\Omega) \rightarrow H^{r+\frac{3}{2}}(\Omega) \text{ continuous, } \forall r \in \mathbb{R}, \\ \text{there exists } \tilde{C}_r > 0, \text{ such that} \\ \|\tilde{S}_\lambda\eta\|_{H^{r+\frac{3}{2}}(\Omega)} \leq \tilde{C}_r \|\eta\|_{H^r(\partial\Omega)}, \quad \forall \eta \in H^r(\partial\Omega). \end{cases}$$

Also, from (3.13), for each $r \in \mathbb{R}$, there exists a $\tilde{\tilde{C}}_r > 0$ such that

$$(4.7) \quad \tilde{\tilde{C}}_r^{-1} \|\eta\|_{H^r(\partial\Omega)} \leq \|S_\lambda\eta\|_{H^{r+1}(\partial\Omega)} \leq \tilde{\tilde{C}}_r \|\eta\|_{H^r(\partial\Omega)}, \quad \forall \eta \in H^r(\partial\Omega).$$

Let $\{\mathcal{S}_h \mid 0 < h \leq h_0\}$ be a 1-parameter family of finite-dimensional boundary element approximation spaces that form an (ℓ, m) -system $S_h^{\ell,m}(\partial\Omega)$, with $\ell, m \in \mathbb{Z}^+ \equiv \{0, 1, 2, \dots\}$, $\ell \geq m + 1$, on $\partial\Omega$, in the sense of Babuška and Aziz [4], satisfying

(1) **Approximation property:**

For each $\phi \in H^t(\partial\Omega)$, there exists a $\phi_h \in \mathcal{S}_h$ such that

$$(4.8) \quad \|\phi - \phi_h\|_{H^s(\partial\Omega)} \leq C_{t,s} h^{t-s} \|\phi\|_{H^t(\partial\Omega)}, \quad \forall h: 0 < h \leq h_0,$$

where $-\ell \leq s \leq t \leq \ell; |s|, |t| \leq m$, and $C_{t,s}$ is a constant independent of h and ϕ .

(2) **Inverse property:**

There exists a constant $M_{s,t} > 0$ such that

$$(4.9) \quad \|\phi_h\|_{H^t(\partial\Omega)} \leq M_{s,t} h^{s-t} \|\phi_h\|_{H^s(\partial\Omega)}, \quad \forall \phi_h \in \mathcal{S}_h, \quad \forall h: 0 < h \leq h_0,$$

where $s \leq t$, and $|s|, |t| \leq m$.

We now implement the monotone iteration scheme according to Corollary 3.3. For definiteness, we will only discretize the (supersolution) systems (3.47) and (3.49). The subsolution systems (3.43) and (3.46) are analogous and thus can

be handled in a nearly identical way. The Galerkin boundary element method can be formulated as follows:

(i) Set

$$(4.10) \quad u_h^0(x) = \bar{u}(x), x \in \bar{\Omega}, \text{ cf. the first equation in (3.43).}$$

(ii) For each $n = 0, 1, 2, \dots$, and $h: 0 < h \leq h_0$, find $\eta_h^{n+1} \in \mathcal{S}_h$ such that

$$(4.11) \quad \langle S_\lambda \eta_h^{n+1}, \phi_h \rangle = \langle g + \mathcal{V}_\lambda F(\cdot, u_h^n), \phi_h \rangle, \quad \forall \phi_h \in \mathcal{S}_h,$$

where $\langle \cdot, \cdot \rangle$ signifies the $L^2(\partial\Omega)$ inner product.

(iii) Define

$$(4.12) \quad u_h^{n+1}(x) = (\tilde{S}_\lambda \eta_h^{n+1})(x) - (\mathcal{V}_\lambda F(\cdot, u_h^n))(x), \quad x \in \bar{\Omega}, \quad n = 0, 1, 2, \dots .$$

Note that (4.11) and (4.12) correspond, respectively, to the discretization of (3.42) and (3.45). We also see that in (4.11) and (4.12), quadrature evaluations are involved in the determination of $\mathcal{V}_\lambda F(\cdot, u_h^n)$ and $\mathcal{V}_{\lambda,\Gamma} F(\cdot, u_h^n)$. Since, in principle, these quadrature evaluations may be carried out to as high an accuracy as desired, we assume that $\mathcal{V}_\lambda F(\cdot, u_h^n)$ and $\mathcal{V}_{\lambda,\Gamma} F(\cdot, u_h^n)$ are *exact* (although in practice, these quadrature evaluations are rather tedious because the integration is carried out on the domain, and numerical errors are involved). From now on, without further mention, we also maintain all the assumptions and notation used in Theorem 2.2.

Our main objective is to estimate $\|u_\infty - u_h^n\|_{H^r(\Omega)}$, where u_∞ is a stable solution as announced in Theorem 2.2(ii). The following working assumption is crucial; it enables us to obtain rates of convergence:

$$[H] \quad \left\{ \begin{array}{l} \text{There exist constants } \gamma_1 \text{ and } \gamma_2 \text{ such that} \\ -\lambda_1 < \gamma_1 \leq \frac{\partial F(x,u)}{\partial u} \leq \gamma_2, \quad \forall (x,u) \in \bar{Q}, \quad \bar{Q} \equiv \bar{\Omega} \times [a,b], \\ \text{where } \lambda_1 > 0 \text{ is the smallest eigenvalue of } -\Delta \text{ on } \Omega \text{ subject} \\ \text{to the homogeneous Dirichlet boundary condition. } \square \end{array} \right.$$

Because $\frac{\partial F}{\partial u}$ is continuous, it easily follows from [H] that for a sufficiently small $\delta > 0$, there exist $\tilde{\gamma}_1$ and $\tilde{\gamma}_2$ such that

$$(4.13) \quad -\lambda_1 < \tilde{\gamma}_1 \leq \frac{\partial F(x,u)}{\partial u} \leq \tilde{\gamma}_2, \quad \forall (x,u) \in \bar{Q}_\delta, \quad \bar{Q}_\delta \equiv \bar{\Omega} \times [a-\delta, b+\delta].$$

This small δ will provide the “breathing room” to accommodate the errors in the numerical operation for the iterates to stay within the range of monotone iteration.

Lemma 4.1. *Let u_{n+1} be iterated according to (3.43) and let u_∞ be the limit as in (3.48). Denote $e^n = u_\infty - u_n$. Then we have*

$$(4.14) \quad \|e^{n+1}\|_{L^2(\Omega)} \leq \alpha_\lambda^{n+1} \|e^0\|_{L^2(\Omega)}, \quad n = 0, 1, 2, \dots ,$$

where

$$(4.15) \quad 0 < \alpha_\lambda \equiv \frac{\lambda - \tilde{\gamma}_1}{\lambda + \lambda_1} < 1.$$

Proof. Since u_∞ and u_{n+1} satisfy, respectively, (2.6) (with $Bu \equiv u$) and (3.43), we obtain by subtraction

$$(4.16) \quad \begin{cases} (\Delta - \lambda)e^{n+1}(x) = F_\lambda(x, u_\infty(x)) - F_\lambda(x, u_n(x)), & x \in \Omega, \\ e^{n+1}(x) = 0, & x \in \partial\Omega. \end{cases}$$

Multiplying the above by $-e^{n+1}$, integrating by parts and using $\lambda_1 \|e^{n+1}\|^2 \leq \|\nabla e^{n+1}\|^2$, we get

$$(4.17) \quad \|e^{n+1}\| \leq \frac{1}{\lambda + \lambda_1} \|F_\lambda(\cdot, u_\infty) - F_\lambda(\cdot, u_n)\|,$$

where all the norms without subscripts are $L^2(\Omega)$. Let

$$(4.18) \quad M_\lambda \equiv \sup_{(x,u) \in \bar{Q}_\delta} \left[\lambda - \frac{\partial F(x, u)}{\partial u} \right].$$

Then

$$(4.19) \quad 0 < M_\lambda < \lambda - \tilde{\gamma}_1.$$

(Actually, $\tilde{\gamma}_1$ can be taken to be γ_1 here.) We get

$$(4.20) \quad \begin{aligned} \|F_\lambda(\cdot, u_\infty) - F_\lambda(\cdot, u_n)\| &= \left\| - \left[\lambda - \frac{F(\cdot, u_\infty) - F(\cdot, u_n)}{u_\infty - u_n} \right] (u_\infty - u_n) \right\| \\ &\leq M_\lambda \|u_\infty - u_n\|. \end{aligned}$$

Substituting (4.20) into (4.17), we obtain

$$\|e^{n+1}\| \leq \frac{M_\lambda}{\lambda + \lambda_1} \|e^n\| \leq \alpha_\lambda \|e^n\|, \quad \alpha_\lambda \equiv \frac{\lambda - \tilde{\gamma}_1}{\lambda + \lambda_1} < 1.$$

Hence (4.14) follows. □

Lemma 4.2. Let η_h^{n+1} and u_h^{n+1} be defined as in (4.11) and (4.12) for $n = 0, 1, 2, \dots$. Let ζ_h^{n+1} , defined on $\partial\Omega$, be the (unique) solution of the BIE

$$(4.21) \quad (S_\lambda \zeta_h^{n+1})(x) = g(x) + \mathcal{V}_{\lambda, \Gamma} F(x, u_h^n(x)), \quad x \in \partial\Omega.$$

Then there exists $\tilde{C}_{t,s} > 0$, depending only on s and t , such that

$$(4.22) \quad \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^s(\partial\Omega)} \leq \tilde{C}_{t,s} h^{t-s} \|\zeta_h^{n+1}\|_{H^t(\partial\Omega)},$$

where $m \leq \ell - 1$, $-(\ell + 1) \leq s \leq t \leq \ell$, $s \leq m$, $-\ell \leq -\frac{1}{2} \leq m$ and $-m \leq -\frac{1}{2} \leq t$.

Proof. Since S_λ satisfies (4.7), according to Hsiao and Wendland [10], the order of S_λ is $2\alpha = -1$, i.e., $\alpha = -\frac{1}{2}$. Invoking [10, Corollary 2.1], we obtain the conclusion. □

Lemma 4.3. *Let $N = 2$ or 3 , and let*

$$(4.23) \quad \varepsilon_0(N) = \begin{cases} 0, & \text{if } N = 2, \\ \text{an arbitrarily small positive number,} & \text{if } N = 3. \end{cases}$$

Then for $\ell \geq 2$ and $m \geq 1$, there exist positive constants $K_{\varepsilon_0(N)}$ and K such that

$$(4.24) \quad \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{\varepsilon_0(N)}(\partial\Omega)} \leq K_{\varepsilon_0(N)} h^{\frac{1}{2}-\varepsilon_0(N)} [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} + \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}],$$

$$(4.25) \quad \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{-\frac{3}{2}}(\partial\Omega)} \leq Kh^2 [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} + \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}],$$

$\forall n = 0, 1, 2, \dots, 0 < h \leq h_0.$

Proof. We have from (4.22) and property (4.7) of S_λ ,

$$(4.26) \quad \begin{aligned} \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{\varepsilon_0(N)}(\partial\Omega)} &\leq \tilde{C}_{\frac{1}{2}, \varepsilon_0(N)} h^{\frac{1}{2}-\varepsilon_0(N)} \cdot \tilde{C}_{\frac{1}{2}} \|S_\lambda \zeta_h^{n+1}\|_{H^{\frac{3}{2}}(\partial\Omega)} \\ &\leq (\tilde{C}_{\frac{1}{2}, \varepsilon_0(N)} \tilde{C}_{\frac{1}{2}}) h^{\frac{1}{2}-\varepsilon_0(N)} [\|g\|_{H^{\frac{3}{2}}(\partial\Omega)} + \|\mathcal{V}_{\lambda, \Gamma} F_\lambda(\cdot, u_h^n)\|_{H^{\frac{3}{2}}(\partial\Omega)}] \quad (\text{by (4.21)}) \\ &\leq (\tilde{C}_{\frac{1}{2}, \varepsilon_0(N)} \tilde{C}_{\frac{1}{2}}) h^{\frac{1}{2}-\varepsilon_0(N)} [\|g\|_{H^{\frac{3}{2}}(\partial\Omega)} + \bar{C}_0 \|F_\lambda(\cdot, u_h^n)\|_{L^2(\Omega)}] \quad (\text{by (4.4)}) \\ &\leq (\tilde{C}_{\frac{1}{2}, \varepsilon_0(N)} \tilde{C}_{\frac{1}{2}}) h^{\frac{1}{2}-\varepsilon_0(N)} [\|g\|_{H^{\frac{3}{2}}(\partial\Omega)} + \bar{C}_0 (\|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)} \\ &\quad + \|F_\lambda(\cdot, u_n) - F_\lambda(\cdot, u_\infty)\|_{L^2(\Omega)} + \|F_\lambda(\cdot, u_\infty)\|_{L^2(\Omega)})] \\ &\leq (\tilde{C}_{\frac{1}{2}, \varepsilon_0(N)} \tilde{C}_{\frac{1}{2}}) h^{\frac{1}{2}-\varepsilon_0(N)} [C \|u_\infty\|_{H^2(\Omega)} + \bar{C}_0 M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} \\ &\quad + \bar{C}_0 \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}], \end{aligned}$$

where in the last inequality we have utilized the properties that u_∞ satisfies (2.6), the trace theorem, (4.14) and (4.20). Therefore, (4.24) has been established. From (4.22), we have

$$\begin{aligned} \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{-\frac{3}{2}}(\partial\Omega)} &\leq \tilde{C}_{\frac{1}{2}, -\frac{3}{2}} h^2 \|\zeta_h^{n+1}\|_{H^{\frac{1}{2}}(\partial\Omega)} \\ &\leq \tilde{C}_{\frac{1}{2}, -\frac{3}{2}} h^2 \cdot \tilde{C}_{\frac{1}{2}} \|S_\lambda \zeta_h^{n+1}\|_{H^{\frac{3}{2}}(\partial\Omega)}. \end{aligned}$$

The rest is the same as in (4.26). □

Remark 4.4. Lemma 4.3 remains valid even if $\varepsilon_0(N) = 0$ when $N = 3$, where the constant $K_{\varepsilon_0(N)}$ remains bounded as $\varepsilon_0(N) \downarrow 0$. But in the discussions henceforth we will only utilize the part when $\varepsilon_0(N)$ is positive yet nonvanishingly small. □

From (4.21), we now define

$$(4.27) \quad w_h^{n+1}(x) = (\tilde{S}_\lambda \zeta_h^{n+1})(x) - (\mathcal{V}_\lambda F(\cdot, u_h^n))(x), \quad x \in \Omega, n = 0, 1, 2, \dots, 0 < h \leq h_0.$$

It is readily seen that w_h^{n+1} satisfies

$$(4.28) \quad \begin{cases} (\Delta - \lambda)w_h^{n+1} = F_\lambda(\cdot, u_h^n) & \text{on } \Omega, \\ w_h^{n+1} = g & \text{on } \partial\Omega. \end{cases}$$

Lemma 4.5. *Let $N = 2$ or 3 . Let $C^{0,\beta}(\overline{\Omega})$ be the Hölder space of continuous functions with exponent β , $0 < \beta < 1$, on $\overline{\Omega}$. Then for $\varepsilon_0(N)$ satisfying (4.23), and for β sufficiently small, we have a positive constant $K'_{\varepsilon_0(N)}$ such that*

$$(4.29) \quad \|u_h^{n+1} - w_h^{n+1}\|_{C^{0,\beta}(\overline{\Omega})} \leq K'_{\varepsilon_0(N)} h^{\frac{1}{2} - \varepsilon_0(N)} [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} + \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u^n)\|_{L^2(\Omega)}],$$

for $n = 0, 1, 2, \dots, 0 < h \leq h_0$, where $K'_{\varepsilon_0(3)}$ may grow unbounded as $\varepsilon_0(3) \downarrow 0$ when $N = 3$.

Proof. We use the Sobolev imbedding theorem (cf. [6, Theorem 2.1.3], e.g.)

$$(4.30) \quad \|f\|_{C^{0,\beta}(\overline{\Omega})} \leq C \|f\|_{H^s(\Omega)}, \quad \text{for } 0 < \beta = s - \frac{N}{2} < 1,$$

to get

$$(4.31) \quad \|u_h^{n+1} - w_h^{n+1}\|_{C^{0,\beta}(\overline{\Omega})} \leq c_{\varepsilon_0} \|u_h^{n+1} - w_h^{n+1}\|_{H^{\frac{3}{2} + \varepsilon_0(N)}(\Omega)},$$

where

$$(4.32) \quad \left. \begin{array}{l} 0 < \beta < \frac{1}{2} \\ \beta = \varepsilon_0(3) \end{array} \right\} \quad \text{if} \quad \begin{cases} N = 2, \\ N = 3. \end{cases}$$

Note that $c_{\varepsilon_0} \uparrow \infty$ as $\varepsilon_0(3) \downarrow 0$. Continuing from the right-hand side of (4.31), we have

$$\begin{aligned} \|u_h^{n+1} - w_h^{n+1}\|_{C^{0,\beta}(\overline{\Omega})} &\leq c_{\varepsilon_0} \|\tilde{S}_\lambda(\eta_h^{n+1} - \zeta_h^{n+1})\|_{H^{\frac{3}{2} + \varepsilon_0(N)}(\Omega)} \quad (\text{by (4.12) and (4.27)}) \\ &\leq c_{\varepsilon_0} \cdot \tilde{C}_{\varepsilon_0(N)} \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{\varepsilon_0(N)}} \quad (\text{by (4.6)}). \end{aligned}$$

The rest of the proof follows from Lemma 4.3. \square

Lemma 4.6. *Let $N = 2$ or 3 . Then there exists a constant $K > 0$ such that*

$$(4.33) \quad \|w_h^{n+1} - u_{n+1}\|_{C^{0,\beta}(\overline{\Omega})} \leq K \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)} \\ \forall n = 0, 1, 2, \dots, 0 < h \leq h_0, \text{ where } 0 < \beta < 1/2.$$

Proof. From (3.45), (4.27), (4.30) and (4.32), we get

$$\begin{aligned} \|w_h^{n+1} - u_{n+1}\|_{C^{0,\beta}(\overline{\Omega})} &\leq c \|w_h^{n+1} - u_{n+1}\|_{H^2(\Omega)} \quad (\text{for } 0 < \beta < \frac{1}{2}, \text{ by (4.30)}) \\ &\leq c [\|\tilde{S}_\lambda(\zeta_h^{n+1} - \mu_{n+1})\|_{H^2(\Omega)} + \|\mathcal{V}_\lambda(F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n))\|_{H^2(\Omega)}] \\ &\leq c [\tilde{C}_{\frac{1}{2}} \tilde{C}_{\frac{1}{2}} \|S_\lambda(\zeta_h^{n+1} - \mu_{n+1})\|_{H^{\frac{3}{2}}(\partial\Omega)} \\ &\quad + \|\mathcal{V}_\lambda(F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n))\|_{H^2(\Omega)}] \quad (\text{by (4.6), (4.7)}) \\ &\leq c [\tilde{C}_{\varepsilon_0(N)} \tilde{C}_{\varepsilon_0(N)} \|\mathcal{V}_{\lambda,\Gamma}(F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n))\|_{H^{\frac{3}{2}}(\partial\Omega)} \\ &\quad + \|\mathcal{V}_\lambda(F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n))\|_{H^2(\Omega)}] \\ &\leq c' \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)} \quad (\text{by (4.2) and (4.4)}). \square \end{aligned}$$

Similar to Lemmas 4.5 and 4.6, we give the parallel estimates in the $L^2(\Omega)$ norms in Lemma 4.7.

Lemma 4.7. *We have*

$$(4.34) \quad \|u_h^{n+1} - w_h^{n+1}\|_{L^2(\Omega)} \leq Kh^2[\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} \\ + \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}], \text{ for some } K > 0,$$

$$(4.35) \quad \|w_h^{n+1} - u_{n+1}\|_{L^2(\Omega)} \leq \frac{1}{\lambda + \lambda_1} \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}, \\ \forall n = 0, 1, 2, 3, \dots, 0 < h \leq h_0.$$

Proof. From (4.12) and (4.27),

$$\|u_h^{n+1} - w_h^{n+1}\|_{L^2(\Omega)} = \|\tilde{S}_\lambda(\eta_h^{n+1} - \zeta_h^{n+1})\|_{L^2(\Omega)} \\ \leq \tilde{C}_{\frac{3}{2}} \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{-\frac{3}{2}}(\partial\Omega)} \quad (\text{by (4.6)}).$$

The rest follows from (4.32).

To show (4.35), we use that u_{n+1} and w_h^{n+1} satisfy, respectively, (3.43) and (4.28). Subtracting (4.28) from (3.43) and estimating as in (4.16) and (4.17), we obtain (4.35). \square

Corollary 4.8. (i) *Let $N = 2$ or 3 . Then for $\varepsilon_0(N)$ satisfying (4.23) and for $\beta > 0$ sufficiently small, with β being related to $\varepsilon_0(N)$ through (4.32), we have a constant $K'_{\varepsilon_0(N)} > 0$ such that*

$$(4.36) \quad \|u_h^{n+1} - u_{n+1}\|_{C^{0,\beta}(\bar{\Omega})} \leq K'_{\varepsilon_0(N)} h^{\frac{1}{2} - \varepsilon_0(N)} [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)}] \\ + \left[\frac{1}{\lambda + \lambda_1} + K'_{\varepsilon_0(N)} h^{\frac{1}{2} - \varepsilon_0(N)} \right] \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)},$$

$\forall n = 0, 1, 2, \dots, 0 < h \leq h_0$, where $K'_{\varepsilon_0(N)}$ may grow unbounded as $\varepsilon_0(3) \downarrow 0$.

(ii) *There exists a constant $K > 0$ such that*

$$(4.37) \quad \|u_h^{n+1} - u_{n+1}\|_{L^2(\Omega)} \leq Kh^2[\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)}] + \left(\frac{1}{\lambda + \lambda_1} + Kh^2 \right) \\ \cdot \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}, \\ \forall n = 0, 1, 2, \dots, 0 < h \leq h_0.$$

Proof. The inequality (4.36) follows from an application of the triangle inequality to (4.29) and (4.33), while (4.37) follows likewise from (4.34) and (4.35). \square

Theorem 4.9. *Let $N = 2$ or 3 and let $\varepsilon_0(N)$ be given by (4.23) and $\beta > 0$ be sufficiently small, with β being related to $\varepsilon_0(N)$ through (4.32). Then there exist $\bar{h}_0 : 0 < \bar{h}_0 \leq h_0$ and $\rho_\lambda : 0 < \rho_\lambda < 1$, independent of h and n , such that*

$$(4.38) \quad \|u_h^{n+1} - u_{n+1}\|_{C^{0,\beta}(\bar{\Omega})} \leq K''_{\varepsilon_0(N)} h^{\frac{1}{2} - \varepsilon_0(N)} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^{n+1} \|e^0\|_{L^2(\Omega)}],$$

$$(4.39) \quad \|u_h^{n+1} - u_{n+1}\|_{L^2(\Omega)} \leq \bar{K} h^2 [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^{n+1} \|e^0\|_{L^2(\Omega)}],$$

and

$$(4.40) \quad a - \delta \leq u_h^{n+1}(x) \leq b + \delta, \quad \forall n = 0, 1, 2, \dots, 0 < h \leq \bar{h}_0,$$

where $K''_{\varepsilon_0(N)}$ and \bar{K} are constants independent of h and n .

Proof. Since $0 < \alpha_\lambda < 1$, we can choose $\bar{h}_0 > 0$ so small that

$$(4.41) \quad 0 < \alpha_\lambda + KM_\lambda \bar{h}_0^2 \leq \rho_\lambda < 1,$$

$$(4.42) \quad [K'_{\varepsilon_0(N)} \bar{h}_0^{\frac{1}{2} - \varepsilon_0(N)} + (\alpha_\lambda + K'_{\varepsilon_0(N)} M_\lambda \bar{h}_0^{\frac{1}{2} - \varepsilon_0(N)}) \left(\frac{K}{1 - \rho_\lambda} \right) \bar{h}_0^2] \\ \cdot [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \|e^0\|_{L^2(\Omega)}] \leq \delta,$$

for some ρ_λ : $0 < \rho_\lambda < 1$. By induction, we may assume

$$u_h^k(x) \in [a - \delta, b + \delta], \quad \forall x \in \bar{\Omega}, \quad k = 0, 1, 2, \dots, n, \quad 0 < h \leq \bar{h}_0.$$

Hence, similar to (4.18)–(4.20), we have

$$(4.43) \quad \|F(\cdot, u_h^k) - F(\cdot, u_k)\|_{L^2(\Omega)} \leq M_\lambda \|u_h^k - u_k\|_{L^2(\Omega)}, \quad k = 0, 1, 2, \dots, n, \quad 0 < h \leq \bar{h}_0.$$

Using (4.41) and (4.43) in (4.37), we obtain

$$(4.44) \quad \|u_h^{n+1} - u_{n+1}\|_{L^2(\Omega)} \\ \leq Kh^2[\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)}] \\ + \rho_\lambda \{Kh^2[\|u_\infty\|_{H^2(\Omega)} + M_\lambda \alpha_\lambda^{n-1} \|e^0\|_{L^2(\Omega)}] + \rho_\lambda \|u_h^{n-1} - u_{n-1}\|_{L^2(\Omega)}\} \\ \leq \dots \\ \leq Kh^2(1 + \rho_\lambda + \dots + \rho_\lambda^n) \|u_\infty\|_{H^2(\Omega)} \\ + KM_\lambda h^2(\alpha_\lambda^n + \rho_\lambda \alpha_\lambda^{n-1} + \dots + \rho_\lambda^{n-1} \alpha_\lambda + \rho_\lambda^n) \|e^0\|_{L^2(\Omega)},$$

where in the last inequality, we have used $u_h^0 = u_0 = \bar{u}$. Since $\alpha_\lambda < 1$, we derive

$$(4.45) \quad \|u_h^{n+1} - u_{n+1}\|_{L^2(\Omega)} \leq \left(\frac{K}{1 - \rho_\lambda} \right) h^2 [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \|e^0\|_{L^2(\Omega)}], \\ \forall n = 0, 1, 2, \dots, \quad 0 < h \leq \bar{h}_0.$$

By applying (4.43) and (4.45) to (4.36), we then obtain

$$(4.46) \quad \|u_h^{n+1} - u_{n+1}\|_{C^{0,\beta}(\bar{\Omega})} \leq \bar{K}_{\varepsilon_0(N)} h^{\frac{1}{2} - \varepsilon_0(N)} [\|u_\infty\|_{H^2(\Omega)} + M_\lambda \|e^0\|_{L^2(\Omega)}], \\ \forall n = 0, 1, 2, \dots, \quad 0 < h \leq \bar{h}_0,$$

where

$$(4.47) \quad \bar{K}_{\varepsilon_0(N)} = K'_{\varepsilon_0(N)} + (\alpha_\lambda + K'_{\varepsilon_0(N)} M_\lambda \bar{h}_0^{\frac{1}{2} - \varepsilon_0(N)}) \left(\frac{K}{1 - \rho_\lambda} \right) \bar{h}_0^{\frac{3}{2} + \varepsilon_0(N)}.$$

Thus, (4.40) is obtained from (4.42), (4.46), (4.47) and the induction hypothesis.

To get (4.39), we note that $\alpha_\lambda < \rho_\lambda$ from (4.41). Then, instead of (4.45), we obtain from (4.44)

$$(4.48) \quad \begin{aligned} \|u_h^{n+1} - u_{n+1}\|_{L^2(\Omega)} &\leq \left(\frac{K}{1 - \rho_\lambda} \right) h^2 \|u_\infty\|_{H^2(\Omega)} + K M_\lambda \left(\frac{\rho_\lambda^{n+1}}{\rho_\lambda - \alpha_\lambda} \right) h^2 \|e^0\|_{L^2(\Omega)}, \\ &\leq \left(\frac{K}{1 - \rho_\lambda} \right) h^2 \|u_\infty\|_{H^2(\Omega)} + \left(\frac{1}{\bar{h}_0^2} \right) \rho_\lambda^{n+1} h^2 \|e^0\|_{L^2(\Omega)} \\ &\quad \forall n = 0, 1, 2, \dots, \quad 0 < h \leq \bar{h}_0, \end{aligned}$$

where we used (4.41). Thus, (4.39) follows from (4.48) by setting $\bar{K} = \max\{\frac{K}{1 - \rho_\lambda}, \frac{1}{\bar{h}_0^2}\}$.

Finally, by using (4.40), (4.41) and then (4.48), we obtain (4.38) from (4.36). The proof is complete. \square

The inequality (4.40) is important because it guarantees that each numerical iterate $u_h^n(x)$ does not fall out of the range of validity, $\Omega \times [a - \delta, b + \delta]$, of the monotone iteration, and for the applicability of (4.18)–(4.20). It also partially explains why we need an assumption like [H].

Theorem 4.10. *Let $\{\mathcal{S}_h \mid 0 < h \leq h_0\}$ be a family of (ℓ, m) -systems on $\partial\Omega$ with $\ell \geq 2$ and $m \geq 1$, and let $N = 2$ or 3 . Assume [H], so that there exists a small number $\delta > 0$ for which (4.13) holds. Then there exist constants $K_1 > 0$, $K_2 > 0$, and an \bar{h}_0 : $0 < \bar{h}_0 \leq h_0$, such that for any r : $0 \leq r \leq 2$, we have*

$$(4.49) \quad \|u_h^n - u_\infty\|_{H^r(\Omega)} \leq K_1 h^{2-r} \|u_\infty\|_{H^2(\Omega)} + K_2 \rho_\lambda^n \|e^0\|_{L^2(\Omega)}$$

and

$$a - \delta \leq u_h^n(x) \leq b + \delta,$$

for all $n = 0, 1, 2, \dots$, h : $0 < h \leq \bar{h}_0$, where $e^0 = u_\infty - \bar{u}$ and $0 < \rho_\lambda < 1$. Therefore, u_h^n converges to u_∞ as n tends to ∞ and h tends to 0.

Proof. Similar to the procedures in (4.26), by using (4.39) and (4.43), we have

$$(4.50) \quad \begin{aligned} \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^s(\partial\Omega)} &\leq \tilde{C}_{\frac{1}{2},s} h^{\frac{1}{2}-s} \|\zeta_h^{n+1}\|_{H^{\frac{1}{2}}(\partial\Omega)} \\ &\leq (\tilde{C}_{\frac{1}{2},s} \tilde{C}_{\frac{1}{2}}) h^{\frac{1}{2}-s} [C \|u_\infty\|_{H^2(\Omega)} + \bar{C}_0 M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} \\ &\quad + \bar{C}_0 \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)}] \\ &\leq (\tilde{C}_{\frac{1}{2},s} \tilde{C}_{\frac{1}{2}}) h^{\frac{1}{2}-s} [C \|u_\infty\|_{H^2(\Omega)} + \bar{C}_0 M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} \\ &\quad + \bar{C}_0 M_\lambda \bar{K} h^2 (\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)})] \\ &\leq C_1 h^{\frac{1}{2}-s} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}], \end{aligned}$$

where $-(\ell + 1) \leq s \leq 1/2$. From here we then have, cf. the proof of Lemma 4.7,

$$(4.51) \quad \begin{aligned} \|u_h^{n+1} - w_h^{n+1}\|_{H^{s+\frac{3}{2}}(\Omega)} &= \|\tilde{S}_\lambda(\eta_h^{n+1} - \zeta_h^{n+1})\|_{H^{s+\frac{3}{2}}(\Omega)} \leq \tilde{C}_s \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^s(\partial\Omega)} \\ &\leq C_2 h^{\frac{1}{2}-s} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}] \quad (\text{by (4.50)}). \end{aligned}$$

Let $r = s + \frac{3}{2}$ in (4.51). Then

$$(4.52) \quad \|u_h^{n+1} - w_h^{n+1}\|_{H^r(\Omega)} \leq C_2 h^{2-r} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}],$$

where $-\ell + \frac{1}{2} \leq r \leq 2$.

For $0 \leq r \leq 2$, we have

$$(4.53) \quad \begin{aligned} \|w_h^{n+1} - u_{n+1}\|_{H^r(\Omega)} &\leq \|w_h^{n+1} - u_{n+1}\|_{H^2(\Omega)} \\ &\leq \bar{C}_0 \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)} \quad (\text{by (2.13) and (4.28)}) \\ &\leq \bar{C}_0 M_\lambda \|u_h^n - u_n\|_{L^2(\Omega)} \quad (\text{by (4.43)}) \\ &\leq (\bar{C}_0 M_\lambda K) h^2 [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}] \quad (\text{by (4.39)}). \end{aligned}$$

Combining (4.52) and (4.53), we obtain

$$(4.54) \quad \begin{aligned} \|u_h^{n+1} - u_{n+1}\|_{H^r(\Omega)} &\leq C_3 h^{2-r} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}], \quad 0 \leq r \leq 2, \\ &\quad \forall n = 0, 1, 2, \dots, 0 < h \leq \bar{h}_0. \end{aligned}$$

Also,

$$(4.55) \quad \begin{aligned} \|u_{n+1} - u_\infty\|_{H^r(\Omega)} &\leq \|u_{n+1} - u_\infty\|_{H^2(\Omega)} \\ &\leq C_\lambda \|F_\lambda(\cdot, u_h) - F_\lambda(\cdot, u_\infty)\|_{L^2(\Omega)} \quad (\text{by (4.16)}) \\ &\leq C_\lambda M_\lambda \|u_n - u_\infty\|_{L^2(\Omega)} \\ &\leq C_\lambda M_\lambda \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} \quad (\text{by (4.14)}). \end{aligned}$$

Then, upon noting $\alpha_\lambda < \rho_\lambda$, we see that (4.49) follows from (4.54) and (4.55). \square

Let μ_∞ be the limit of the simple-layer densities $\{\mu_n\}_1^\infty$ given in Corollary 3.3. We give the error estimate $\|\eta_h^n - \mu_\infty\|$ in the following theorem.

Theorem 4.11. *Let $N = 2$ or 3 . Let $\{\mathcal{S}_h \mid 0 < h \leq h_0\}$ be a family of (ℓ, m) -systems with $\ell \geq 2$ and $m \geq 1$. Then for s : $-\frac{3}{2} \leq s \leq \frac{1}{2}$, there are constants $K_1 > 0$ and $K_2 > 0$ such that*

$$(4.56) \quad \begin{aligned} \|\eta_h^n - \mu_\infty\|_{H^s(\partial\Omega)} &\leq K_1 h^{\frac{1}{2}-s} \|u_\infty\|_{H^2(\Omega)} + K_2 \rho_\lambda^n \|e^0\|_{L^2(\Omega)}, \\ &\quad \forall n = 0, 1, 2, \dots, \quad h: 0 < h \leq \bar{h}_0. \end{aligned}$$

Proof. First, from (4.21) and (3.47), we have that for $s \leq 1/2$,

$$\begin{aligned}
 (4.57) \quad & \|\zeta_h^{n+1} - \mu_n\|_{H^s(\partial\Omega)} = \|S_\lambda^{-1}[\mathcal{V}_{\lambda,\Gamma}(F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n))]\|_{H^s(\partial\Omega)} \\
 & \leq \tilde{C}_{s+1} \|\mathcal{V}_{\lambda,\Gamma}[F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_n)]\|_{H^{\frac{3}{2}}(\partial\Omega)} \\
 & \hspace{15em} \text{(for any } s \in \mathbb{R}, \text{ by (4.7))} \\
 & \leq K \|u_h^n - u_n\|_{L^2(\Omega)} \quad \text{(by (4.2) and (4.43))} \\
 & \leq K'h^2[\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}], \text{ from (4.39).}
 \end{aligned}$$

Also, from (4.50),

$$(4.58) \quad \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^s(\partial\Omega)} \leq K'' h^{\frac{1}{2}-s} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)}], \quad -(\ell + 1) \leq s \leq \frac{1}{2}.$$

Similarly, from (3.47), (3.48), and the proofs of (4.57) and Lemma 4.1, we have

$$(4.59) \quad \|\mu_n - \mu_\infty\|_{H^s(\partial\Omega)} \leq K''' \alpha_\lambda^n \|e^0\|_{L^2(\Omega)} \quad \text{for } s \leq \frac{1}{2}.$$

From (4.57)–(4.59) and the triangle inequality, (4.56) follows. \square

In our proofs above, for the $S_h^{\ell,m}(\partial\Omega)$ -systems we have used, because of the need of the inverse inequality we require that $|\alpha| \leq m$ (cf. the proof of Lemma 4.2 for α), $\alpha = -\frac{1}{2}$, leading to $m \geq 1$. Thus piecewise constant boundary element spaces $S_h^{1,0}(\partial\Omega)$ are excluded. However, this restriction may be relaxed for $N = 2$ if we use S_h^d , the space of smoothest splines of degree d with respect to a quasi-uniform mesh on $\partial\Omega$, which is a smooth Jordan curve in \mathbb{R}^2 . In this case the approximation property (4.8) holds for all $s \leq t \leq d + 1$, $s < d + \frac{1}{2}$, and the inverse property (4.9) holds for $s \leq t < d + \frac{1}{2}$ [12]. By [14, Corollary 4], in (4.24) we now have

$$\begin{aligned}
 (4.60) \quad & \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^s(\partial\Omega)} \leq Ch^{t-s} \|\zeta_h^{n+1}\|_{H^t(\partial\Omega)} - (d + 2) \leq s \leq t \leq d + 1, \\
 & \hspace{10em} s < d + \frac{1}{2}, \quad t > -\left(d + \frac{3}{2}\right), \quad d \geq 0.
 \end{aligned}$$

Therefore, all of the lemmas and theorems after Lemma 4.2 in this section remain valid for these S_h^d systems with a quasiuniform mesh on $\partial\Omega$ when $N = 2$. In particular, we state the following.

Theorem 4.12. *Let $N = 2$ and let $\{\mathcal{S}_h \mid 0 < h \leq h_0\}$ be a family of S_h^d spaces of smoothest splines of degree $d \in \mathbb{Z}^+$ with respect to a quasiuniform mesh on $\partial\Omega$. Under otherwise the same assumptions as in Theorems 4.10 and 4.11, we have $K_i > 0$, $i = 1, 2, 3, 4$, and $\bar{h}_0: 0 < \bar{h}_0 \leq h_0, 0 < \rho_\lambda < 1$, such that*

$$\begin{aligned}
 \|u_h^n - u_\infty\|_{H^2(\Omega)} & \leq K_1 h^{2-r} \|u_\infty\|_{H^2(\Omega)} + K_2 \rho_\lambda^n \|e^0\|_{L^2(\Omega)}, \\
 \|\eta_h^n - u_\infty\|_{H^s(\partial\Omega)} & \leq K_3 h^{\frac{1}{2}-s} \|u_\infty\|_{H^2(\Omega)} + K_4 \rho_\lambda^n \|e^0\|_{L^2(\Omega)} \\
 & \hspace{15em} (K_3 \text{ and } K_4 \text{ depend on } s),
 \end{aligned}$$

$\forall r: 0 \leq r \leq 2, \quad n = 0, 1, 2, \dots, \quad h: 0 < h \leq \bar{h}_0; \quad s: -(d + 2) \leq s < 1/2. \quad \square$

Remark 4.13. The error estimates obtained in Theorems 4.10–4.12 are all of the separable form

$$K_1 h^\gamma \|u_\infty\|_{H^2(\Omega)} + K_2 \rho_\lambda^n \|e^0\|_{L^2(\Omega)}, \quad \gamma > 0.$$

It is obvious that these estimates are asymptotically tight. \square

Remark 4.14. Error estimates of $\|u_h^n - u_\infty\|_{H^r(\Omega)}$ with respect to higher-order Sobolev space norms $H^r(\Omega)$, with $r > 2$, and for space dimension $N > 3$, are possible if we make an assumption like

$$(4.61) \quad \|F_\lambda(\cdot, u_1(\cdot)) - F_\lambda(\cdot, u_2(\cdot))\|_{H^m(\Omega)} \leq \widetilde{M}_\lambda \|u_1 - u_2\|_{H^m(\Omega)}, \quad m \geq 1, \quad \forall u_1, u_2 \in H^m(\Omega),$$

for some sufficiently small \widetilde{M}_λ , to supersede (4.20). But (4.61) appears unnatural, and leads to cumbersome notation. So we do not go into this. \square

5. HIGHER THAN OPTIMAL-ORDER ERROR ESTIMATES
FOR NONLINEARITIES THAT ARE SEPARABLE

A special feature of boundary elements for nonlinear PDEs is that for an important class of nonlinearities, one can obtain higher than “optimal order” error estimates with respect to the h -parameter. Let us assume that $F(x, u)$ in (1.1) is separable, of the form

$$(5.1) \quad F(x, u) = a_0(x)f(u) + f_0(x),$$

where $a_0, f_0 \in C^\infty(\overline{\Omega})$, and $f \in C^\infty(\mathbb{R})$, and that $g(x) \equiv 0$ on $\partial\Omega$.

Lemma 5.1. *Let $\{\mathcal{S}_h \mid 0 < h \leq h_0\}$ be $S_h^{\ell,m}(\partial\Omega)$ -systems with $\ell \geq 2$ and $m \geq 1$ when $N = 2$ or 3 , or be S_h^d -systems of smoothest splines of degree d with quasi-uniform mesh when $N = 2$. Then there are $\bar{h}_0: 0 < \bar{h}_0 \leq h_0$, $\rho_\lambda: 0 < \rho_\lambda < 1$, and constants $K_1 > 0$, $K_2 > 0$ such that*

$$(5.2) \quad \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{-3/2}(\partial\Omega)} \leq K_1 h^{2+\sigma} [\|u\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)} + K_2], \quad \forall \sigma: 0 < \sigma < 1,$$

for all $0 < h \leq \bar{h}_0, n = 0, 1, 2, \dots$, provided that $F(x, u)$ is of the form (5.1), and $g \equiv 0$. In particular, $K_2 = 0$ if $a_0(x)$ and $f_0(x)$ in (5.1) are constant functions.

Proof. We first recall the Sloboditskii norm $\|u\|_\sigma^2 \equiv |u|_\sigma^2 + \|u\|_{L^2(\Omega)}^2$, where

$$|u|_\sigma^2 = \int_{\Omega \times \Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{N+2\sigma}} dx dy$$

for $0 < \sigma < 1$. It is known that the Sloboditskii norm is equivalent to the Sobolev space norm $\|\cdot\|_{H^\sigma(\Omega)}^2$. From (5.1), we have, for $a - \delta \leq u(x) \leq b + \delta, x \in \Omega$,

$$\begin{aligned} & |F_\lambda(x, u(x)) - F_\lambda(y, u(y))| \\ &= |\lambda[u(x) - u(y)] - [a_0(x) - a_0(y)]f(u(x)) - a_0(y)[f(u(x)) - f(u(y))] \\ & \quad + [f_0(x) - f_0(y)]|, \end{aligned}$$

from which it follows that

$$(5.3) \quad \begin{aligned} |F_\lambda(\cdot, u)|_\sigma &\leq \lambda|u|_\sigma + \|f\|_{C[a-\delta, b+\delta]}|a_0|_\sigma + \|a_0\|_{C^0(\bar{\Omega})}\|f'\|_{C^0[a-\delta, b+\delta]}|u|_\sigma + |f_0|_\sigma \\ &\equiv C_1|u|_\sigma + C_2, \quad 0 < \sigma < 1. \end{aligned}$$

Because of (4.48), we can get

$$(5.4) \quad \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_\infty)\|_{L^2(\Omega)} \leq M_\lambda \|u_h^n - u_\infty\|_{L^2(\Omega)}.$$

Similarly as in the proof of Lemma 4.5, we obtain

$$\begin{aligned} \|\eta_h^{n+1} - \zeta_h^{n+1}\|_{H^{-3/2}(\partial\Omega)} &\leq C_3 h^{2+\sigma} \|\zeta_h^{n+1}\|_{H^{\frac{1}{2}+\sigma}(\partial\Omega)} \\ &\leq C_4 h^{2+\sigma} \|F_\lambda(\cdot, u_h^n)\|_{H^\sigma(\Omega)} \quad (\text{because } g = 0 \text{ in (4.21)}) \\ &\leq C_4 h^{2+\sigma} [\|F_\lambda(\cdot, u_h^n)|_\sigma + \|F_\lambda(\cdot, u_h^n) - F_\lambda(\cdot, u_\infty)\|_{L^2(\Omega)} + \|F_\lambda(\cdot, u_\infty)\|_{L^2(\Omega)}] \\ &\leq C_4 h^{2+\sigma} [C_1|u_h^n|_\sigma + C_2 + M_\lambda \|u_h^n - u_\infty\|_{L^2(\Omega)} + C_5 \|u_\infty\|_{H^2(\Omega)}] \\ &\hspace{15em} (\text{by (5.3) and (5.4)}) \\ &\leq C_4 h^{2+\sigma} [C_6(\|u_h^n - u_\infty\|_{H^\sigma(\Omega)} + C_7 \|u_\infty\|_{H^2(\Omega)}) + C_2] \\ &\leq C_8 h^{2+\sigma} [\|u_\infty\|_{H^2(\Omega)} + \rho_\lambda^n \|e^0\|_{L^2(\Omega)} + C_9] \quad (\text{by Theorem 4.10}). \square \end{aligned}$$

Now, the following is obvious.

Theorem 5.2. *Assume the same conditions as in Lemma 5.1. Then there are three constants $C_i > 0$, $i = 1, 2, 3$, such that*

$$\begin{aligned} \|u_h^n - u_\infty\|_{H^r(\Omega)} &\leq C_1 \rho_\lambda^n \|e^0\|_{L^2(\Omega)} + C_2 h^{2-r+\sigma} (\|u_\infty\|_{H^2(\Omega)} + C_3) \\ \forall h: \quad 0 < h \leq \bar{h}_0, \quad n = 0, 1, 2, \dots, \quad 0 \leq \sigma < 1, \quad 0 \leq r \leq 2. \end{aligned}$$

In particular, $C_3 = 0$ if $a_0(x)$ is a constant function and $f_0(x) \equiv 0$ in (5.1). □

6. NEUMANN AND ROBIN BOUNDARY CONDITIONS

The treatments in §4 and §5 deal with the Dirichlet boundary condition. We now consider the case when the boundary condition is

$$(6.1) \quad \frac{\partial}{\partial n} u(x) + \alpha(x)u(x) = g(x), \quad x \in \partial\Omega,$$

where $\alpha(x) \geq 0$, $\alpha \in C^\infty(\partial\Omega)$. The condition (6.1) corresponds to a Neumann or a Robin boundary value problem depending on whether $\alpha(x)$ is or is not identically equal to zero.

For (6.1), we formulate a corresponding assumption similar to [H]. Let λ'_1 be the smallest eigenvalue of the operator $(-\Delta, [\frac{\partial}{\partial n} + \alpha(\cdot)]|_\Gamma)$. Then $\lambda'_1 = 0$ if $\alpha(x) \equiv 0$ and $\lambda'_1 > 0$ if $\alpha(x) \geq 0$ but $\alpha(x) \not\equiv 0$. We assume:

[H]' There exist constants γ'_1 and γ'_2 such that

$$-\lambda'_1 < \gamma'_1 \leq \frac{\partial F(x, u)}{\partial u} \leq \gamma'_2, \quad \forall (x, u) \in \bar{Q} \equiv \bar{\Omega} \times [a, b].$$

Similar to (4.13), we now have that for a sufficiently small $\delta > 0$, there exist $\tilde{\gamma}'_1$ and $\tilde{\gamma}'_2$ such that

$$(6.2) \quad -\lambda'_1 < \tilde{\gamma}'_1 \leq \frac{\partial F(x, u)}{\partial u} \leq \tilde{\gamma}'_2, \quad \forall (x, u) \in \overline{Q}_\delta \equiv \overline{\Omega} \times [a - \delta, b + \delta].$$

Let $\lambda > 0$ be chosen sufficiently large such that

$$(6.3) \quad M'_\lambda \equiv \sup_{(x, u) \in \overline{Q}_\delta} \left[\lambda - \frac{\partial F(x, u)}{\partial u} \right] > 0.$$

Then

$$(6.4) \quad 0 < M'_\lambda < \lambda - \tilde{\gamma}'_1.$$

Lemma 6.1. *Let the assumptions in Theorem 2.2 and (6.1)–(6.4) hold. Let u_{n+1} be iterated according to*

$$(6.5) \quad \begin{cases} (\Delta - \lambda)u_{n+1}(x) = F_\lambda(x, u_n(x)), & x \in \Omega, \\ \frac{\partial}{\partial n}u_{n+1}(x) + \alpha(x)u_{n+1}(x) = g(x), & x \in \partial\Omega, \quad g \in C^\infty(\partial\Omega), \end{cases}$$

and let u_∞ be the limit as in (2.16). Denote $e^n = u_\infty - u_n$. Then we have

$$\|e^{n+1}\|_{L^2(\Omega)} \leq \alpha'_\lambda{}^{n+1} \|e^0\|_{L^2(\Omega)}, \quad n = 0, 1, 2, \dots,$$

where

$$0 < \alpha'_\lambda \equiv \frac{\lambda - \tilde{\gamma}'_1}{\lambda + \lambda'_1} < 1.$$

Proof. Refer to the proof of Lemma 4.1. Instead of (4.16), we now have

$$\begin{cases} (\Delta - \lambda)e^{n+1}(x) = F_\lambda(x, u_\infty(x)) - F_\lambda(x, u_n(x)), & x \in \Omega, \\ \frac{\partial}{\partial n}e^{n+1}(x) + \alpha(x)e^{n+1}(x) = 0, & x \in \partial\Omega. \end{cases}$$

Multiplying the above by $-e^{n+1}$ and integrating by parts, we get

$$\begin{aligned} \int_{\partial\Omega} \alpha |e^{n+1}|^2 d\sigma + \int_{\Omega} |\nabla e^{n+1}|^2 dx + \lambda \int_{\Omega} |e^{n+1}|^2 dx \\ \leq \|F_\lambda(\cdot, u_\infty) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)} \|e^{n+1}\|_{L^2(\Omega)}. \end{aligned}$$

But

$$\inf_{v \in H^1(\Omega)} \frac{\left[\int_{\partial\Omega} \alpha |v|^2 d\sigma + \int_{\Omega} |\nabla v|^2 dx \right]}{\|v\|_{L^2(\Omega)}^2} = \lambda'_1.$$

Thus,

$$(\lambda + \lambda'_1) \|e^{n+1}\|_{L^2(\Omega)}^2 \leq \|F_\lambda(\cdot, u_\infty) - F_\lambda(\cdot, u_n)\|_{L^2(\Omega)} \|e^{n+1}\|_{L^2(\Omega)}.$$

The rest of the arguments are the same as in the proof of Lemma 4.1. \square

Proceeding as in §4, we now note where the places are that need to be adjusted to accommodate the boundary condition (6.1). The iterate $u_{n+1}(\cdot)$ satisfying (6.5) is still represented as in (3.8), but η_{n+1} is determined from the BIE

$$\begin{aligned}
 (6.6) \quad \mathcal{B}_\lambda u_{n+1}(x) &\equiv \frac{1}{2} \eta_{n+1}(x) + \int_{\partial\Omega} \left[\frac{\partial}{\partial n_x} E(x-y; \lambda) + \alpha(x) E(x-y; \lambda) \right] \eta_{n+1}(y) d\sigma_y \\
 &= g(x) + \alpha(x) (\mathcal{V}_{\lambda, \Gamma} F(\cdot, u_n))(x) + \frac{\partial}{\partial n} (\mathcal{V}_\lambda F(\cdot, u_n))(x), \quad x \in \partial\Omega,
 \end{aligned}$$

instead of (3.47). Note that \mathcal{B}_λ is a strongly elliptic pseudodifferential operator of order 0 ($= 2\alpha$ according to the notation in [10]) with principal symbol $\frac{1}{2}$ such that

$$(6.7) \quad \mathcal{B}_\lambda: H^s(\partial\Omega) \longrightarrow H^s(\partial\Omega) \quad \text{isomorphically,} \quad \forall s \in \mathbb{R}.$$

Therefore, (6.6) is uniquely solvable.

The Galerkin boundary element method for (6.5) is almost the same as in (4.10)–(4.12), except that (4.11) is replaced by

(ii)' For each $n = 0, 1, 2, \dots$, and $h: 0 < h \leq h_0$, find $\eta_h^{n+1} \in \mathcal{S}_h$ such that

$$\langle \mathcal{B}_\lambda \eta_h^{n+1}, \phi_h \rangle_{L^2(\partial\Omega)} = \langle *, \phi_h \rangle_{L^2(\partial\Omega)}, \quad \forall \phi_h \in \mathcal{S}_h,$$

where $*$ denotes the sum in the last equality of (6.6), but with u_h^n taking the place of u_h .

Next, we let ζ_h^{n+1} be the unique solutions of the BIE

$$\begin{aligned}
 (\mathcal{B}_\lambda \zeta_h^{n+1})(x) &= g(x) + \alpha(x) (\mathcal{V}_{\lambda, \Gamma} F(\cdot, u_h^n))(x) + \frac{\partial}{\partial n} (\mathcal{V}_\lambda F(\cdot, u_h^n))(x), \quad x \in \partial\Omega, \\
 &\text{(instead of (4.21))}
 \end{aligned}$$


from which we define $w_h^{n+1}(x)$ just as in (4.27).

It is now routine to check that beginning from (4.21) in Lemma 4.2 and ending at (4.60), all the estimates remain valid. Indeed, we obtain the same results under a *less restrictive assumption* on $S_h^{\ell, m}(\partial\Omega)$ —here we only require that $\ell \geq 1$ (and $m \geq 0$) because 2α , the order of \mathcal{B}_λ , is zero. Thus the requirement that $|\alpha| \leq m$ made in [10, Corollary 2.1] and [14, Corollary 3] is now automatically satisfied with $m \geq 0$.

7. NUMERICAL EXAMPLES

The theory developed so far was molded by computations of concrete examples of semilinear PDEs. In the following, we will present several such examples featuring various situations, with numerical solutions illustrated by computer graphics.

The steps that we take are indicated by a flowchart next. For definiteness, we only consider the Dirichlet boundary condition for $\Omega \equiv D_1$, the unit open disk in



\mathbb{R}^2 . The boundary value problem is of the form

$$(7.1) \quad \begin{cases} \Delta u(x) - F(x, u(x)) = 0, & x \in D_1, \\ u(x) = g(x), & x \in \partial D_1. \end{cases}$$

Note that at Steps 2 and 4 of the flowchart, the volume potential $V_{\lambda,j}(x)$ and the iterate $w_{j+1}(x)$ are computed by quadratures of high accuracy with 864 Gauss points placed on the unit disk. The Gauss points are selected by using the Cartesian Gauss rule for the polar coordinate variables (r, θ) : dividing the unit disk into $4_r \times 24_\theta$ elements and using $3_r \times 3_\theta$ Gauss points in each element. The key step is the solution of the boundary integral equation for the simple-layer density η_{j+1} in Step 3. In our numerical work, η_{j+1} is solved *by point collocation* using piecewise constant boundary elements (i.e., $S_h^{1,0}(\partial\Omega)$ -systems) with uniform mesh on $\partial\Omega$, where $\partial\Omega$ is divided into 64 equal pieces of arc (i.e., $h = 2\pi/64$). Here collocation is taken as a measure of convenience, because the Galerkin boundary element method as stated in (4.10) and (4.12) involve significantly more programming work. It is also for this reason that in this section we have not attempted to confirm the error estimates given in §5 and §6 experimentally. We nevertheless hope that the examples furnished here will serve as useful benchmarks for researchers doing similar work, and that in the future we will also be able to derive and numerically verify error estimates for the boundary element point-collocation scheme.

In all our numerical computations, $\varepsilon = 10^{-4}$ has been used as the relative L_2 error in solutions to terminate the iteration, see Flow-chart Step 5.

We first test the accuracy of our computer program this way. Choose a known (smooth) function $y(x), x \in \bar{\Omega}$, and for a somewhat arbitrary function F , where F

is nonlinear in u , we compute the solution of

$$\begin{cases} \Delta u(x) - F(x, u(x)) = 0, & x \in \Omega, \\ u(x) = y(x), & x \in \partial\Omega, \end{cases}$$

where

$$F(x, u(x)) = F(u(x)) + [\Delta y(x) - F(y(x))], \quad x \in \Omega.$$

Obviously, $y(x)$ is an *exact solution* of the (artificially set example) above. We then use $y(x)$ as a benchmark to test whether we can numerically recover $u(x) = y(x)$. The computer results are affirmative: the output data have shown high accuracy.

In the following, we provide numerical examples which do not have exact solutions available. The accompanying theoretical discussions are to a large extent motivated by visualization of the numerical results.

Example 7.1. Multiple solutions—two stable solutions bounding an unstable one.

Consider

$$(7.2) \quad \begin{cases} \Delta u(x) - \gamma u(x)[u(x) - a][u(x) - 1] = 0, & x \in D_1, \\ u(x) = 0, & x \in \partial D_1, \end{cases}$$

where $\gamma > 0$, $0 < a < 1/2$. It is easy to see that $u(x) = \bar{v}(x) \equiv 0$ is a solution. One can further establish that $\bar{v}(x) \equiv 0$ is *stable*. This solution is of less interest because it is *trivial*.

Substituting $\bar{u}(x) \equiv 1$ for u into (7.2), we also easily verify that $\bar{u}(x)$ is a supersolution. (Actually, $\bar{u}(x) \equiv C$ for any $C \geq 1$ will do.)




FIGURE 7.1. A bell-shaped stable positive boundary element numerical solution $u(x)$ satisfying $\Delta u - u(u - \frac{1}{4})(u - 1) = 0$ and the zero Dirichlet boundary condition for Example 7.1.

In (7.2), we have

$$(7.3) \quad \begin{aligned} F(x, u) &\equiv F(u) = \gamma u(u - a)(u - 1), \\ \frac{\partial F}{\partial u} &= \gamma[(u - a)(u - 1) + u(u - 1) + u(u - a)] \\ &= \gamma[3u^2 - 2(a + 1)u + a]. \end{aligned}$$

In our computation, we choose

$$\gamma = 100, \quad a = \frac{1}{4}, \quad \text{yielding} \quad \max \frac{\partial F}{\partial u} = 75, \quad u \in [0, 1].$$

Thus, if we choose $\bar{u}(x) \equiv 1$, $\lambda = 75.69$, then (3.10) is satisfied.

The monotone scheme beginning from $u_0(x) = \bar{u}(x) \equiv 1$ takes 30 iterations, yielding a bell-shaped function $u_\infty(x)$ as shown in Fig. 7.1. According to Theorem 2.2, this is a stable solution. Since $u_\infty(x) \not\equiv 0 \equiv \bar{v}(x) = v_\infty(x)$, we know from Theorem 2.2 that there is at least one unstable solution of (7.2), called a “spike-layer” solution, sandwiched between v_∞ and u_∞ .

Generally, it is known that for (7.2), the following are true:

- (i) Any solution u of (7.2) must satisfy $0 \leq u(x) \leq 1$;
- (ii) For $\gamma > 0$ sufficiently small, (7.2) has the trivial solution as its only solution;
- (iii) The spike-layer solution exists only when $\gamma > 0$ is large.

Meanwhile, the antimonotone convergence of the simple-layer densities, guaranteed by Theorem 3.1, is reflected in Fig. 7.2. □




FIGURE 7.2. The antimonotone convergence of the simple-layer densities for Example 7.1. While the solutions $u_h^n(x)$ decrease monotonically on the unit disk, the simple-layer densities $\eta_h^n(x)$ increase monotonically for $n = 1$ (bottom curve) to $n = 10$ (top curve).

Example 7.2. A monotone dissipative operator equation.

Consider

$$(7.4) \quad \begin{cases} \Delta u(x) - 4u^3(x) = \gamma x_1^2, & x = (x_1, x_2) \in D_1, \quad \gamma > 0, \\ u(x) = 0, & x \in \partial D_1. \end{cases}$$

The nonlinear operator \mathcal{N} on the left-hand side of the PDE in (7.1) is known to be a monotone dissipative operator as it satisfies

$$\begin{aligned} \langle \mathcal{N}u_1 - \mathcal{N}u_2, u_1 - u_2 \rangle_{L^2(D_1)} &= \langle (\Delta u_1 - 4u_1^3) - (\Delta u_2 - 4u_2^3), u_1 - u_2 \rangle \\ &= \int_{D_1} [\Delta(u_1 - u_2) - 4(u_1^3 - u_2^3)](u_1 - u_2) dx \\ &= - \int [|\nabla(u_1 - u_2)|^2 + 4(u_1^3 - u_2^3)(u_1 - u_2)] dx \leq 0, \end{aligned}$$

for $u_1, u_2 \in H_0^2(D_1)$. From the theory of monotone dissipative operators it is straightforward ([11], e.g.) to establish that (7.4) has a *unique* solution $u \in C^\infty(\overline{D_1})$.

From the way (7.4) is given, we can further prove that the solution satisfies $u(x) < 0$, $x \in D_1$. Assume the contrary. Then by the usual maximum principle [13] $G \equiv \{x \in D_1 \mid u(x) > 0\}$ is nonempty. Thus, on G , u satisfies

$$\begin{cases} \Delta u(x) = 4u^3(x) + \gamma x_1^2 > 0, & x \in G, \\ u(x) > 0, & x \in G, \\ u(x) = 0, & x \in \partial G. \end{cases}$$

But for u satisfying $\Delta u \geq 0$ on G and $u = 0$ on ∂G , the maximum principle implies that $u(x) \leq 0$ on G , a contradiction.

For (7.4), we have $F(x, u) = 4u^3 + \gamma x_1^2$. For D_1 , it is known that λ_1 , the smallest eigenvalue of $-\Delta$ corresponding to the homogeneous Dirichlet condition, is (cf. [7, p. 188], e.g.)

$$\lambda_1 \approx (2.4048)^2 \approx 5.783063.$$

Therefore,

$$(7.5) \quad \frac{\partial F}{\partial u} = 12u^2 \geq 0 > -\lambda_1 \approx -5.783063.$$

We may take $\bar{u}(x) \equiv 0$ as a supersolution, and $\bar{v}(x) \equiv -C$, $C > 0$, as a subsolution, if C is chosen such that $4C^3 - \gamma x_1^2 \geq 0$, $\forall x \in \overline{D_1}$.

Because of (7.5), we easily see that [H] and (4.15) are satisfied for $\tilde{\gamma}_1 < 0$ if $|\tilde{\gamma}_1|$ is small.

In our numerical computations, we choose $\gamma = 10$ and $\lambda = 4$. The computation takes 8 iterations to converge. The numerical solution is plotted in Fig. 7.3. The reader can easily see that the solution $u(x)$ is nonpositive. \square

Note that Theorem 5.2 applies to Examples 7.1 and 7.2.




FIGURE 7.3. Boundary element numerical solution of $\Delta u - 4u^3 \equiv 10x_1^2$ subject to the zero Dirichlet boundary condition, Example 7.2.

Example 7.3. A nonlinear PDE of nonmonotone type whose existence or nonexistence of stable solutions of one sign depends on the size of forcing.

Consider

$$(7.6) \quad \begin{cases} \Delta u(x) + 4u^3(x) = \gamma x_1^2, & x \in D_1, \quad \gamma > 0, \\ u(x) = 0, & x \in \partial D_1. \end{cases}$$

Suppose we are interested in finding nonpositive solutions

$$(7.7) \quad u(x) \leq 0 \quad \text{on } D_1.$$

This condition is now appended to (7.6). Later on, we will be able to remove (7.7) provided that $\gamma > 0$ is sufficiently small, and that proper supersolutions and subsolutions are chosen; see (iii) below.

We now show the following.

(i) *The system (7.6) and (7.7) has no solution if γ is large.*

Let ϕ_1 be the first eigenfunction of $-\Delta$ satisfying

$$(7.8) \quad \begin{cases} \Delta \phi_1 + \lambda_1 \phi_1 = 0 & \text{on } D_1 \quad (\lambda_1 \approx 5.783063), \\ \phi_1(x) > 0 & \text{on } D_1, \\ \phi_1|_{\partial D_1} = 0. \end{cases}$$

Here, $\phi_1(x) > 0$ because we know that ϕ_1 is of one sign on D_1 (cf. [6, §3.1], e.g.).

From (7.6)–(7.8), we get

$$\begin{aligned}
 \gamma \int_{D_1} x_1^2 \phi_1(x) dx &= \int_{D_1} [\Delta u(x) + 4u^3(x)] \phi_1(x) dx \\
 &= \int_{D_1} [-\lambda_1 u(x) \phi_1(x) + 4u^3(x)] \phi_1(x) dx \\
 &= \int_{D_1} u(x) \phi_1(x) [4u^2(x) - \lambda_1] dx \\
 &= \int_{D_{1,1}} u(x) \phi_1(x) [4u^2(x) - \lambda_1] dx \\
 &\quad + \int_{D_{1,2}} u(x) \phi_1(x) [4u^2(x) - \lambda_1] dx,
 \end{aligned}$$

where

$$D_{1,1} = \{x \in D_1 \mid -\sqrt{\lambda_1}/2 \leq u(x) \leq 0\}, \quad D_{1,2} = \{x \in D_1 \mid u(x) < -\sqrt{\lambda_1}/2\}.$$

Hence,

$$(7.9) \quad 0 < \gamma \int_{D_1} x_1^2 \phi_1(x) dx \leq \int_{D_{1,1}} u(x) \phi_1(x) [4u^2(x) - \lambda_1] dx.$$

On $D_{1,1}$, we have $\sqrt{\lambda_1}/2 \geq -u(x)$, and $\lambda_1 \geq \lambda_1 - 4u^2(x)$. Using the above in (7.9), we obtain

$$\begin{aligned}
 &\frac{1}{2} \int_{D_{1,1}} \sqrt{\lambda_1} \phi_1(x) \lambda_1 dx \geq \gamma \int_{D_1} x_1^2 \phi_1(x) dx, \\
 (7.10) \quad \gamma &\leq \frac{\lambda_1^{3/2} \int_{D_{1,1}} \phi_1(x) dx}{2 \int_{D_1} x_1^2 \phi_1(x) dx} \leq \frac{\lambda_1^{3/2} \int_{D_1} \phi_1(x) dx}{2 \int_{D_1} x_1^2 \phi_1(x) dx}.
 \end{aligned}$$

Thus, if $\gamma > 0$ is chosen large enough to violate (7.10), the system (7.6) and (7.7) will not have a solution.

(ii) *If the system (7.6) and (7.7) has a solution for $\gamma = \gamma_1 > 0$, then for all $\gamma: 0 < \gamma < \gamma_1$, the system (7.6) and (7.7) also has a solution.*

Let \tilde{u} be the solution corresponding to γ_1 :

$$(7.11) \quad \begin{cases} \Delta \tilde{u}(x) + 4\tilde{u}^3(x) = \gamma_1 x_1^2, & x \in D_1, \\ \tilde{u}(x) \leq 0 & \text{on } D_1, \\ \tilde{u}(x) = 0 & \text{on } \partial D_1. \end{cases}$$

Consider u , satisfying

$$(7.12) \quad \begin{cases} \Delta u(x) + 4u^3(x) = \gamma x_1^2, & x \in D_1, \quad 0 < \gamma < \gamma_1, \\ u(x) \leq 0 & \text{on } D_1, \\ u(x) = 0 & \text{on } \partial D_1. \end{cases}$$

Thus, it is easy to check that $\bar{u}(x) \equiv 0$ is a supersolution for (7.12). We also claim that $\bar{v}(x) = \tilde{u}(x)$ is a subsolution, since

$$\begin{aligned} \Delta \bar{v}(x) + 4\bar{v}^3(x) - \gamma x_1^2 &= [\Delta \tilde{u}(x) + 4\tilde{u}^3(x) - \gamma x_1^2] + (\gamma_1 - \gamma)x_1^2 \\ &= (\gamma_1 - \gamma)x_1^2 \geq 0 \quad \text{on } D_1, \end{aligned}$$

and $\bar{v}(x) = \tilde{u}(x) = 0$ on ∂D_1 . By Theorem 2.3, a solution $u(x)$ of (7.12) exists, satisfying $\tilde{u}(x) \leq u(x) \leq 0$ on D_1 . \square

(iii) *The system (7.6) and (7.7) possesses a stable solution $\tilde{u}_1: \tilde{u}_1(x) \leq 0$ as well as a second solution $\tilde{u}_2: \tilde{u}_2(x) < \tilde{u}_1(x), x \in \Omega$, if $\gamma > 0$ is sufficiently small.*

We already know that $\bar{u}(x) \equiv 0$ is a supersolution of (7.6) and (7.7). We now construct a subsolution. Let $w(x)$ satisfy the Lane-Emden equation

$$(7.13) \quad \begin{cases} \Delta w(x) + 4w^3(x) = 0, & \text{on } D_1, \\ w(x) > 0, & \text{on } D_1, \\ w(x) = 0, & \text{on } \partial D_1. \end{cases}$$

It is known (cf. [12], e.g.) that (7.14) has a unique solution (that is, also radially symmetric and monotonically decreasing in $|x|$). Thus, we write $w(x) = w(r)$, $r = |x|$, $x \in D_1$. Define

$$(7.14) \quad w_\alpha(x) = \alpha w(\alpha r), \quad 0 < \alpha < 1.$$

Then it is routine to verify that $w_\alpha(x)$ satisfies

$$\Delta w_\alpha(x) + 4w_\alpha^3(x) = 0, \quad w_\alpha(x) \geq \varepsilon > 0, \text{ for } \varepsilon = \alpha w(\alpha) \text{ and } x \in D_1.$$

Let

$$(7.15) \quad \bar{v}(x) = -\beta w_\alpha(x), \quad 1 > \beta > 0, \text{ for some fixed } \beta.$$

Then

$$\begin{aligned} \Delta \bar{v}(x) + 4\bar{v}^3(x) - \gamma x_1^2 &= -\beta \Delta w_\alpha(x) - 4\beta^3 w_\alpha^3(x) - \gamma x_1^2 \\ &= 4\beta w_\alpha^3(x) - 4\beta^3 w_\alpha^3(x) - \gamma x_1^2 \\ &= 4\beta(1 - \beta^2)w_\alpha^3(x) - \gamma x_1^2 \\ &\geq 4\beta(1 - \beta^2)\varepsilon^3 - \gamma x_1^2 \geq 0, \end{aligned}$$

if $\gamma > 0$ is sufficiently small. Also,

$$\bar{v}(x) = -\beta w_\alpha(x) = -\beta \alpha w(\alpha) \leq -\beta \alpha \varepsilon, \quad \text{if } x \in \partial D_1.$$

Thus $\bar{v}(x)$ is a subsolution, if $\gamma > 0$ is sufficiently small. Hence the system (7.6) and (7.7) admit a stable solution for $\gamma > 0$ sufficiently small. Call this solution \tilde{u}_1 . From (ii), we further see that $\tilde{u}_1 \uparrow 0$ uniformly as $\gamma \downarrow 0$.

To establish the existence of the second solution $\tilde{u}_2 < \tilde{u}_1$ (on Ω), we use a variational approach. For ease of discussion, set $z = -u$ in (7.6) and (7.7), giving

$$(7.16) \quad \begin{cases} \Delta z(x) + 4z^3(x) + \gamma x_1^2 = 0, & \text{on } D_1, \\ z(x) > 0, & \text{on } D_1, \\ z(x) = 0, & \text{on } \partial D_1. \end{cases}$$

Note that $z_1 = -\tilde{u}_1$ is a solution of (7.16). Setting $w = z - z_1$ in (7.16), we obtain

$$(7.17) \quad \begin{cases} \Delta w(x) + [12z_1^2(x)]w(x) + [12z_1(x)]w^2(x) + 4w^3(x) = 0, & x \in D_1, \\ w(x) = 0, & x \in \partial D_1. \end{cases}$$

We now look for a solution of (7.17) that further satisfies

$$(7.18) \quad w(x) > 0, \quad x \in D_1.$$

When $\gamma > 0$ is sufficiently small, $\|z_1\|_{L^\infty(\Omega)}$ is small. Thus $\|12z_1^2\|_{L^\infty(\Omega)} < \lambda_1$ (cf. (7.8) for λ_1) is satisfied for small $\gamma > 0$. Then an application of the usual Mountain-Pass Lemma [3, 12] yields a classical solution w of (7.17) and (7.18). Hence, $\tilde{u}_2 \equiv -(z_1 + w) < \tilde{u}_1 < 0$ on Ω is a second solution.

For (7.6), we have

$$F(x, u) = -4u^3 + \gamma x_1^2, \quad \frac{\partial F}{\partial u} = -12u^2.$$

Although the subsolution as given in (7.14) and (7.15) is *not known explicitly*, it can be verified that the conditions [H] and (4.15) are satisfied if $\lambda \geq 0$.

In our numerical experiments, we have chosen $\gamma = 10$ and $\lambda = 4$. The numerical iterates converge after 16 iterations. The graph of the numerical solution \tilde{u}_1 can be found in Fig. 7.4. \square




FIGURE 7.4. Boundary element numerical solution of $\Delta u + 4u^3 = 10x_1^2$ subject to the zero Dirichlet boundary condition, Example 7.3.

Example 7.4. A nonlinear PDE of nonmonotone type with a unique stable non-positive solution.

Consider

$$(7.19) \quad \begin{cases} \Delta u(x) + 4u^2(x) = \gamma x_1^2, & x \in D_1, \quad \gamma > 0, \\ u(x) = 0, & x \in \partial D_1. \end{cases}$$

It is easy to see that $\bar{u}(x) \equiv 0$ is a supersolution, and $\bar{v}(x) \equiv -C$, $C > 0$, is a subsolution if C is chosen such that $4C^2 - \gamma x_1^2 \geq 0, \forall x_1: |x_1| \leq 1$. Thus, by Theorem 2.2, there exists a stable solution $u(x)$ such that $-C \leq u(x) \leq 0, \forall x \in D_1$. We now show that *nonpositive* solutions of (7.19) are unique. Suppose $u^{(1)}$ and $u^{(2)}$ both are nonpositive and satisfy (7.19). We may assume that

$$(7.20) \quad u^{(1)}(x) \geq u^{(2)}(x) \quad \forall x \in D_1, \quad u^{(1)} \not\equiv u^{(2)}.$$

Otherwise, by Theorem 2.2 and Remark 2.4(a), $\bar{v}(x) = \min\{u^{(1)}(x), u^{(2)}(x)\}, x \in D_1$, is a subsolution of (7.19) and monotone iteration with $\bar{v}(x)$ as the initial state will yield a limit solution $u_\infty(x)$ such that $u^{(1)}(x) \geq u_\infty(x), \forall x \in D_1, u^{(1)} \not\equiv u_\infty$. From (7.19) and (7.20), we have

$$\begin{aligned} 0 &= \int_{\partial D_1} \left[u^{(2)} \frac{\partial u^{(1)}}{\partial n} - u^{(1)} \frac{\partial u^{(2)}}{\partial n} \right] d\sigma = \int_{D_1} [u^{(2)} \Delta u^{(1)} - u^{(1)} \Delta u^{(2)}] dx \\ &= - \int_{D_1} [u^{(2)}(4u^{(1)2} - \gamma x_1^2) - u^{(1)}(4u^{(2)2} - \gamma x_1^2)] dx \\ &= - \int_{D_1} [4u^{(1)}u^{(2)}(u^{(1)} - u^{(2)}) + \gamma x_1^2(u^{(1)} - u^{(2)})] dx < 0, \end{aligned}$$

a contradiction. □

Thus, in contrast to Example 7.3, the size of γ here does not affect the existence or nonexistence of the unique stable nonpositive solution. (Nevertheless, we do not rule out that some *other stable solutions not of one sign may still exist.*)

For (7.19), we have

$$(7.21) \quad F(x, u) = -4u^2 + \gamma x_1^2, \quad \frac{\partial F}{\partial u} = -8u.$$

Note that we need only consider $u \leq 0$. Here, to ensure that [H] and (4.15) are satisfied, we need only use $\lambda = 0$.

In our numerical computation, we choose $\gamma = 10$ in (7.19) and use $\lambda = 4$. The numerical solutions converge after 9 iterations. The limiting solution is plotted in Fig. 7.5.

Although the PDEs in Examples 7.2, 7.3 and 7.4 seem to have vastly different nonlinearities in a qualitative sense, we nevertheless notice that the solution graphics in Figs. 7.3, 7.4 and 7.5 seem to display similar profiles. □

Example 7.5. A nonlinear PDE of nonmonotone type, whose stable solutions, not of one sign, may be obtained through symmetry.

Consider

$$(7.22) \quad \begin{cases} \Delta u(x) + 4u^3(x) = \gamma x_1^3, & x \in D_1, \quad \gamma > 0, \\ u(x) = 0, & x \in \partial D_1. \end{cases}$$




FIGURE 7.5. Boundary element numerical solution of $\Delta u + 4u^2 = 10x_1^2$ subject to the zero Dirichlet boundary condition, Example 7.4.

Here the forcing term γx_1^3 changes sign as x moves from $D_{1,-}$ into $D_{1,+}$:

$$D_{1,-} \equiv \{x \in D_1 \mid x_1 < 0\}, \quad D_{1,+} \equiv \{x \in D_1 \mid x_1 > 0\}.$$

Note that, if $u(x)$ is a solution of (7.22), then so is $-u(-x_1, x_2)$, i.e., u is antisymmetric with respect to the x_2 -axis, and thus we deduce that $u(x) = 0$, $x \in \overline{D}_{1,-} \cap \overline{D}_{1,+}$. So we need only consider the boundary value problem

$$(7.23) \quad \begin{cases} \Delta \tilde{u}(x) + 4\tilde{u}^3(x) = \gamma x_1^3, & x \in D_{1,+}, \\ \tilde{u}(x) = 0, & x \in \partial D_{1,+}, \end{cases}$$

because on $D_{1,-}$ we simply have $u(x_1, x_2) = -\tilde{u}(-x_1, x_2)$ for $x \in D_{1,-}$. The nonlinear boundary value problem (7.23) is similar to the one considered in (7.6) because the forcing terms γx_1^3 and γx_1^2 are all nonnegative on each domain. So essentially all the arguments in parts (i)–(iii) of Example 7.3 apply. The only concern one might have for (7.23) may be that $\partial D_{1,+}$ is not a C^∞ curve. Nevertheless, $\partial D_{1,+}$ is Lipschitz continuous. With suitable modifications, the arguments still go through and there will be no problems.

In our numerical computations, we again choose $\gamma = 10$ in (7.22) and use $\lambda = 4$. The numerical solutions converge after 13 iterations, using $u_0(x) \equiv 0$. The graphics is given in Fig. 7.6. The reader will find that the solution changes sign across the x_2 -axis. \square




FIGURE 7.6. Boundary element numerical solution of $\Delta u + 4u^3 = 10x_1^3$ subject to the zero Dirichlet boundary condition, Example 7.5.

Remark 7.6. How restrictive is the working assumption [H] made in §4? From Examples 7.1–7.5, we see that for the monotone equation (7.4) in Example 7.2, [H] is always satisfied. The BVP (7.4) also has the least trouble with existence and uniqueness. For the BVP in Example 7.1, when γ is sufficiently small, condition [H] holds. However, [H] fails when γ is large, because the system (7.1) has multiple solutions that can be separated only by nonconstant sup- and subsolutions. So it appears that an alternative assumption to [H] is required. Nevertheless, computationally we are still able to obtain convergent numerical solutions.

For Examples 7.3 and 7.4, even though the existence/nonexistence and uniqueness/multiplicity of solutions are harder to determine, condition [H] can still largely be confirmed. In any case, regardless of whether [H] can be confirmed a priori or a posteriori, numerical solutions can always be generated by the computer and motivate us to establish further theoretical properties of solutions as given in this section. This is an indication of the usefulness of the development of a combination of theory (i.e., monotone iterations) and numerical methods (i.e., BEM). \square

REFERENCES

1. M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions*, Dover, New York, 1965. MR **34**:8607
2. H. Amann, *Supersolution, monotone iteration and stability*, J. Diff. Eq. **21** (1976), 367-377. MR **53**:11226
3. A. Ambrosetti and P. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. **14** (1973), 349-381. MR **51**:6412

4. I. Babuška and A.K. Aziz, *The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations*, Academic Press, New York, 1972. MR **54**:9111
5. C.A. Brebbia and S. Walker, *Boundary Element Techniques in Engineering*, Newnes-Butterworths, London, 1980.
6. G. Chen and J. Zhou, *Boundary Element Methods*, Academic Press, London, 1992. MR **93e**:65143
7. G. Chen and J. Zhou, *Vibration and Damping in Distributed Systems*, Vol. II: WKB and Wave Methods, Visualization and Experimentation CRC Press, Boca Raton, Florida, 1993.
8. Y. Deng, *Boundary element methods for nonlinear partial differential equations*, Ph.D. dissertation, Math. Dept., Texas A&M Univ., College Station, Texas, August 1994.
9. G. Fichera, *Proc. of Symposium on Partial Differential Equations and Continuum Mechanics* (R.E. Langer, ed.), 55-80, Univ. of Wisconsin Press, Madison, Wisconsin, 1961. MR **27**:6016
10. G.C. Hsiao and W.L. Wendland, *The Aubin-Nitsche lemma for integral equations*, J. Integral Eq. 3 (1981), 299-315. MR **83j**:45019
11. J.L. Lions, *Quelques Méthodes de Résolution des Problèmes aux Limites Non Linéaires*, Dunod, Paris, 1969. MR **41**:4326
12. W.M. Ni, *Lecture Notes published by Institute of Mathematics*, National Tsing Hua Univ., Hsinchu, Taiwan, Rep. of China, May, 1987.
13. M. Protter and H. Weinberger, *Maximum Principles in Differential Equations*, Prentice-Hall, Englewood Cliffs, New Jersey, 1967. MR **36**:2935
14. K. Ruotsalainen and J. Saranen, *On the convergence of the Galerkin method for nonsmooth solutions of integral equations*, Num. Math. **54** (1988), 295-302. MR **90a**:65281
15. K. Ruotsalainen and W.L. Wendland, *On the boundary element method for a nonlinear boundary value problem*, Boundary Elements IX (C.A. Brebbia, ed.), Springer-Verlag, New York, 1987, pp. 385-393. MR **90e**:65161
16. M. Sakakihara, *An iterative boundary integral equation method for mildly nonlinear elliptic partial differential equations*, Boundary Elements VII (C.A. Brebbia and G. Maier, ed.), vol. . II, Springer-Verlag, Berlin-Heidelberg, 1985, pp. 13.49-13.58. CMP 20:14
17. D. Sattinger, *Topics in Stability and Bifurcation Theory*, Lecture Notes in Mathematics, vol. . 309, Springer-Verlag, New York, 1973. MR **57**:3569

(Y. Deng, G. Chen, and J. Zhou) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843

E-mail address: `ydeng@cs.tamu.edu`

E-mail address: `gchen@math.tamu.edu`

E-mail address: `jzhou@math.tamu.edu`

(W.-M. Ni) SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455