

## ON THE DISCRETE LOGARITHM IN THE DIVISOR CLASS GROUP OF CURVES

HANS-GEORG RÜCK

ABSTRACT. Let  $X$  be a curve which is defined over a finite field  $k$  of characteristic  $p$ . We show that one can evaluate the discrete logarithm in  $Pic_0(X)_{p^n}$  by  $O(n^2 \log p)$  operations in  $k$ . This generalizes a result of Semaev for elliptic curves to curves of arbitrary genus.

Let  $k$  be a finite field of characteristic  $p$ . We consider a projective irreducible nonsingular curve  $X$  of genus  $g \geq 1$  which is defined over  $k$ . We assume that the curve  $X$  has a  $k$ -rational point  $P_0$ . Let  $Pic_0(X)_m$  be the subgroup of the  $m$ -torsion points in the group of divisor classes of degree 0 on  $X$ .

In [1] it is shown that one can reduce the evaluation of the discrete logarithm in  $Pic_0(X)_m$  by  $O(\log m)$  operations to the evaluation of the discrete logarithm in  $k(\zeta_m)^*$ , where  $\zeta_m$  is a primitive  $m$ -th root of unity, if the integer  $m$  is prime to  $p$ . If  $m = p$  and if the curve  $X$  is an elliptic curve (i.e.,  $g = 1$ ), then it is proved in [2] that the discrete logarithm in  $Pic_0(X)_p$  can be evaluated by  $O(\log p)$  operations in  $k$ . We want to extend this result to curves  $X$  of arbitrary genus  $g$ , and we will see that its proof is based on the connection between  $Pic_0(X)_p$  and logarithmic holomorphic differentials on  $X$ .

**Theorem.** *The discrete logarithm in  $Pic_0(X)_{p^n}$  can be evaluated by  $O(n^2 \log p)$  operations in  $k$ .*

*Proof.* Let  $x \in Pic_0(X)_{p^n}$  be an element of order  $p^n$  and let  $y$  be contained in the cyclic group generated by  $x$ . We have to show that  $\lambda \in \mathbb{Z}/p^n\mathbb{Z}$  with  $y = \lambda \cdot x$  can be evaluated by  $O(n^2 \log p)$  operations. It is a standard argument to reduce the evaluation of  $\lambda = \sum_{i=0}^{n-1} \lambda_i p^i$  with  $0 \leq \lambda_i < p$  to the evaluation of  $\lambda_i$  (by multiplication with  $p^i$ ,  $0 \leq i \leq n-1$ ) as solutions of  $n$  discrete logarithms in  $Pic_0(X)_p$ . Hence we can assume that  $n = 1$ .

The key point of the proof is the following result of Serre ([3], Proposition 10). Let  $\Omega^1(X)$  be the  $k$ -vector space of holomorphic differentials on  $X$ . Then there is an isomorphism from  $Pic_0(X)_p$  into  $\Omega^1(X)$  given by the following rule: Choose a divisor  $D$  of degree 0 with  $p \cdot D = (f)$ , where  $f$  is a function on  $X$ , then the divisor class  $\overline{D} \in Pic_0(X)_p$  is mapped to the holomorphic differential  $df/f$ .

---

Received by the editor August 8, 1997.

1991 *Mathematics Subject Classification.* Primary 11T71; Secondary 94A60.

©1999 American Mathematical Society

Now let  $t$  be a local parameter of  $P_0$ , then we get  $df/f = \frac{\partial f/\partial t}{f} dt$ . We evaluate the power series expansion

$$\frac{\partial f/\partial t}{f} = \sum_{i=0}^{\infty} a_i t^i \quad \text{with } a_i \in k.$$

We denote by  $(a_0, a_1, \dots, a_{2g-2})(f)$  the vector of the coefficients at  $1, t, \dots, t^{2g-2}$  of  $f^{-1}(\partial f/\partial t)$ . The Riemann-Roch theorem says that these coefficients determine the holomorphic differential  $df/f$  uniquely. Hence we get an isomorphism  $\phi$  from  $Pic_0(X)_p$  into  $k^{2g-1}$  which is defined by  $\phi(\overline{D}) = (a_0, a_1, \dots, a_{2g-2})(f)$ .

For elliptic curves this is the isomorphism in Lemma 2 of [2].

It remains to evaluate  $\phi(\overline{D})$  by  $O(\log p)$  operations, because the discrete logarithm in  $k^{2g-1}$  can be evaluated by this complexity. For this we modify the ideas of Chapter 3 in [1]. Since the addition in  $Pic_0(X)$  should be given explicitly, it is possible to solve the following problem:

- (\*) Let  $A^{(1)}$  and  $A^{(2)}$  be positive divisors of degree  $g$ ; find a positive divisor  $A^{(3)}$  of degree  $g$  and a function  $h$  such that the divisor of  $h$  is equal to  $A^{(1)} + A^{(2)} - A^{(3)} - gP_0$ .

Let  $S$  be a finite subgroup of  $Pic_0(X)_p$ . We suppose that  $S$  has a set of representatives  $\{A_s\}$  under  $c_g$  which are prime to  $P_0$  (here  $c_g$  is given by  $c_g(A_s) = \overline{A_s - gP_0}$ ).

We define the following group law on  $S \times k^{2g-1}$ :

$$(s_1, v_1) \odot (s_2, v_2) = (c_g(A_{s_3}), v_1 + v_2 + (a_0, \dots, a_{2g-2})(h)),$$

where  $A_{s_3} = A^{(3)}$  is the divisor and  $h$  is the function in (\*) corresponding to  $A^{(1)} = A_{s_1}$  and  $A^{(2)} = A_{s_2}$ ;  $(a_0, \dots, a_{2g-2})(h)$  is defined as above, even if the differential  $h^{-1}(\partial h/\partial t) dt$  has a pole at  $P_0$ . Furthermore  $s_3$  is the sum of  $s_1$  and  $s_2$  in  $S$ .

In other words we use the 2-cocycle  $S \times S \rightarrow k^{2g-1}$  with  $(s_1, s_2) \mapsto (a_0, \dots, a_{2g-2})(h)$  to define the group law. This is the additive version of the Tate pairing.

It can be shown easily by induction that  $(\overline{D}, 0) \odot \dots \odot (\overline{D}, 0)$  ( $p$ -times) is equal to  $(0, \phi(\overline{D}))$ .

Hence using repeated doubling in the group  $(\langle \overline{D} \rangle \times k^{2g-1}, \odot)$  we can evaluate  $\phi(\overline{D})$  by  $O(\log p)$  operations in the field  $k$ .

#### REFERENCES

- [1] G. Frey and H.-G. Rück, *A remark concerning  $m$ -divisibility and the discrete logarithm in the divisor class group of curves*, Math. Comp. **62** (1994), 865-874. MR **94h**:11056
- [2] I. A. Semaev, *Evaluation of discrete logarithms in a group of  $p$ -torsion points of an elliptic curve in characteristic  $p$* , Math. Comp. **67** (1998), 353-356. MR **98c**:94017
- [3] J. P. Serre, *Sur la topologie des variétés algébriques en caractéristique  $p$* , Sympos. Internat. Topologia Algebraica, Mexico City 1956, 24-53. MR **20**:4559

INSTITUT FÜR EXPERIMENTELLE MATHEMATIK, UNIVERSITÄT GH ESSEN, ELLERNSTR. 29,  
D-45326 ESSEN, GERMANY

*E-mail address*: rueck@exp-math.uni-essen.de