

NONTRIVIAL GALOIS MODULE STRUCTURE OF CYCLOTOMIC FIELDS

MARC CONRAD AND DANIEL R. REPLOGLE

ABSTRACT. We say a tame Galois field extension L/K with Galois group G has trivial Galois module structure if the rings of integers have the property that \mathcal{O}_L is a free $\mathcal{O}_K[G]$ -module. The work of Greither, Replogle, Rubin, and Srivastav shows that for each algebraic number field other than the rational numbers there will exist infinitely many primes l so that for each there is a tame Galois field extension of degree l so that L/K has nontrivial Galois module structure. However, the proof does not directly yield specific primes l for a given algebraic number field K . For K any cyclotomic field we find an explicit l so that there is a tame degree l extension L/K with nontrivial Galois module structure.

1. INTRODUCTION TO CYCLOTOMIC SWAN SUBGROUPS AND GALOIS MODULE THEORY

Let G be a group of finite order m . Let L/K be a tame (i.e., at most tamely ramified) Galois extension of algebraic number fields with finite Galois group $\text{Gal}(L/K) \cong G$. Let \mathcal{O}_L and \mathcal{O}_K denote the respective rings of algebraic integers. We say L/K has a trivial Galois module structure if \mathcal{O}_L is a free $\mathcal{O}_K[G]$ -module. Equivalently, one says in this case that L/K has a normal integral basis. (Note: One always has that \mathcal{O}_L is a rank one locally free $\mathcal{O}_K[G]$ -module whenever L/K is a tame Galois extension with Galois group G .)

The classical Hilbert-Speiser theorem proves that any abelian extension of \mathbb{Q} , the field of rational numbers, has a trivial Galois module structure. Call a field K Hilbert-Speiser if each tame abelian extension has a trivial Galois module structure. In [2] tame elementary abelian extensions and Swan modules are considered to find conditions a Hilbert-Speiser field must satisfy. Let $V_l = (\mathcal{O}_K/l\mathcal{O}_K)^*/\text{Im}(\mathcal{O}_K^*)$, where for any ring S we let S^* denote its group of multiplicative units, and Im denotes the image of \mathcal{O}_K^* under the canonical surjection $\psi : \mathcal{O}_K \rightarrow \mathcal{O}_K/l\mathcal{O}_K$. Then we have the following theorem.

Theorem 1.0 ([4, Theorem 1]). *Let K be a Hilbert-Speiser number field. Then:*

- (i) *The class number of K is one.*
- (ii) *For each odd prime l the group V_l has exponent dividing $\frac{(l-1)^2}{2}$.*
- (iii) *The group V_2 is trivial.*

Received by the editor November 6, 2000 and, in revised form, July 15, 2001.

2000 *Mathematics Subject Classification*. Primary 11R33, 11R29; Secondary 11R27, 11R18.

Key words and phrases. Swan subgroups, cyclotomic units, Galois module structure, tame extension, normal integral basis.

This theorem and a Galois theoretic argument are used to show that for any $K \neq \mathbb{Q}$ there is some odd prime l for which condition (ii) is violated. Thus we have the following theorem.

Theorem 1.1 ([4, Theorem 2]). *Among all algebraic number fields, only the rational numbers are Hilbert-Speiser.*

For the convenience of the reader we outline the ideas in the proofs of these two results. The overall idea is that V_l^{l-1} will be seen to be a lower bound on the Swan subgroup, and $V_l^{(l-1)^2/2}$ will be seen to be a lower bound on the group of realisable classes. Let $\Lambda = \mathcal{O}_K[G]$. Then Λ is an order in the group algebra $K[G]$. For each s in \mathcal{O}_K so that s and m are relatively prime, one defines the Swan module $\langle s, \Sigma \rangle$ by $\langle s, \Sigma \rangle = s\Lambda + \Lambda\Sigma$, where $\Sigma = \sum_{g \in G} g$. It is easily shown that each $\langle s, \Sigma \rangle$ is a rank one locally free Λ -module. (See [15, Proposition 2] for example.) Hence each Swan module determines a class in the locally free class group $Cl(\Lambda)$. We denote the class of $\langle s, \Sigma \rangle$ by $[s, \Sigma]$. Denote the set (at this point) of all classes of Swan modules over Λ by $T(\Lambda)$. Let $D(\Lambda)$, the kernel group, denote the subgroup of $Cl(\Lambda)$ of all classes that become trivial upon extension of scalars to the maximal order in $K[G]$ containing Λ . Let $\Gamma = \Lambda/\Sigma\Lambda$, let $\psi : \mathcal{O}_K \rightarrow \mathcal{O}_K/m\mathcal{O}_K = \overline{\mathcal{O}_K}$ denote the canonical quotient map, let $\epsilon : \Lambda \rightarrow \mathcal{O}_K$ denote the augmentation map, and let $\bar{\epsilon} : \Gamma \rightarrow \overline{\mathcal{O}_K}$ be induced from ϵ . The main result of [9] shows that if the group algebra $K[G]$ satisfies an Eichler condition (which is always satisfied if G is abelian), then there exists a Mayer-Vietoris exact sequence

$$(1.0) \quad \mathcal{O}_K^* \times \Gamma^* \xrightarrow{h} \overline{\mathcal{O}_K}^* \xrightarrow{\delta} D(\Lambda) \rightarrow D(\Gamma) \oplus D(\mathcal{O}_K) \rightarrow 0,$$

where for any ring S we have been denoting its group of multiplicative units by S^* . It is further shown in [9] that the map h is given by $h[(u, v)] = \psi(u)\bar{\epsilon}(v)^{-1}$. From [15] we have that the map δ is given by $\delta(s) = [s, \Sigma]$. Hence we have from (1.0) that $T(\Lambda)$ is a subgroup of $D(\Lambda)$ and

$$(1.1) \quad T(\Lambda) \cong (\overline{\mathcal{O}_K}^*)/h(\mathcal{O}_K^* \times \Gamma^*).$$

Now let $V_l = (\mathcal{O}_K/l\mathcal{O}_K)^*/Im(\mathcal{O}_K^*)$ and set $G \cong C_l$, the cyclic group of prime order l . Then in [4] a lower bound on $T(\Lambda)$ is deduced from (1.1). Namely, one may show that for $G \cong C_l$ and $\Lambda = \mathcal{O}_K[C_l]$ there is a natural surjective map

$$(1.2) \quad T(\Lambda) \rightarrow V_l^{l-1}.$$

See [4, Theorem 5], which handles the elementary abelian group case.

Now consider tame Galois extensions of K with $Gal(L/K) \cong C_l$, where C_l is the cyclic group of order l . Of course each such extension L has the property that \mathcal{O}_L is a locally free rank one $\mathcal{O}_K[G]$ -module. Hence each \mathcal{O}_L determines a tame Galois module class $[\mathcal{O}_L]$ in the locally free class group $Cl(\Lambda)$. Denote by $R(\Lambda)$ the set of all such classes. Observe that $R(\Lambda)$ measures to what extent a given base field K fails to have Galois extensions with Galois group G having nontrivial Galois module structure. In [7] McCulloh gives an explicit description of $R(\Lambda)$ as a subgroup of $Cl(\Lambda)$ for the case considered here. We note that in [8] McCulloh shows that $R(\Lambda)$ is in fact a subgroup of $Cl(\Lambda)$ for all abelian G ; however, that result is not used here. Using the description of $R(\Lambda)$ from [7], we have the following relationship between $R(\Lambda)$, $D(\Lambda)$, and $T(\Lambda)$, stated for the case we are considering. If $G \cong C_l$, where $l > 2$ is prime, then by [4, Proposition 4] the Swan subgroup has the property

that

$$(1.3) \quad T(\Lambda)^{(l-1)/2} \subseteq R(\Lambda) \cap D(\Lambda).$$

Observe that condition (ii) of Theorem 1.0 follows from (1.2) and (1.3).

The proof of Theorem 1.1 uses the Chebotarev density theorem to establish the existence of infinitely many primes for which condition (ii) of Theorem 1.0 is violated. It is not clear whether one can use the proof of Theorem 1.1 to actually write down any explicit such primes. In fact, as the proof of [4, Theorem 2] uses the Chebotarev density theorem to show the existence of such a prime by showing infinitely many must exist, it is unlikely that one could use that argument to write down any such primes. We adopt the following terminology. If, given a field K , we have an explicit prime l so that there is a tame Galois extension L/K with $Gal(L/K) \cong C_l$, the cyclic group of order l , so that \mathcal{O}_L is not a free $\mathcal{O}_K[C_l]$ -module, we say K is not Hilbert-Speiser for l . Then Theorem 1.1 says that, for any algebraic number field K , there is some l for which K is not Hilbert-Speiser for l .

In this article we show that for any cyclotomic field one may in fact find a specific prime l for which the field is not Hilbert-Speiser for l . That is, let $K_n = \mathbb{Q}(\zeta_n)$, where ζ_n is a primitive n th root of unity. Note that without loss of generality we may assume $n \not\equiv 2 \pmod{4}$, as if $n \equiv 2 \pmod{4}$ then $K_n = K_{\frac{n}{2}}$.

For K_n the ring of algebraic integers is $\mathbb{Z}[\zeta_n]$, which we denote by \mathcal{O}_n . The class number of an algebraic number field K is denoted by h_K and for K_n is denoted by h_n . In light of Theorem 1.0, to accomplish our goal it suffices to prove the following theorem, the main result for this article.

Theorem 1.2 (Main Theorem). *Let K_n be as above and assume $h_n = 1$. One may find an explicit prime l so that $V_{l,n} = (\mathcal{O}_n/l\mathcal{O}_n)^*/Im(\mathcal{O}_n^*)$ does not have exponent dividing $\frac{(l-1)^2}{2}$. Hence for every cyclotomic field one may find an explicit prime l so that there is a tame degree l field extension with nontrivial Galois module structure.*

We begin our study by taking care of two special cases. We first treat the case $h_n \neq 1$ by providing some details about condition (i) of Theorem 1.0. Using [6], one can show that any field of class number not equal to one has a quadratic extension which does not possess a relative integral basis. Thus we have the following restatement of this in our language.

Theorem 1.3. *Let K be an algebraic number field with class number $h_K \neq 1$. Then K is not Hilbert-Speiser for $l = 2$.*

While this result is decisive for fields of class number not equal to 1, we must note that much more recent studies of Galois module theory have been made. For example, the work of Fröhlich [3] gives an excellent account of Galois module classes. The results in [1] which we state below provide an important Galois module structure result for extensions of cyclotomic fields.

Next we treat the case when n is prime. We note that for this case one need not make any assumptions on the class number.

Proposition 1.4. *For $n > 3$ prime, K_n is not Hilbert-Speiser for n .*

Proof. Let $n > 3$ be prime. We show that $V_n = (\mathcal{O}_n/n\mathcal{O}_n)^*/Im(\mathcal{O}_n)$ is an elementary abelian group of exponent n , from which the result follows from Theorem 1.0 (ii). As n totally ramifies, we have $(\mathcal{O}_n/l\mathcal{O}_n)^* \cong C_{n-1} \times C_n^{n-2}$, and from Dirichlet's unit theorem we have $\mathcal{O}_n^* \cong \langle -\zeta_n \rangle \times \langle \epsilon_1 \rangle \times \cdots \times \langle \epsilon_{(n-3)/2} \rangle$, where the ϵ_i are a system

of fundamental units. Since ζ_n is not congruent to 1 mod l , $V_n^{n-1} \cong C_n^{l-2-j}$ for some integer j with $1 \leq j \leq (n-1)/2$. Now, as $n \geq 5$, we have that V_n^{n-1} is a nontrivial elementary abelian group of exponent n . \square

Notes. 1. Proposition 1.4 is essentially [10, Proposition 15].

2. In the case when both n is prime and $n \nmid h_n^+$, where h_n^+ is the class number of the maximal real subfield of K_n , one may in fact explicitly compute $T(\mathbb{Z}[\zeta_n]C_n)$. See [11, Theorem 1].

3. For applications to Hopf orders in $\mathbb{Z}[\zeta_n]C_n$ when n is a prime power, see [12].

So, we need to consider only the case where either $n = 3$ or n is composite so that $h_n = 1$. This reduces the question to the following list. (See [16, Theorem 11.1], for example, for a listing of those cyclotomic fields K_n with $n \not\equiv 2 \pmod 4$ so that $h_n = 1$.)

List 1.5. *Let n be not congruent to 2 mod 4 and not a prime greater than 3. Then $h_n = 1$ only if $n = 3, 4, 8, 9, 12, 15, 16, 20, 21, 24, 25, 27, 28, 32, 33, 35, 36, 40, 44, 45, 48, 60$, or 84.*

In the next section we will provide several primes l for each n in this list so that K_n is not Hilbert-Speiser for l . That is, for each n we will exhibit several primes l so that $V_{l,n}$ does not have exponent dividing $\frac{(l-1)^2}{2}$. This will complete the proof of the theorem.

Before beginning we introduce the important related result of Chan and Lim, [1]. For any Galois extension of number fields L/K with Galois group G one may define the associated order by $\mathcal{A}_{L/K} = \{\alpha \in K[G] \mid \alpha \mathcal{O}_L \subseteq \mathcal{O}_L\}$. Then a classical result is L/K is at most tamely ramified if and only if $\mathcal{A}_{L/K} = \mathcal{O}_K[G]$. So, one may generalize the notion of trivial Galois module structure to say that a Galois extension of number fields L/K has trivial Galois module structure if \mathcal{O}_L is a free $\mathcal{A}_{L/K}$ -module. Chan and Lim show in [1] that any Galois extension L/K with L and K both cyclotomic fields has a trivial Galois module structure. Hence, for the primes l for which we detect the existence of degree l extensions with nontrivial Galois module structure, we know these extension fields are not cyclotomic fields.

2. CYCLOTOMIC UNITS AND NONTRIVIAL GALOIS MODULE STRUCTURE

To complete the proof of Theorem 1.2 it suffices to exhibit a prime l for each n in List 1.5 so that $V_{l,n} \cong (\mathcal{O}_n/l\mathcal{O}_n)^*/\text{Im}(\mathcal{O}_n^*)$ is not of exponent dividing $\frac{(l-1)^2}{2}$. Of course the issues are finding an l that splits nicely in \mathcal{O}_n ; then representing the quotient $\mathcal{O}_n/l\mathcal{O}_n$ in a way amenable to computation; then finding the image of the units of \mathcal{O}_n in this representation; then finally developing a computer algorithm to solve the problem. We will not use the results in [5] here, but we must note that our method is somewhat analogous.

The first two parts of our task are completed by the following lemma. Let \mathbb{F}_l denote the field of l elements and $\mathbb{F}_l[x]$ its polynomial ring. Further, let ϕ be the usual Euler ϕ -function.

Lemma 2.0. *For an l that does not divide n let f be minimal such that $n|k := l^f - 1$, and $g = \phi(n)/f$. Then $(\mathcal{O}_n/l\mathcal{O}_n)^* \cong (\mathbb{Z}/k\mathbb{Z})^g$, where the isomorphism is*

explicitly given by the following maps κ , λ_i , and μ_i :

$$(2.0) \quad \kappa : \mathcal{O}_n/l\mathcal{O}_n \longrightarrow \prod_{i=1}^g \mathcal{O}_n/P_i\mathcal{O}_n,$$

$$a + l\mathcal{O}_n \mapsto (a + P_i\mathcal{O}_n)_{i=1, \dots, g},$$

where each P_i is an ideal generated by two generators $P_i = \langle l, p_i \rangle$, and p_i is a polynomial of degree f , which we obtain by factoring the n -th cyclotomic polynomial $\Phi_n(x)$ as a polynomial in $\mathbb{F}_l[x]$ into irreducible polynomials. Also,

$$(2.1) \quad \lambda_i : \mathcal{O}_n/P_i\mathcal{O}_n \rightarrow \mathbb{F}_l[x]/(p_i),$$

$$a + P_i\mathcal{O}_n \mapsto a + (p_i),$$

for $i = 1, \dots, g$; and, finally,

$$(2.2) \quad \mu_i : (\mathbb{F}_l[x]/(p_i))^* \rightarrow \mathbb{Z}/k\mathbb{Z},$$

$$a + (p_i) \mapsto b \pmod k,$$

where $a + (p_i) = w_i^b$ and w_i is a (fixed) generator of the cyclic group $(\mathbb{F}_l[x]/(p_i))^*$.

Proof. The isomorphism (2.0) follows by [16, Theorem 2.13], which describes the splitting behaviour of a prime in K_n , and the Chinese remainder theorem for ideals in an algebraic number field. The isomorphism (2.1) can be proved straightforwardly as $\mathcal{O}_n \cong \mathbb{Z}[x]/(\Phi_n(x))$, and (2.2) is the well known fact that the multiplicative group of a finite field is cyclic. \square

The larger issue is the units. We show first that it is sufficient to consider cyclotomic units. We introduce some mostly standard notation at this point: Let W_n denote the roots of unity in \mathcal{O}_n . Let E_n denote the units of \mathcal{O}_n . Last, let E_n^+ denote the units in the ring of algebraic integers of the maximal real subfield of K_n . Then we have $[E_n : W_n E_n^+] = Q$, where Q equals one if n is a prime power and equals two otherwise. For details, see chapters 4 and 8 of [16].

Let C_n^+ denote the cyclotomic units in E_n^+ . Let h_n^+ denote the class number of the maximal real subfield of K_n . Next we have the following result, due to Sinnott [14], stated only for what we are considering here.

Theorem 2.1 ([14]). *The group E_n^+/C_n^+ is finite, and $[E_n^+ : C_n^+] = 2^b h_n^+$. The integer b is defined by $b = 0$ if $r = 1$ and $b = 2^{r-2} + 1 - r$ if $r > 1$, where r is the number of distinct primes dividing n .*

This theorem has two immediate corollaries which we prove.

Corollary 2.2. *If $h_n = 1$, then $E_n^+ = C_n^+$.*

Proof. For each K_n of class number one, three or fewer primes divide n . Hence $b = 0$ in Theorem 2.1. \square

Corollary 2.3. *For each n of List 1.5 we have $C_n = E_n$.*

Proof. Note first that $E_n^+ = C_n^+$ by Corollary 2.2, and obviously we have $C_n \supseteq W_n C_n^+$. If n is a prime power we have $Q = 1$, and therefore $E_n = W_n E_n^+ = W_n C_n^+ = C_n$.

In the case $Q = 2$ the result is proven when we show that $[C_n : W_n C_n^+] > 1$. This follows because $1 - \zeta_n \in C_n$ but $\zeta_n^\nu(1 - \zeta_n) = \zeta_n^\nu - \zeta_n^{\nu+1} \notin \mathbb{R} \supseteq C_n^+$ for each $\nu = 1, \dots, n$. (A necessary condition for $\zeta_n^\nu - \zeta_n^\mu \in \mathbb{R}$ is $2(\nu + \mu) = n$, which is impossible for $\mu = \nu + 1$ as $n \not\equiv 2 \pmod{4}$.) \square

From this it follows we may consider just cyclotomic units. That is, the group of cyclotomic units generates the full group of units. From [2] we have an explicit description of a basis of the cyclotomic units in K_n .

Proposition 2.4 ([2]). *A basis of the group of cyclotomic units is explicitly given as a subset of the set*

$$(2.3) \quad \left\{ \frac{1 - \zeta_q^a}{1 - \zeta_q} \mid q|n, 1 < a < q, (a, q) = 1, q \text{ is a prime power} \right\} \\ \cup \{1 - \zeta_d^a \mid d|n, 1 < a < d, (a, d) = 1, d \text{ is not a prime power} \} \\ \cup \{\pm \zeta_n\}.$$

An algorithm which computes a basis according to [2] is implemented in SIMATH [13]. Now we work toward developing an algorithm to compute $V_{i,n}$. We will need the following lemma.

Lemma 2.5. *Let $k, s, g \in \mathbb{N}$ and $G = (\mathbb{Z}/k\mathbb{Z})^g$. Further, let $H = \langle v_1, \dots, v_g \rangle$ be a subgroup of G which is generated by the g -tuples $v_i = (v_{1,i}, \dots, v_{g,i})$ and $V = (v_{i,j})_{1 \leq i, j \leq g}$, the $g \times g$ -matrix which is made up by the v_i . Then we have:*

If there exists a prime p such that $p \mid k$, $p \mid \det(V)$ and $p \nmid s$, then the exponent of G/H does not divide s .

Proof. “The exponent of G/H does not divide s ” means that $\exists w \in G$ such that $sw \notin H$, or—more concretely— $\exists w \in G \forall \alpha := (\alpha_1, \dots, \alpha_g) \in \mathbb{Z}^g$:

$$(2.4) \quad \sum_{i=1}^g \alpha_i v_i \not\equiv sw \pmod{k}.$$

A sufficient condition for (2.4) is obviously that there exists a prime $p \mid k$ with

$$(2.5) \quad \sum_{i=1}^g \alpha_i v_i \not\equiv sw \pmod{p}.$$

The only primes p which are in question are those p with $p \nmid s$; otherwise $(0, \dots, 0)$ would be a solution. For those p the number s^{-1} is well defined mod p , and we obtain from (2.5) that we have to show: $\exists w \in G$ with $\alpha s^{-1} V \not\equiv w \pmod{p}$. So in fact we have to show that the endomorphism

$$\varphi : (\mathbb{Z}/p\mathbb{Z})^g \rightarrow (\mathbb{Z}/p\mathbb{Z})^g, \\ x \mapsto xs^{-1}V,$$

is not surjective, which is equivalent to $\det(s^{-1}V) \equiv 0 \pmod{p}$, or $p \mid \det(V)$. \square

Remark 2.6. Of course the subgroup H can be generated by fewer than g vectors. Then we obtain the lemma by dropping the condition $p \mid \det(V)$.

Remark 2.7. Note that we do not need to factor k in order to check whether a p as required in Lemma 2.5 exists. We can use the following algorithm:

- i) Let $c := \gcd(k, \det(V))$.
- ii) do $t := c$; $c := c / \gcd(c, s)$ while $c \neq t$.
- iii) if $c = 1$ then $\nexists p$ else $\exists p$.

Except for the proof that the group of cyclotomic units already is the full unit group we have not used the fact that $h_n = 1$. So we define

$$V_{l,n}^{cyc} \cong (\mathcal{O}_n / l\mathcal{O}_n)^* / \text{Im}(C_n),$$

where we have replaced the full unit group \mathcal{O}_n^* by the group of cyclotomic units C_n . With the results of Lemmata 2.0 and 2.5 we obtain the following algorithm.

Algorithm 2.8. *If for a given pair (l, n) with $l \nmid n$ the following algorithm returns “yes”, then the exponent of $V_{l,n}^{cyc}$ does not divide $(l - 1)^2/2$.*

- i) Factor the n -th cyclotomic polynomial Φ_n modulo l . Let $L = \{p_1, \dots, p_g\}$ be the list of irreducible factors of Φ_n , let f be the degree of one of the p_i (which is in fact the same for all p_i), and set $k = l^f - 1$.
- ii) For each polynomial $p_i \in L$ determine a generator w_i of the finite field $\mathbb{F}_l[x]/(p_i(x))$.
- iii) Determine a (finite) set $B = \{b_1, \dots, b_s\}$ of generators of cyclotomic units. This may be done e.g. with the function `lcycbas()` of SIMATH, [13].
- iv) Compute the matrix $V \in (\mathbb{Z}/k\mathbb{Z})^{s \times g}$, where each entry $v_{i,j}$ of V is given by solving the discrete logarithm problem $b_i \equiv w_j^{v_{i,j}} \pmod{p_j}$ in $\mathbb{F}_l[x]$.
- v) If $s > g$, eliminate rows of V by Gauss elimination over the ring $\mathbb{Z}/k\mathbb{Z}$, until V is a square matrix. If $s < g$, generate a square matrix by adding zero rows (or set $\det(V) = 0$ in the next step).
- vi) Check (via Remark 2.7) if there exists a prime p such that $p \mid \gcd(\det(V), k)$ and $p \nmid ((l - 1)^2/2)$. If such a p exists, return “yes”.

This algorithm has been implemented and is available as function `iscynfHS()` in Version 4.5 of SIMATH.

For each n in List 1.5 we now in fact exhibit several primes l for which $V_{l,n}^{cyc}$ is not of exponent dividing $\frac{(l-1)^2}{2}$. For these n one has $V_{l,n}^{cyc} = V_{l,n}$, as $h_n = 1$. So, we provide a table, obtained using the above algorithm, such that for each n in List 1.5 we have $K_n = \mathbb{Q}(\zeta_n)$ and for the l associated to n we have that there is a tame Galois field extension L/K_n of degree l for which l splits in K_n/\mathbb{Q} and the extension L/K_n is so that L/K_n is not Hilbert-Speiser, i.e., does not have a normal integral basis. This together with Theorem 1.3 and Proposition 1.4 proves Theorem 1.2. We note that the algorithm presented applies to those n treated in Proposition 1.4 also. For example, for $n = 7$ one obtains the list $l = 5, 11, 13, 17, 23, 37, 41, 53, 67, 79, 83, 97$. Of course, these l split, whereas 7 totally ramifies in K_7 .

TABLE 2.9.

n	l
3	17, 29, 41, 53, 59, 71, 83, 89
4	11, 19, 23, 43, 47, 59, 67, 71, 79, 83
8	5, 11, 13, 19, 23, 29, 37, 43, 47, 53, 59, 61, 67, 71, 79, 83
9	5, 7, 11, 13, 17, 31, 43, 53, 61, 67, 71, 79, 89, 97
12	11, 17, 19, 23, 29, 41, 43, 47, 53, 59, 67, 71, 79, 83, 89
15	7, 11, 13, 17, 19, 23, 29, 37, 41, 43, 47, 53, 59, 67, 71, 73, 79, 89
16	3, 5, 11, 13, 19, 23, 29, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 89
20	7, 11, 13, 17, 19, 23, 29, 37, 43, 47, 53, 59, 67, 71, 73, 79, 89
21	5, 11, 13, 17, 29, 37, 41, 67, 71, 79, 83, 97
24	5, 11, 13, 17, 19, 23, 29, 37, 41, 43, 47, 53, 59, 61, 67, 71, 79, 83, 89
25	7, 11, 31, 43
27	17, 19, 37, 53, 73
28	3, 5, 11, 13, 17, 37, 41, 43, 53, 71, 83, 97
32	3, 5, 7, 17, 23, 41, 47, 71, 73, 79
33	5, 23, 43, 89
35	3, 11, 13, 29, 41, 43
36	5, 7, 11, 13, 17, 19, 53, 61, 71, 89, 97
40	3, 7, 11, 13, 17, 19, 23, 29, 37, 43, 47, 53, 59, 61, 67, 71, 73, 79, 89
44	3, 5, 23, 43, 67
45	11, 17, 19, 31, 37, 53, 61, 71, 73, 89
48	5, 11, 13, 17, 19, 23, 29, 37, 41, 43, 47, 53, 59, 71, 73, 79, 89
60	7, 11, 13, 17, 19, 23, 29, 37, 41, 43, 47, 53, 59, 67, 71, 73, 79, 89
84	5, 11, 13, 17, 29, 37, 41, 43, 71, 83, 97

ACKNOWLEDGMENTS

The authors would wish to thank L. Washington for conversations regarding this project and for his excellent text ([16]) that both of the present authors use frequently.

REFERENCES

- [1] S-P Chan and C-H Lim, *Relative Galois module structure of rings of integers of cyclotomic fields*, J. Reine Angew. Math. **434** (1993), 205-230. MR **93i**:11127
- [2] M. Conrad, *Construction of bases for the group of cyclotomic units*, J. Num. Theory **81** (2000), 1-15. MR **2001f**:11182
- [3] A. Fröhlich, *Galois Module Structure of Algebraic Integers*, Springer-Verlag, Berlin, 1983. MR **85h**:11067
- [4] C. Greither, D. R. Replogle, K. Rubin, and A. Srivastav, *Swan Modules and Hilbert-Speiser number fields*, J. Num. Theory **79** (1999), 164-173. MR **2000m**:11111
- [5] T. Kohl and D. R. Replogle, *Computation of several Cyclotomic Swan Subgroups*, Math. Comp., **71** (2002), 343-348.
- [6] H. B. Mann, *On integral bases*, Proc. Amer. Soc. Math. **9** (1958), 162-172. MR **20**:26
- [7] L. R. McCulloh, *Galois Module Structure of Elementary Abelian Extensions*, J. Alg. **82** (1983), 102-134. MR **85d**:11093
- [8] L. R. McCulloh, *Galois Module Structure of Abelian Extensions*, J. Reine Angew. Math. **375/376** (1987), 259-306. MR **88k**:11080
- [9] I. Reiner and S. Ullom, *A Mayer-Vietoris sequence for class groups*, J. Alg. **31** (1974), 305-342. MR **50**:2321

- [10] D. R. Replegle, *Swan Modules and Realisable Classes for Kummer Extensions of Prime Degree*, *J. Alg.* **212** (1999), 482-494. MR **2000a**:11161
- [11] D. R. Replegle, *Cyclotomic Swan subgroups and irregular indices*, *Rocky Mountain Journal Math.* **31** (Summer 2001), 611-618.
- [12] D. R. Replegle and R. G. Underwood, *Nontrivial tame extensions over Hopf orders*, *Acta Arithmetica* (to appear).
- [13] The SIMATH group/H. G. Zimmer, *SIMATH: A computer algebra system for algebraic number theory*, www.simath.info.
- [14] W. Sinnott, *On the Stickelberger ideal and the circular units of a cyclotomic field*, *Annals Math.* **108** (1978), 107-134. MR **58**:5585
- [15] S. V. Ullom, *Nontrivial lower bounds for class groups of integral group rings*, *Illinois Journal Mathematics* **20** (1976), 361-371. MR **52**:14024
- [16] L. Washington, *Introduction to Cyclotomic Fields*, Graduate Texts in Mathematics 83, Springer-Verlag, New York, 1982. MR **85g**:11001

FACULTY OF TECHNOLOGY, SOUTHAMPTON INSTITUTE, EAST PARK TERRACE, SOUTHAMPTON,
S014 0YN GREAT BRITAIN

E-mail address: `marc@pension-perisic.de`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, COLLEGE OF SAINT ELIZABETH, 2
CONVENT ROAD, MORRISTOWN, NEW JERSEY 07960

E-mail address: `dreplegle@cse.edu`