

MAXIMUM NORM STABILITY OF DIFFERENCE SCHEMES FOR PARABOLIC EQUATIONS ON OVERTSET NONMATCHING SPACE-TIME GRIDS

T. P. MATHEW AND G. RUSSO

ABSTRACT. In this paper, theoretical results are described on the maximum norm stability and accuracy of finite difference discretizations of parabolic equations on overset nonmatching space-time grids. We consider parabolic equations containing a linear reaction term on a space-time domain $\Omega \times [0, T]$ which is decomposed into an overlapping collection of cylindrical subregions of the form $\Omega_l^* \times [0, T]$, for $l = 1, \dots, p$. Each of the space-time domains $\Omega_l^* \times [0, T]$ are assumed to be independently grided (in parallel) according to the local geometry and space-time regularity of the solution, yielding space-time grids with mesh parameters h_l and τ_l . In particular, the different space-time grids need not match on the regions of overlap, and the time steps τ_l can differ from one grid to the next. We discretize the parabolic equation on each local grid by employing an explicit or implicit θ -scheme in time and a finite difference scheme in space satisfying a discrete maximum principle. The local discretizations are coupled together, without the use of Lagrange multipliers, by requiring the boundary values on each space-time grid to match a suitable interpolation of the solution on adjacent grids. The resulting global discretization yields a large system of coupled equations which can be solved by a parallel Schwarz iterative procedure requiring some communication between adjacent subregions. Our analysis employs a contraction mapping argument.

Applications of the results are briefly indicated for reaction-diffusion equations with contractive terms and *heterogeneous* hyperbolic-parabolic approximations of parabolic equations.

1. INTRODUCTION

In this paper, theoretical bounds are described (extending results in [32, 10]) for the maximum norm stability and convergence of discretizations of parabolic equations on nonmatching, overset space-time grids. Nonmatching overset spatial grids are popular in several fluid dynamics computations involving complex geometries [33, 13]. They permit independent (parallel) generation of local grids adapted to the local geometry (without the restriction of matching the grids on the regions of overlap) at the cost of increased computations in coupling the various local discretizations. For evolution problems, additional flexibility can be obtained

Received by the editor July 25, 2000.

2000 *Mathematics Subject Classification*. Primary 65N20, 65F10.

Key words and phrases. Nonmatching overset space-time grids, maximum norm stability, composite grids, parallel Schwarz alternating method, parabolic equations, discrete maximum principle, discrete barrier functions.

by permitting different time steps and choice of explicit or implicit schemes on each of the different space-time subregions [33, 13, 17, 18, 7, 23, 5, 7, 20, 22].

In the computational literature, several approaches have been proposed for coupling discretizations on nonmatching grids. These include Lagrange multipliers (including mortar methods) and least squares based techniques (see [33, 13, 32, 5, 4, 9, 23, 20, 1, 2, 10, 19]). The method considered in this paper does not use either Lagrange multipliers or least squares to couple the various local problems. It is simpler to implement (see [32, 10]); however, it applies only to a certain class of parabolic equations exhibiting a contraction property, and it requires overlap amongst adjacent grids.

Our study will be restricted to a small class of parabolic equations of the form

$$(1.1) \quad \begin{cases} u_t - a\Delta u + \vec{b}(x) \cdot \nabla u + c(x)u &= f(x, t), & \text{in } \Omega \times [0, T] \\ u(x, t) &= 0, & \text{on } \partial\Omega \times [0, T], \\ u(x, 0) &= u_0(x), & \text{on } \Omega, \end{cases}$$

where $f(x, t)$, $\vec{b}(x)$, $c(x)$ and $u_0(x)$ are sufficiently smooth functions and $a > 0$. Here $\Omega \subset R^d$ for $d = 1, 2, \dots$. In order to have a contraction property for homogeneous solutions, we will require that

$$c(x) \geq c_0 > 0,$$

for some positive constant c_0 .

Given the cylindrical space-time domain $\Omega \times [0, T]$, we decompose it into an overlapping collection of *cylinders* of the form $\{\Omega_l^* \times [0, T]\}_{l=1}^p$ that form a covering of $\Omega \times [0, T]$. Each cylinder $\Omega_l^* \times [0, T]$ will be assumed to be triangulated by a space-time grid with mesh and time parameters h_l and τ_l (see Figure 1.1). We employ finite difference methods in space and implicit or explicit θ -schemes in time, independently on each space-time grid. On each subdomain boundary $\partial\Omega_l^* \times [0, T]$, we require the local solution to match some suitably chosen interpolant of the solution from adjacent grids (see [33, 32, 13, 10]).

Our main result in the paper, stated in Theorem 4.4, concerns the accuracy of the global discretization. Let $u_{h,\tau}$ denote the restriction of the exact solution u of the parabolic equation to all the space-time gridpoints, and let $U_{h,\tau}$ denote the computed solution of the global discretization. Suppose the truncation and intergrid interpolation errors for the discretization and boundary conditions, respectively, on the l th space-time grid $\Omega_l^* \times [0, T]$ satisfy

$$\text{Local truncation error on } \Omega_l^* \times [0, T] = \|u\|_{q_l+2, q_l+2, \infty, \Omega_l^* \times [0, T]} (h_l^{q_l+1} + \tau_l^{q_l+2}),$$

$$\text{Local interpolation error on } \partial\Omega_l^* \times [0, T] = \|u\|_{r_l+1, r_l+2, \infty, B_2^{l,*} \times [0, T]} (h_l^{r_l+1} + \tau_l^{r_l+2}),$$

where $\|\cdot\|_{q-l;1, q-l;2, \infty, \Omega_l^* \times [0, T]}$ and $\|\cdot\|_{r-l;1, r-l;2, \infty, B_2^{l,*} \times [0, T]}$ denote Sobolev norms. Here $B_2^{l,*}$ is a small spatial region covering the the boundary segment $B_2^l = (\partial\Omega_l^* \cap \Omega)$ of the l th spatial subdomain. Theorem 4.4 states that, under suitable assumptions, the maximum norm of the global error $u_{h,\tau} - U_{h,\tau}$ satisfies the bound

$$\begin{aligned} \|u_{h,\tau} - U_{h,\tau}\| \leq C \max_l \{ & \|u\|_{q-l;1+2, q-l;2+2, \infty, \Omega_l^* \times [0, T]} (h_l^{q_l+1} + \tau_l^{q_l+2}) \\ & + \|u\|_{r_l+1, r_l+2, \infty, B_2^{l,*} \times [0, T]} (h_l^{r_l+1} + \tau_l^{r_l+2}) \}, \end{aligned}$$

where $C > 0$ is independent of the mesh sizes. From this we deduce that, ideally, the local grid sizes h_l and τ_l should be chosen so that all the local error terms are “balanced”. This would mean smaller h_l and τ_l on regions where the solution is




FIGURE 1.1. Sample nonmatching overset space-time grid.

less regular. Additionally, the intergrid interpolation maps for determining local boundary data should be chosen so that the interpolation errors are balanced with the local truncation errors. Then, the resulting accuracy of the global discretization will be of optimal order.

The rest of the paper is outlined as follows. In Section 2, we introduce notation for the overlapping space-time subregions, discuss explicit and implicit local discretizations on each space-time grid, intergrid interpolation maps, the global discretization, and a parallel Schwarz iterative procedure for solving the resulting system of equations. In Section 3, we discuss theoretical properties of the local schemes, such as *a priori* estimates, maximum principles, comparison theorems, barrier functions, and contraction properties of homogeneous solutions. In Section 4, we analyze the stability and accuracy of our global space-time discretization by employing Picard's contraction mapping theorem.

2. GLOBAL DISCRETIZATION ON NONMATCHING OVERLAPPING SPACE-TIME GRIDS

In this section, we describe the construction of a global discretization of (1.1) and a parallel Schwarz iterative method for solving the resulting large system of equations.

2.1. Space-time subdomains. Let $\Omega \times [0, T]$ denote the space-time region on which the parabolic equation (1.1) is posed. We will describe here the construction of an overlapping collection of space-time subregions that covers the above region.

Let $\{\Omega_l\}_{l=1}^p$ denote a partition or a covering of the spatial domain Ω :

$$\bar{\Omega} \subset \bigcup_{l=1}^p \bar{\Omega}_l.$$

In practice, the subregions Ω_l may be chosen according to the geometry of Ω or the regularity of the solution (if known, or else by estimating the regularity from prior numerical approximations). For each subregion Ω_l , choose a parameter $\beta_l > 0$ and enlarge Ω_l to Ω_l^* as

$$\Omega_l^* \equiv \{x \in \Omega : \text{dist}(x, \Omega_l) < \beta_l\}.$$

The collection of subregions $\{\Omega_l^*\}_{l=1}^p$ will form an *overlapping covering* of Ω (with overlap parameters β_l). An overlapping covering of the space-time region $\Omega \times [0, T]$ can be immediately constructed:

$$\Omega \times [0, T] = \bigcup_{l=1}^p (\Omega_l^* \times [0, T]),$$

where each space-time subregion $\Omega_l^* \times [0, T]$ is cylindrical.

We will denote the boundary of each spatial subregion Ω_l^* by $B^l \equiv \partial\Omega_l^*$. It will be convenient to further partition each boundary $B^l = \partial\Omega_l^*$ into two segments B_1^l and B_2^l (we will omit the superscript l when the subregion is clear from the context):

$$B_1^l \equiv \partial\Omega_l^* \cap \partial\Omega \quad \text{and} \quad B_2^l \equiv \partial\Omega_l^* \cap \Omega.$$

Corresponding to this, the space-time boundary $B^l \times [0, T]$ of each local subregion can be decomposed into $B_1^l \times [0, T]$ and $B_2^l \times [0, T]$.

2.2. Local space-time grids. On each of the local space-time cylinders $\Omega_l^* \times [0, T]$, we assume that a space-time grid $\Omega_{h_l}^* \times \{0, \tau_l, 2\tau_l, \dots, T - \tau_l, T\}$ is constructed, taking into account the geometry of Ω_l^* and the regularity of the solution on this space-time region (see Figure 1.1). Here h_l denotes the mesh size on Ω_l^* and τ_l denotes the time step with

$$\tau_l = \frac{T}{N_l},$$

for some integer $N_l \geq 1$. Throughout the paper, $x_i^{h_l}$ will denote the i th gridpoint in $\overline{\Omega}_{h_l}^*$.

We will use I^{h_l} to denote the *interior* nodes in the grid $\Omega_{h_l}^*$ and B^{h_l} to denote its *boundary* nodes. Since B^l is decomposed into B_1^l and B_2^l , we denote by $B_1^{h_l}$ and $B_2^{h_l}$, the gridpoints on B_1^l and B_2^l , respectively. We will denote a grid function on $\overline{\Omega}_{h_l}^*$ by w_{h_l} . Corresponding to the partition of gridpoints in $\overline{\Omega}_{h_l}^*$ into I^{h_l} and B^{h_l} , we obtain the block vector

$$w_{h_l} = (w_{h_l, I}, w_{h_l, B})^T.$$

When it is necessary to distinguish the block components of w_{h_l} corresponding to the boundary subgrids $B_1^{h_l}$ and $B_2^{h_l}$, we will use the notation w_{h_l, B_1} and w_{h_l, B_2} , respectively:

$$w_{h_l, B} = (w_{h_l, B_1}, w_{h_l, B_2})^T.$$

We will denote a grid function on $\overline{\Omega}_{h_l}^*$ at time $k\tau_l$ by $w_{h_l}^k$. A space-time grid function on $\overline{\Omega}_{h_l}^* \times \{0, \tau_l, \dots, T\}$ will be denoted by w_{h_l, τ_l} with

$$w_{h_l, \tau_l} = \{w_{h_l}^k\}_{k=0}^{N_l}.$$

A grid function on the entire family of space-time grids will be denoted by $w_{h,\tau}$, where

$$\begin{aligned} w_{h,\tau} &\equiv (w_{h_l,\tau_l})_{l=1}^p \\ &= \left(\{w_{h_l}^k\}_{k=0}^{N_l} \right)_{l=1}^p. \end{aligned}$$

Given a continuous spatial function $w(x)$, we will use $\pi_{h_l}w$ to denote its spatial interpolation onto the gridpoints in $\bar{\Omega}_{h_l}^*$

$$(\pi_{h_l}w)_i \equiv w(x_i^{h_l}), \quad x_i^{h_l} \in \bar{\Omega}_{h_l}^*.$$

Similarly, $\pi_{h_l,I}w$ will denote interpolation of $w(x)$ onto the interior gridpoints in $\Omega_{h_l}^*$. If $w(x,t)$ is a continuous space-time function, we will use π_{h_l,τ_l} to denote the interpolation of $w(.,.)$ onto the l th space-time grid

$$\pi_{h_l,\tau_l}w \equiv \{\pi_{h_l}w(.,k\tau_l)\}_{k=0}^{N_l}.$$

If $w(x,t)$ is a continuous space-time function, we will use $\pi_{h,\tau}$ to denote the interpolation of $w(.,.)$ onto all the space-time grids

$$\pi_{h,\tau}w \equiv (\pi_{h_l,\tau_l}w)_{l=1}^p.$$

2.3. Local subproblems and local discretizations. On each of the space-time subdomains $\Omega_l^* \times [0, T]$, the original parabolic equation (1.1) will be replaced by the following local parabolic initial boundary value problem with suitably chosen boundary conditions $g_l(x,t)$ that couple the adjacent problems:

$$\begin{aligned} u_t + Lu &= f(x,t), & (x,t) \in \Omega_l^* \times [0, T], \\ u &= 0, & (x,t) \in B_1^l \times [0, T], \\ u &= g_l(x,t), & (x,t) \in B_2^l \times [0, T], \\ u(x,0) &= u_0(x), & t = 0, \end{aligned}$$

where L denotes the elliptic operator

$$Lu \equiv -a\Delta u + \vec{b} \cdot \nabla u + c(x)u.$$

Here, the choice of local initial data is $u_0(x)$ restricted to Ω_l^* since the exact solution restricted to $\Omega_l^* \times [0, T]$ would satisfy this initial condition and since $u_0(x)$ is assumed to be known. The boundary data on $B_1^l \times [0, T]$ is zero, since the exact solution satisfies this boundary condition. The boundary data $g_l(x,t)$ will play a crucial role, as it is not known. We will require (see subsection 2.5) that $g_l(x,t)$ equals a suitable interpolation of the solutions from adjacent regions. This will couple the various local problems and require an iterative process to compute $g_l(x,t)$.

Each local parabolic equation will be discretized on the space-time grid $\Omega_{h_l}^* \times \{0, \tau_l, \dots, T\}$ by a finite difference scheme in space and an implicit or explicit θ -scheme in time. The elliptic operator L will be discretized on each spatial grid $\Omega_{h_l}^*$ by a finite difference scheme with coefficient matrix A^{h_l} . If $x_i^{h_l}$ is the i th interior gridpoint in $\Omega_{h_l}^*$, then the discretization of L at this gridpoint will be denoted by

$$Lw(x_i^{h_l}) = \sum_j A_{ij}^{h_l} w(x_j^{h_l}) + \mathcal{C}_{h_l}(w, x_i^{h_l}),$$

where $\mathcal{C}(w, x_i^{h_l})$ is the local truncation error at $x_i^{h_l}$ for an arbitrary smooth function $w(x)$. The matrix A^{h_l} will be rectangular; the first index (i in the preceding

equation) corresponds to interior gridpoints, while the second index (j above) corresponds to interior and/or boundary gridpoints. Corresponding to the partition $w_{h_i} = (w_{h_i,I}, w_{h_i,B})^T$, the rectangular matrix A^{h_i} can be block partitioned

$$A^{h_i} = \begin{bmatrix} A_{II}^{h_i} & A_{IB}^{h_i} \end{bmatrix}, \quad \text{with} \quad A^{h_i} w_{h_i} = A_{II}^{h_i} w_{h_i,I} + A_{IB}^{h_i} w_{h_i,B}.$$

Assumption A1. We will assume the following about the entries of A^{h_i} .

1. $A_{ii}^{h_i} > 0$ for all $i \in I^{h_i}$.
2. $A_{ij}^{h_i} \leq 0$ when $i \neq j$ with $i \in I^{h_i}$ and $j \in I^{h_i} \cup B^{h_i}$.
3. $\sum_j A_{ij}^{h_i} = c_i^{h_i} \geq c_0 > 0$.

Remark 1. Finite difference discretizations satisfying Assumption A1 can be constructed in many ways. If the grid $\Omega_{h_i}^*$ is uniform, then the standard second-order five-point stencil may be applied to approximate $\sum_i \frac{\partial}{\partial x_i} \left(a \frac{\partial w}{\partial x_i} \right)$. If a is not small in relation to $\|\vec{b}\|_{\infty, \Omega_i^*}$, then centered finite differences may be applied to obtain a second order approximation to $\vec{b}(x) \cdot \nabla w$, provided a local cell Peclet restriction (of the form $\|\vec{b}\|_{\infty, \Omega_i^*} h < 2a$) is satisfied. If a is small in relation to $\|\vec{b}\|_{\infty, \Omega_i^*}$ (or zero), then a first order upwind discretization can be applied to approximate $\vec{b}(x) \cdot \nabla w$. The term $c(x)w(x)$ can be approximated by a one-point stencil at each gridpoint. If the grid is nonuniform, finite volume based finite differences may be applied to construct the desired approximations. For instance, if $\Omega \subset R^2$, then a Delaunay triangulation need first to be constructed for the grid and finite volume based finite differences can be applied (see [6]).

Remark 2. If matrix A^{h_i} satisfies Assumption A1, then A^{h_i} will be strictly diagonally dominant and $((A_{II}^{h_i})^{-1})_{ij} \geq 0$ for all i, j . In particular, $A_{II}^{h_i}$ will be an M -matrix (see, for instance [31]).

Semi-discretization of the local parabolic initial boundary value problem on the spatial grid $\Omega_{h_i}^*$ yields

$$\begin{aligned} \frac{dU_{h_i,I}}{dt} + A_{II}^{h_i} U_{h_i,I} + A_{IB}^{h_i} U_{h_i,B} &= f_{h_i,I}(t), \\ U_{h_i,B_1}(t) &= 0, \\ U_{h_i,B_2}(t) &= g_{h_i,B_2}(t), \\ U_{h_i}(0) &= \pi_{h_i} u_0(x), \end{aligned} \tag{2.1}$$

where $f_{h_i,I}(t) \equiv \pi_{h_i,I} f(\cdot, t)$. The boundary conditions $g_{h_i,B_2}(t)$ will be specified in subsections 2.4 and 2.5.

To discretize (2.1) in time, we choose $0 \leq \theta_l \leq 1$, for $l = 1, \dots, p$ independently, and apply a θ -scheme to obtain

$$\begin{aligned}
 (2.2) \quad & \frac{U_{h_l, I}^{k+1} - U_{h_l, I}^k}{\tau_l} + \theta_l A_{II}^{h_l} U_{h_l, I}^{k+1} + \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} \\
 & + \tilde{\theta}_l A_{II}^{h_l} U_{h_l, I}^k + \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k = \theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k, \\
 & U_{h_l, B_1}^{k+1} = 0, \\
 & U_{h_l, B_2}^{k+1} = g_{h_l, B_2}^{k+1}, \\
 & U_{h_l, 0}^0 = \pi_{h_l, I} u_0,
 \end{aligned}$$

for $k = 0, \dots, N_l - 1$, where $\tilde{\theta}_l \equiv 1 - \theta_l$. The discrete boundary conditions $\{g_{h_l, B_2}^k\}$ for $k = 1, \dots, N_l$ and $l = 1, \dots, p$ are crucial for coupling the various local parabolic discretizations and will be described in the next section.

2.4. Intergrid interpolation. The local space-time discretizations (2.2) will be coupled together by requiring that the local boundary data g_{h_l, B_2}^k match a suitable interpolation $\mathcal{I}_{h_l}^k U_{h, \tau}$ of the discrete solution $U_{h, \tau}$ from adjacent grids. The linear map $\mathcal{I}_{h_l}^k$ is described below.

Let $x_i^{h_l}$ (from $\bar{\Omega}_{h_l}^*$) be a gridpoint on B_2^l . At time $k\tau_l$ the boundary data $(g_{h_l, B_2}^k)_i = (U_{h_l}^k)_i \approx u(x_i^{h_l}, k\tau_l)$ will be approximated by *linear combinations* of nodal values of $(U_{h_{\tilde{l}}}^k)_j \approx u(x_j^{h_{\tilde{l}}}, k\tau_{\tilde{l}})$ from adjacent space-time grids $\bigcup_{\tilde{l} \neq l} \bar{\Omega}_{\tilde{l}} \times [0, T]$ with

$$(U_{h_l}^k)_i = (\mathcal{I}_{h_l}^k U_{h, \tau})_i,$$

where the *interpolation map* $\mathcal{I}_{h_l}^k$ is defined in terms of a tensor $\alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i}$:

$$(2.3) \quad (\mathcal{I}_{h_l}^k U_{h, \tau})_i \equiv \sum_{\tilde{l}=1}^p \sum_{\tilde{k}=0}^{N_{\tilde{l}}} \sum_{\tilde{i}} \alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i} (U_{h_{\tilde{l}}}^{\tilde{k}})_{\tilde{i}}.$$

Below we list assumptions about the weights $\alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i}$ used to define $(\mathcal{I}_{h_l}^k U_{h, \tau})_i$.

1. *Assumption A2.* The intergrid interpolation map $\mathcal{I}_{h_l}^k$ must use only values from adjacent grids $\tilde{l} \neq l$. In terms of the coefficients, this places the following requirement on the weights $\alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i}$:

$$\alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i} = 0, \quad \text{when } l = \tilde{l}.$$

2. *Assumption A3.* Given l , the intergrid interpolation map $\mathcal{I}_{h_l}^k$ should involve only nodal values from gridpoints in the *unextended* subregions $\bar{\Omega}_{\tilde{l}} \times [0, T]$ for $\tilde{l} \neq l$, i.e., it should not involve nodal values from gridpoints in the *extended* regions $\bigcup_{\tilde{l} \neq l} (\Omega_{\tilde{l}}^* \setminus \bar{\Omega}_{\tilde{l}}) \times [0, T]$. In terms of the coefficients $\alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i}$ this requirement is

$$\alpha_{\tilde{l}, \tilde{k}, \tilde{i}}^{l, k, i} = 0, \quad \text{when } x_j^{h_{\tilde{l}}} \in \Omega_{\tilde{l}}^* \setminus \bar{\Omega}_{\tilde{l}} \text{ or } \tilde{l} = l.$$

Given B_2^l , let $B_2^{l,*} \subset \Omega$ denote the smallest region such that $B_2^{l,*} \times [0, T]$ contains all the cells and gridpoints from adjacent subregions used in defining the interpolation map $\mathcal{I}_{h_l}^k$ for $k = 0, \dots, N_l$.

3. *Assumption A4.* Let $w(x, t)$ be a smooth space-time function which is zero on $\partial\Omega \times [0, T]$. For each space-time gridpoint $(x_j^{h_l}, k\tau_l)$ on the boundary segment $B_2^l \times [0, T]$ we assume that the interpolation is chosen so that the error satisfies

$$w(x_j^{h_l}, k\tau_l) - (\mathcal{I}_{h_l}^k w_{h,\tau})_j = \mathcal{D}_{h_l}^k(w, x_j^{h_l}),$$

where the local interpolation error can be estimated by Taylor series expansion

$$|\mathcal{D}_{h_l}^k(w, x_j^{h_l})| \leq C \|w\|_{r_{l,1}, r_{l,2}, \infty, B_2^{l,*} \times [0, T]} (h_l^{r_{l,1}} + \tau_l^{r_{l,2}}),$$

for some integers $r_{l,1} \geq 1$ and $r_{l,2} \geq 1$, where C is independent of h_l, τ_l and $\|w\|_{r_{l,1}, r_{l,2}, \infty, B_2^{l,*} \times [0, T]}$ denotes a Sobolev norm of the space-time function $w(\cdot, \cdot)$ on the region $B_2^{l,*} \times [0, T]$.

Definition. Throughout the paper, we will use $\sigma_{h,\tau}$ to denote the maximum norm of the intergrid interpolation map

$$\sigma_{h,\tau} \equiv \max_{l,k,i} \sum_{\tilde{l}=1}^p \sum_{\tilde{k}=0}^{N_{\tilde{l}}} \sum_{\tilde{i}} |\alpha_{\tilde{l},\tilde{k},\tilde{i}}^{l,k,i}|.$$

Example. We include a simple example to illustrate the intergrid interpolation map for a one-dimensional region $\Omega = (0, 4)$ with $T = 1$ and a two-subdomain decomposition with $\Omega_1 = (0, 2)$ and $\Omega_2 = (2, 4)$. Let the overlap parameters be $\beta_l = 1$ for $l = 1, 2$. Then $\Omega_1^* = (0, 3)$ and $\Omega_2^* = (1, 4)$. Let the space-time grids be chosen with $h_1 = 3/10, \tau_1 = 1/10, h_2 = 3/4$ and $\tau_2 = 1/3$. Let the gridpoints be $x_i^{h_1} = ih_1$ for $i = 0, \dots, 10$ in $\Omega_{h_1}^*$ and $x_i^{h_2} = 1 + ih_2$ for $i = 0, \dots, 4$. We will consider a second-order accurate interpolation scheme. In this example, $B_1^{h_1} = 0, B_2^{h_1} = 3, B_1^{h_2} = 4$ and $B_2^{h_2} = 1$. The space-time gridpoints on $B_2^1 \times [0, 1]$ are $\{(3, k/10) : k = 0, \dots, 10\}$. We will describe how the entries of the map $\mathcal{I}_{h_l}^k$ can be constructed for defining the interpolated values at $(x_i^{h_l}, k\tau_l)$ for $k = 1, l = 1$ and $i = 10$. The other grid values can be constructed similarly. The boundary gridpoint $(3, 1/10)$ on boundary $B_2^1 \times [0, 1]$ is enclosed in the cell with vertices $(2.5, 0), (3.25, 0), (2.5, 1/3)$ and $(3.25, 1/3)$ (whose vertices are all gridpoints in the space-time subdomain $\Omega_2^* \times [0, 1]$). Note that these four nodes are contained in $\overline{\Omega_2} \times [0, 1]$. We will define our approximation to $(U_{h_1}^1)_{10}$ by using (second-order) bilinear interpolation:

$$(U_{h_1}^1)_{10} = \frac{23}{90}U_{h_2}(2.5, 0) + \frac{46}{90}U_{h_2}(3.25, 0) + \frac{7}{90}U_{h_2}(2.5, 1/3) + \frac{14}{90}U_{h_2}(3.25, 1/3).$$

The interpolation map can be defined similarly for the other gridpoints on $B_2^{h_1} \times [0, 1]$ so that Assumptions A2 and A3 are satisfied. If all the stencils involve convex combinations (as in the above stencil), then the interpolation map will have maximum norm $\sigma_{h,\tau} = 1$. In this example, the interpolation error is second order in the mesh parameters h_2 and τ_2 of the space-time grid on $\Omega_2^* \times [0, T]$ with coefficients

that depend on second derivatives of the function being interpolated (in the convex hull of the four nodes involved in the stencil):

$$\begin{aligned} & \left| w(x_{10}^{h_1}, \tau_1) - \left(\frac{23}{90}W_{h_2}(2.5, 0) + \frac{46}{90}W_{h_2}(3.25, 0) \right. \right. \\ & \quad \left. \left. + \frac{7}{90}W_{h_2}(2.5, 1/3) + \frac{14}{90}W_{h_2}(3.25, 1/3) \right) \right| \\ & \leq C \|w\|_{2,2,\infty,[2.5,3.25] \times [0,1/3]} (h_2^2 + \tau_2^2). \end{aligned}$$

If the exact solution is less smooth near $B_2^{h_1} \times [0, T]$, then higher accuracy stencils should be applied. However, determining the region where the solution is less smooth would require some estimates (see [24]), for the smoothness of the solution.

2.5. Global discretization. A global discretization of (1.1) can be obtained from the local discretizations (2.2) once the boundary data g_{h_l, B_2}^k are specified using the intergrid interpolation map $\mathcal{I}_{h_l}^k$.

For each $l = 1, \dots, p$, let $0 \leq \theta_l \leq 1$ be the choice of the θ -scheme on $\Omega_l^* \times [0, T]$. Multiplying each equation in (2.2) by π and rearranging terms, the global discretization of (1.1) becomes

$$(2.4) \quad \left\{ \begin{aligned} \left(I + \pi \theta_l A_{II}^{h_l} \right) U_{h_l, I}^{k+1} &= \left(I - \pi \tilde{\theta}_l A_{II}^{h_l} \right) U_{h_l, I}^k \\ &+ \pi \left(-\theta_l A_{IB}^{h_l} U_{h_l, I}^{k+1} + \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, I}^k \right) \\ &+ \pi \left(\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k \right), \\ U_{h_l, B_1}^{k+1} &= 0, \\ U_{h_l, B_2}^{k+1} &= \mathcal{I}_{h_l}^{k+1} \left(U_{h_1, \tau_1}, \dots, U_{h_p, \tau_p} \right)^T, \\ U_{h_l, I}^0 &= \pi_{h_l, I} u_0, \end{aligned} \right.$$

for $l = 1, \dots, p$ and $k = 0, \dots, N_l - 1$.

The above system couples the p local parabolic discretizations through the $U_{h_l, B_2}^{k+1} = \mathcal{I}_{h_l}^{k+1} U_{h, \tau}$ terms. It yields a very large system of linear equations whose parallel iterative solution will be described next. Techniques will also be described for reducing the size of the system and the local memory requirements.

2.6. A parallel Schwarz iterative method. The linear system (2.4) can be solved by a parallel version of the Schwarz iterative method [26, 15, 16, 35, 28, 12] which will (under assumptions stated in Section 4) converge geometrically. On a parallel architecture, each processor can in principle be assigned to a different space-time grid. Some communication between processors will, however, be necessary (to compute the boundary conditions g_{h_l, B_2}^k involving the intergrid interpolation maps $\mathcal{I}_{h_l}^k$). The loads may be well balanced if the subproblems are of comparable size.

System (2.4) will be very large in general, involving all the unknowns on all the space-time grids. However, with some care, the number of unknowns and the memory requirements can be reduced.

1. First, suppose m is a common factor of the number of time steps N_l on each grid for $l = 1, \dots, p$ (i.e., $m = \text{gcd}(N_1, \dots, N_p)$). Then, T can be reduced by a factor m by defining $\tilde{T} \equiv T/m$ and repeatedly applying the global scheme on the time intervals $[0, \tilde{T}]$, $[\tilde{T}, 2\tilde{T}]$, \dots , $[(m-1)\tilde{T}, m\tilde{T}]$.

For example, suppose there are *two* subregions (i.e., $p = 2$) and the time steps are $\tau_1 = T/100$ and $\tau_2 = T/200$. Then choose $\tilde{T} = \tau_1 = 2\tau_2$ and repeatedly apply the global discretization scheme $m = 100$ times on $[0, \tau_1]$, $[\tau_1, 2\tau_1], \dots, [99\tau_1, 100\tau_1]$ to obtain a solution on the time interval $[0, T]$. This procedure will reduce the size of the linear system by a factor 100.

2. Within each space-time grid $\Omega_{h_l}^* \times \{0, \tau_1, \dots, N_l\tau_1\}$ the discrete solution $U_{h_l}^k$ need not be stored for $k = 0, \dots, N_l$. Store the *initial data* $U_{h_l, I}^0$ and the *boundary data* g_{h_l, B_2}^k for $k = 0, \dots, N_l$. Using these, the local discrete solution can be generated by solving the local equations. For the parallel Schwarz algorithm, described next, it would also be necessary to store the nodal values of $(U_{h_l}^k)_i$ that will be used to compute the intergrid interpolation maps.

Once the size of the global system has been reduced by reducing N_l so that $\gcd(N_1, \dots, N_p) = 1$, then system (2.4) can be solved by a parallel Schwarz iterative algorithm. To distinguish the different *iterates* in the Schwarz procedure, we introduce the following notation: $U_{h_l}^{k;(n)}$ will denote the n th Schwarz *iterate* at time $k\tau_l$ on the grid $\Omega_{h_l}^*$.

Parallel Schwarz iteration. Let $\{\{U_{h_l}^{k;(0)}\}_{k=0}^{N_l}\}_{l=1}^p$ be a given starting guess.

1. **For** $n = 0, 1, \dots$ until convergence **do**
2. **For** $l = 1, \dots, p$ in parallel **do**
3. Compute the local boundary conditions for $k = 1, \dots, N_l$:

$$g_{h_l, B_2}^{k;(n+1)} = \mathcal{I}_{h_l}^k \left(U_{h_1, \tau_1}^{(n)}, \dots, U_{h_p, \tau_p}^{(n)} \right)^T.$$

4. **EndFor**
5. **For** $l = 1, \dots, p$ in parallel **do**
6. Let

$$U_{h_l, I}^{0;(n+1)} = \pi_{h_l, I} u_0.$$

7. **For** $k = 0, \dots, N_l - 1$ **solve**

$$(2.5) \quad \begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) U_{h_l, I}^{k+1;(n+1)} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) U_{h_l, I}^{k;(n+1)} \\ &+ \tau_l \left(-\theta_l A_{IB}^{h_l} U_{h_l, I}^{k+1;(n)} + \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, I}^{k;(n+1)} \right) \\ &+ \tau_l \left(\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k \right), \end{aligned}$$

$$U_{h_l, B_1}^{k+1;(n+1)} = 0,$$

$$U_{h_l, B_2}^{k+1;(n+1)} = g_{h_l, B_2}^{k+1;(n+1)}.$$

8. **EndFor**
9. **Endfor**
10. **Endfor**

Under suitable assumptions, the iterates $\{U_{h_l}^{k;(n)}\}$ above can be shown to converge geometrically to the exact solution of (2.4) as $n \rightarrow \infty$ (see Section 4).

3. MAXIMUM NORM PROPERTIES OF THE LOCAL DISCRETIZATIONS

In this section we describe several background results. For *local discretizations* of parabolic equations, we describe maximum norm stability, maximum principles, comparison theorems, barrier functions, and contraction properties. These results will be used in Section 4 to study the maximum norm stability of the global discretization (2.4). The proofs are presented here for the convenience of the reader, though most of the results are scattered in the literature [34, 14, 25, 21, 28, 10].

3.1. Maximum norm stability and a priori estimates for local discretizations. The following preliminary result provides the basis for maximum norm estimates of solutions to θ -schemes.

Lemma 3.1. *Suppose the following hold.*

1. Let $A^{h_l} = \begin{bmatrix} A_{II}^{h_l} & A_{IB}^{h_l} \end{bmatrix}$ satisfy Assumption A1.
2. Let $0 \leq \theta_l \leq 1$ and let $0 < \tau_l$.
3. Let $w_{h_l} = [w_{h_l,I}, w_{h_l,B}]^T$ satisfy

$$\begin{aligned} (I + \tau_l \theta_l A_{II}^{h_l}) w_{h_l,I} + \tau_l \theta_l A_{IB}^{h_l} w_{h_l,B} &= \tilde{f}_{h_l,I}, \\ w_{h_l,B} &= \tilde{g}_{h_l,B}. \end{aligned}$$

Then the following holds:

$$|w_{h_l}|_\infty \leq \max \left\{ \frac{1}{1 + \tau_l \theta_l c_0} |\tilde{f}_{h_l,I}|_\infty, |\tilde{g}_{h_l,B}|_\infty \right\}.$$

Proof. Without loss of generality (if needed multiply w_{h_l} by -1), let $(w_{h_l})_i = |w_{h_l}|_\infty$ (i.e., $(w_{h_l})_i \geq |(w_{h_l})_j|$ for all j). If $i \in B^{h_l}$, then since $w_{h_l,B} = \tilde{g}_{h_l,B}$ the desired result holds. Therefore, in the following, suppose $i \in I^{h_l}$. Then the following holds:

$$\begin{aligned} (\tilde{f}_{h_l,I})_i &= (1 + \tau_l \theta_l A_{ii}^{h_l}) (w_{h_l})_i + \tau_l \theta_l \sum_{j \neq i} A_{ij}^{h_l} (w_{h_l})_j \\ &= (1 + \tau_l \theta_l A_{ii}^{h_l}) |w_{h_l}|_\infty + \tau_l \theta_l \sum_{j \neq i} A_{ij}^{h_l} (w_{h_l})_j \\ &\geq (1 + \tau_l \theta_l A_{ii}^{h_l}) |w_{h_l}|_\infty + \tau_l \theta_l \sum_{j \neq i} A_{ij}^{h_l} |w_{h_l}|_\infty, \quad \text{since } A_{ij}^{h_l} \leq 0 \text{ for } i \neq j \\ &= \left(1 + \tau_l \theta_l A_{ii}^{h_l} + \sum_{j \neq i} \tau_l \theta_l A_{ij}^{h_l} \right) |w_{h_l}|_\infty \\ &= (1 + \tau_l \theta_l c_i^{h_l}) |w_{h_l}|_\infty \\ &\geq (1 + \tau_l \theta_l c_0) |w_{h_l}|_\infty. \end{aligned}$$

Thus, $|w_{h_l}|_\infty \leq |(\tilde{f}_{h_l,I})_i| / (1 + \tau_l \theta_l c_0) \leq |\tilde{f}_{h_l,I}|_\infty / (1 + \tau_l \theta_l c_0)$. □

The preceding result can be applied to the linear system occurring at each time step in the θ -scheme, provided a *stability constraint* is satisfied by τ_l .

Lemma 3.2. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumption A1.
2. Let $0 \leq \theta_l \leq 1$ and define $\tilde{\theta}_l \equiv 1 - \theta_l$.

3. Let $0 < \tau_l$ satisfy the stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0.$$

4. Let $U_{h_l}^{k+1}$ satisfy

$$\begin{aligned} (I + \tau_l \theta_l A_{II}^{h_l}) U_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} &= (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k \\ &\quad + \tau_l (\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k), \\ U_{h_l, B}^{k+1} &= g_{h_l, B}^{k+1}, \end{aligned}$$

where $U_{h_l, I}^k, U_{h_l, B}^k, g_{h_l, B}^{k+1}, f_{h_l, I}^{k+1}$ and $f_{h_l, I}^k$ are given.

Then the following holds:

$$\begin{aligned} |U_{h_l}^{k+1}|_\infty \leq \max \left\{ |g_{h_l, B}^{k+1}|_\infty, \frac{\tau_l \theta_l}{1 + \tau_l \theta_l c_0} |f_{h_l, I}^{k+1}|_\infty \right. \\ \left. + \frac{\tau_l \tilde{\theta}_l}{1 + \tau_l \theta_l c_0} |f_{h_l, I}^k|_\infty + \frac{1 - \tau_l \tilde{\theta}_l c_0}{1 + \tau_l \theta_l c_0} |U_{h_l}^k|_\infty \right\}. \end{aligned}$$

Proof. We apply the preceding lemma with

$$\begin{aligned} \tilde{f}_{h_l, I} &\equiv (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k + \tau_l (\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k), \\ \tilde{g}_{h_l, B} &\equiv g_{h_l, B}^{k+1}. \end{aligned}$$

We need to estimate $|\tilde{f}_{h_l, I}|_\infty$. Since $1 - \tau_l \theta_l A_{ii} \geq 0$ for all i (by the stability criterion for τ_l) and $-\tau_l \theta_l A_{ij}^{h_l} \geq 0$ for $i \neq j$, we obtain

$$\begin{aligned} |(\tilde{f}_{h_l, I})_i| &\leq (1 - \tau_l \theta_l A_{ii}^{h_l}) |(U_{h_l}^k)_i| - \tau_l \theta_l \sum_{j \neq i} A_{ij}^{h_l} |(U_{h_l}^k)_j| \\ &\quad + \tau_l \left(\theta_l |(f_{h_l, I}^{k+1})_i| + \tilde{\theta}_l |(f_{h_l, I}^k)_i| \right) \\ &\leq (1 - \tau_l \theta_l \sum_j A_{ij}^{h_l}) |U_{h_l}^k|_\infty \\ &\quad + \tau_l \left(\theta_l |f_{h_l, I}^{k+1}|_\infty + \tilde{\theta}_l |f_{h_l, I}^k|_\infty \right) \\ &\leq (1 - \tau_l \theta_l c_0) |U_{h_l}^k|_\infty \\ &\quad + \tau_l \left(\theta_l |f_{h_l, I}^{k+1}|_\infty + \tilde{\theta}_l |f_{h_l, I}^k|_\infty \right), \end{aligned}$$

where we used that $-\tau_l \theta_l \sum_j A_{ij}^{h_l} \leq -\tau_l \theta_l c_0$. The desired result now follows by an application of the preceding lemma. \square

By recursively applying the preceding result for $k = 0, 1, \dots, N_l - 1$, one can obtain an *a priori* estimate for the solution to the discretized local parabolic equations.

Lemma 3.3. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumption A1.
2. Let $0 \leq \theta_l \leq 1$ and define $\tilde{\theta}_l \equiv 1 - \theta_l$.

3. Let $0 < \tau_l$ satisfy the stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0.$$

4. Let $U_{h_l, I}^{k+1}$ satisfy

$$\begin{aligned} (I + \tau_l \theta_l A_{II}^{h_l}) U_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} &= (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k \\ &\quad + \tau_l (\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k), \\ U_{h_l, B}^{k+1} &= g_{h_l, B}^{k+1}, \end{aligned}$$

for $k = 0, 1, \dots, N_l - 1$, where $\{g_{h_l, B}^k\}_{k=0}^{N_l}$, $\{f_{h_l, I}^k\}_{k=0}^{N_l}$ and $U_{h_l, I}^0$ are given.

Then the following holds:

$$\begin{aligned} |U_{h_l}^{N_l}|_\infty &\leq \max \left\{ |g_{h_l, B}^{N_l}|_\infty, \frac{1 - \tau_l \tilde{\theta}_l c_0}{1 + \tau_l \theta_l c_0} |g_{h_l, B}^{N_l-1}|_\infty, \dots, \left(\frac{1 - \tau_l \tilde{\theta}_l c_0}{1 + \tau_l \theta_l c_0} \right)^{N_l} |g_{h_l, B}^0|_\infty \right\} \\ &\quad + \tau_l \sum_{k=0}^{N_l-1} \left(\theta_l \left(\frac{1 - \tau_l \tilde{\theta}_l c_0}{1 + \tau_l \theta_l c_0} \right)^{N_l-k} |f_{h_l, I}^{k+1}|_\infty \right. \\ &\quad \quad \left. + \tilde{\theta}_l \left(\frac{1 - \tau_l \tilde{\theta}_l c_0}{1 + \tau_l \theta_l c_0} \right)^{N_l-k} |f_{h_l, I}^k|_\infty \right) \\ &\quad + \left(\frac{1 - \tau_l \tilde{\theta}_l c_0}{1 + \tau_l \theta_l c_0} \right)^{N_l} |U_{h_l, I}^0|_\infty. \end{aligned}$$

Proof. The proof follows by a recursive application of the preceding lemma. \square

Remark 3. By setting $c_0 = 0$ in the preceding result, we may obtain a less sharp result:

$$\begin{aligned} |U_{h_l}^{N_l}|_\infty &\leq \max \left\{ |g_{h_l, B}^{N_l}|_\infty, |g_{h_l, B}^{N_l-1}|_\infty, \dots, |g_{h_l, B}^0|_\infty \right\} \\ &\quad + \tau \theta |f_{h_l, I}^{N_l}|_\infty + \tau \left(\sum_{k=1}^{N_l-1} |f_{h_l, I}^k|_\infty \right) + \tau \tilde{\theta} |f_{h_l, I}^0|_\infty \\ &\quad + |U_{h_l, I}^0|_\infty. \end{aligned}$$

Remark 4. If $f_{h_l, I}^k \equiv 0$ for $k = 0, \dots, N_l$ and $U_{h_l, I}^0 = 0$, then the above result yields

$$|U_{h_l}^{N_l}|_\infty \leq \max \left\{ |g_{h_l, B}^{N_l}|_\infty, |g_{h_l, B}^{N_l-1}|_\infty, \dots, |g_{h_l, B}^0|_\infty \right\},$$

which is a form of the discrete maximum principle for homogeneous solutions of the local discretized parabolic equation.

3.2. Maximum principles and comparison theorems. The following is a maximum principle for the locally discretized parabolic equations.

Lemma 3.4. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumption A1.
2. Let $0 \leq \theta_l \leq 1$ and define $\tilde{\theta}_l \equiv 1 - \theta_l$.

3. Let $0 < \tau_l$ satisfy the stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0.$$

4. Let $U_{h_l, I}^{k+1}$ satisfy

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) U_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k \\ &\quad + \tau_l \left(\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k \right), \end{aligned}$$

$$U_{h_l, B}^{k+1} = g_{h_l, B}^{k+1},$$

for $k = 0, 1, \dots, N_l - 1$, where $\{g_{h_l, B}^k\}_{k=0}^{N_l}$, $\{f_{h_l, I}^k\}_{k=0}^{N_l}$ and U_I^0 are given.

5. Let the initial and boundary data satisfy

$$\begin{aligned} \left(U_{h_l}^0 \right)_i &\geq 0, \quad i \in I^{h_l} \\ \left(f_{h_l, I}^k \right)_i &\geq 0, \quad i \in I^{h_l}, \quad k = 0, \dots, N_l \\ \left(g_{h_l, B}^k \right)_j &\geq 0, \quad j \in B^{h_l}, \quad k = 0, \dots, N_l. \end{aligned}$$

Then, the following holds:

$$\left(U_{h_l, I}^k \right)_i \geq 0, \quad i \in I^{h_l} \text{ and } k = 1, \dots, N_l.$$

Proof. Let $k_0 \geq 1$ denote the smallest integer such that there is an i_0 with $\left(U_{h_l, I}^{k_0} \right)_{i_0} < 0$ (for, if there does not exist such a k_0 and i_0 , then $\left(U_{h_l, I}^k \right)_i \geq 0$ for all k and i , and the desired conclusion holds). Without loss of generality, suppose that

$$\left(U_{h_l, I}^{k_0} \right)_{i_0} = \min_{i \in I^{h_l}} \left(U_{h_l, I}^{k_0} \right)_i < 0.$$

Consider the local discretized equation at time $k_0 \tau_l$

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) U_{h_l, I}^{k_0} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k_0} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) U_{h_l, I}^{k_0-1} - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^{k_0-1} \\ &\quad + \tau_l \left(\theta_l f_{h_l, I}^{k_0} + \tilde{\theta}_l f_{h_l, I}^{k_0-1} \right), \\ U_{h_l, B}^{k_0} &= g_{h_l, B}^{k_0}. \end{aligned}$$

At the i_0 -th gridpoint this becomes

$$\begin{aligned} \left(1 + \tau_l \theta_l A_{i_0 i_0}^{h_l} \right) \left(U_{h_l}^{k_0} \right)_{i_0} + \tau_l \theta_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0} \right)_j \\ = \left(I - \tau_l \tilde{\theta}_l A_{i_0 i_0}^{h_l} \right) \left(U_{h_l}^{k_0-1} \right)_{i_0} - \tau_l \tilde{\theta}_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0-1} \right)_j \\ + \tau_l \left(\theta_l \left(f_{h_l, I}^{k_0} \right)_{i_0} + \tilde{\theta}_l \left(f_{h_l, I}^{k_0-1} \right)_{i_0} \right), \end{aligned}$$

where $\left(U_{h_l, I}^{k_0} \right)_j = \left(g_{h_l, B}^k \right)_j$ for $j \in B^{h_l}$. Using the nonnegativity of $\left(f_{h_l, I}^{k_0} \right)_i$, we obtain

$$\begin{aligned} \left(1 + \tau_l \theta_l A_{i_0 i_0}^{h_l} \right) \left(U_{h_l}^{k_0} \right)_{i_0} + \tau_l \theta_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0} \right)_j &\geq \left(I - \tau_l \tilde{\theta}_l A_{i_0 i_0}^{h_l} \right) \left(U_{h_l}^{k_0-1} \right)_{i_0} \\ &\quad - \tau_l \tilde{\theta}_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0-1} \right)_j. \end{aligned}$$

Since $(U_{h_l}^{k_0-1})_j \geq 0$ for all j , $(1 - \tau_l \tilde{\theta}_l A_{i_0 i_0}^{h_l}) \geq 0$ and $-\tau_l \tilde{\theta}_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \geq 0$, we obtain

$$\begin{aligned} \left(1 + \tau_l \theta_l A_{i_0 i_0}^{h_l}\right) \left(U_{h_l}^{k_0}\right)_{i_0} + \tau_l \theta_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0}\right)_j &\geq \left(1 - \tau_l \tilde{\theta}_l A_{i_0 i_0}^{h_l}\right) \left(U_{h_l}^{k_0-1}\right)_{i_0} \\ &\quad - \tau_l \tilde{\theta}_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0-1}\right)_j \\ &\geq 0. \end{aligned}$$

By assumption $-(U_{h_l}^{k_0})_{i_0} \geq -(U_{h_l}^{k_0})_j$ and $A_{i_0 j}^{h_l} \leq 0$ for $j \neq i_0$. Rearranging terms in the left hand side above, we obtain

$$\begin{aligned} 0 &\leq \left(1 + \tau_l \theta_l A_{i_0 i_0}^{h_l}\right) \left(U_{h_l}^{k_0}\right)_{i_0} + \tau_l \theta_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0}\right)_j \\ &\leq \left(1 + \tau_l \theta_l A_{i_0 i_0}^{h_l}\right) \left(U_{h_l}^{k_0}\right)_{i_0} + \tau_l \theta_l \sum_{j \neq i_0} A_{i_0 j}^{h_l} \left(U_{h_l}^{k_0}\right)_{i_0} \\ &= \left(1 + \tau_l \theta_l \sum_j A_{i_0 j}^{h_l}\right) \left(U_{h_l}^{k_0}\right)_{i_0} \\ &= \left(1 + \tau_l \theta_l c_{i_0}^{h_l}\right) \left(U_{h_l}^{k_0}\right)_{i_0} \\ &< 0, \end{aligned}$$

since $1 + \tau_l \theta_l c_{i_0}^{h_l} \geq 1$ and $(U_{h_l}^{k_0})_{i_0} < 0$ (by assumption). We have thus arrived at a contradiction. Therefore, our assumption was incorrect and we must have

$$\left(U_{h_l}^k\right)_i \geq 0, \quad \text{for } k = 0, \dots, N_l \text{ and } i \in I^{h_l},$$

which is the desired conclusion. □

Remark 5. A similar result can be shown to hold when $c_0 = 0$, provided A^{h_l} is an M -matrix.

As an immediate application of the preceding lemma, we obtain the following *comparison principle*.

Lemma 3.5. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumption A1.
2. Let $0 \leq \theta_l \leq 1$ and define $\tilde{\theta}_l \equiv 1 - \theta_l$.
3. Let $0 < \tau_l$ satisfy the stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0.$$

4. Let $U_{h_l, I}^{k+1}$ satisfy

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l}\right) U_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l}\right) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k \\ &\quad + \tau_l \left(\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k\right), \end{aligned}$$

$$U_{h_l, B}^{k+1} = g_{h_l, B}^{k+1},$$

for $k = 0, 1, \dots, N_l - 1$, where $\{f_{h_l, I}^k\}_{k=0}^{N_l}$, $\{g_{h_l, B}^k\}_{k=0}^{N_l}$ and $U_{h_l, I}^0$ are given.

5. Let \tilde{U}_I^{k+1} satisfy

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) \tilde{U}_{h_l, I}^{k+1} + \tau \theta A_{IB}^{h_l} \tilde{U}_{h_l, B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) \tilde{U}_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} \tilde{U}_{h_l, B}^k \\ &\quad + \tau_l \left(\theta_l \tilde{f}_{h_l, I}^{k+1} + \tilde{\theta}_l \tilde{f}_{h_l, I}^k \right), \\ \tilde{U}_{h_l, B}^{k+1} &= \tilde{g}_{h_l, B}^{k+1}, \end{aligned}$$

for $k = 0, 1, \dots, N_l - 1$, where $\{\tilde{f}_{h_l, I}^k\}_{k=0}^{N_l}$, $\{\tilde{g}_{h_l, B}^k\}_{k=0}^{N_l}$ and $\tilde{U}_{h_l, I}^0$ are given.

6. Let the forcing terms, the initial and boundary data satisfy:

$$\begin{aligned} \left(U_{h_l, I}^0 \right)_i &\geq \left(\tilde{U}_{h_l, I}^0 \right)_i, \quad i \in I^{h_l}, \\ \left(f_{h_l, I}^k \right)_i &\geq \left(\tilde{f}_{h_l, I}^k \right)_i, \quad i \in I^{h_l}, \quad k = 0, \dots, N_l, \\ \left(g_{h_l, B}^k \right)_j &\geq \left(\tilde{g}_{h_l, B}^k \right)_j, \quad j \in B^{h_l}, \quad k = 0, \dots, N_l. \end{aligned}$$

Then, the following will hold:

$$\left(U_{h_l, I}^k \right)_i \geq \left(\tilde{U}_{h_l, I}^k \right)_i \quad \forall i \in I^{h_l} \text{ and } k = 1, \dots, N_l.$$

Proof. Subtract the two sets of discretized parabolic equations to obtain

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) \left(U_{h_l, I}^{k+1} - \tilde{U}_{h_l, I}^{k+1} \right) + \tau_l \theta A_{IB}^{h_l} \left(U_{h_l, B}^{k+1} - \tilde{U}_{h_l, B}^{k+1} \right) \\ = \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) \left(U_{h_l, I}^k - \tilde{U}_{h_l, I}^k \right) - \tau_l \tilde{\theta}_l A_{IB}^{h_l} \left(U_{h_l, B}^k - \tilde{U}_{h_l, B}^k \right) \\ + \tau_l \theta_l \left(f_{h_l, I}^{k+1} - \tilde{f}_{h_l, I}^{k+1} \right) + \tau_l \tilde{\theta}_l \left(f_{h_l, I}^k - \tilde{f}_{h_l, I}^k \right), \\ U_{h_l, B}^{k+1} - \tilde{U}_{h_l, B}^{k+1} = g_{h_l, B}^{k+1} - \tilde{g}_{h_l, B}^{k+1}, \end{aligned}$$

for $k = 0, 1, \dots, N_l - 1$ for the differences $\{U_{h_l, I}^k - \tilde{U}_{h_l, I}^k\}$. Since $U_{h_l, I}^0 - \tilde{U}_{h_l, I}^0$, $g_{h_l, B}^k - \tilde{g}_{h_l, B}^k$, and $f_{h_l, I}^k - \tilde{f}_{h_l, I}^k$ all have nonnegative entries by assumption, we can apply the preceding maximum principle to obtain

$$\left(U_{h_l, I}^k - \tilde{U}_{h_l, I}^k \right)_i \geq 0, \quad \forall i \in I^{h_l} \text{ for } k = 1, \dots, N_l,$$

which is the desired result. \square

Our primary application of the comparison principle for discretized parabolic equations will be to estimate, on each local grid, the modulus of discrete homogeneous solutions \tilde{U}_{h_l, τ_l} with trivial initial data, using a suitable *comparison* or *barrier* grid function w_{h_l, τ_l} . Accordingly, given a discrete homogeneous solution \tilde{U}_{h_l, τ_l} satisfying trivial initial conditions $\tilde{U}_{h_l}^0 = 0$ and nontrivial boundary conditions $\tilde{U}_{h_l, B_2}^k = \tilde{g}_{h_l, B_2}^k$ for $k = 0, \dots, N_l$, we would need to construct (or guarantee the existence of) a suitable grid function w_{h_l, τ_l} satisfying the requirements of the preceding lemma. For convenience, we will seek a comparison (or barrier) grid function w_{h_l, τ_l}^k to be a stationary grid function w_{h_l} , i.e.,

$$w_{h_l}^k = w_{h_l}, \quad k = 0, \dots, N_l.$$

To ensure applicability of the preceding comparison principle, the requirements on w_{h_l} of the preceding lemma will be reduced to

$$\begin{aligned} \left(A_{II}^{h_l} w_{h_l, I} + A_{IB}^{h_l} w_{h_l, B} \right)_i &\geq 0, & i \in I^{h_l}, \\ (w_{h_l, B})_i &\geq \max_{k=0, \dots, N_l} \left| \left(\tilde{g}_{h_l, B}^k \right)_i \right|, & i \in B^{h_l}, \\ (w_{h_l})_i &\geq 0, & i \in I^{h_l}, \end{aligned}$$

since $\tilde{f}_{h_l}^k = 0$ and $\tilde{U}_{h_l}^0 = 0$. We have the following result.

Lemma 3.6. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumption A1.
2. Let $0 \leq \theta_l \leq 1$ and define $\tilde{\theta}_l \equiv 1 - \theta_l$.
3. Let $0 < \tau_l$ satisfy the stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\theta_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0.$$

4. Let $\tilde{U}_{h_l, I}^{k+1}$ solve

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) \tilde{U}_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} \tilde{U}_{h_l, B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) \tilde{U}_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} \tilde{U}_{h_l, B}^k, \\ \tilde{U}_{h_l, B}^{k+1} &= \tilde{g}_{h_l, B}^{k+1}, \end{aligned}$$

for $k = 0, 1, \dots, N_l - 1$ with $\tilde{U}_{h_l, I}^0 = 0$, where $\{\tilde{g}_{h_l, B}^k\}_{k=0}^{N_l}$ are given.

5. Let $w_{h_l} = (w_{h_l, I}, w_{h_l, B})^T$ satisfy

$$\begin{aligned} \left(A_{II}^{h_l} w_{h_l, I} + A_{IB}^{h_l} w_{h_l, B} \right)_i &\geq 0, & i \in I^{h_l}, \\ (w_{h_l, B})_j &\geq \max_{k=0, \dots, N_l} \left| \left(\tilde{g}_{h_l, B}^k \right)_j \right|, & j \in B^{h_l}. \end{aligned}$$

Then, the following holds:

$$(w_{h_l, I})_i \geq \left| \left(\tilde{U}_{h_l, I}^k \right)_i \right|, \quad \forall i \in I^{h_l}, \quad \text{for } k = 0, \dots, N_l.$$

Proof. We will apply the comparison principle (Lemma 3.5) using \tilde{U}_{h_l, τ_l} as defined above and

$$U_{h_l}^k \equiv w_{h_l}, \quad \text{for } k = 0, \dots, N_l.$$

Due to the stationarity of $U_{h_l}^k$, the finite difference approximation of its time derivative is zero and

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) U_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} \\ \geq \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k + \tau_l \tilde{f}_{h_l, I}, \\ \left(U_{h_l, B}^{k+1} \right)_j \geq \left| \left(\tilde{g}_{h_l, B}^{k+1} \right)_j \right|, \quad j \in B^{h_l}, \end{aligned}$$

holds for $k = 0, \dots, N_l - 1$.

In order to apply Lemma 3.5, we also need to verify that

$$\left(U_{h_l, I}^0 \right)_i \geq \left(\tilde{U}_{h_l}^0 \right)_i = 0, \quad \forall i \in I^{h_l},$$

which in our application corresponds to $(w_{h_l, I})_i \geq 0$ for all i . From the definition of w_{h_l} , we obtain

$$w_{h_l, I} = \left(A_{II}^{h_l}\right)^{-1} \left(\tilde{f}_{h_l, I} - A_{IB}^{h_l} w_{h_l, B}\right).$$

Since A^{h_l} satisfies Assumption A1, it is a strictly diagonally dominant (rectangular) M -matrix and satisfies

$$\left(\left(A_{II}^{h_l}\right)^{-1}\right)_{ij} \geq 0, \quad \forall i, j \in I^{h_l}.$$

Since $(A^{h_l})_{ij} \leq 0$ for $i \neq j$, $\tilde{f}_{h_l}^k = 0$ and $(w_{h_l, B})_i \geq 0$ in the above, we conclude that $(w_{h_l, I})_i \geq 0$ for $i \in I^{h_l}$. By the comparison principle

$$(U_{h_l}^k)_i = (w_{h_l})_i \geq \left(\tilde{U}_{h_l}^k\right)_i, \quad \forall i, k.$$

To complete the proof, we observe that all of the above inequalities remain valid if $\tilde{U}_{h_l, I}^k$ is replaced by $-\tilde{U}_{h_l, I}^k$. Applying identical arguments yields

$$(w_{h_l})_i \geq -\left(\tilde{U}_{h_l}^k\right)_i, \quad \forall i, k.$$

Combining these two results yields

$$(w_{h_l})_i \geq \left|\tilde{U}_{h_l}^k\right|_i, \quad \forall i, k,$$

which is the desired result. \square

In the next section, we describe a technique for constructing grid functions w_{h_l} , for *sufficiently small* mesh size h_l (with $h_l \leq h_l^*$) when $c_0 > 0$. Such grid functions (actually, standardized versions of them where the boundary values are suitably scaled) will be referred to as discrete *barrier* or *comparison* functions. Once barrier functions are constructed (or their properties are known), they can be applied to derive a contraction property for homogeneous solutions.

3.3. Existence of continuous and discrete barrier functions. In this section we prove the existence of a discrete barrier grid function w_{h_l} when $c_0 > 0$ on each local grid and describe some of its properties. We state below the precise requirements that a grid function must satisfy in order to be called a barrier function.

Definition. A grid function $w_{h_l} = (w_{h_l, I}, w_{h_l, B})^T$ that satisfies

$$\begin{aligned} \left(A_{II}^{h_l} w_{h_l, I} + A_{IB}^{h_l} w_{h_l, B}\right)_i &\geq 0, \quad i \in I^{h_l}, \\ (3.1) \quad (w_{h_l, B_1})_i &\geq 0, \quad i \in B_1^{h_l}, \\ (w_{h_l, B_2})_i &\geq 1, \quad i \in B_2^{h_l}, \end{aligned}$$

will be referred to as a discrete *barrier* (or *comparison*) grid function.

The existence of discrete barrier grid functions will be proved in two stages. First, results on the existence of a continuous barrier function $w_l(x)$ associated with the continuous analog of (3.1) on Ω_l^* will be described. Second, a grid function $w_{h_l} = \pi_{h_l} w_l$ will be defined by nodal interpolation of $w_l(x)$ onto the grid $\Omega_{h_l}^*$. For *sufficiently small* h_l (i.e., for $h_l \leq h_l^*$ for some $h_l^* > 0$), it will be shown that the resulting grid function w_{h_l} will satisfy the requirements (3.1). We have the following result for continuous barrier functions.

Lemma 3.7. *Suppose the following hold.*

1. Let $0 \leq d_l(x)$ denote the distance from x to the boundary segment $B_2^l = \partial\Omega_l^* \cap \Omega$:

$$d_l(x) \equiv \text{dist}(x, B_2^l).$$

2. Let $w_l(x)$ be defined as

$$w_l(x) \equiv e^{-\alpha d_l(x)}, \quad x \in \overline{\Omega}_l^*,$$

for $\alpha > 0$.

Then there exists a choice $\alpha = \alpha_l > 0$ so that $w_l(x)$ defined above satisfies

$$Lw_l \geq \frac{c_0}{2}, \quad x \in \Omega_l^*,$$

$$w_l(x) \geq 0, \quad x \in B_1^l,$$

$$w_l(x) = 1, \quad x \in B_2^l.$$

Proof. We follow the construction in Lions [27] (see also [28]). Direct computation of $Le^{-\alpha d_l(x)}$ yields

$$\begin{aligned} Le^{-\alpha d_l(x)} &= e^{-\alpha d_l(x)} \left(a\alpha \Delta d_l(x) - a\alpha^2 |\nabla d_l(x)|^2 - \alpha \vec{b}(x) \cdot \nabla d_l(x) + c(x) \right) \\ &\geq e^{-\alpha d_l(x)} \left(a\alpha \Delta d_l(x) - a\alpha^2 |\nabla d_l(x)|^2 - \alpha \vec{b}(x) \cdot \nabla d_l(x) + c_0 \right). \end{aligned}$$

If we choose $\alpha = \alpha_l > 0$

$$\alpha_l \equiv \min \left\{ 1, \frac{c_0}{2 \left(\|\vec{b}\|_{\infty, \overline{\Omega}_l^*} \|\nabla d_l\|_{\infty, \overline{\Omega}_l^*} + a \|\nabla d_l\|_{\infty, \overline{\Omega}_l^*}^2 + a \|\Delta d_l\|_{\infty, \overline{\Omega}_l^*} \right)} \right\},$$

then

$$\frac{c_0}{2} + a\alpha \Delta d_l(x) - a\alpha^2 |\nabla d_l(x)|^2 - \alpha \vec{b}(x) \cdot \nabla d_l(x) \geq 0,$$

and we obtain $Le^{-\alpha_l d_l(x)} \geq c_0/2 > 0$. Since $d_l(x) = 0$ on B_2^l , it follows that $e^{-\alpha_l d_l(x)} = 1$ on B_2^l . Since the exponential $e^{-\alpha_l d_l(x)}$ is always nonnegative, it follows that $0 \leq e^{-\alpha_l d_l(x)}$ on B_1^l . Additionally, since $0 \leq d_l(x)$ for all x , it follows that $0 \leq e^{-\alpha_l d_l(x)} \leq 1$. \square

Remark 6. In the above construction, we tacitly assumed smoothness of the distance function $d_l(x)$. Unfortunately, this may not be the case in general, even if $B_2^l = (\partial\Omega_l^* \cap \Omega)$ is smooth. However, given any $0 < \epsilon_l \ll \beta_l$, for our applications we may replace $d_l(x)$ by any smooth function $0 \leq \tilde{d}_{l,\epsilon_l}(x)$ satisfying

$$\begin{aligned} \tilde{d}_{l,\epsilon_l}(x) &\geq 0, & \forall x, \\ \tilde{d}_{l,\epsilon_l}(x) &= 0, & x \in B_2^l, \\ \tilde{d}_{l,\epsilon_l}(x) &\leq d_l(x) + \epsilon_l, & \forall x, \\ \tilde{d}_{l,\epsilon_l}(x) &\geq d_l(x) - \epsilon_l, & \forall x. \end{aligned}$$

Such a ‘‘pseudo-distance’’ function $\tilde{d}_{l,\epsilon_l}(x)$ can be constructed as follows. For any $\gamma > 0$, let Ω_l^γ denote

$$\Omega_l^\gamma \equiv \{x : \text{dist}(x, \Omega_l) < \gamma\}.$$

Then $\Omega_l^* = \Omega_l^{\beta_l}$.

1. Let $S_{l;\epsilon_l}$ denote a region with *smooth* boundaries satisfying

$$\left(\Omega_l^{\beta_l + \frac{\epsilon_l}{2}} \setminus \Omega_l^{\beta_l - \frac{\epsilon_l}{2}}\right) \subset S_{l,\epsilon_l} \subset \left(\Omega_l^{\beta_l + \epsilon_l} \setminus \Omega_l^{\beta_l - \epsilon_l}\right).$$

Then $B_2^l \subset S_{l,\epsilon_l}$.

2. Given S_{l,ϵ_l} , let $d_{l;\epsilon_l}(x)$ denote

$$d_{l;\epsilon_l}(x) \equiv \text{dist}(x, S_{l,\epsilon_l}).$$

Then $d_{l;\epsilon_l}(x)$ will have the following properties:

$$\begin{aligned} d_{l;\epsilon_l}(x) &\geq 0, & \forall x, \\ d_{l;\epsilon_l}(x) &= 0, & x \in S_{l,\epsilon_l}, \\ d_{l;\epsilon_l}(x) &\leq d_l(x) + \frac{\epsilon_l}{2}, & \forall x, \\ d_{l;\epsilon_l}(x) &\geq d_l(x) - \frac{\epsilon_l}{2}, & \forall x. \end{aligned}$$

Unfortunately, $d_{l;\epsilon_l}(x)$ will not be smooth in the regions where the level sets of S_{l,ϵ_l} intersect.

3. Let $0 \leq \psi_{\epsilon_l}(x)$ denote a smooth probability density function having compact support of diameter $\epsilon_l/4$ centered at the origin. Define $\tilde{d}_l(x)$ as the convolution (mollification) of $d_{l;\epsilon_l}(x)$ with $\psi_{\epsilon_l}(x)$ as

$$\tilde{d}_l(x) \equiv \int_y \psi_{\epsilon_l}(y) d_{l;\epsilon_l}(x - y) dy.$$

By construction $\tilde{d}_l(x)$ will be smooth. Due to the nonnegativity and compact support of $\psi_{\epsilon_l}(x)$ of diameter $\epsilon_l/4$, it will further satisfy

$$\begin{aligned} \tilde{d}_l(x) &\geq 0, & \forall x, \\ \tilde{d}_l(x) &= 0, & x \in B_2^l, \\ \tilde{d}_l(x) &\leq d_l(x) + \epsilon_l, & \forall x, \\ \tilde{d}_l(x) &\geq d_l(x) - \epsilon_l, & \forall x. \end{aligned}$$

Thus, given a suitable small but fixed choice of ϵ_l , $e^{-\alpha \tilde{d}_l(x)} \leq e^{-\alpha(d_l(x) + \epsilon_l)}$ will satisfy the requirements of a barrier function for the value of α_l given in the preceding lemma (with $\tilde{d}_l(x)$ replacing $d_l(x)$). For convenience, however, we will henceforth assume that $d_l(x)$ is smooth.

Given the continuous barrier function $w_l(x)$, we will interpolate it onto the grid $\Omega_{h_l}^*$ to construct a discrete barrier function. To ensure that the resulting grid function satisfies (3.1), we will require that the discretization A^{h_l} be at least first order accurate and that h_l be sufficiently small.

Assumption A5. Let $x_i^{h_l}$ denote an interior gridpoint in $\Omega_{h_l}^*$. Then for any sufficiently smooth test function $v(x)$ we assume that

$$(Lv)(x_i^{h_l}) = (A^{h_l} \pi_{h_l} v)_i + \mathcal{C}_{h_l}(v, x_i^{h_l}) h_l^{s_l}$$

holds, where the coefficient $\mathcal{C}_{h_l}(v, x_i^{h_l})$ involves higher order derivatives of $v(x)$ in the convex hull of the gridpoints of the local stencil

$$(3.2) \quad |\mathcal{C}_{h_l}(v, x_i^{h_l})| \leq C \|v\|_{s_l+2, \infty, \Omega_l^*},$$

where C is a positive constant independent of h_l and $v(\cdot)$ and $1 \leq s_l$ is an integer.

Lemma 3.8. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumptions A1 and A5.
2. Let $w_l(x) = e^{-\alpha_l d_l(x)}$ be a smooth continuous barrier function satisfying

$$\begin{aligned} Lw_l &\geq \frac{c_0}{2}, & x \in \Omega_l^*, \\ w_l(x) &\geq 0, & x \in B_1^l, \\ w_l(x) &= 1, & x \in B_2^l. \end{aligned}$$

3. Define $w_{h_l} \equiv \pi_{h_l} w_l$ where

$$(w_{h_l})_i = w_l(x_i^{h_l}), \quad \forall x_i^{h_l} \in \overline{\Omega}_{h_l}^*.$$

4. Let h_l satisfy

$$h_l \leq h_l^* \equiv \left(\frac{c_0}{2C \|w_l\|_{s_l+2, \infty, \Omega_l^*}} \right)^{1/r_l},$$

where C is defined in (3.2).

Then, the following will hold:

$$\begin{aligned} A_{II}^{h_l} w_{h_l, I} + A^{h_l} w_{h_l, B} &\geq 0, & \text{componentwise,} \\ w_{h_l, B_2} &\geq 1, & \text{componentwise,} \\ w_{h_l, B_1} &\geq 0, & \text{componentwise,} \end{aligned}$$

i.e., w_{h_l} is a discrete barrier function.

Proof. By Assumption A5 on the local consistency of the finite difference discretization matrix A^{h_l} , we obtain

$$\begin{aligned} (A^{h_l} w_{h_l})_i &= (Lw_l)(x_i^{h_l}) + \mathcal{C}_{h_l}(w_l, x_i^{h_l}) h_l^{s_l}, \\ &\geq \frac{c_0}{2} + \mathcal{C}_{h_l}(w_l, x_i^{h_l}) h_l^{s_l}, & \text{since } Lw_l \geq \frac{c_0}{2} \\ &\geq \frac{c_0}{2} - C \|w_l\|_{s_l+2, \infty, \Omega_l^*} h_l^{s_l}, & \text{using A5} \\ &\geq 0, & \text{if } h_l \leq h_l^*, \end{aligned}$$

where

$$h_l^* = \left(\frac{c_0}{2C \|w_l\|_{s_l+2, \infty, \Omega_l^*}} \right)^{1/s_l}.$$

The desired result now holds due to the properties of $w_l(x)$ and since $w_{h_l} = \pi_{h_l} w_l$. \square

In the next section, we use barrier grid functions to prove a contraction property of homogeneous solutions to discretized parabolic equations.

3.4. Contraction property of homogeneous solutions. Suppose $\{W_{h_l}^k\}_{k=0}^{N_l}$ denotes a homogeneous solution of the discretized parabolic equation on the l th local grid with trivial initial conditions $W_{h_l, I}^0 = 0$ and nontrivial boundary data $W_{h_l, B_2}^k = g_{h_l, B_2}^k$ on $B_2^{h_l}$. By the discrete maximum principle for parabolic equations, we obtain the bound

$$\left| (W_{h_l}^k)_j \right| \leq \max_{k=0, \dots, N_l} \max_{\{i \in B_2^{h_l}\}} \left| (g_{h_l, B_2}^k)_i \right|,$$

at any interior gridpoint $x_j^{h_l}$ at time $k\tau_l$ for $k = 0, \dots, N_l$. However, when $c_0 > 0$ (as we have assumed), a stronger property will hold in the *interior* region $\bar{\Omega}_l \times [0, T]$

$$\max_{j \in \bar{\Omega}_{h_l}} \left| (W_{h_l}^k)_j \right| \leq \rho_{h_l} \max_{k=0, \dots, N_l} \max_{i \in B_2^{h_l}} \left| (g_{h_l, B_2}^k)_i \right|,$$

for some $\rho_{h_l} < 1$. This will be referred to as the local *contraction property* and will be essential in establishing the stability of the global discretization.

We define now the normalized contraction factor $0 \leq \rho_{h_l} \leq 1$ from a domain Ω_l^* to a subregion Ω_l for discrete homogeneous solutions of the discretized parabolic equation on the l th grid.

Definition. Let $\{W_{h_l}^k\}_{k=0}^{N_l}$ denote a homogeneous solution of the following discretized parabolic equation with trivial initial conditions:

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) W_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} W_{h_l, B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) W_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} W_{h_l, B}^k, \\ W_{h_l, B_1}^{k+1} &= 0, \\ W_{h_l, B_2}^{k+1} &= g_{h_l, B_2}^k, \\ W_{h_l, I}^0 &= 0, \end{aligned}$$

where $(g_{h_l, B_2}^k)_i = 1$ for $i \in B_2^{h_l}$. We define the normalized contraction factor ρ_{h_l} (with $0 \leq \rho_{h_l} \leq 1$) as

$$(3.3) \quad \rho_{h_l} \equiv \max_{k=0, \dots, N_l} \max_{i \in \bar{\Omega}_{h_l}} \left| (W_{h_l, I}^k)_i \right|.$$

Our first result in this section provides an upper bound for ρ_{h_l} , in terms of the contraction factor for the continuous barrier function $e^{-\alpha_l \bar{d}_l(x)}$ from the preceding section.

Lemma 3.9. *Suppose the following hold.*

1. Let $0 < \beta_l$ denote the overlap parameter from subdomain Ω_l to Ω_l^* .
2. Let matrix A^{h_l} satisfy Assumptions A1 and A5, and let τ_l satisfy the stability criterion.
3. Let $c_0 > 0$.
4. Let $0 < \alpha_l$ be chosen so that the grid function $w_{h_l} = \pi_{h_l} e^{-\alpha_l \bar{d}_l(x)}$ is a discrete barrier grid function for $h_l \leq h_l^*$.
5. Let ρ_{h_l} denote the normalized contraction factor on the l th grid with W_{h_l} as employed in (3.3).

Then the following holds:

$$\begin{aligned} \rho_{h_l} &\leq \max_{i \in \bar{\Omega}_{h_l}} (w_{h_l})_i \\ &\leq \max_{i \in \bar{\Omega}_{h_l}} e^{-\alpha_l \tilde{d}_l(x_i^{h_l})}. \end{aligned}$$

Consequently

$$\begin{aligned} \rho_{h_l} &\leq e^{-\alpha_l \beta_l}, && \text{if } \tilde{d}_l(x) = d_l(x) \\ &\leq e^{-\alpha_l(\beta_l - \epsilon_l)}, && \text{if } d_l(x) - \epsilon \leq \tilde{d}_l(x) \leq d_l(x), \end{aligned}$$

for some $\epsilon_l > 0$. In either case, $\rho_{h_l} < 1$ if $\beta_l \geq \epsilon_l$.

Proof. By assumption on the mesh size h_l , the preceding lemma yields that $w_{h_l} \equiv \pi_{h_l} e^{-\alpha_l \tilde{d}_l(x)}$ is a discrete barrier grid function. By applying comparison principle 3.5 from subsection 3.2, with $U_{h_l}^k = w_{h_l}$ and $\tilde{U}_{h_l}^k = W_{h_l}^k$ above, we obtain

$$\begin{aligned} \left| (W_{h_l}^k)_i \right| &\leq (w_{h_l})_i && \forall k \\ &\leq e^{-\alpha_l \tilde{d}_l(x_i^{h_l})} && \forall k \\ &\leq e^{-\alpha_l(\beta_l - \epsilon_l)} && \forall k, \end{aligned}$$

which is the desired result. □

The above estimates for the contraction factor ρ_{h_l} are qualitative, and involve an unknown constant α_l . We indicate below how more quantitative theoretical bounds can be obtained for ρ_{h_l} on uniform grids (see [21]).

Example. Consider a parabolic equation $u_t + Lu = f$ in one space dimension (i.e., $\Omega \subset R$) where $Lu = -u'' + bu' + cu$ and where $0 < b$ and $0 < c$ are constants. For convenience, consider a subdomain $\Omega_l^* = (0, 1)$ with $\Omega_l = (\tilde{a}, \tilde{b})$ where $0 < \tilde{a} < \tilde{b} < 1$. Suppose that a uniform grid is constructed on Ω_l^* with mesh size $h_l = 1/M_l$ and gridpoints $x_i^{h_l} = ih_l$ for $0, \dots, M_l$. Discretize $-u''$ by three point finite differences, bu' by upwind finite differences and cu by a one point approximation on the above uniform grid and suppose 0 and 1 are interior points in Ω . To estimate the discrete contraction factor ρ_{h_l} , we solve the difference equations

$$\begin{cases} -(bh + 1)u_{i-1} + (2 + bh + ch^2)u_i - 1u_{i+1} = 0, & \text{for } i = 1, \dots, M_l - 1, \\ u_0 = 1, \\ u_{M_l} = 1, \end{cases}$$

whose general solution has the form

$$u_i = c_1 \sigma_1^i + c_2 \sigma_2^i,$$

where σ_1 and σ_2 are roots of the quadratic

$$\sigma^2 - (2 + bh + ch^2)\sigma + (bh + 1) = 0.$$

In terms of b , c and h_l the roots are

$$\begin{aligned}\sigma_1 &= 1 + \frac{bh}{2} + \frac{ch^2}{2} + h \left(\frac{4c + b^2 + 2bhc + c^2h^2}{4} \right)^{1/2}, \\ \sigma_2 &= 1 + \frac{bh}{2} + \frac{ch^2}{2} - h \left(\frac{4c + b^2 + 2bhc + c^2h^2}{4} \right)^{1/2}.\end{aligned}$$

The constants c_1 and c_2 can be computed by enforcing the boundary conditions

$$c_1 + c_2 = 1,$$

$$\sigma_1^{M_l} c_1 + \sigma_2^{M_l} c_2 = 1,$$

which yields

$$\begin{aligned}c_1 &= \frac{\sigma_2^{M_l} - 1}{\sigma_2^{M_l} - \sigma_1^{M_l}}, \\ c_2 &= \frac{1 - \sigma_1^{M_l}}{\sigma_2^{M_l} - \sigma_1^{M_l}}.\end{aligned}$$

The contraction factor ρ_{h_l} can be estimated as

$$\rho_{h_l} \equiv \max_{\{i: x_i^{h_l} \in (\tilde{a}, \tilde{b})\}} \left| \frac{(\sigma_2^{M_l} - 1)\sigma_1^i + (1 - \sigma_1^{M_l})\sigma_2^i}{\sigma_2^{M_l} - \sigma_1^{M_l}} \right|.$$

Given b , c , h_l , \tilde{a} and \tilde{b} , these can be determined quantitatively.

Our main result in this section relates the discrete contraction factor ρ_{h_l} , which was defined for the normalized Dirichlet boundary conditions, to the case of general Dirichlet boundary conditions.

Lemma 3.10. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumptions A1 and A5.
2. Let $0 \leq \theta_l \leq 1$ and define $\tilde{\theta}_l \equiv 1 - \theta_l$.
3. Let $0 < \tau_l$ satisfy the stability condition

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0.$$

4. Let $\tilde{V}_{h_l}^k$ satisfy

$$\begin{aligned}(I + \tau_l \theta_l A_{II}^{h_l}) \tilde{V}_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} \tilde{V}_{h_l, B}^{k+1} &= (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) \tilde{V}_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} \tilde{V}_{h_l, B}^k, \\ \tilde{V}_{h_l, B_1}^{k+1} &= 0, \\ \tilde{V}_{h_l, B_2}^{k+1} &= \tilde{g}_{h_l, B_2}^{k+1}, \\ \tilde{V}_{h_l, I}^0 &= 0,\end{aligned}$$

for $k = 0, \dots, N_l - 1$.

5. Let ρ_{h_l} denote the normalized contraction factor defined by (3.3) in subsection 3.4 with associated grid function $\{W_{h_l}^k\}_{k=0}^{N_l}$.

Then the following holds:

$$\max_{\{i: x_i^{h_l} \in \bar{\Omega}_{h_l}, k=0, \dots, N_l\}} \left| \left(\tilde{V}_{h_l}^k \right)_i \right| \leq \rho_{h_l} \max_{i \in \bar{\Omega}_{h_l}^*} \max_{k=0, \dots, N_l} \left| \left(\tilde{V}_{h_l}^k \right)_i \right|.$$

Proof. Let $\|\tilde{V}_{B_2^{h_l}}\|$ denote the number

$$\|\tilde{V}_{B_2^{h_l}}\| \equiv \max_{i \in B_2^{h_l}} \max_{k=0, \dots, N_l} \left| \left(\tilde{V}_{h_l}^k \right)_i \right|.$$

Define a grid function $\tilde{W}_{h_l}^k$ by scaling $\tilde{V}_{h_l}^k$ so that its boundary values have maximum modulus *one*:

$$\left(\tilde{W}_{h_l}^k \right)_i \equiv \frac{\left(\tilde{V}_{h_l}^k \right)_i}{\|\tilde{V}_{B_2^{h_l}}\|}, \quad \forall i, k.$$

Apply comparison principle 3.5 from subsection 3.2, employing $U_{h_l}^k = W_{h_l}^k$, where $W_{h_l}^k$ is the grid function associated with the normalized contraction factor ρ_{h_l} defined in (3.3) of subsection 3.4, and employing $\tilde{U}_{h_l}^k = \pm \tilde{W}_{h_l}^k$, where $\tilde{W}_{h_l}^k$ is defined above. The desired result follows immediately from the normalized contraction property. \square

Remark 7. Due to the stationarity of the discrete barrier function w_{h_l} , the upper bound for the contraction factor ρ_{h_l} will be independent of τ_l .

4. MAXIMUM NORM STABILITY AND ACCURACY OF THE GLOBAL DISCRETIZATION

In this section, we prove that the global discretization (2.4) is stable in the maximum norm and analyze its accuracy. We also show that the parallel Schwarz algorithm is geometrically convergent. The proofs are motivated by [32, 10] and employ Picard's contraction mapping theorem. In the first section, we describe the contraction mapping theorem and existence and uniqueness results for (2.4). We also discuss the geometric convergence of the parallel Schwarz iterates. In the second section we describe the stability of the global discretization. In the third section, we apply the stability theorem to estimate the accuracy of the global non-matching grid discretization in terms of the local discretization and interpolation errors.

4.1. Contraction mapping theorem. The existence and uniqueness of solutions to the global discretized system (2.4) will be proved by applying Picard's contraction mapping theorem [3], which we summarize below for convenience. In addition, the geometric convergence of the parallel Schwarz iterates will follow from properties of the contraction mapping.

Theorem 4.1. *Suppose the following hold.*

1. Let \mathcal{H} be a complete metric space with metric $d(\cdot, \cdot)$.
2. Let $T : \mathcal{H} \rightarrow \mathcal{H}$ be a contractive mapping, i.e., for any $X, Y \in \mathcal{H}$ we have

$$d(TX, TY) \leq \delta d(X, Y),$$

where $\delta < 1$.

Then the following hold.

1. There exists a unique fixed point $U \in \mathcal{H}$ of \mathcal{T} satisfying

$$U = \mathcal{T}U.$$

2. Let $U^{(0)}$ be any element of \mathcal{H} . Then $U^{(n)} \equiv \mathcal{T}^n U^{(0)} \rightarrow U$ geometrically with

$$\begin{aligned} d(U^{(n+1)}, U) &\leq \delta d(U^{(n)}, U) \\ &\leq \delta^n d(U^{(0)}, U). \end{aligned}$$

3. For any $U^{(0)} \in \mathcal{H}$ we have

$$d(U^{(0)}, U) \leq \frac{1}{1-\delta} d(\mathcal{T}U^{(0)}, U^{(0)}).$$

Proof. See [3]. □

In our applications, we will choose the metric space \mathcal{H} and mapping \mathcal{T} so that global discretization (2.4) is a fixed point equation for \mathcal{T} . Furthermore, the parallel Schwarz iterates $\{U^{(n)}\}$, given a starting guess $U^{(0)}$, will correspond to $U^{(n+1)} = \mathcal{T}U^{(n)}$. For convenience, we will consider a system of equations more general than (2.4). Let $\tilde{f}_{h_l, I}^k$ and \tilde{g}_{h_l, B_2}^k be given forcing terms for $k = 0, \dots, N_l$ and $l = 1, \dots, p$. Let $\tilde{u}_{h_l, I}^0$ be given initial data for $l = 1, \dots, p$. We consider the following general system of equations for unknowns $U_{h, \tau} = \{\{U_{h_l}^k\}_{k=0}^{N_l}\}_{l=1}^p$:

$$\begin{aligned} (4.1) \quad (I + \tau_l \theta_l A_{II}^{h_l}) U_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} U_{h_l, B}^{k+1} &= (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) U_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} U_{h_l, B}^k \\ &\quad + \tau_l (\theta_l \tilde{f}_{h_l, I}^{k+1} + \tilde{\theta}_l \tilde{f}_{h_l, I}^k), \\ U_{h_l, B_1}^{k+1} &= 0, \\ U_{h_l, B_2}^{k+1} &= \mathcal{I}_{h_l}^{k+1} U_{h, \tau} + \tilde{g}_{h_l, B_2}^{k+1}, \\ U_{h_l, I}^0 &= \tilde{u}_{h_l, I}^0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$.

Our choice for \mathcal{H} will be based on the linear system (4.1). Given $\tilde{f}_{h_l, I}^k, \tilde{g}_{h_l, B_2}^k$ for $k = 0, \dots, N_l$ and $l = 1, \dots, p$ and $\tilde{u}_{h_l, I}^0$, we define

$$\begin{aligned} \mathcal{H} &\equiv \{X_{h, \tau} : X_{h_l}^k \text{ satisfy following constraints}\} \\ (I + \tau_l \theta_l A_{II}^{h_l}) X_{h_l, I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} X_{h_l, B}^{k+1} &= (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) X_{h_l, I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} X_{h_l, B}^k \\ &\quad + \tau_l (\theta_l \tilde{f}_{h_l, I}^{k+1} + \tilde{\theta}_l \tilde{f}_{h_l, I}^k), \\ X_{h_l, B_1}^{k+1} &= 0, \\ X_{h_l, I}^0 &= \tilde{u}_{h_l, I}^0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$. \mathcal{H} is not a vector space due to the linear *inhomogeneous* constraints. However, \mathcal{H} is closed in the vector space of all grid functions endowed with the maximum norm, and consequently, \mathcal{H} will be a complete metric space if the metric $d(\cdot, \cdot)$ defined on \mathcal{H} is inherited from the standard maximum norm. Given $X_{h, \tau}, Y_{h, \tau} \in \mathcal{H}$, we define

$$d(X_{h, \tau}, Y_{h, \tau}) \equiv \| \|X_{h, \tau} - Y_{h, \tau}\| \|,$$

where $\|\cdot\|$ denotes the maximum norm on the vector space of all space-time grid functions

$$\|X_{h,\tau} - Y_{h,\tau}\| \equiv \max_{l=1,\dots,p} \max_{k=0,\dots,N_l} \max_{i \in \bar{\Omega}_{h_l}^*} |(X_{h_l}^k - Y_{h_l}^k)_i|.$$

For the above \mathcal{H} , we define a mapping $\mathcal{T} : \mathcal{H} \rightarrow \mathcal{H}$ so that linear system (4.1) is a fixed point equation of \mathcal{T} on \mathcal{H} . Given $X_{h,\tau} \in \mathcal{H}$ define $\tilde{X}_{h,\tau} = \mathcal{T}X_{h,\tau}$ as

$$(4.2) \quad \begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) \tilde{X}_{h_l,I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} \tilde{X}_{h_l,B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) \tilde{X}_{h_l,I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} \tilde{X}_{h_l,B}^k \\ &\quad + \tau_l \left(\theta_l \tilde{f}_{h_l,I}^{k+1} + \tilde{\theta}_l \tilde{f}_{h_l,I}^k \right), \\ \tilde{X}_{h_l,B_1}^{k+1} &= 0, \\ \tilde{X}_{h_l,B_2}^{k+1} &= \mathcal{I}_{h_l}^{k+1} X_{h,\tau} + \tilde{g}_{h_l,B_2}^{k+1}, \\ \tilde{X}_{h_l,I}^0 &= \tilde{u}_{h_l,I}^0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$. It immediately follows that system (4.1) is a fixed point equation of \mathcal{T} . Additionally, the parallel Schwarz iterates from Section 2 can be described in terms of the mapping \mathcal{T} : Given a starting guess $U_{h,\tau}^{(0)}$ to (4.1), the subsequent parallel Schwarz iterates $\{U_{h,\tau}^{(n)}\}$ are

$$U_{h,\tau}^{(n)} \equiv \mathcal{T}U_{h,\tau}^{(n-1)}, \quad n = 1, 2, \dots$$

The existence and uniqueness of solutions to system (4.1) and (2.4) will be guaranteed by Picard’s contraction mapping theorem, provided \mathcal{T} is a *contraction* mapping in the metric space \mathcal{H} . This contraction property of \mathcal{T} is pivotal to the stability analysis in this paper and is proved next.

Theorem 4.2. *Suppose the following hold.*

1. Let matrix A^{h_l} satisfy Assumptions A1 and A5 for $l = 1, \dots, p$.
2. Let $0 < \tau_l$ satisfy the local stability criterion

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0,$$

for $l = 1, \dots, p$.

3. Let the intergrid interpolation maps $\{\mathcal{I}_{h_l}^k\}_{k,l}$ satisfy Assumptions A2 and A3.
4. Let the local overlap parameters β_l be chosen large enough, and the local mesh size $h_l \leq h_l^*$ small enough so that the contraction factor ρ_{h_l} satisfies

$$\delta_h \equiv \sigma_{h,\tau} \max_l \rho_{h_l} \leq \delta < 1,$$

where $\sigma_{h,\tau}$ denotes the maximum norm of the map $\{\mathcal{I}_{h_l}^k\}_{k,l}$.

Then the following hold.

1. The mapping \mathcal{T} will be a contraction mapping on \mathcal{H} satisfying

$$d(\mathcal{T}X_{h,\tau}, \mathcal{T}Y_{h,\tau}) \leq \delta d(X_{h,\tau}, Y_{h,\tau}), \quad \forall X_{h,\tau}, Y_{h,\tau} \in \mathcal{H}.$$

2. System (4.1) will be uniquely solvable with a solution $U_{h,\tau}$.

3. Given any starting guess $U_{h,\tau}^{(0)} \in \mathcal{H}$, the iterates $U_{h,\tau}^{(n)} = \mathcal{T}^n U_{h,\tau}^{(0)}$ (which correspond to the parallel Schwarz iterates) converge geometrically to the unique fixed point $U_{h,\tau}$:

$$\|\mathcal{T}^n U_{h,\tau}^{(0)} - U_{h,\tau}\| \leq \delta^n \|U_{h,\tau}^{(0)} - U_{h,\tau}\|.$$

Proof. The proof is similar to [10] and relies on the maximum principle and the contraction property. Given $X_{h,\tau}, Y_{h,\tau} \in \mathcal{H}$, we need to estimate $d(\mathcal{T}X_{h,\tau}, \mathcal{T}Y_{h,\tau})$ in terms of $d(X_{h,\tau}, Y_{h,\tau})$. For convenience, denote $\tilde{X}_{h,\tau} = \mathcal{T}X_{h,\tau}$ and $\tilde{Y}_{h,\tau} = \mathcal{T}Y_{h,\tau}$. By applying the definition of \mathcal{T} , we note that $\tilde{X}_{h,\tau} - \tilde{Y}_{h,\tau}$ will satisfy the homogeneous system

$$\begin{aligned} (4.3) \quad & (I + \tau_l \theta_l A_{II}^{h_l}) (\tilde{X}_{h_l, I}^{k+1} - \tilde{Y}_{h_l, I}^{k+1}) + \tau_l \theta_l A_{IB}^{h_l} (\tilde{X}_{h_l, B}^{k+1} - \tilde{Y}_{h_l, B}^{k+1}) \\ & = (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) (\tilde{X}_{h_l, I}^k - \tilde{Y}_{h_l, I}^k) + \tau_l \tilde{\theta}_l A_{IB}^{h_l} (\tilde{X}_{h_l, B}^k - \tilde{Y}_{h_l, B}^k), \\ & (\tilde{X}_{h_l, B_1}^{k+1} - \tilde{Y}_{h_l, B_1}^{k+1}) = 0, \\ & (\tilde{X}_{h_l, B_2}^{k+1} - \tilde{Y}_{h_l, B_2}^{k+1}) = \mathcal{I}_{h_l}^{k+1} (X_{h,\tau} - Y_{h,\tau}), \\ & (\tilde{X}_{h_l, I}^0 - \tilde{Y}_{h_l, I}^0) = 0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$. By the maximum principle from Section 3 we obtain

$$\begin{aligned} \|\tilde{X}_{h,\tau} - \tilde{Y}_{h,\tau}\| & \leq \max_{k=0, \dots, N_l} \max_{l=1, \dots, p} |\mathcal{I}_{h_l}^k (X_{h,\tau} - Y_{h,\tau})| \\ & \leq \sigma_{h,\tau} \max_{k=0, \dots, N_l} \max_{l=1, \dots, p} \max_{i \in \bar{\Omega}_l} |(X_{h_l}^k - Y_{h_l}^k)_i|. \end{aligned}$$

Here $\sigma_{h,\tau}$ is the maximum norm of the intergrid interpolation map $\mathcal{I}_{h_l}^k$.

Since $X_{h,\tau}, Y_{h,\tau} \in \mathcal{H}$, their difference $X_{h,\tau} - Y_{h,\tau}$ satisfies a discretized *homogeneous* parabolic equation, and by the contraction property, we obtain

$$\max_{i \in \bar{\Omega}_l} |(X_{h_l}^k - Y_{h_l}^k)_i| \leq \rho_{h_l} \|X_{h,\tau} - Y_{h,\tau}\|,$$

and consequently

$$\max_{k=0, \dots, N_l} \max_{l=1, \dots, p} \max_{i \in \bar{\Omega}_l} |(X_{h_l}^k - Y_{h_l}^k)_i| \leq \max_{l=1, \dots, p} \rho_{h_l} \|X_{h,\tau} - Y_{h,\tau}\|.$$

Combining the two bounds, we obtain

$$\begin{aligned} \|\tilde{X}_{h,\tau} - \tilde{Y}_{h,\tau}\| & \leq \max_{k=0, \dots, N_l} \max_{l=1, \dots, p} |\mathcal{I}_{h_l}^k (X_{h,\tau} - Y_{h,\tau})| \\ & \leq \sigma_{h,\tau} \max_{k=0, \dots, N_l; l=1, \dots, p} \max_{i \in \bar{\Omega}_l} |(X_{h_l}^k - Y_{h_l}^k)_i| \\ & \leq \sigma_{h,\tau} \max_{l=1, \dots, p} \rho_{h_l} \|X_{h,\tau} - Y_{h,\tau}\| \\ & = \delta \|X_{h,\tau} - Y_{h,\tau}\|. \end{aligned}$$

Since $d(\tilde{X}_{h,\tau}, \tilde{Y}_{h,\tau}) = \|\tilde{X}_{h,\tau} - \tilde{Y}_{h,\tau}\|$ and $d(X_{h,\tau}, Y_{h,\tau}) = \|X_{h,\tau} - Y_{h,\tau}\|$, and since by assumption $\delta < 1$, we obtain that \mathcal{T} is a contraction

$$d(\tilde{V}_{h,\tau}, \tilde{W}_{h,\tau}) \leq \delta d(V_{h,\tau}, W_{h,\tau}).$$

The unique solvability of the linear system follows from the uniqueness of the fixed point of a contraction mapping. The geometric convergence of the parallel Schwarz iterates follows by the geometric convergence of the Picard iterates. \square

We note that the preceding result does not directly provide an *a priori* bound for the maximum norm of the solution. This will be done in the next section.

4.2. Stability of the global discretization. In this section, we will derive an *a priori* bound for the maximum norm of the solution to (4.1).

Theorem 4.3. *Suppose the following hold.*

1. Let matrices A^{h_l} satisfy Assumptions A1 and A5 for $l = 1, \dots, p$.
2. Let $0 < \tau_l$ satisfy the local stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0,$$

for $l = 1, \dots, p$.

3. Let $\mathcal{I}_{h_l}^k$ satisfy Assumptions A2 and A3.
4. Let the overlap parameters β_l be chosen large enough, and the mesh sizes small enough with $h_l \leq h_l^*$ so that $\delta \equiv \sigma_{h,\tau} \max_l \rho_{h_l} \leq \delta < 1$.
5. Let $U_{h,\tau} = \{U_{h_l}^k\}_{k,l}$ denote the unique solution of system (4.1).

Then, the maximum norm of the solution $U_{h,\tau}$ satisfies

$$\begin{aligned} \|U_{h,\tau}\| \leq & \left(1 + \frac{\sigma_{h,\tau}}{1-\delta}\right) \max_{l=1,\dots,p} \left(\|\tilde{u}_{h_l,I}^0\|_{\infty,\Omega_{h_l}^*} + \tau_l \theta_l \|\tilde{f}_{h_l,I}^{N_l}\|_{\infty,\Omega_{h_l}^*} \right. \\ & \left. + \tau_l \sum_{k=1}^{N_l-1} \|\tilde{f}_{h_l,I}^k\|_{\infty,\Omega_{h_l}^*} + \tau_l \tilde{\theta}_l \|\tilde{f}_{h_l,I}^0\|_{\infty,\Omega_{h_l}^*} + \sum_{k=0}^{N_l} \|\tilde{g}_{h_l,B_2}^k\|_{\infty,B_2^{h_l}} \right). \end{aligned}$$

Proof. Choose any suitable grid function $X_{h,\tau} \in \mathcal{H}$ and use it as an initial guess in the Picard fixed point iteration. By Theorem 4.1 (the contraction mapping)

$$d(X_{h,\tau}, U_{h,\tau}) \leq \frac{1}{1-\delta} d(X_{h,\tau}, \mathcal{T}X_{h,\tau}).$$

Using that the metric in \mathcal{H} was inherited from the maximum norm, we obtain

$$\begin{aligned} \|U_{h,\tau}\| & \leq \|X_{h,\tau}\| + \|U_{h,\tau} - X_{h,\tau}\|, && \text{triangle inequality} \\ & = \|X_{h,\tau}\| + d(U_{h,\tau}, X_{h,\tau}) \\ & = \|X_{h,\tau}\| + \frac{1}{1-\delta} d(X_{h,\tau}, \mathcal{T}X_{h,\tau}), && \text{from above.} \end{aligned}$$

Therefore, to obtain a bound for $\|U_{h,\tau}\|$ we only need to choose $X_{h,\tau} \in \mathcal{H}$ and estimate $\|X_{h,\tau}\|$ and $d(X_{h,\tau}, \mathcal{T}X_{h,\tau})$.

Accordingly, choose $X_{h,\tau} = \{X_{h_l}^k\}_{k,l}$ as the solutions to the local discretized parabolic equations with trivial boundary conditions

$$(4.4) \quad \begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) X_{h_l,I}^{k+1} + \tau_l \theta_l A_{IB}^{h_l} X_{h_l,B}^{k+1} &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) X_{h_l,I}^k - \tau_l \tilde{\theta}_l A_{IB}^{h_l} X_{h_l,B}^k \\ &\quad + \tau_l \left(\theta_l \tilde{f}_{h_l,I}^{k+1} + \tilde{\theta}_l \tilde{f}_{h_l,I}^k \right), \\ X_{h_l,B_1}^{k+1} &= 0, \\ X_{h_l,B_2}^{k+1} &= 0, \\ X_{h_l,I}^0 &= \tilde{u}_{h_l,I}^0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$. Since the local problems for $\{X_{h_l}^k\}_k$ are *decoupled*, we can estimate the maximum norm of each local component X_{h_l,τ_l} independently by using the local *a priori* estimates from Section 3. This yields

$$\begin{aligned} \| \| X_{h,\tau} \| \| \leq \max_{l=1,\dots,p} \left(\| \tilde{u}_{h_l,I}^0 \|_\infty + \tau_l \theta_l \| \tilde{f}_{h_l,I}^{N_l} \|_{\infty, \Omega_{h_l}^*} + \tau_l \sum_{k=1}^{N_l-1} \| \tilde{f}_{h_l,I}^k \|_{\infty, \Omega_{h_l}^*} \right. \\ \left. + \tau_l \tilde{\theta}_l \| \tilde{f}_{h_l,I}^0 \|_{\infty, \Omega_{h_l}^*} + \sum_{k=0}^{N_l} \| \tilde{g}_{h_l,B_2}^k \|_{\infty, B_2^{h_l}} \right). \end{aligned}$$

We next estimate $\| \| X_{h,\tau} - \mathcal{T} X_{h,\tau} \| \|$. For convenience, let $\tilde{X}_{h,\tau} = \mathcal{T} X_{h,\tau}$. We note that $X_{h,\tau} - \tilde{X}_{h,\tau}$ will satisfy the discretized homogeneous parabolic equation

$$(4.5) \quad \begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) \left(X_{h_l,I}^{k+1} - \tilde{X}_{h_l,I}^{k+1} \right) + \tau_l \theta_l A_{IB}^{h_l} \left(X_{h_l,B}^{k+1} - \tilde{X}_{h_l,B}^{k+1} \right) \\ = \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) \left(X_{h_l,I}^k - \tilde{X}_{h_l,I}^k \right) - \tau_l \tilde{\theta}_l A_{IB}^{h_l} \left(X_{h_l,B}^k - \tilde{X}_{h_l,B}^k \right), \\ \left(X_{h_l,B_1}^{k+1} - \tilde{X}_{h_l,B_1}^{k+1} \right) = 0, \\ \left(X_{h_l,B_2}^{k+1} - \tilde{X}_{h_l,B_2}^{k+1} \right) = -\mathcal{I}_{h_l}^{k+1} X_{h,\tau}, \\ \left(X_{h_l,I}^0 - \tilde{X}_{h_l,I}^0 \right) = 0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$. By applying the discrete maximum principle and using $\sigma_{h,\tau} = \| \| \mathcal{I}_{h_l}^k \| \|$, we obtain

$$\| \| X_{h,\tau} - \tilde{X}_{h,\tau} \| \| \leq \sigma_{h,\tau} \| \| X_{h,\tau} \| \|.$$

Substituting these in our expression for $\| \| U_{h,\tau} \| \|$, we obtain

$$\begin{aligned} \| \| X_{h,\tau} \| \| \leq \left(1 + \frac{\sigma_{h,\tau}}{1-\delta} \right) \max_{l=1,\dots,p} \left(\| \tilde{u}_{h_l,I}^0 \|_{\infty, \Omega_{h_l}^*} + \tau_l \theta_l \| \tilde{f}_{h_l,I}^{N_l} \|_{\infty, \Omega_{h_l}^*} \right. \\ \left. + \tau_l \sum_{k=1}^{N_l-1} \| \tilde{f}_{h_l,I}^k \|_{\infty, \Omega_{h_l}^*} + \tau_l \tilde{\theta}_l \| \tilde{f}_{h_l,I}^0 \|_{\infty, \Omega_{h_l}^*} + \sum_{k=0}^{N_l} \| \tilde{g}_{h_l,B_2}^k \|_{\infty, B_2^{h_l}} \right), \end{aligned}$$

which is the desired result. \square

The above result depends critically on the contraction factor $\delta < 1$. If the overlap β_l of the local subregions are sufficiently large so that $\sigma_{h,\tau} \max_l \rho_{h_l} \leq \delta < 1$ (uniformly in h_l) then the global discretization will be stable. In the next section, we apply the above stability result to estimate the accuracy of the nonmatching grid discretization (2.4).

4.3. Accuracy of the global discretization. From the general theory for discretization of linear evolution equations [30], we expect a stable and consistent scheme to be convergent. The same holds for the nonmatching overlapping grid discretization scheme (2.4) considered here. We will now consider the consistency of the global scheme (2.4), which is measured by the magnitude of the residual when $\pi_{h,\tau}u$ (i.e., the exact solution $u(x, t)$ restricted to the collection of space-time grids) is substituted into the scheme.

Definition. Given the restriction $u_{h,\tau} = \pi_{h,\tau}u$ of the exact solution $u(x, t)$ to the space-time grids, we define the grid function $\mathcal{E}_{h_l}^k(u)$ to represent the *local discretization error* on the grid $\Omega_{h_l}^*$ at time $k\tau$

$$\begin{aligned} \mathcal{E}_{h_l}^{k+1}(u) \equiv & \left(I + \tau_l \theta_l A_{II}^{h_l} \right) u_{h_l, I}^{k+1} - \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) u_{h_l, I}^k \\ & + \tau_l \left(\theta_l A_{IB}^{h_l} u_{h_l, B}^{k+1} + \tilde{\theta}_l A_{IB}^{h_l} u_{h_l, B}^k \right) \\ & - \tau_l \left(\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k \right). \end{aligned}$$

We use $\mathcal{D}_{h_l}^k(u)$ to denote the boundary grid function representing the local intergrid interpolation error

$$\mathcal{D}_{h_l}^k(u) \equiv u_{h_l, B_2}^{k+1} - \mathcal{I}_{h_l}^{k+1} u_{h,\tau}.$$

At each gridpoint $(x_i^{h_l}, k\tau_l)$, the local discretization error $\mathcal{E}_{h_l}^k(u)$ and the interpolation error $\mathcal{D}_{h_l}^k(u)$ can be estimated by expanding the stencils using Taylor series expansions centered at the gridpoint. The resulting estimate will involve the local mesh parameters h_l and τ_l , and higher order derivatives of $u(\cdot, \cdot)$ at one or more points in the convex hull of the gridpoints involved in that stencil. For convenience suppose that the discretization and interpolation errors satisfy

$$(4.6) \quad \begin{aligned} \left| (\mathcal{E}_{h_l}^k(u))_i \right| & \leq C \|u\|_{q_l, 1+2, q_l, 2+1, \infty, \overline{\Omega}_l^* \times [0, T]} (h_l^{q_l, 1} + \tau_l^{q_l, 2}), \\ \left| (\mathcal{D}_{h_l}^k(u))_i \right| & \leq C \|u\|_{r_l, 1, r_l, 2, \infty, B_2^{h_l, *}} \times [0, T] (h_l^{r_l, 1} + \tau_l^{r_l, 2}), \end{aligned}$$

where $B_2^{h_l, *}$ is a neighborhood of the boundary B_2^l containing the union of all cells involved in the local intergrid interpolation.

We will now estimate the accuracy $\|u_{h,\tau} - U_{h,\tau}\|$ of the global discretization (2.4) in terms of the local discretization and interpolation errors, $\mathcal{E}_{h_l}^k(u)$ and $\mathcal{D}_{h_l}^k(u)$, respectively.

Theorem 4.4. *Suppose the following hold.*

1. Let matrices A^{h_l} satisfy Assumptions A1 and A5 for $k = 1, \dots, p$.

2. Let $0 < \tau_l$ satisfy the following local stability constraint

$$\tau_l \leq \min_{i \in I^{h_l}} \frac{1}{\tilde{\theta}_l A_{ii}^{h_l}}, \quad \text{if } \tilde{\theta}_l \neq 0,$$

for $l = 1, \dots, p$.

3. Let $\mathcal{I}_{h_l}^k$ satisfy Assumptions A2 and A3.

4. Let the overlap parameter β_l be chosen sufficiently large, and the mesh size sufficiently small with $h_l \leq h_l^*$ so that

$$\delta \equiv \sigma_{h,\tau} \max_l \rho_{h_l} < 1.$$

5. Let $U_{h,\tau}$ denote the solution to (2.4).

6. Let $u_{h,\tau} = \pi_{h,\tau} u$, the exact solution restricted to the grid, satisfy

$$\begin{aligned} (I + \tau_l \theta_l A_{II}^{h_l}) u_{h_l, I}^{k+1} &= (I - \tau_l \tilde{\theta}_l A_{II}^{h_l}) u_{h_l, I}^k \\ &\quad + \tau_l \left(-\theta_l A_{IB}^{h_l} u_{h_l, I}^{k+1} + \tilde{\theta}_l A_{IB}^{h_l} u_{h_l, I}^k \right) \\ &\quad + \tau_l \left(\theta_l f_{h_l, I}^{k+1} + \tilde{\theta}_l f_{h_l, I}^k \right) \\ &\quad + \mathcal{E}_{h_l}^{k+1}(u) \end{aligned} \tag{4.7}$$

$$u_{h_l, B_1}^{k+1} = 0,$$

$$u_{h_l, B_2}^{k+1} = \mathcal{I}_{h_l}^{k+1} u_{h,\tau} + \mathcal{D}_{h_l}^{k+1}(u),$$

$$u_{h_l, I}^0 = \pi_{h_l} u_0,$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$, where $\mathcal{E}_{h_l}^k(u)$ and $\mathcal{D}_{h_l}^k(u)$ denote the local discretization and interpolation errors, respectively.

7. Let the local discretization and interpolation errors satisfy bounds (4.6).

Then $u_{h,\tau} - U_{h,\tau}$ satisfies the bounds

$$\begin{aligned} \|u_{h,\tau} - U_{h,\tau}\| &\leq \left(1 + \frac{\sigma_{h,\tau}}{1 - \delta} \right) \max_{l=1, \dots, p} \left(\tau_l \theta_l \|\mathcal{E}_{h_l}^{N_l}(u)\|_{\infty, \bar{\Omega}_{h_l}^*} \right. \\ &\quad \left. + \tau_l \sum_{k=1}^{N_l-1} \|\mathcal{E}_{h_l}^k(u)\|_{\infty, \bar{\Omega}_{h_l}^*} + \tau_l \tilde{\theta}_l \|\mathcal{E}_{h_l}^0(u)\|_{\infty, \bar{\Omega}_{h_l}^*} \right. \\ &\quad \left. + \max_{k=0, \dots, N_l} \|\mathcal{D}_{h_l}^k(u)\|_{\infty, B_{l,2}^*} \right) \\ &\leq \left(1 + \frac{\sigma_{h,\tau}}{1 - \delta} \right) \max_{l=1, \dots, p} \left(C \|u\|_{q_1+2, q_2+1, \infty, \bar{\Omega}_l^* \times [0, T]} (h_l^{q_1+1} + \tau_l^{q_1+2}) \right. \\ &\quad \left. + C \|u\|_{r_{l,1}, r_{l,2}, \infty, B_2^{h_l, *}} \times [0, T] (h_l^{r_{l,1}} + \tau_l^{r_{l,2}}) \right). \end{aligned} \tag{4.8}$$

Proof. Subtract equation (4.7) satisfied by $u_{h,\tau}$ from (2.4) satisfied by $U_{h,\tau}$. We obtain

$$\begin{aligned} \left(I + \tau_l \theta_l A_{II}^{h_l} \right) \left(u_{h_l,I}^{k+1} - U_{h_l,I}^{k+1} \right) &= \left(I - \tau_l \tilde{\theta}_l A_{II}^{h_l} \right) \left(u_{h_l,I}^k - U_{h_l,I}^k \right) \\ &\quad + \tau_l \left(-\theta_l A_{IB}^{h_l} \left(u_{h_l,I}^{k+1} - U_{h_l,I}^{k+1} \right) \right. \\ &\quad \quad \left. + \tilde{\theta}_l A_{IB}^{h_l} \left(u_{h_l,I}^k - U_{h_l,I}^k \right) \right) \\ &\quad + \mathcal{E}_{h_l}^{k+1}(u) \\ \left(u_{h_l,B_1}^{k+1} - U_{h_l,B_1}^{k+1} \right) &= 0, \\ \left(u_{h_l,B_2}^{k+1} - U_{h_l,B_2}^{k+1} \right) &= \mathcal{I}_{h_l}^{k+1} \left(u_{h,\tau} - U_{h,\tau} \right) + \mathcal{D}_{h_l}^{k+1}(u), \\ \left(u_{h_l,I}^0 - U_{h_l,I}^0 \right) &= 0, \end{aligned}$$

for $k = 0, \dots, N_l - 1$ and $l = 1, \dots, p$.

Since the hypothesis of the stability theorem from the preceding section is satisfied, we may apply it with $\tilde{f}_{h_l,I}^k = \mathcal{E}_{h_l}^k(u)$ and $\tilde{g}_{h_l}^k = \mathcal{D}_{h_l}^k(u)$ to obtain

$$\begin{aligned} \|u_{h,\tau} - U_{h,\tau}\| \leq \left(1 + \frac{\sigma_{h,\tau}}{1 - \delta} \right) \max_{l=1,\dots,p} \left(\tau_l \theta_l \|\mathcal{E}_{h_l}^{N_l}(u)\|_{\infty, \overline{\Omega}_{h_l}^*} \right. \\ \quad \left. + \tau_l \sum_{k=1}^{N_l-1} \|\mathcal{E}_{h_l}^k(u)\|_{\infty, \overline{\Omega}_{h_l}^*} + \tau_l \tilde{\theta}_l \|\mathcal{E}_{h_l}^0(u)\|_{\infty, \overline{\Omega}_{h_l}^*} \right. \\ \quad \left. + \max_{k=0,\dots,N_l} \|\mathcal{D}_{h_l}^k(u)\|_{\infty, B_{l,2}^*} \right). \end{aligned}$$

Substituting the bounds (4.6) for the local discretization and interpolation errors, we obtain

$$\begin{aligned} \|u_{h,\tau} - U_{h,\tau}\| \\ \leq \left(1 + \frac{\sigma_{h,\tau}}{1 - \delta} \right) \max_{l=1,\dots,p} \left(C \|u\|_{q_{l,1}+2, q_{l,2}+1, \infty, \overline{\Omega}_l^* \times [0,T]} \left(h_l^{q_{l,1}} + \tau_l^{q_{l,2}} \right) \right. \\ \quad \left. + C \|u\|_{r_{l,1}, r_{l,2}, \infty, B_2^{h_l,*} \times [0,T]} \left(h_l^{r_{l,1}} + \tau_l^{r_{l,2}} \right) \right), \end{aligned}$$

which is the desired result. \square

Remark 8. The above global error bound provides some guidance on the selection of the local mesh sizes. If a global error of $O(\epsilon)$ is desired, then h_l , τ_l and $\mathcal{I}_{h_l}^k$ should be chosen so that

$$\begin{aligned} \epsilon &\approx \|u\|_{r_{l,1}, r_{l,2}, \infty, B_2^{h_l,*} \times [0,T]} \left(h_l^{r_{l,1}} + \tau_l^{r_{l,2}} \right) \\ &\approx \|u\|_{q_{l,1}+2, q_{l,2}+1, \infty, \overline{\Omega}_l^* \times [0,T]} \left(h_l^{q_{l,1}} + \tau_l^{q_{l,2}} \right), \end{aligned}$$

for $l = 1, \dots, p$. Thus, the mesh sizes should be smaller (or the interpolation stencil, of higher order) in regions where the solution is less smooth.

Remark 9. We assumed throughout this paper that $c(x) \geq c_0 > 0$, in order to guarantee the stability of the global discretization (2.4). In practice, however, the discretization considered should be stable and convergent even if this condition is

violated, though the authors do not have rigorous results on this. We illustrate this by a simple example. Consider the parabolic equation

$$\left\{ \begin{array}{ll} u_t - u_{xx} = f(x, t), & \text{in } (0, 1) \times [0, T], \\ u(x, t) = 0, & \text{on } \{0\} \times [0, T], \\ u(x, t) = 0, & \text{on } \{1\} \times [0, T], \\ u(x, 0) = u_0(x), & \text{on } (0, 1). \end{array} \right.$$

Choose the following subdomains $\Omega_1 = (0, 1/3)$, $\Omega_2 = (1/3, 2/3)$, $\Omega_3 = (2/3, 1)$ and the following extended subdomains $\Omega_1^* = (0, 1/2)$, $\Omega_2^* = (1/4, 3/4)$ and $\Omega_3^* = (1/2, 1)$. Since the associated elliptic operator is $Lu = -u_{xx}$, we can solve local homogeneous problems explicitly to obtain the estimates for the local contraction factors

$$\rho_1 = \frac{1}{6}, \quad \rho_2 = 1, \quad \rho_3 = \frac{1}{6}.$$

We note that $\rho_2 = 1$ since on subdomain Ω_2^* , $u(x) \equiv 1$ is the solution to the homogeneous elliptic equation $-u_{xx} = 0$ with boundary conditions $u(1/4) = 1$ and $u(3/4) = 1$. Thus, the condition $\delta = \sigma_{h,\tau} \max_l \rho_l < 1$ will be violated, and the mapping \mathcal{T} will not be a contraction. However, it can easily be verified that the mapping \mathcal{T}^2 is a contraction with contraction factor $1/6$, and the global scheme should be stable. More generally, when $c(x) = 0$, we expect the subdomains adjacent to the boundary $\partial\Omega$ to have contraction factors less than 1 and interior (floating) subdomains to have contraction factors of 1. Repeated applications of \mathcal{T} should however “propagate” the contraction property to the interior subdomains, and \mathcal{T}^{k_0} may be contractive for some $k_0 > 1$. Unfortunately, rigorous results along these lines are not known to the authors.

Remark 10. We indicate briefly how the results of this paper can be extended to certain *semilinear* reaction diffusion equations. Consider the equation

$$\left\{ \begin{array}{ll} u_t - \Delta u + \vec{b}(x) \cdot \nabla u + c(x, u) = f(x, t), & \text{in } \Omega \times [0, T], \\ u(x, t) = 0, & \text{on } \partial\Omega \times [0, T], \\ u(x, 0) = u_0(x), & \text{on } (0, 1), \end{array} \right.$$

where $c(x, u)$ is a smooth function and satisfies

$$\frac{\partial c}{\partial u}(x, u) \geq c_0 > 0, \quad \forall x, u.$$

On each local grid, we will discretize the semilinear elliptic operator

$$L(u) \equiv -\Delta u + \vec{b}(x) \cdot \nabla u + c(x, u)$$

using finite difference schemes, and discretize the parabolic equation in time using a θ -scheme. Due to the nonlinearity $c(x, u)$, the following approximation will be used to avoid solving a nonlinear equation at each time step

$$c(x, U_{h_i}^{k+1}) \approx c(x, U_{h_i}^k) + c_u(x, U_{h_i}^k) (U_{h_i}^{k+1} - U_{h_i}^k).$$

This will correspond to applying one-step of a Newton approximation and the resulting θ -scheme approximation of the semilinear terms on $[k\tau_l, (k+1)\tau_l]$ will be:

$$\begin{aligned} & \theta_l c(x_i^{h_l}, (U_{h_l}^{k+1})_i) \\ & + \tilde{\theta}_l c(x_i^{h_l}, (U_{h_l}^k)_i) \approx \theta_l \left\{ c(x_i^{h_l}, (U_{h_l}^k)_i) + c_u(x_i^{h_l}, (U_{h_l}^k)_i) ((U_{h_l}^{k+1} - U_{h_l}^k)_i) \right\} \\ & \quad + \tilde{\theta}_l c(x_i^{h_l}, (U_{h_l}^k)_i) \\ & = c(x_i^{h_l}, (U_{h_l}^k)_i) \\ & \quad + \tilde{\theta}_l c_u(x_i^{h_l}, (U_{h_l}^k)_i) ((U_{h_l}^{k+1} - U_{h_l}^k)_i). \end{aligned}$$

Due to the above linearization and the sign properties of $c_u(\cdot, \cdot)$, the existence of solutions of the local θ -discretizations are guaranteed. Provided the local time steps τ_l are small enough to satisfy the local stability conditions (which would depend on the $c_u(\cdot, \cdot)$ terms), the theory developed in this paper can be extended to analyze the accuracy of the global discretization of the semilinear parabolic equation. The local truncation error terms will be different from the linear case and the local interpolation error will be identical to the linear case. We omit the details.

Remark 11. To simplify our discussions, we had assumed throughout the paper that the diffusion term was $a\Delta u$. More generally, the results of this paper are valid even if the diffusion term is of the form $\nabla \cdot (a(x)\nabla u)$, where $a(x)$ is a diagonal matrix with positive diagonal entries, provided the spatial discretization yields an M -matrix. In particular, if $a(x)$ is zero on a subregion (or on all of Ω) the results would still hold (provided the discretization yields an M -matrix, which, due to the hyperbolic nature of the equation, would necessitate that a first order upwind discretization be used to discretize $\vec{b}(x) \cdot \nabla u$). Consider now the original parabolic equation with $a \ll 1$ (i.e., a singularly perturbed parabolic equation). Suppose Ω_1^* and Ω_2^* are two overlapping subregions covering Ω , such that

$$|a\Delta u| \leq \eta, \quad \text{on } \Omega_1^* \times [0, T], \quad \text{where } \eta \ll 1.$$

If $a\Delta u$ is dropped on $\Omega_1^* \times [0, T]$, the resulting hyperbolic equation may be discretized by an explicit scheme (with suitable restriction on the time step) without requiring the solution of a linear system. A coupled hyperbolic-parabolic problem may be constructed to approximate the original parabolic equation (see for instance [29, 11]), as indicated next. Let $B_1^i = \partial\Omega_i^* \cap \partial\Omega$ and $B_2^i = \partial\Omega_i^* \cap \Omega$ for $i = 1, 2$, and let $B_{j,in}^1$ denote the inflow segment of B_j^1 . Additionally, let v and w denote the approximations of u on $\Omega_1^* \times [0, T]$ and $\Omega_2^* \times [0, T]$, respectively. A coupled hyperbolic-parabolic system for v and w can be posed as follows:

$$\left\{ \begin{array}{ll} v_t + \vec{b}(x) \cdot \nabla v + c(x)v = f(x), & \text{in } \Omega_1^* \times [0, T], \\ v = 0, & \text{on } B_{1,in}^1 \times [0, T], \\ v = w, & \text{on } B_{2,in}^1 \times [0, T], \\ v = u_0(x), & \text{for } t = 0 \end{array} \right.$$

and

$$\left\{ \begin{array}{ll} w_t - a\Delta w + \vec{b}(x) \cdot \nabla w + c(x)w = f(x), & \text{in } \Omega_2^* \times [0, T], \\ w = 0, & \text{on } B_1^2 \times [0, T], \\ w = v, & \text{on } B_2 \times [0, T], \\ w = u_0(x), & \text{for } t = 0. \end{array} \right.$$

Provided the M -matrix assumption holds for the local discretizations, Theorem 4.4 would guarantee the maximum norm stability of the global nonmatching overset grid discretization of the above system. If u denotes the solution of the original parabolic equation (with $u_{h,\tau}$ denoting its restriction to the space-time grids) and if $(v_{h_1,\tau_1}, w_{h_2,\tau_2})$ denotes the discrete hyperbolic-parabolic solution, then the error $u_{h,\tau} - (v_{h_1,\tau_1}, w_{h_2,\tau_2})$ will satisfy (4.8) with an additional term of magnitude η on $\Omega_1^* \times [0, T]$:

$$\begin{aligned} & \| \| u_{h,\tau} - (v_{h_1,\tau_1}, w_{h_2,\tau_2}) \| \| \\ & \leq \left(1 + \frac{\sigma_{h,\tau}}{1 - \delta} \right) \max_{l=1,\dots,p} \left(C \| u \|_{q_{l,1}+2, q_{l,2}+1, \infty, \overline{\Omega}_l^* \times [0, T]} (h_l^{q_{l,1}} + \tau_l^{q_{l,2}}) \right. \\ & \qquad \qquad \qquad \left. + C \| u \|_{r_{l,1}, r_{l,2}, \infty, B_2^{h_l, \tau_l} \times [0, T]} (h_l^{r_{l,1}} + \tau_l^{r_{l,2}}) + \eta \right). \end{aligned}$$

Thus, if η is of the same magnitude as the local truncation and interpolation errors, then

$$\| \| u_{h,\tau} - (v_{h_1,\tau_1}, w_{h_2,\tau_2}) \| \| \approx \| \| u_{h,\tau} - (U_{h_1,\tau_1}, U_{h_2,\tau_2}) \| \|,$$

where $(U_{h_1,\tau_1}, U_{h_2,\tau_2})$ is the nonmatching overset grid solution of the full parabolic equation. The domain Ω_1^* may be adaptively determined as in the χ -formulation (see [8]). We omit further details.

ACKNOWLEDGMENTS

The authors wish to thank the *Consiglio Nazionale delle Ricerche* of Italy for support during the visit of the first author to the Università dell'Aquila.

REFERENCES

1. G. Abdoulaev, Y. Achdou, J. Hontand, Y. Kuznetsov, O. Pironneau, and C. Prud'homme, *Nonmatching grids for fluids*, The tenth international conference on domain decomposition methods for partial differential equations (Providence, R.I) (J. Mandel, C. Farhat, and X.-C. Cai, eds.), AMS, 1998. MR **99i**:76103
2. Y. Achdou, Y. Maday, and O. Widlund, *Iterative substructuring preconditioners for mortar element methods in two dimensions*, Tech. report, TR-735, Department of Computer Science, Courant Institute of Mathematical Sciences, 1997; SIAM J. Numer. Anal. **36** (1999), 551–580. MR **99m**:65233
3. V. I. Arnold, *Ordinary differential equations*, Springer-Verlag, New York, 1992. MR **93b**:34001
4. F. Ben Belgacem, *The mortar finite element method with Lagrange multipliers*, Numer. Math. **84** (1999), 173–197.
5. C. Bernardi, Y. Maday, and A. Patera, *A new nonconforming approach to domain decomposition: The mortar element method*, Colloque de France Seminar (H. Brezis and J. L. Lions, eds.), Pitman, 1990. MR **95a**:65201
6. C. Börgers and C. S. Peskin, *A Lagrangian fractional step method for the incompressible Navier-Stokes equations on a periodic domain*, J. Comp. Phys. **70** (1987), no. 2, 397–438. MR **89c**:76007
7. J. H. Bramble, R. E. Ewing, J.E. Pasciak, and A. H. Schatz, *A preconditioning technique for the efficient solution of problems with local grid refinement*, Comput. Meth. Appl. Mech. Engrg. **67** (1988), 149–159.
8. F. Brezzi, C. Canuto, and A. Russo, *A self-adaptive formulation for the Euler/Navier-Stokes coupling*, Comp. Meth. in Appl. Mech. Engrg. **73** (1989), 317–330. MR **90h**:76075
9. X. C. Cai, M. Dryja, and M. Sarkis, *Overlapping nonmatching grids mortar element methods for elliptic problems I: Error analysis*, SIAM J. Numer. Anal. (2000), **36** (1999), 581–606. MR **2000a**:65142

10. X. C. Cai, T. P. Mathew, and M. Sarkis, *Maximum norm analysis of overlapping nonmatching grid discretizations of elliptic equations*, SIAM J. Numer. Anal. **37** (2000), no. 5, 1709–1728. MR **2001c**:65130
11. C. Canuto and A. Russo, *Self-adaptive coupling of mathematical models and/or numerical methods*, Contemporary Mathematics **157** (1994), 35–44. MR **94m**:35085
12. T. F. Chan and T. P. Mathew, *Domain decomposition algorithms*, Acta Numerica (1994), 61–143. MR **95f**:65214
13. G. Chesshire and W. D. Henshaw, *Composite overlapping meshes for the solution of partial differential equations*, J. Comp. Phys. **90** (1990), 1–64.
14. P. G. Ciarlet, *Discrete maximum principle for finite-difference operators*, Aequationes Math. **4** (1970), 338–352. MR **45**:1404
15. M. Dryja and O. B. Widlund, *An additive variant of the Schwarz alternating method for the case of many subregions*, Tech. Report 339, also Ultracomputer Note 131, Department of Computer Science, Courant Institute, 1987.
16. ———, *Domain decomposition algorithms with small overlap*, SIAM J. Sci. Comp. **15** (1994), no. 3, 604–620. MR **95d**:65102
17. R. E. Ewing, *Domain decomposition techniques for efficient adaptive local grid refinement*, Domain Decomposition Methods (Philadelphia, PA) (Tony Chan, Roland Glowinski, Jacques Périaux, and Olof Widlund, eds.), SIAM, 1989. MR **90c**:65141
18. R. E. Ewing, R. D. Lazarov, and P. S. Vassilevski, *Local refinement techniques for elliptic problems on cell-centered grids*, Tech. report, University of Wyoming, 1988.
19. C. Farhat, J. Mandel, and F. X. Roux, *Optimal convergence properties of the FETI domain decomposition method*, Comp. Meth. Appl. Mech. Engrg. **115** (1994), 365–385. MR **95d**:65091
20. P. J. J. Ferket and A. A. Reusken, *A finite difference discretization method for elliptic problems on composite grids*, Computing **56** (1996), 343–369. MR **97c**:65174
21. M. Garbey, Y. Kuznetsov, and Y. Vassilevski, *Parallel Schwarz methods for advection-diffusion equations*, SIAM J. Sci. Comp. **22**–3, (2000), 891–916.
22. E. Giladi and H. B. Keller, *Space-time domain decomposition for parabolic problems*, Proceedings of IMACS conference (J. Wang, M. Allen, B. Chen, and T. Mathew, eds.), 1997, Also appeared as technical report from Center for Research on Parallel Computation, CRPC-TR97701.
23. W. D. Henshaw, *Automatic grid generation*, Acta Numerica (1996), 121–148. CMP 98:14
24. C. Johnson, *Numerical solutions of partial differential equations by the finite element method*, Cambridge University Press, Cambridge, 1987. MR **89b**:65003a and MR **89b**:65003b
25. A. Karafiat, *Discrete maximum principle in parabolic boundary value problems*, Ann. Polon. Math. **53** (1991), 253–265. MR **92d**:35052
26. P. L. Lions, *On the Schwarz alternating method. I.*, First International Symposium on Domain Decomposition Methods for Partial Differential Equations (Philadelphia, PA) (Roland Glowinski, Gene H. Golub, Gérard A. Meurant, and Jacques Périaux, eds.), SIAM, 1988. MR **90a**:65248
27. ———, *On the Schwarz alternating method. II.*, Domain Decomposition Methods (Philadelphia, PA) (Tony Chan, Roland Glowinski, Jacques Périaux, and Olof Widlund, eds.), SIAM, 1989. MR **90e**:65140
28. T. P. Mathew, *Uniform convergence of the Schwarz alternating method for solving singularly perturbed advection diffusion equations*, SIAM J. Numer. Anal. **35** (1998), no. 4, 1663–1683. MR **99f**:65156
29. A. Quarteroni and A. Valli, *Theory and applications of Steklov-Poincaré operators for boundary-value problems: the heterogeneous operator case*, Proceedings of 4th International Conference on Domain Decomposition Methods, Moscow (Philadelphia) (Tony Chan, Roland Glowinski, Jacques Périaux, and Olof Widlund, eds.), SIAM, 1990. MR **92g**:65132
30. R. D. Richtmyer and K. W. Morton, *Difference methods for initial-value problems*, Wiley Interscience, New York, 1967. MR **36**:3515
31. Y. Saad, *Iterative methods for sparse linear systems*, PWS Publishing Company, Boston, MA, U.S.A., 1996.
32. G. Starius, *Composite mesh difference methods for elliptic boundary value problems*, Numer. Math. **28** (1977), 243–258. MR **57**:1923
33. J. Steger and J. Benek, *On the use of composite grid schemes in computational aerodynamics*, Comp. Meth. Appl. Mech. Engrg. **64** (1987), 301–320. MR **88i**:65146

34. R. S. Varga, *Matrix iterative analysis*, Prentice-Hall, 1962. MR **28**:1725
35. J. Xu, *Iterative methods by space decomposition and subspace correction*, SIAM Review **34** (1992), 581–613. MR **93k**:65029

115 SEAL ROCK DRIVE, SAN FRANCISCO, CALIFORNIA 94121
E-mail address: `tmathew@mindspring.com`

DIPARTIMENTO DI MATEMATICA ED INFORMATICA, UNIVERSITÀ DI CATANIA, VIALE ANDREA
DORIA 6, 95125 CATANIA, ITALY
E-mail address: `russo@dmi.unict.it`