

NEW AMICABLE FOUR-CYCLES

KARSTEN BLANKENAGEL, WALTER BORHO, AND AXEL VOM STEIN

ABSTRACT. Fifty new amicable four-cycles are discovered by the constructive method invented in 1969 by the second author.

1. Let $\tau(n)$ denote the sum of proper divisors of a natural number n , and let $\sigma(n) = n + \tau(n)$. We consider when the sequence $n, \tau(n), \tau^{(2)}(n) := \tau(\tau(n)), \dots$ becomes periodic. If $n = \tau^{(k)}(n)$ with k minimal, then $n_1 = n, n_2 = \tau(n), n_3 = \tau^{(2)}(n), \dots, n_k = \tau^{(k-1)}(n)$ is called an *amicable k -cycle*. The study of amicable 1-cycles (*perfect numbers*) and of amicable 2-cycles (*amicable pairs*) has a thousand-year-old history. Here we study amicable four-cycles. The smallest example is

$$\begin{aligned} n_1 &= 2^2 \cdot 5 \cdot 17 \cdot 3719, & n_3 &= 2^2 \cdot 521 \cdot 829, \\ n_2 &= 2^2 \cdot 5 \cdot 193 \cdot 401, & n_4 &= 2^5 \cdot 40787, \end{aligned}$$

discovered by H. Cohen [5] in 1970 by an exhaustive trial and error search below 60,000,000.

2. Alternatively, one may try to *construct* amicable four-cycles of a special form. This can be done by means of the following theorem, due to the second author.

Theorem 1 ([3]). *Let a_1 and a_2 be natural numbers, $a_1 \neq a_2$, and let $D := a_1 a_2 - \tau(a_1) \tau(a_2)$. Let $d_1 d_2 = a_1 a_2$ be any factorization into two natural numbers d_1, d_2 . Consider the six numbers p_{ij}, r_i ($i, j = 1, 2$)*

$$\begin{aligned} (1) \quad p_{ij} &:= \frac{1}{D} (\tau(a_{i+1}) \sigma(a_i) + d_j \sigma(a_{i+1})), & \text{where } a_3 &:= a_1, \\ (2) \quad r_i &:= \frac{1}{a_i} \tau(a_i p_{i1} p_{i2}). \end{aligned}$$

If all six are primes, and $p_{ij} \nmid a_i, r_i \nmid a_i, p_{i1} \neq p_{i2}$ ($i, j = 1, 2$), then the following is an amicable four-cycle:

$$\begin{aligned} n_1 &= a_1 p_{11} p_{12}, & n_3 &= a_2 p_{21} p_{22}, \\ n_2 &= a_1 r_1, & n_4 &= a_2 r_2. \end{aligned}$$

The smallest example is

$$\begin{aligned} n_1 &= 3^3 \cdot 5 \cdot 7 \cdot 83 \cdot 359, & n_3 &= 3^3 \cdot 5 \cdot 11 \cdot 79 \cdot 263, \\ n_2 &= 3^3 \cdot 5 \cdot 7 \cdot 31643, & n_4 &= 3^3 \cdot 5 \cdot 11 \cdot 20183, \end{aligned}$$

which was found in [3] in 1969 without use of a computer.

Received by the editor September 19, 2001 and, in revised form, February 18, 2002.
 2000 *Mathematics Subject Classification*. Primary 11A25.
Key words and phrases. Number theory, sociable numbers.

To specify these four numbers, it is enough to give a_1, a_2 and d_1 , because the rest is then given by formulas (1) and (2) of the theorem. In the example

$$3^3 \cdot 5 \cdot \begin{cases} 7 & = a_1, \\ 11 & = a_2, \end{cases} \quad d_1 = 3^2 \cdot 5 \cdot 7.$$

3. By a constructive search based on the above theorem, we produced 50 new (amicable) four-cycles specified below. Of the 60 (amicable) four-cycles previously known, 30 were found by this same constructive method, while 30 were found by trial and error (below 10^{12}). Here is the list of the discoverers:

years	Discoverers	number of discoveries	method	references
1969	Borho	1	constructive	[3]
1970	Cohen	8	trial and error	[5]
1972	David / Root	4 / 5	trial and error	[6] / [2]
1990	Yuanhua	3	constructive	[10]
1990	Flammenkamp	7	trial and error	[7]
1990-95	Moews, Moews	9	trial and error	[8] / [9]
1990-98	Moews, Moews	8	constructive	[8]
1997-00	Pedersen	9	constructive	
2000	Baader	1	trial and error	
2000-01	Ren Yuanhua	9	constructive	
2000-01	new	50	constructive	

4. In the constructive search with the theorem above, the choice of a_1 and a_2 is of course essential. It is useful to check the following necessary condition for a_1, a_2 .

Theorem 2 (Four-cycle condition). *In order that the p_{ij} in Theorem 1 can be integers, it is necessary that*

$$(a_1 - a_2) f g \equiv 0 \pmod{D},$$

where f resp. g is the greatest common divisor of $\sigma(a_1), \sigma(a_2)$ resp. of a_1, a_2 .

We call a four-cycle given by Theorem 1 “regular of type (r, s) ” if a_1, a_2 have the form

$$a_1 = a p_1 p_2 \dots p_r, \quad a_2 = a q_1 q_2 \dots q_s,$$

with some greatest common divisor a , and different primes $p_1, \dots, p_r, q_1, \dots, q_s$ not dividing a . The 50 new amicable four-cycles discovered here are all “regular” in this sense. Nonregular or “exotic” examples were found by Ren Yuanhua and Pedersen.

5. For our constructive search for four-cycles, we must make the denominator $D := D(a_1, a_2) = a_1 a_2 - \tau(a_1) \tau(a_2) = a_1 \sigma(a_2) + a_2 \sigma(a_1) - \sigma(a_1) \sigma(a_2)$ small in order to increase the chance that the p_{ij} become integers. But we can cancel by $g_i := \gcd(a_i, \sigma(a_i))$ for $i = 1, 2$, and in addition by $\gcd(\sigma(a_1)/g_1, \sigma(a_2)/g_2)$. Let

$\bar{D}(a_1, a_2)$ denote the result. What we really have to make small is this $\bar{D}(a_1, a_2)$. (One might take d_j a multiple of g_j to make p_{ij} integer.)

Let us describe, for example, our search for regular four-cycles of type (1, 1). So let $a_1 = a p_1, a_2 = a q_1$ as in section 4. For a we took all values $\leq 50,000,000$ with $\bar{D}(a, a) < 100$. These are 6224 numbers. Next, p_1 must be chosen greater than $\tau(a) \sigma(a) / D(a, a)$ (to make $D(a p_1, a)$ positive). All primes p_1 greater than this lower bound were taken in increasing order as long as $\bar{D}(a p_1, a) < 100$. Finally, q_1 was chosen in such a way that the four-cycle condition (Theorem 2) was satisfied. To all pairs $a_1 = a p_1, a_2 = a q_1$ so chosen, Theorem 1 was applied; that is, all factorizations $d_1 d_2 = a_1 a_2$ were checked to see whether they give an amicable four-cycle.

This search produced seven of the thirteen known four-cycles of type (1, 1), plus in addition two new ones, numbers 2), 3) of the list below.

Searches for four-cycles of types (1, 0), (2, 1), (2, 2), (3, 1), (3, 2), (3, 3) were performed in a similar manner. The new results are listed below. Note that no four-cycles of type (1, 0) are known so far. It is an interesting open problem to find a four-cycle of this type.

An interesting phenomenon that occurs so far only in four-cycles of type (1, 1) is that p_{11} may equal p_{21} . This actually occurs in 9 of the 16 known cases of type (1, 1).

For our search for amicable four-cycles we spent a total of about 3 years of computation on Sun desk calculators of 333 MHz.

6. The following table specifies all 50 new amicable four-cycles found by the searches described above. In fact, for each of the 50 four-cycles we give the triples a_1, a_2 and d_1 in prime factorization. From these data, the full four-cycle itself can be easily computed by the formulae given in Theorem 1. For instance, for four-cycle number 4 below, one obtains the 14 decimal digit numbers (in prime decomposition):

$$n_1 = 2 \cdot 7 \cdot 223 \cdot 5 \cdot 11 \cdot 3121 \cdot 73589, \quad n_3 = 2 \cdot 7 \cdot 223 \cdot 23 \cdot 4013 \cdot 215417,$$

$$n_2 = 2 \cdot 7 \cdot 223 \cdot 5 \cdot 11 \cdot 288231067, \quad n_4 = 2 \cdot 7 \cdot 223 \cdot 23 \cdot 689238587.$$

We let each line of the following table begin with an ordinal number of the corresponding four-cycle, followed by the number of decimal digits of the smallest member of the four-cycle, with a “D” for “decimal digits”. Next, a_1 and a_2 are given, with the common divisor given only once. Finally, d_1 is given. The four-cycles are ordered according to their type.

type (1, 1)			
1)	23D	$2 \cdot 7^2 \cdot 13 \cdot 19 \cdot 101 \cdot$	$\begin{cases} 118169 \\ 236339 \end{cases} \quad d_1 = 2 \cdot 7^2 \cdot 13 \cdot 19 \cdot 101$
2)	26D	$3^2 \cdot 5^2 \cdot 19 \cdot 29 \cdot 41 \cdot$	$\begin{cases} 491 \\ 294461 \end{cases} \quad d_1 = 3^2 \cdot 5^2 \cdot 19 \cdot 41 \cdot 294461$
3)	27D	$3^3 \cdot 7^2 \cdot 13 \cdot 17 \cdot 67 \cdot$	$\begin{cases} 179 \\ 49139 \end{cases} \quad d_1 = 3^3 \cdot 7^2 \cdot 13 \cdot 17 \cdot 67$

type (2, 1)

- | | | | |
|-----|-----|---|--|
| 4) | 14D | $2 \cdot 7 \cdot 223 \cdot \begin{cases} 5 \cdot 11 \\ 23 \end{cases}$ | $d_1 = 2 \cdot 7 \cdot 223$ |
| 5) | 15D | $3^3 \cdot 5^3 \cdot \begin{cases} 7 \cdot 59 \\ 89 \end{cases}$ | $d_1 = 3^3 \cdot 5^3$ |
| 6) | 16D | $3^2 \cdot 5^2 \cdot 31 \cdot \begin{cases} 7 \cdot 59 \\ 89 \end{cases}$ | $d_1 = 3^2 \cdot 5^2 \cdot 31$ |
| 7) | 17D | $3^2 \cdot 5 \cdot 13 \cdot 19 \cdot \begin{cases} 53 \cdot 379 \\ 71 \end{cases}$ | $d_1 = 3^2 \cdot 5 \cdot 13 \cdot 19 \cdot 53$ |
| 8) | 22D | $3^5 \cdot 5 \cdot 13 \cdot \begin{cases} 17 \cdot 109 \\ 17929 \end{cases}$ | $d_1 = 3^7 \cdot 5 \cdot 13 \cdot 109$ |
| 9) | 23D | $3^5 \cdot 7^2 \cdot 13 \cdot 83 \cdot \begin{cases} 11 \cdot 53 \\ 1871 \end{cases}$ | $d_1 = 3^5 \cdot 7^2 \cdot 13 \cdot 1871$ |
| 10) | 24D | $3^2 \cdot 5 \cdot 13 \cdot 23 \cdot 137 \cdot \begin{cases} 29 \cdot 547 \\ 349 \end{cases}$ | $d_1 = 3^2 \cdot 5 \cdot 13 \cdot 23 \cdot 29 \cdot 137$ |
| 11) | 24D | $2^4 \cdot 101 \cdot \begin{cases} 29 \cdot 797 \\ 113 \end{cases}$ | $d_1 = 2^4 \cdot 29$ |
| 12) | 26D | $3^7 \cdot 5^2 \cdot 41 \cdot \begin{cases} 11 \cdot 79 \\ 7529 \end{cases}$ | $d_1 = 3^{11} \cdot 5^2 \cdot 41$ |
| 13) | 31D | $3^2 \cdot 5 \cdot 13 \cdot 19 \cdot \begin{cases} 571 \cdot 162749 \\ 29 \end{cases}$ | $d_1 = 3^2 \cdot 5 \cdot 13 \cdot 19 \cdot 571$ |
| 14) | 32D | $3^4 \cdot 7^2 \cdot 11 \cdot 19 \cdot \begin{cases} 379 \cdot 523 \\ 5177119 \end{cases}$ | $d_1 = 3^4 \cdot 11 \cdot 19 \cdot 5177119$ |
| 15) | 36D | $2^8 \cdot 619 \cdot \begin{cases} 1487 \cdot 105229 \\ 391455599 \end{cases}$ | $d_1 = 2^8 \cdot 619 \cdot 105229$ |
| 16) | 36D | $3^2 \cdot 7 \cdot 11 \cdot 17 \cdot 23 \cdot \begin{cases} 101 \cdot 397493 \\ 2039 \end{cases}$ | $d_1 = 3^2 \cdot 7 \cdot 17 \cdot 23^2 \cdot 101$ |

type (2, 2)

- | | | | |
|-----|-----|---|---|
| 17) | 16D | $3^2 \cdot 5 \cdot 13 \cdot \begin{cases} 11 \cdot 1279 \\ 23 \cdot 127 \end{cases}$ | $d_1 = 3^2 \cdot 5 \cdot 13 \cdot 1279$ |
| 18) | 17D | $2^3 \cdot \begin{cases} 13 \cdot 3079 \\ 19 \cdot 293 \end{cases}$ | $d_1 = 2^3 \cdot 13$ |
| 19) | 19D | $3^4 \cdot 11^2 \cdot 17 \cdot \begin{cases} 5 \cdot 373 \\ 13 \cdot 43 \end{cases}$ | $d_1 = 3^4 \cdot 11^2 \cdot 13 \cdot 17 \cdot 43$ |
| 20) | 20D | $2 \cdot 5 \cdot 13 \cdot \begin{cases} 29 \cdot 139 \\ 47 \cdot 2099 \end{cases}$ | $d_1 = 2 \cdot 5 \cdot 13 \cdot 47 \cdot 139$ |
| 21) | 21D | $3^4 \cdot 7^2 \cdot 11^2 \cdot \begin{cases} 17 \cdot 97 \\ 41 \cdot 1049 \end{cases}$ | $d_1 = 3^5 \cdot 7^2 \cdot 11^2 \cdot 1049$ |
| 22) | 22D | $3^3 \cdot 7^2 \cdot 13 \cdot \begin{cases} 11 \cdot 199 \\ 17 \cdot 1559 \end{cases}$ | $d_1 = 3^3 \cdot 7^2 \cdot 11 \cdot 13 \cdot 199$ |
| 23) | 22D | $2^6 \cdot \begin{cases} 107 \cdot 2099 \\ 179 \cdot 2939 \end{cases}$ | $d_1 = 2^5 \cdot 179 \cdot 2099$ |
| 24) | 24D | $2^5 \cdot 97 \cdot \begin{cases} 193 \cdot 2447 \\ 197 \cdot 2909 \end{cases}$ | $d_1 = 2^5 \cdot 97 \cdot 197 \cdot 2447$ |

25) 25D	$3^2 \cdot 5^3 \cdot 13^2 \cdot \begin{cases} 29 \cdot 193 \\ 103 \cdot 14549 \end{cases}$	$d_1 = 3^2 \cdot 5^3 \cdot 13^2 \cdot 103 \cdot 193$
26) 25D	$3^3 \cdot 5 \cdot 11 \cdot 43 \cdot \begin{cases} 101 \cdot 1031 \\ 251 \cdot 1289 \end{cases}$	$d_1 = 3^3 \cdot 5 \cdot 11 \cdot 43 \cdot 101 \cdot 1289$
27) 25D	$3^2 \cdot 5 \cdot 13 \cdot 41 \cdot \begin{cases} 19 \cdot 233 \\ 29 \cdot 2347 \end{cases}$	$d_1 = 3^2 \cdot 5 \cdot 13 \cdot 41^2 \cdot 233$
28) 26D	$3 \cdot 5 \cdot 7 \cdot 19 \cdot \begin{cases} 13 \cdot 509 \\ 577 \cdot 4787 \end{cases}$	$d_1 = 3 \cdot 5 \cdot 7 \cdot 19 \cdot 509$
29) 26D	$3^2 \cdot 7 \cdot 11 \cdot 13 \cdot 43 \cdot \begin{cases} 101 \cdot 1031 \\ 251 \cdot 1289 \end{cases}$	$d_1 = 3^2 \cdot 7 \cdot 11 \cdot 13 \cdot 43 \cdot 101 \cdot 1289$
30) 28D	$3^3 \cdot 5^4 \cdot 19 \cdot \begin{cases} 37 \cdot 239 \\ 47 \cdot 9803 \end{cases}$	$d_1 = 3^4 \cdot 5^5 \cdot 19 \cdot 47$
31) 28D	$2^6 \cdot \begin{cases} 79 \cdot 2699 \\ 367 \cdot 23819 \end{cases}$	$d_1 = 2^6 \cdot 367 \cdot 2699$
32) 30D	$3^4 \cdot 5 \cdot 11 \cdot \begin{cases} 23 \cdot 11699 \\ 419 \cdot 3167 \end{cases}$	$d_1 = 3^4 \cdot 5 \cdot 11 \cdot 23$
33) 30D	$2^2 \cdot 11 \cdot 79 \cdot \begin{cases} 17 \cdot 13903 \\ 89 \cdot 112507 \end{cases}$	$d_1 = 2^2 \cdot 11 \cdot 79 \cdot 13903$
34) 31D	$3^3 \cdot 5^3 \cdot 13 \cdot \begin{cases} 199 \cdot 4751 \\ 271 \cdot 336599 \end{cases}$	$d_1 = 3^3 \cdot 5^3 \cdot 13 \cdot 271$
35) 31D	$3^2 \cdot 5^2 \cdot 13 \cdot 31 \cdot \begin{cases} 199 \cdot 4751 \\ 271 \cdot 336599 \end{cases}$	$d_1 = 3^2 \cdot 5^2 \cdot 13 \cdot 31 \cdot 271$
36) 37D	$3^2 \cdot 5 \cdot 13 \cdot 23 \cdot \begin{cases} 19 \cdot 101429 \\ 461 \cdot 22271129 \end{cases}$	$d_1 = 3^2 \cdot 5 \cdot 13 \cdot 19 \cdot 23 \cdot 101429$
37) 41D	$3^2 \cdot 5 \cdot 11 \cdot 19 \cdot \begin{cases} 131 \cdot 5651377 \\ 509 \cdot 30557 \end{cases}$	$d_1 = 3^2 \cdot 5 \cdot 11 \cdot 19 \cdot 131$
type (3, 1)		
38) 32D	$3^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot \begin{cases} 71 \cdot 83 \cdot 2851 \\ 245519 \end{cases}$	$d_1 = 3^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot 71 \cdot 83$
39) 48D	$2 \cdot 5 \cdot 11^2 \cdot 241 \cdot \begin{cases} 127 \cdot 181 \cdot 70853 \\ 434571199 \end{cases}$	$d_1 = 2 \cdot 5 \cdot 11^2 \cdot 241$
type (3, 2)		
40) 17D	$2^3 \cdot \begin{cases} 13 \cdot 23 \cdot 109 \\ 167 \cdot 1231 \end{cases}$	$d_1 = 2^3 \cdot 23 \cdot 167$
41) 19D	$3^3 \cdot 5^2 \cdot \begin{cases} 11 \cdot 29 \cdot 167 \\ 23 \cdot 449 \end{cases}$	$d_1 = 3^3 \cdot 5^2 \cdot 11 \cdot 23 \cdot 29$
42) 24D	$2^2 \cdot 11 \cdot \begin{cases} 17 \cdot 439 \cdot 503 \\ 31 \cdot 11087 \end{cases}$	$d_1 = 2^2 \cdot 11 \cdot 439 \cdot 503$
43) 25D	$3^2 \cdot 5^3 \cdot 13 \cdot \begin{cases} 29 \cdot 59 \cdot 389 \\ 149 \cdot 8423 \end{cases}$	$d_1 = 3^2 \cdot 5^3 \cdot 13 \cdot 29 \cdot 389$
44) 35D	$3^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot \begin{cases} 97 \cdot 109 \cdot 569 \\ 197 \cdot 1667959 \end{cases}$	$d_1 = 3^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot 109 \cdot 569$
45) 40D	$3^5 \cdot 5 \cdot 13 \cdot \begin{cases} 17 \cdot 263 \cdot 67103 \\ 197 \cdot 122039807 \end{cases}$	$d_1 = 3^5 \cdot 5 \cdot 13 \cdot 122039807$

$$\begin{array}{ll}
46) 42D & 2^5 \cdot \begin{cases} 47 \cdot 2137 \cdot 6323 \\ 101 \cdot 642640039 \end{cases} & d_1 = 2^5 \cdot 2137 \cdot 6323 \\
47) 47D & 2 \cdot 5 \cdot 11 \cdot \begin{cases} 239 \cdot 1051 \cdot 2763119 \\ 31 \cdot 181324319039 \end{cases} & d_1 = 2 \cdot 5 \cdot 31 \cdot 239 \cdot 1051 \cdot 2763119 \\
48) 52D & 2^2 \cdot 11 \cdot 149 \cdot \begin{cases} 13 \cdot 1297 \cdot 15074627 \\ 167 \cdot 272306275957 \end{cases} & d_1 = 2^2 \cdot 11 \cdot 149 \cdot 272306275957 \\
\text{type (3, 3)} & & \\
49) 28D & 2^3 \cdot \begin{cases} 19 \cdot 23 \cdot 47 \\ 79 \cdot 251 \cdot 31727 \end{cases} & d_1 = 2^3 \cdot 23 \cdot 79 \cdot 251 \\
50) 45D & 2^4 \cdot \begin{cases} 23 \cdot 83 \cdot 227 \\ 191 \cdot 15791 \cdot 10112749 \end{cases} & d_1 = 2^4 \cdot 83
\end{array}$$

7. The two four-cycles 1), 13) were found by a different type of search: a_1, a_2 were chosen as so-called breeders for amicable number pairs, as studied in [4], [1]. These two new cycles are due to all three authors; the other ones are due to the first two authors. A complete list of all 110 known amicable four-cycles is provided by J. O. M. Pedersen on his internet amicable.adsl.dk pages

<http://amicable.adsl.dk/aliquot/sociable.txt>.

REFERENCES

1. S. Battiato and W. Borho, *Breeding amicable numbers in abundance II*, Math. Comp. **70**, (2001), 1329–1333. MR **2002b**:11011
2. M. Beeler, R. W. Gosper and R. Schroepel, *M.I.T. Artificial Intelligence Memo #239*, (29.02.72), Item 61.
3. W. Borho, *Über die Fixpunkte der k -fach iterierten Teilersummenfunktion*, Mitt. Math. Gesellsch. Hamburg 9, (1969), 34–48. MR **40**:7189
4. W. Borho and H. Hoffmann, *Breeding amicable numbers in abundance*, Math. Comp. **46**, (1986), 281–293. MR **87c**:11003
5. H. Cohen, *On amicable and sociable numbers*, Math. Comp. **24**, (1970), 423–429. MR **42**:5887
6. J.S. Devitt, R. K. Guy and J.L. Selfridge, *Third report on aliquot sequences*, Congr. Numer. XVIII, Proc. 6th Manitoba Conf. Numerical Math., (1976), 177–204. MR **80d**:10001
7. A. Flammenkamp, *New sociable numbers*, Math. Comp. **56**, (1991), 871–873. MR **92a**:11004
8. D. Moews and P. C. Moews, *A search for aliquot cycles below 10^{10}* , Math. Comp. **57**, (1991), 849–855. MR **92e**:11151
9. D. Moews and P. C. Moews, *A search for aliquot cycles and amicable pairs*, Math. Comp. **61**, (1993), 935–938. MR **94a**:11198
10. Ren Yuanhua, *On constructing sociable numbers*, (1990), preprint.

KARSTEN BLANKENAGEL, BUGH FB 7, GAUSSSTRASSE 20, 42097 WUPPERTAL, GERMANY
E-mail address: kblanken@math.uni-wuppertal.de

WALTER BORHO, BUGH FB 7, GAUSSSTRASSE 20, 42097 WUPPERTAL, GERMANY
E-mail address: Walter.Borho@math.uni-wuppertal.de

AXEL VOM STEIN, BUGH FB 7, GAUSSSTRASSE 20, 42097 WUPPERTAL, GERMANY
E-mail address: vomstein@uni-wuppertal.de