

DISCRETE FREDHOLM PROPERTIES AND CONVERGENCE ESTIMATES FOR THE ELECTRIC FIELD INTEGRAL EQUATION

SNORRE H. CHRISTIANSEN

ABSTRACT. The Galerkin discretization of the Electric Field Integral Equation is reinvestigated. We prove quasi-optimal convergence estimates at nonresonant frequencies, using orthogonal splittings of the Galerkin space. At resonant frequencies we show that the spurious electric currents radiate only weakly in the exterior domain. This is achieved through the study of some finitely degenerated problems in terms of LBB Inf-Sup estimates and the use of discrete Helmholtz decompositions.

INTRODUCTION

The Electric Field Integral Equation (EFIE) is defined in Section 2.1. It is the basic equation in the theory of integral methods for the scattering of electromagnetic harmonic (i.e., mono-frequential) waves. The discretization of this equation is a widespread method for the computation of radar cross sections, antenna performance and electromagnetic compatibility issues, to mention just a few important fields of application.

At nonresonant frequencies we give a new and rather general proof of the Inf-Sup condition for the Galerkin discretization of the EFIE on a large class of Galerkin spaces. The challenge is that the operator is not a compact perturbation of a coercive form. To compensate for this, we use splittings of the Galerkin spaces into orthogonal sums, where one of the two subspaces is the kernel of the divergence operator. We prove that the discrete Inf-Sup condition holds whenever the *gap* from the other subspace to its continuous analogue (the orthogonal of the kernel of the divergence operator) tends to zero. In turn this condition is shown to hold whenever discrete Helmholtz decompositions of tangent fields (analogues of the decompositions $u = \text{grad } p + v$ with $\text{div } v = 0$) are sufficiently well behaved, which is checked for the standard spaces. We deduce quasi-optimal convergence rates in natural norms, for the approximate solutions in standard surface Finite Element spaces.

At resonant frequencies the integral operator appearing in the EFIE is degenerate. However the operator is of Fredholm type, and for exterior problems the right-hand side is in general compatible—that is, the EFIE has a solution. We address the question as to what happens for the discretized equation. When the right-hand

Received by the editor December 26, 2000 and, in revised form, April 10, 2002.
2000 *Mathematics Subject Classification.* Primary 65N12, 65N38, 78M15.
This work received financial support from Thales Airborne Systems.

side is compatible, it is a straightforward remark that there exist bounded families (indexed by the mesh refinement parameter h) of approximate solutions to the discrete equations, for instance the best approximations in any chosen norm. We prove a converse property, namely that for any bounded family of approximate solutions the *exterior* electromagnetic field is well approximated. Since by the above remark it suffices to consider only the homogeneous equation, an interpretation of this result is that *discrete* spurious currents radiate much less in the exterior domain than their norm would indicate: they are close to the kernel of the *continuous* EFIE operator.

Other integral equations are currently used to solve electromagnetic scattering problems. Some of these, such as the Magnetic Field Integral Equation (MFIE) are also degenerate only for a discrete set of frequencies. However at these, the right-hand side is usually not compatible. Linear combinations of the EFIE and MFIE known as CFIEs avoid the existence of resonances at the expense of introducing a somewhat arbitrary (complex) parameter. This is also known as the Brakhage and Werner trick. The EFIE has the advantage of having an unknown with a physical interpretation: it is the electric current density on the surface and has been perceived to be more robust on singular scatterers; in particular CFIEs cannot treat open surfaces (screens). However a more elaborate comparison is beyond the goals of this paper.

One of the main motivations for constructing the present theory was to provide a rigorous justification for the use of a new nonlocal finite element space, which appeared naturally in a preconditioning technique for the EFIE described in Christiansen-Nédélec [16]. We therefore remain sufficiently general throughout the paper to provide a theory that applies to such nonstandard spaces.

To achieve our goals the three main tools used are

- a reformulation of the EFIE as an equivalent saddle-point problem, following a technique used by Nédélec [29] to prove the Fredholm property of the (continuous) EFIE. It enables us to argue in terms of compact perturbations of coercive forms.
- estimates on discrete Helmholtz decompositions closely related to a discrete compactness result due to Kikuchi [24]. They have recently been found to be important in the discretization of eigenvalue problems by saddle-point formulations.
- a general theory of the discretization of finitely degenerated problems, which we have tried to develop in the spirit of Babuska's Inf-Sup conditions [2].

The most well-known Galerkin space is the one introduced for the EFIE by Rao-Wilton-Glisson in [32]. This space of tangent fields on Γ had previously been used by Nédélec [27] for the computation of eddy currents and, as noted by him, is an adaptation to arbitrary surfaces of the lowest degree Raviart-Thomas (RT) finite element (FE) space developed for planar problems [33]. A numerical analysis of the Galerkin discretization of the EFIE by RT FE of any order was first presented by Bendali [5]. We improve the results of Bendali by getting rid of a parasitic factor of the form $Ch^{-1/2-\epsilon}$ (for arbitrary $\epsilon > 0$, C depending on ϵ) in the error estimates. It should be remarked that we do not need to compute any residual error in the sense of Hsiao-Kleinman [21] to assess the quality of the solution.

The paper is organized as follows:

- Section 1: We develop a theory of discrete Fredholm properties in an abstract setting.
- Section 2: We recall the main results we need on the integral representations of electromagnetic fields and the related Electric Field Integral Equation. In particular we introduce the appropriate splitting of the solution space.
- Section 3: We turn to the discretization and prove the Inf-Sup condition for the EFIE and the other announced results using the theory of the first two sections.

1. DISCRETE FREDHOLM PROPERTIES

The EFIE is an example of an equation $Au = v$ involving a Fredholm operator A and a right-hand side v which in many cases of practical interest is compatible. In this section we develop a tool for studying the discretization of Fredholm operators by Galerkin methods. First we recall the definition of a left semi-Fredholm (LSF) operator and Babuska's Inf-Sup conditions for the solvability of equations involving invertible operators. Then we propose a definition for a discrete LSF condition. It is a generalisation of Babuska's Inf-Sup condition such that

- if an injective operator is discrete LSF (on a given family of Galerkin spaces), then it satisfies a standard (one-sided) Babuska Inf-Sup condition (Corollary 1.11);
- if an operator is discrete LSF, then so is any compact perturbation of it (Corollary 1.13).

The combination of these two properties is practical for proving Inf-Sup conditions for injective operators since, in the course of such a proof, many compact perturbations are sometimes involved, but injectivity is not guaranteed in the intermediate steps. In this paper, one compact perturbation appears once appropriate splittings are considered for the EFIE (Theorem 2.6); others are due to the relative compactness of the frequency part $G_k - G_0$ of the Green kernel.

1.1. Results on left semi-Fredholm operators. We collect some well-known facts that will be useful later and that will serve as references for discrete counterparts. The classical reference on the subject is Kato [23].

Theorem 1.1. *Let X and Y be two Banach spaces and $A : X \rightarrow Y$ be linear and continuous. The following conditions are equivalent:*

- (1) A has closed range and finite dimensional kernel;
- (2) $\dim \ker A < \infty$ and on any closed supplementary M of $\ker A$

$$(1.1) \quad \inf_{u \in M} \|Au\|/\|u\| > 0;$$

- (3) there is a closed subspace M of X with finite codimension such that

$$(1.2) \quad \inf_{u \in M} \|Au\|/\|u\| > 0.$$

Definition 1.2. Let X and Y be two Banach spaces and $A : X \rightarrow Y$ be linear and continuous. We say that A is *left semi-Fredholm* (LSF) if the stated conditions are satisfied. The set of LSF operators from X to Y is denoted $\text{LSF}(X, Y)$.

Theorem 1.3. *Let X and Y be two Banach spaces. For all $n \in \mathbb{N}$ the following sets (whose union is $\text{LSF}(X, Y)$) are open in $L(X, Y)$:*

$$(1.3) \quad \{A \in L(X, Y) : A \text{ has closed range and } \dim \ker A \leq n\}.$$

Moreover $\text{LSF}(X, Y)$ is stable under translation by compact operators.

1.2. Galerkin methods in the presence of Gårding inequalities. Let X be a Hilbert space. The base field \mathbb{K} of X is either \mathbb{R} or \mathbb{C} . Let $a : X \times X \rightarrow \mathbb{K}$ be a \mathbb{K} -bilinear continuous form on X . We say that a is nondegenerate if the induced map $\mathcal{A} : X \rightarrow X^*$, $u \mapsto a(u, \cdot)$ is an isomorphism (since X is reflexive, this is equivalent to $u \mapsto a(\cdot, u)$ being one). When a is nondegenerate, we construct approximations of \mathcal{A}^{-1} by choosing a family of closed subspaces (X_h) of X , which is *approximating* in the sense that

$$(1.4) \quad \forall u \in X \quad \lim_h \inf_{u' \in X_h} \|u - u'\| = 0.$$

Then for a given $l \in X^*$ we seek solutions of

$$(1.5) \quad u \in X_h \quad \forall u' \in X_h \quad a(u, u') = l(u').$$

In accordance with usual conventions it is implicit that $(X_h) = (X_h)_{h \in H}$, where H is some subset of \mathbb{R}_+^* accumulating at 0, and that \lim_h means $\lim_{h \rightarrow 0}$.

Let \mathcal{A}_h be the map: $X_h \rightarrow X_h^*$, $u \mapsto a(u, \cdot)$. Then equation (1.5) can be restated as $u \in X_h \quad \mathcal{A}_h u = l|_{X_h}$. If \mathcal{A}_h is invertible and if

$$(1.6) \quad \exists C > 0 \quad \forall l \in X^* \quad \forall h \quad \|\mathcal{A}^{-1}l - \mathcal{A}_h^{-1}l|_{X_h}\| \leq C \inf_{u' \in X_h} \|\mathcal{A}^{-1}l - u'\|,$$

we say that the Galerkin method yields *quasi-optimal convergence*.

A sufficient condition for a to be nondegenerate is that it is coercive in the sense that for some \mathbb{R} -linear isometric involution of X , denoted $u \mapsto \bar{u}$ and called conjugation, we have

$$(1.7) \quad \exists C > 0 \quad \forall u \in X \quad \Re a(u, \bar{u}) \geq (1/C)\|u\|^2.$$

This is the Lax-Milgram Theorem. If a is coercive, Céa's Lemma asserts that if the spaces X_h are stable under conjugation (i.e., $u \mapsto \bar{u}$ maps X_h into X_h), the Galerkin method yields quasi-optimal convergence.

In the context of boundary integral equations one soon encounters bilinear forms satisfying only a weaker estimate, known as a Gårding inequality:

$$(1.8) \quad \exists C > 0 \quad \forall u \in X \quad \Re a(u, \bar{u}) \geq (1/C)\|u\|_X^2 - C\|u\|_Y^2,$$

where Y is a Hilbert space containing X and such that the canonical injection $X \rightarrow Y$ is compact.

It was soon recognized that if a is nondegenerate and satisfies a Gårding inequality, and if the spaces X_h are stable under conjugation, then there exists $h_0 > 0$ such that the Galerkin method has quasi-optimal convergence on $(X_h)_{h < h_0}$. Early contributions in this direction include Schatz [36]. This fact is naturally stated in terms of the Inf-Sup conditions due to Babuska [2], [3]. For an insightful account on this we refer to Demkowicz [19], who attributes the result to Mikhlin; see also Wendland [40]. For future reference we recall Babuska's Theorem:

Theorem 1.4. *Let X and Y be two reflexive Banach spaces. Let a be a continuous bilinear form on $X \times Y$. If*

$$(1.9) \quad \inf_{u \in X} \sup_{v \in Y} \frac{|a(u, v)|}{\|u\| \|v\|} \geq \alpha > 0,$$

$$(1.10) \quad \forall v \in Y \quad (\forall u \in X \ a(u, v) = 0) \Rightarrow (v = 0),$$

then

$$(1.11) \quad \inf_{v \in Y} \sup_{u \in X} \frac{|a(u, v)|}{\|v\| \|u\|} = \inf_{u \in X} \sup_{v \in Y} \frac{|a(u, v)|}{\|u\| \|v\|},$$

and for all $l \in Y^*$ there is a unique $u \in X$ such that

$$(1.12) \quad \forall v \in Y \quad a(u, v) = l(v).$$

It satisfies $\|u\| \leq \alpha^{-1} \|l\|$.

We will use the following notation whenever it makes sense:

$$(1.13) \quad \sup_Y \mathcal{A} = \sup_{v \in Y} \frac{\|\mathcal{A}v\|}{\|v\|}, \quad \inf_X \sup_Y \mathcal{A} = \inf_X \sup_Y a = \inf_{u \in X} \sup_{v \in Y} \frac{|a(u, v)|}{\|u\| \|v\|},$$

where it is implicit that u and v are nonzero.

1.3. Discrete left semi-Fredholm operators. The following lemma is trivial and stated just for the record.

Lemma 1.5. *Let X and Y be two Banach spaces, and let $\mathcal{A} : X \rightarrow Y^*$ be continuous. Let M be a closed subspace of X . Let (X_h) and (Y_h) be families of closed subspaces of X and Y . Suppose (X_h) is approximating (and $Y_h \neq \{0\}$). If*

$$(1.14) \quad \liminf_h \inf_{X_h \cap M} \sup_{Y_h} \mathcal{A} > 0,$$

then $M \cap \ker \mathcal{A} = \{0\}$. If in addition M has finite codimension, then \mathcal{A} is LSF.

We will need two more lemmas, which are less trivial.

Lemma 1.6. *Let X be a Banach space. Let M and N be closed subspaces of X such that $M \oplus N = X$ and N is finite dimensional. Let $P : X \rightarrow X$ be the projector with range M and kernel N . Let (X_h) be a family of closed subspaces of X which is approximating. Then*

- there are projectors P_h with range M which converge in norm to P and which leave X_h stable for sufficiently small h ;
- if (P_h) is any family of such projectors, then for sufficiently small h , $\ker P_h \subset X_h$ and $X_h = (X_h \cap M) \oplus \ker P_h$.

Proof. Choose a basis (e^i) of N , and pick elements e_h^i of X_h such that

$$(1.15) \quad \lim_h \|e^i - e_h^i\| = 0.$$

For any h , let N_h be the space spanned by the e_h^i . Then there is $h_0 > 0$ such that for $h < h_0$ we have $X = M \oplus N_h$.

(1) For $h < h_0$ let P_h be the projector with range M and kernel N_h , and for $h \geq h_0$ put $P_h = P$. One checks that the family (P_h) has the desired properties.

(2) If (P_h) is a family of projectors with these properties, then $P_h|_{X_h}$ are projectors with range $X_h \cap M$, so for sufficiently small h

$$(1.16) \quad \dim N_h \leq \dim \ker P_h|_{X_h} \leq \dim \ker P_h \leq \dim N.$$

For sufficiently small h these dimensions are all equal. \square

Lemma 1.7. *Let X be a Banach space and Y be a reflexive Banach space. Let $\mathcal{A} : X \rightarrow Y^*$ be continuous and let it have closed range. Then for any $l \in X^*$ that vanishes on $\ker \mathcal{A}$ there is a $v \in Y$ such that*

$$(1.17) \quad \forall u \in X \quad l(u) = (\mathcal{A}u)(v).$$

Proof. Remark first that \mathcal{A} induces an isomorphism of Banach spaces $X/\ker \mathcal{A} \rightarrow \mathcal{A}(X)$. Let \mathcal{Z} be the inverse mapping. For convenience we put $N = \ker \mathcal{A}$.

Pick $l \in X^*$ that vanishes on N . Let \tilde{l} be the canonical image of l in $(X/N)^*$. Put $f = \mathcal{Z}^* \tilde{l} \in (\mathcal{A}(X))^*$. Extend f to a continuous linear form on Y^* , and put $v = (i_Y)^{-1} f$. For all $u \in X$ we have

$$(1.18) \quad (\mathcal{A}u)(v) = f(\mathcal{A}u) = \tilde{l}(\mathcal{Z}\mathcal{A}u) = \tilde{l}(\tilde{u}),$$

where \tilde{u} is the canonical image of u in X/N . The proof is complete. \square

Theorem 1.8. *Let X and Y be two reflexive Banach spaces, and $\mathcal{A} : X \rightarrow Y^*$ be LSF. Let (X_h) and (Y_h) be families of closed subspaces of X and Y which are approximating.*

Put $N_0 = \ker \mathcal{A}$. Suppose we have closed subspaces M_0 and M_1 such that $M_0 \oplus N_0 = X$, $M_1 \oplus N_1 = M_0$ and $\dim N_1 < \infty$. If

$$(1.19) \quad \liminf_h \inf_{X_h \cap M_1} \sup_{Y_h} \mathcal{A} > 0,$$

then

$$(1.20) \quad \liminf_h \inf_{X_h \cap M_0} \sup_{Y_h} \mathcal{A} > 0.$$

Proof. If not, let $(u_n), (h_n)$ be sequences such that

$$(1.21) \quad u_n \in X_{h_n} \cap M_0, \quad \|u_n\| = 1, \quad \lim_n h_n = 0, \quad \limsup_n \sup_{Y_{h_n}} \mathcal{A}u_n = 0.$$

Let P be the projector with range M_1 and kernel $N_1 \oplus N_0$. The sequence $(u_n - Pu_n)$ is a bounded sequence in the finite dimensional space $N_1 \oplus N_0$, so modulo extraction we can suppose that it converges in norm to some $u^N \in N_1 \oplus N_0$. Since $u_n \in M_0$, we actually have $u^N \in N_1$.

Since (Y_{h_n}) is approximating, we have

$$(1.22) \quad \forall v \in Y \quad \lim_n (\mathcal{A}u_n)(v) = 0.$$

Therefore, by Lemma 1.7, for any $l \in X^*$ that vanishes on N_0 we have

$$(1.23) \quad \lim_n l(u_n) = 0.$$

If $l \in X^*$ vanishes on $M_1 \oplus N_0$, we have

$$(1.24) \quad l(u^N) = \lim_n l(u_n - Pu_n),$$

and $Pu_n \in M_1$ so this limit must be 0. Therefore, by the Hahn-Banach theorems, $u^N \in M_1 \oplus N_0$, so $u^N = 0$.

Let $P_h : X \rightarrow X$ be projectors with range M_1 , leaving X_h stable and converging in norm to P , as in Lemma 1.6. Notice that

$$(1.25) \quad \lim_h \|u_n - P_{h_n} u_n\| = 0.$$

One obtains a contradiction using the discrete Inf-Sup estimate on $(X_h \cap M_1) \times Y_h$ for the sequence $(P_{h_n} u_n)$. \square

Theorem 1.9. *Let X and Y be two reflexive Banach spaces and $\mathcal{A} : X \rightarrow Y^*$ be LSF. Let (X_h) and (Y_h) be families of closed subspaces of X and Y which are approximating.*

Let N denote the kernel of \mathcal{A} and let M_1 and M_2 be two supplementaries of N in X . If

$$(1.26) \quad \liminf_h \inf_{X_h \cap M_1} \sup_{Y_h} \mathcal{A} > 0,$$

then

$$(1.27) \quad \liminf_h \inf_{X_h \cap M_2} \sup_{Y_h} \mathcal{A} > 0.$$

Proof. Choose $\alpha > 0$ and h_0 such that for all $h < h_0$

$$(1.28) \quad \inf_{X_h \cap M_1} \sup_{Y_h} \mathcal{A} \geq \alpha.$$

Let P denote the projection with range M_1 and kernel N . Since P induces a continuous bijection $M_2 \rightarrow M_1$, by the standard theorems there is $\beta > 0$ such that

$$(1.29) \quad \forall u \in M_2 \quad \|Pu\| \geq \beta\|u\|.$$

Moreover let $P_h : X \rightarrow X$ be projectors with range M_1 , leaving X_h stable and converging in norm to P , as in Lemma 1.6.

For all $u \in X_h \cap M_2$ and all $v \in Y_h$ we have

$$(1.30) \quad |(\mathcal{A}u)(v)| = |(\mathcal{A}Pu)(v)|$$

$$(1.31) \quad \geq |(\mathcal{A}P_h u)(v)| - |(\mathcal{A}(P - P_h)u)(v)|.$$

Hence for $h < h_0$

$$(1.32) \quad \sup_{Y_h} \mathcal{A}u \geq \alpha\|P_h u\| - \|\mathcal{A}\|\|P - P_h\|\|u\|.$$

Moreover

$$(1.33) \quad \|P_h u\| \geq \|Pu\| - \|(P - P_h)u\| \geq \beta\|u\| - \|P - P_h\|\|u\|,$$

so

$$(1.34) \quad \sup_{Y_h} \mathcal{A}u \geq \alpha\beta\|u\| - (\|\mathcal{A}\| + \alpha)\|P - P_h\|\|u\|.$$

It follows that

$$(1.35) \quad \inf_{X_h \cap M_2} \sup_{Y_h} \mathcal{A} \geq \alpha\beta - (\|\mathcal{A}\| + \alpha)\|P - P_h\|.$$

□

Definition 1.10. Let X and Y be two reflexive Banach spaces, and let $\mathcal{A} : X \rightarrow Y^*$ be continuous. Let (X_h) and (Y_h) be families of closed subspaces of X and Y . We say that \mathcal{A} is *discrete LSF* on $(X_h \times Y_h)$ if there is a closed subspace M of X , with finite codimension in X such that

$$(1.36) \quad \liminf_h \inf_{X_h \cap M} \sup_{Y_h} \mathcal{A} > 0.$$

In this situation, if we wish to be more precise, we say that \mathcal{A} is discrete LSF on $(X_h \times Y_h)$ with respect to (w.r.t.) M .

Corollary 1.11. *With the hypotheses of Theorem 1.8, if \mathcal{A} is discrete LSF on $(X_h \times Y_h)$ w.r.t. M , then it is so w.r.t. any closed subspace whose intersection with $\ker \mathcal{A}$ is $\{0\}$. In particular if in addition \mathcal{A} is injective, then \mathcal{A} satisfies a uniform discrete Inf-Sup condition on $(X_h \times Y_h)$ in the sense of Babuska for small enough parameters h .*

1.4. Compact perturbations of discrete LSF operators.

Theorem 1.12. *Let X and Y be two reflexive Banach spaces, and let $\mathcal{A} : X \rightarrow Y^*$ be LSF. Let $\mathcal{B} : X \rightarrow Y^*$ be compact. Let (X_h) and (Y_h) be families of closed subspaces of X and Y . Suppose (Y_h) is approximating.*

Suppose furthermore that M_0 is a closed subspace of X that satisfies

$$(1.37) \quad \liminf_h \inf_{X_h \cap M_0} \sup_{Y_h} \mathcal{A} > 0.$$

Let N_1 be the kernel of $\mathcal{A} + \mathcal{B}$ restricted to M_0 , and suppose it has a closed supplementary M_1 in M_0 (i.e., $M_0 = M_1 \oplus N_1$). Then

$$(1.38) \quad \liminf_h \inf_{X_h \cap M_1} \sup_{Y_h} \mathcal{A} + \mathcal{B} > 0.$$

Proof. If not, let $(u_n), (h_n)$ be sequences such that

$$(1.39) \quad u_n \in X_{h_n} \cap M_1, \quad \|u_n\| = 1, \quad \lim_n h_n = 0, \quad \limsup_n \sup_{Y_{h_n}} (\mathcal{A} + \mathcal{B})u_n = 0.$$

Since (Y_{h_n}) is approximating, we have

$$(1.40) \quad \forall v \in Y \quad \lim_n ((\mathcal{A} + \mathcal{B})u_n)(v) = 0.$$

Therefore, for any continuous linear form $l \in X^*$ that vanishes on $\ker(\mathcal{A} + \mathcal{B})$ we have

$$(1.41) \quad \lim_n l(u_n) = 0.$$

Any continuous linear form on M_1 has a continuous extension to X that vanishes on $\ker(\mathcal{A} + \mathcal{B})$; therefore (u_n) converges weakly to 0 in M_1 , hence also in X . Since \mathcal{B} is compact, $(\mathcal{B}u_n)$ converges in norm to 0. One obtains a contradiction using the discrete Inf-Sup estimate for \mathcal{A} on $((X_h \cap M_0) \times Y_h)$. \square

Corollary 1.13. *With the hypotheses of the above Theorem 1.12, if \mathcal{A} is discrete LSF on $(X_h \times Y_h)$, then so is $\mathcal{A} + \mathcal{B}$.*

Example 1.14. If X is a Hilbert space equipped with a conjugation and $\mathcal{A} : X \rightarrow X^*$ is continuous and satisfies a Gårding inequality, then \mathcal{A} is discrete LSF on any approximating family of conjugation-stable and closed subspaces.

1.5. Error control of discrete LSF operators. The following property of discrete LSF operators will be important to us:

Theorem 1.15. *Let X and Y be two reflexive Banach spaces, and let $\mathcal{A} : X \rightarrow Y^*$ be LSF. Let (X_h) and (Y_h) be approximating families of closed subspaces of X and Y . If \mathcal{A} is discrete LSF on $(X_h \times Y_h)$, then there is $C > 0$, such that for all $\epsilon > 0$ there is $h_\epsilon > 0$ such that for all $h < h_\epsilon$ and all $u \in X_h$*

$$(1.42) \quad \sup_Y \mathcal{A}u \leq C \sup_{Y_h} \mathcal{A}u + \epsilon \|u\|.$$

Proof. Let N be the kernel of \mathcal{A} on X and let M be a closed supplementary of N in X such that \mathcal{A} is discrete LSF w.r.t. M . Let P be the projector with range M and kernel N , and let P_h be projectors as in Lemma 1.6. Pick $\alpha > 0$ and h_0 such that for all $h < h_0$, X_h is stable under P_h and

$$(1.43) \quad \inf_{X_h \cap M} \sup_{Y_h} \mathcal{A} \geq \alpha.$$

Note first that for all $u \in X$,

$$(1.44) \quad \mathcal{A}u = \mathcal{A}Pu.$$

In particular,

$$(1.45) \quad \sup_Y \mathcal{A}u = \sup_Y \mathcal{A}Pu \leq \|\mathcal{A}\| \|Pu\|.$$

Moreover,

$$(1.46) \quad \|Pu\| \leq \|P_h u\| + \|P - P_h\| \|u\|,$$

and if $u \in X_h$ and $h < h_0$,

$$(1.47) \quad \|P_h u\| \leq \alpha^{-1} \sup_{Y_h} \mathcal{A}P_h u,$$

and

$$(1.48) \quad \sup_{Y_h} \mathcal{A}P_h u \leq \sup_{Y_h} \mathcal{A}u + \|\mathcal{A}\| \|P - P_h\| \|u\|.$$

All in all, if $u \in X_h$ and $h < h_0$,

$$(1.49) \quad \sup_Y \mathcal{A}u \leq \alpha^{-1} \|\mathcal{A}\| \sup_{Y_h} \mathcal{A}u + \|\mathcal{A}\| (1 + \alpha^{-1} \|\mathcal{A}\|) \|P - P_h\| \|u\|.$$

The theorem follows. □

Remark 1.16. Equation (1.49) together with the construction of projectors P_h shown in point (1) of the proof of Lemma 1.6 yield rather explicit bounds on the constant C and the threshold h_ϵ appearing in the theorem.

In particular under the hypotheses of the theorem, \mathcal{A} has the following property:

$$(1.50) \quad \forall \epsilon > 0 \exists \delta > 0 \exists h_0 > 0 \forall h < h_0 \forall u \in X_h \\ (\sup_{Y_h} \mathcal{A}u \leq \delta \|u\|) \Rightarrow (\sup_Y \mathcal{A}u \leq \epsilon \|u\|).$$

In other words, in a sense, small discrete residual error indicates small continuous residual error. This property is easily extended to hold uniformly for some families of operators, as follows:

Corollary 1.17. *Let X and Y be two reflexive Banach spaces, and let \mathfrak{A} be a compact subset of $\text{LSF}(X, Y^*)$ (in the norm topology). Let (X_h) and (Y_h) be approximating families of closed subspaces of X and Y . If all \mathcal{A} in \mathfrak{A} are discrete LSF on $(X_h \times Y_h)$, then*

$$(1.51) \quad \forall \epsilon > 0 \exists \delta > 0 \exists h_0 > 0 \forall h < h_0 \forall u \in X_h \forall \mathcal{A} \in \mathfrak{A} \\ (\sup_{Y_h} \mathcal{A}u \leq \delta \|u\|) \Rightarrow (\sup_Y \mathcal{A}u \leq \epsilon \|u\|).$$

Proof. Choose a $\epsilon > 0$. For each $\mathcal{A} \in \mathfrak{A}$ choose a $\delta(\mathcal{A})$ and a $h_0(\mathcal{A})$ so that (1.50) holds for \mathcal{A} relative to $\epsilon/2$.

The family of balls with center \mathcal{A} and radius $\min\{\delta(\mathcal{A}), \epsilon\}/2$, for $\mathcal{A} \in \mathfrak{A}$, covers \mathfrak{A} , so we can extract a finite subcover indexed by, say, $\mathcal{A}_1, \dots, \mathcal{A}_n$.

Put $\delta = \min_i \delta(\mathcal{A}_i)/2$ and $h_0 = \min_i h_0(\mathcal{A}_i)$. Take $\mathcal{A} \in \mathfrak{A}$ and choose i so that $\|\mathcal{A} - \mathcal{A}_i\|$ is less than both $\delta(\mathcal{A}_i)/2$ and $\epsilon/2$.

For any $h < h_0$ and any $u \in X_h$ we have: If $\sup_{Y_h} \mathcal{A}u \leq \delta\|u\|$, then $\sup_{Y_h} \mathcal{A}u \leq (\delta(\mathcal{A}_i)/2)\|u\|$, so $\sup_{Y_h} \mathcal{A}_i u \leq \delta(\mathcal{A}_i)\|u\|$. This implies $\sup_Y \mathcal{A}_i u \leq (\epsilon/2)\|u\|$, and hence $\sup_Y \mathcal{A}u \leq \epsilon\|u\|$. \square

1.6. Extensions to quasi-conforming Galerkin approximations.

Definition 1.18. Let X be a Banach space and X_0 a closed subspace. Let (X_h) be a family of closed subspaces of X . We say that

(1) (X_h) is *approximating* in X_0 if

$$(1.52) \quad \forall u \in X_0 \quad \lim_h \inf_{u' \in X_h} \|u - u'\| = 0;$$

(2) (X_h) is *quasi-conforming* in X_0 if

$$(1.53) \quad \lim_h \sup_{u \in X_h} \inf_{u' \in X_0} \|u - u'\|/\|u\| = 0.$$

The quantity

$$(1.54) \quad \delta(X_h, X_0) = \sup_{u \in X_h} \inf_{u' \in X_0} \|u - u'\|/\|u\|$$

is called the *gap* from X_h to X_0 (see Kato [23]).

Suppose X_0 has a closed supplementary X_1 in X (one says that X_0 *splits*), and let $P : X \rightarrow X$ be the projection with range X_0 and kernel X_1 . For all $u \in X$ one has

$$(1.55) \quad \forall u' \in X_0 \quad \|u - Pu\| = \|(u - u') - (Pu - u')\| = \|(u - u') - P(u - u')\|,$$

hence

$$(1.56) \quad \left| \|u\| - \|Pu\| \right| \leq \|u - Pu\| \leq (1 + \|P\|)\|u\| \inf_{u' \in X_0} \|u - u'\|/\|u\|.$$

In particular if (X_h) is a family of closed subspaces which is quasi-conforming in X_0 , then for sufficiently small h the spaces PX_h are closed in X_0 and P induces isomorphisms $X_h \rightarrow PX_h$ which are arbitrarily close in norm to the identity mapping on X_h . Also if in addition (X_h) is approximating in X_0 , then so is (PX_h) . From this we deduce the following lemma.

Lemma 1.19. *Let X and Y be Banach spaces and let $\mathcal{A} : X \rightarrow Y^*$ be continuous. Let X_0 and Y_0 be closed subspaces of X and Y that split—yielding projectors P_X in X and P_Y in Y as above—and $\mathcal{A}_0 : X_0 \rightarrow Y_0^*$ be the map induced by \mathcal{A} . Let (X_h) and (Y_h) be families of closed subspaces of X and Y , which are quasi-conforming in X_0 and Y_0 (i.e., the corresponding gaps tend to 0). For any closed M_0 in X_0 , one has*

$$(1.57) \quad \liminf_h \inf_{M_0 \cap PX_h} \sup_{P_Y Y_h} \mathcal{A}_0 = \liminf_h \inf_{(M_0 \oplus \ker P_X) \cap X_h} \sup_{Y_h} \mathcal{A}.$$

With this lemma most results from the preceding sections carry over to the quasi-conforming setting in some form or another. However—as will be shown in the next section—the lemma is sufficient for our needs, so we will not develop this possibility here.

2. THE ELECTRIC FIELD INTEGRAL EQUATION

2.1. Integral representation of interior and exterior waves. Let Ω_- be an open bounded subset of \mathbb{R}^3 , and let Γ be its boundary. We suppose that $\overline{\Omega_-} = \Omega_- \cup \Gamma$ is a C^∞ smooth submanifold with boundary, though this hypothesis could be relaxed to C^p smoothness for some $p < \infty$ without much extra effort. More interesting would have been an extension of the theory to Lipschitz manifolds, but even though scalar equations are well understood (see Costabel [18]), the case of Maxwell's equations is still an active research direction (see Buffa et al. [13]) which we will not pursue here. We denote by Ω_+ the complement of $\Omega_- \cup \Gamma$ and by n the outward pointing normal on Γ . We suppose throughout that Ω_+ is *connected*, which ensures uniqueness of solutions to exterior problems.

The free-space harmonic Maxwell equations for the electromagnetic field (E, H) in an open region $\Omega \subset \mathbb{R}^3$ are

$$(2.1) \quad \operatorname{curl} E = +i\omega\mu H,$$

$$(2.2) \quad \operatorname{curl} H = -i\omega\epsilon E,$$

where $\mu > 0$ is the magnetic permeability and $\epsilon > 0$ is the electric permittivity. If the pulsation ω is zero, one also has to add explicitly that the fields are divergence free. In the sequel these equations will be referred to as Maxwell's equations. It is convenient to introduce the *wavenumber* k and the impedance Z defined by

$$(2.3) \quad k = \omega(\mu\epsilon)^{1/2},$$

$$(2.4) \quad Z = (\mu/\epsilon)^{1/2}.$$

Then we have $+i\omega\mu = +ikZ$ and $-i\omega\epsilon = -ik/Z$.

We will consider here only nonhomogeneous boundary value problems for the free-space harmonic Maxwell equations, with *real nonzero positive* pulsations. Referring to Colton-Kress [17], Cessenat [14] and Nédélec [29] for proofs, we briefly state the main results which will be of interest to us.

For any open domain Ω in \mathbb{R}^3 we use the notation

$$(2.5) \quad H_{\operatorname{curl}}^0(\Omega) = \{u \in L_T^2(\Omega) : \operatorname{curl} u \in L_T^2(\Omega)\},$$

where $L_T^2(\Omega)$ denotes the space of square summable tangential fields. On Γ the usual Sobolev spaces of scalar and tangential fields of regularity order $s \in \mathbb{R}$ are denoted $H^s(\Gamma)$ and $H_T^s(\Gamma)$, respectively, and the corresponding norms are both written

$$(2.6) \quad u \mapsto |u|_s.$$

On Γ we denote by div the surface divergence and introduce the Hilbert spaces $H_{\operatorname{div}}^s(\Gamma)$ of tangent fields on Γ

$$(2.7) \quad H_{\operatorname{div}}^s(\Gamma) = \{u \in H_T^s(\Gamma) : \operatorname{div} u \in H^s(\Gamma)\}.$$

They are equipped with the norms

$$(2.8) \quad u \mapsto \|u\|_s : \|u\|_s^2 = |u|_s^2 + |\operatorname{div} u|_s^2.$$

The surface rotational and the spaces $H_{\text{rot}}^s(\Gamma)$ are defined in a similar way but we do not introduce any notation for the corresponding norm. Notice that $u \mapsto u \times n$ induces isomorphisms $H_{\text{rot}}^s(\Gamma) \rightarrow H_{\text{div}}^s(\Gamma)$ and $H_{\text{div}}^s(\Gamma) \rightarrow H_{\text{rot}}^s(\Gamma)$.

Recall that we have well-defined continuous tangential trace operators ([29], Theorem 5.4.2 p. 209)

$$(2.9) \quad \gamma_{\text{T}}^- : \begin{cases} H_{\text{curl}}^0(\Omega_-) & \rightarrow H_{\text{rot}}^{-1/2}(\Gamma), \\ v & \mapsto v_{\text{T}} = v - (v \cdot n)n, \end{cases}$$

and, for arbitrary large enough $R > 0$ (with $B_R = \{x \in \mathbb{R}^3 : |x| < R\}$)

$$(2.10) \quad \gamma_{\text{T}}^+ : \begin{cases} H_{\text{curl}}^0(\Omega_+ \cap B_R) & \rightarrow H_{\text{rot}}^{-1/2}(\Gamma), \\ v & \mapsto v_{\text{T}} = v - (v \cdot n)n. \end{cases}$$

For simplicity we denote by $H_{\text{curl}}^0(\Omega_+)_{\text{loc}}$ the Fréchet space of vector fields in Ω_+ whose restrictions are in $H_{\text{curl}}^0(\Omega_+ \cap B_R)$ for all $R > 0$.

The Silver-Müller radiation condition at infinity for an electromagnetic field $(E, H) \in H_{\text{curl}}^0(\Omega_+)_{\text{loc}}^2$ is

$$(2.11) \quad \mu^{1/2}H \times x/|x| - \epsilon^{1/2}E = o(1/|x|).$$

For *exterior* problems we have the existence and uniqueness result ([29], Theorem 5.4.6, p. 220):

Theorem 2.1. *For all $k > 0$, all $v \in H_{\text{rot}}^{-1/2}(\Gamma)$ there is a unique $(E, H) \in H_{\text{curl}}^0(\Omega_+)_{\text{loc}}^2$ solving Maxwell's equations in Ω_+ , satisfying the Silver-Müller radiation condition, and such that $\gamma_{\text{T}}^+ E = v$. The corresponding solution operator is continuous.*

For *interior* problems we have

Theorem 2.2. *There is a unique real positive strictly increasing and unbounded sequence (k_n) such that with $\mathcal{K} = \{k_n : n \in \mathbb{N}\}$ we have*

- for all $k \notin \mathcal{K}$, for all $v \in H_{\text{rot}}^{-1/2}(\Gamma)$ there is a unique $(E, H) \in H_{\text{curl}}^0(\Omega_-)^2$ solving Maxwell's equations in Ω_- and such that $\gamma_{\text{T}}^- E = v$;
- for all $k \in \mathcal{K}$, the space of solutions $(E, H) \in H_{\text{curl}}^0(\Omega_-)^2$ to Maxwell's equations in Ω_- , such that $\gamma_{\text{T}}^- E = 0$, is a nonzero finite dimensional space.

The elements of \mathcal{K} are called *resonant wavenumbers*. For any k , the vector space of electric fields E in Ω_- , such that with $H = 1/(ikZ) \text{curl} E$, (E, H) is in $H_{\text{curl}}^0(\Omega_-)^2$ and solves Maxwell's equations in Ω_- with the *perfect conductor* boundary condition $\gamma_{\text{T}}^- E = 0$, is denoted \mathcal{E}_k . It is finite dimensional and it is nonzero only if $k \in \mathcal{K}$.

Recall the Sommerfeld radiation condition

$$(2.12) \quad \partial_r u - iku = o(1/r) \quad \text{with } r = |x|,$$

and let G_k be the fundamental solution of the Helmholtz operator $-\Delta - k^2$ satisfying it

$$(2.13) \quad G_k(x, y) = \frac{e^{ik|x-y|}}{4\pi|x-y|}.$$

Let Φ_k be the potential, mapping any sufficiently smooth tangent field u on Γ to the field in \mathbb{R}^3 defined away from Γ by

$$(2.14) \quad (\Phi_k u)(y) = \int_{\Gamma} G_k(x, y) u(x) dx.$$

Of fundamental importance to us will be the representation theorem ([29], Theorem 5.5.1, p. 234):

Theorem 2.3. *Suppose (E, H) is a field whose restrictions to Ω_- and Ω_+ are in $H_{\text{curl}}^0(\Omega_-)^2$ and $H_{\text{curl}}^0(\Omega_+)^2_{\text{loc}}$ and solve Maxwell's equations for a given wavenumber k . Suppose also that it verifies the Silver-Müller radiation condition. Define the electric and magnetic currents j and m on Γ by the jump formulas*

$$(2.15) \quad j = [H \times n] = (\gamma_{\Gamma}^- H - \gamma_{\Gamma}^+ H) \times n,$$

$$(2.16) \quad m = [E \times n] = (\gamma_{\Gamma}^- E - \gamma_{\Gamma}^+ E) \times n.$$

Then in Ω_- and Ω_+ we have

$$(2.17) \quad E = (+ikZ)(1 + (1/k^2) \text{grad div})\Phi_k j + \text{curl } \Phi_k m,$$

$$(2.18) \quad H = (-ik/Z)(1 + (1/k^2) \text{grad div})\Phi_k m + \text{curl } \Phi_k j.$$

Definition 2.4. For $k \neq 0$ we define the Electric Field Integral Operator, on tangent fields on Γ , by (the interior and exterior traces are equal)

$$(2.19) \quad A_k u = \gamma_{\Gamma}(1 + (1/k^2) \text{grad div})\Phi_k u.$$

One shows that A_k is continuous $H_{\text{div}}^s(\Gamma) \rightarrow H_{\text{rot}}^s(\Gamma)$. The EFIE for a given tangent field v is the equation $A_k u = v$. From the preceding equations it follows that if $k \in \mathbb{R}_+^* \setminus \mathcal{K}$, $A_k : H_{\text{div}}^{-1/2}(\Gamma) \rightarrow H_{\text{rot}}^{-1/2}(\Gamma)$ is invertible ([29], Theorem 5.6.2, p. 247). But more interestingly we have the following results even if k is a resonant wavenumber:

- for any $v \in H_{\text{rot}}^{-1/2}(\Gamma)$, if the interior problem with respect to v has a solution (E_-, H_-) , then letting (E_+, H_+) be the exterior solution and putting $u = (\gamma_{\Gamma}^- H_- - \gamma_{\Gamma}^+ H_+) \times n$, we have $u \in H_{\text{div}}^{-1/2}(\Gamma)$ and $A_k u = v$;
- for any $v \in H_{\text{rot}}^{-1/2}(\Gamma)$, if we have a solution $u \in H_{\text{div}}^{-1/2}(\Gamma)$ to $A_k u = v$, then the solution (E, H) of the exterior problem with respect to v is given by

$$(2.20) \quad E = (1 - (1/k^2) \text{grad div})\Phi_k u, \quad H = 1/(ikZ) \text{curl } \Phi_k u;$$

- for any $u \in H_{\text{div}}^{-1/2}(\Gamma)$, $A_k u = 0$ if and only if $u \in (\gamma_{\Gamma}^- \text{curl } \mathcal{E}_k) \times n$.

Thus for instance if v is the tangential trace of a planar wave—which is the case in RCS computations—then the EFIE has a solution even at resonances, which is determined up to a finite dimensional space, and for any such solution the corresponding potential solves the exterior problem. One of the main goals of this paper is to determine to which extent an analogous property holds true for the discretized EFIE.

2.2. Variational formulation and discretization. Recall that the \mathbb{C} bilinear form on tangent fields

$$(2.21) \quad (u, v) \mapsto \langle u, v \rangle = \int_{\Gamma} u \cdot v$$

induces a duality between $H_{\text{div}}^{-1/2}(\Gamma)$ and $H_{\text{rot}}^{-1/2}(\Gamma)$ ([29], Lemma 4.5.1, p. 208).

For sufficiently smooth tangent fields we have (all integrals are on Γ)

$$(2.22) \quad \langle A_k u, u' \rangle = \iint G_k(x, y) u(x) \cdot u'(y) dx dy \\ - (1/k^2) \iint G_k(x, y) \operatorname{div} u(x) \operatorname{div} u'(y) dx dy.$$

In order to have a highest order term which is positive we will sometimes consider the opposite bilinear form (associated with $-A_k$). This leads to a variational formulation of the EFIE, also known as the Rumsey reaction principle. For a given $v \in \mathbf{H}_{\text{rot}}^{-1/2}(\Gamma)$ solve

$$(2.23) \quad u \in \mathbf{H}_{\text{div}}^{-1/2}(\Gamma) \quad \text{and} \quad \forall u' \in \mathbf{H}_{\text{div}}^{-1/2}(\Gamma) \quad \langle A_k u, u' \rangle = \langle v, u' \rangle.$$

Whenever we have a family (X_h) of subspaces of $\mathbf{H}_{\text{div}}^{-1/2}(\Gamma)$, the Galerkin method consists in considering the equations

$$(2.24) \quad u \in X_h \quad \forall u' \in X_h \quad \langle A_k u, u' \rangle = \langle v, u' \rangle$$

and studying the convergence of the corresponding solutions (when they exist), with respect to the parameter h .

2.3. Some strategies for the analysis of the EFIE. As is clear from equation (2.22), the EFIE has dominant terms with different signs when restricted to gradients and divergence-free tangent fields. Therefore the theory of compact perturbations of coercive operators is not by itself enough to study the discretization of the EFIE.

It seems that the only numerical analysis available for the variational discretization of the EFIE is the original approach of Bendali [5], who introduces the charge density $q = \operatorname{div} u$ into the formulation, with Lagrange multipliers. More precisely the EFIE is formulated in the following way:

$$(2.25) \quad \begin{cases} u \in \mathbf{H}_{\text{div}}^{-1/2}(\Gamma) \\ q \in \mathbf{H}^{-1/2}(\Gamma)^\bullet \end{cases} \quad \begin{cases} \forall u' \in \mathbf{H}_{\text{div}}^{-1/2}(\Gamma) \\ \forall q' \in \mathbf{H}^{-1/2}(\Gamma)^\bullet \end{cases} \quad \begin{cases} a(u, u') + b(q, u') = l(u'), \\ c(q', u) + d(q, q') = 0, \end{cases}$$

where for any space W of scalar functions on Γ , W^\bullet denotes the subspace of W whose elements are L^2 orthogonal to the functions that are constant on each connected component of Γ , and

$$(2.26) \quad l(u') = -\langle v, u' \rangle,$$

$$(2.27) \quad a(u, u') = -\iint G_k(x, y) u(x) \cdot u'(y) dx dy,$$

$$(2.28) \quad b(q, u') = (1/k^2) \iint G_k(x, y) q(x) \operatorname{div} u'(y) dx dy,$$

$$(2.29) \quad c(q', u) = (1/k^2) \iint G_0(x, y) q'(x) \operatorname{div} u(y) dx dy,$$

$$(2.30) \quad d(q, q') = -(1/k^2) \iint G_0(x, y) q(x) q'(y) dx dy.$$

This formulation leads to an intricate mathematical analysis relying heavily upon the fact that the operator with kernel $G_k - G_0$ is of order -3 . Its main advantage is that for the usual Galerkin spaces $X_h \subset \mathbf{H}_{\text{div}}^{-1/2}(\Gamma)$ the spaces $W_h = \operatorname{div} X_h$ are simple subspaces of the usual Galerkin spaces of piecewise polynomial functions

(with no continuity requirements). Thus the formulation (2.25) has a discrete counterpart on $W_h \times X_h$.

Nédélec [29] has used the following splitting of $X = \mathbf{H}_{\text{div}}^{-1/2}(\Gamma)$ to prove the Fredholm property of the EFIE. Let a be the bilinear form induced by the EFIE. Put $W = \mathbf{H}^{1/2}(\Gamma)$. Then $\text{rot } W$ is a closed subspace of X . Let V be the orthogonal of $\text{rot } W$ with respect to a , i.e.,

$$(2.31) \quad V = \{u \in X : \forall w \in W \quad a(\text{rot } w, u) = 0\}.$$

Since the restriction of a to $\text{rot } W$ is a compact perturbation of a coercive form, $V \cap W$ is finite dimensional, and $V + W$ is closed and has finite codimension in X . Thus, up to finite dimensional spaces, one can search for the solution of the EFIE in the form $u = v + \text{rot } w$, with (v, w) solving

$$(2.32) \quad \begin{cases} v \in X \\ w \in W \end{cases} \quad \begin{cases} \forall v' \in X & a(v, v') + a(\text{rot } w, v') = l(v'), \\ \forall w' \in W & a(\text{rot } w', v) = 0. \end{cases}$$

The conclusive remark is that on V , a satisfies a Gårding inequality, thus this saddle-point problem is Fredholm of index 0.

2.4. Reformulation of the EFIE as a saddle-point. We will use a nonsymmetric variant of this last technique which has the advantage of not introducing the parasitic finite dimensional subspaces. In particular the analysis we propose does not need the fact that the operator with kernel $G_k - G_0$ is of order -3 , only that is it compact $\mathbf{H}^{-1/2}(\Gamma) \rightarrow \mathbf{H}^{1/2}(\Gamma)$.

In order to be more precise in our statements, we will require the theory of saddle-point problems. We assume familiarity with the theory of Inf-Sup conditions as presented for instance in Roberts-Thomas [34], and in particular the way *discrete* Inf-Sup conditions lead to convergence estimates for Galerkin methods. However, for completeness we include a result of Nicolaides [30], which generalizes the classical theorem of Brezzi [10].

Theorem 2.5. *Let X_i and W_i for $i = 1, 2$ be Hilbert spaces. Let a , b and c be continuous bilinear forms on $X_1 \times X_2$, $W_1 \times X_2$ and $W_2 \times X_1$. Let V_1 (resp. V_2) denote the right-hand kernel of c on $W_2 \times X_1$ (resp. b on $W_1 \times X_2$). Suppose that*

$$(2.33) \quad \inf_{V_1} \sup_{V_2} a \geq \alpha > 0,$$

$$(2.34) \quad \inf_{W_1} \sup_{X_2} b \geq \beta > 0,$$

$$(2.35) \quad \inf_{W_2} \sup_{X_1} c \geq \gamma > 0,$$

$$(2.36) \quad \forall v_2 \in V_2 \quad (\forall v_1 \in V_1 \quad a(v_1, v_2) = 0) \Rightarrow (v_2 = 0).$$

Then for all $(g, f) \in W_2^ \times X_2^*$ there is a unique $(w_1, u_1) \in W_1 \times X_1$ such that*

$$(2.37) \quad \begin{cases} u_1 \in X_1 \\ w_1 \in W_1 \end{cases} \quad \begin{cases} \forall u_2 \in X_2 & a(u_1, u_2) + b(w_1, u_2) = f(u_2), \\ \forall w_2 \in W_2 & c(w_2, u_1) = g(w_2). \end{cases}$$

Moreover one has the continuity estimate for the solution operator

$$(2.38) \quad \|u_1\| \leq \alpha^{-1} \|f\| + \gamma^{-1} (1 + \alpha^{-1} \|a\|) \|g\|,$$

$$(2.39) \quad \|w_1\| \leq \beta^{-1} (1 + \alpha^{-1} \|a\|) (\|f\| + \gamma^{-1} \|a\| \|g\|).$$

If for all (g, f) there is a unique solution (w_1, u_1) , then the above Inf-Sup conditions are satisfied for some α, β and γ , as noted by Bernardi et al. [6].

We return now to the special case of interest, the EFIE. We put $X = \mathbf{H}_{\text{div}}^{-1/2}(\Gamma)$ and let a be the bilinear form on X induced by the EFIE

$$(2.40) \quad \begin{aligned} a(u, u') &= (1/k^2) \iint G_k(x, y) \operatorname{div} u(x) \operatorname{div} u'(y) dx dy \\ &\quad - \iint G_k(x, y) u(x) \cdot u'(y) dx dy. \end{aligned}$$

Let d be the bilinear form on X defined by

$$(2.41) \quad \begin{aligned} d(u, u') &= -(1/k^2) \iint G_0(x, y) \operatorname{div} u(x) \operatorname{div} u'(y) dx dy \\ &\quad - \iint G_0(x, y) u(x) \cdot u'(y) dx dy. \end{aligned}$$

The involution of X induced by the complex conjugation is denoted $u \mapsto \bar{u}$. Then it follows from the positivity of the operator with kernel G_0 that $-d(\cdot, \bar{\cdot})$ is a Hermitian scalar product on X . Thus for any closed subspace X_0 of X and any closed subspace W_0 of X_0 , if X_0 and W_0 are stable under conjugation (*conjugation-stable* for short) one has the orthogonal splitting $X_0 = V_0 \oplus W_0$, with

$$(2.42) \quad V_0 = \{u \in X_0 : \forall w \in W_0 \quad d(w, u) = 0\}.$$

Put $W = \{u \in X : \operatorname{div} u = 0\}$. Now let X_0 be any closed and conjugation-stable subspace of X , and put $W_0 = X_0 \cap W$, which is also closed and conjugation-stable. Then, for all $l \in X_0^*$, u solves

$$(2.43) \quad u \in X_0 \quad \text{and} \quad \forall u' \in X_0 \quad a(u, u') = l(u'),$$

if and only if $u = v + w$, with (v, w) solving

$$(2.44) \quad \begin{cases} v \in X_0 \\ w \in W_0 \end{cases} \quad \begin{cases} \forall v' \in X_0 & a(v, v') + b(w, v') = l(v'), \\ \forall w' \in W_0 & c(w', v) = 0, \end{cases}$$

where b is the restriction of a to $W \times X$ and c the restriction of d to $W \times X$.

Consider first the case $X_0 = X$. Let $\Theta : W \times X \rightarrow W^* \times X^*$ be the operator associated with the left-hand side of the above saddle-point (2.44). If the kernels G_k are replaced by G_0 in a and b (keeping the outside coefficient $(1/k^2)$ untouched), the corresponding operator $\Theta_0 : W \times X \rightarrow W^* \times X^*$ is symmetric. It is the saddle-point mapping associated with the bilinear form a_0 defined by

$$(2.45) \quad \begin{aligned} a_0(u, u') &= (1/k^2) \iint G_0(x, y) \operatorname{div} u(x) \operatorname{div} u'(y) dx dy \\ &\quad - \iint G_0(x, y) u(x) \cdot u'(y) dx dy. \end{aligned}$$

For Θ_0 the Brezzi compatibility estimates (2.34) and (2.35) are trivial. The cornerstone of the argument is

Theorem 2.6. *On the right kernel of c on $W \times X$, the bilinear form a_0 satisfies a Gårding inequality.*

Proof. Let V denote the right kernel of c on $W \times X$. Remark first that V is a supplementary of $\ker_X \operatorname{div}$ and that $\operatorname{div} : X \rightarrow \mathbf{H}^{-1/2}$ has closed range. It follows

that div determines an isomorphism from V to its range, so there is $C > 0$ such that

$$(2.46) \quad \forall v \in V \quad \|v\|_X \leq C |\operatorname{div} v|_{-1/2}.$$

Hence the first term in the right-hand side of equation (2.45) is coercive on V .

Concerning the second term, notice that for elements v of V we have

$$(2.47) \quad \operatorname{rot} \pi_T \int G_0(x, y) v(x) dx = 0,$$

where π_T denotes the orthogonal projection onto the tangent space. As shown in Nédélec [29], it follows that rot maps V continuously into $H^{-1/2}(\Gamma)$, hence V is continuously imbedded in $H_T^{1/2}(\Gamma)$. In turn the injection $H_T^{1/2}(\Gamma) \rightarrow H_T^{-1/2}(\Gamma)$ is compact so the proof is complete. \square

From this it follows that Θ_0 is Fredholm, and by symmetry its index must be 0. Since Θ is a compact perturbation of Θ_0 , Θ is also Fredholm of index 0.

If $u \mapsto a(u, \cdot)$ is injective, the above splitting, in the case $X_0 = X$ shows that the map Θ is also injective. Therefore it is an isomorphism.

Remark 2.7. We could also have made the choice $W = \operatorname{rot} H^{1/2}(\Gamma)$, which differs from the adopted one only by a finite dimensional space, whose elements are known as the Neumann fields (or harmonic fields). Some proofs would have to be modified.

3. INF-SUP ESTIMATES FOR THE EFIE

We now turn to the discretization of the EFIE. Recall notation from Section 2.4. We consider the case where X_0 is a Galerkin space, one among a family (X_h) . Putting $W_h = X_h \cap W$, our strategy will be to first study the saddle-point problem associated with the EFIE on $W_h \times X_h$ and to give sufficient conditions on (X_h) for it to satisfy uniform Inf-Sup estimates. From these we easily deduce Inf-Sup estimates for the original problem on X_h .

3.1. Sufficient conditions for uniform Inf-Sup estimates.

Lemma 3.1. *Let X be a Hilbert space. The scalar product is denoted $(\cdot|\cdot)$, and orthogonality is denoted \perp . Let $X = V \oplus W$ be an orthogonal splitting. Let (X_h) be an approximating family of closed subspaces. Put $W_h = W \cap X_h$, and $V_h = \{u \in X_h : u \perp W_h\}$. Then (V_h) is approximating in V , and if (V_h) is quasi-conforming in V , then (W_h) is approximating in W .*

Proof. Indeed the orthogonal projection P_h onto X_h maps V into V_h , therefore (V_h) is approximating. For any $w \in W$, put $P_h w = v_h + w_h$ with $v_h \in V_h$ and $w_h \in W_h$. We have

$$(3.1) \quad \forall v' \in V \quad (v_h|v_h) = (v_h|P_h w) = (v_h|w) = (v_h - v'|w).$$

Hence

$$(3.2) \quad \|v_h\|^2 \leq \|w\|^2 \sup_{v \in V_h} \inf_{v' \in V} \|v - v'\|/\|v\|,$$

and since

$$(3.3) \quad \|w - w_h\| \leq \|w - P_h w\| + \|v_h\|,$$

it follows that if (V_h) is quasi-conforming in V , then $\|w - w_h\|$ tends to 0. \square

Theorem 3.2. *Recall notation from Section 2.4. Let (X_h) be an approximating family of closed and conjugation-stable subspaces of X . Let V be the right kernel of c on $W \times X$ and V_h the right kernel of c on $W_h \times X_h$. If $\delta(V_h, V) \rightarrow 0$, then Θ is discrete LSF on $(W_h \times X_h) \times (W_h \times X_h)$, and $u \mapsto a(u, \cdot)$ is discrete LSF on $(X_h \times X_h)$.*

Proof. We prove the result for Θ_0 . Then the theorem follows for Θ by Corollary 1.13, for $u \mapsto a_0(u, \cdot)$ by the equivalent splitting and for $u \mapsto a(u, \cdot)$ likewise.

We will use on X the scalar product induced by $-d$. By Theorem 2.6, Example 1.14 and Lemma 1.19 it follows that if N_0 is the kernel of a_0 on $V \times V$, we have, for some $\alpha_0 > 0$ and $h_0 > 0$

$$(3.4) \quad \forall h < h_0 \quad \inf_{N_0^\perp \cap V_h} \sup_{V_h} a_0 > \alpha_0,$$

where N_0^\perp is the orthogonal complement of N_0 in X .

For any h , let V_h^\perp be the orthogonal complement of V_h in X_h . Since c is the restriction of d , we can choose a $\gamma > 0$ such that

$$(3.5) \quad \forall h \quad \gamma < \inf_{V_h^\perp} \sup_{W_h} c^\dagger = \inf_{W_h} \sup_{V_h^\perp} c = \inf_{W_h} \sup_{X_h} c.$$

For any Banach space Y with norm denoted $\|\cdot\|$, any continuous linear form l on Y and any subspace Y_h of Y we use the notation

$$(3.6) \quad \|l\|_{Y_h} = \sup_{u \in Y_h} |l(u)|/\|u\|.$$

Choose $h < h_0$, $w \in W_h$, $v \in N_0^\perp \cap V_h$ and $u \in V_h^\perp$. Put $(g, f) = \Theta_0(w, u + v)$. Following Nicolaides, one checks that

$$(3.7) \quad \|u\| \leq \gamma^{-1}(\|g\|_{W_h}),$$

$$(3.8) \quad \|v\| \leq \alpha_0^{-1}(\|f\|_{X_h} + \|a_0\| \|u\|),$$

$$(3.9) \quad \|w\| \leq \gamma^{-1}(\|f\|_{X_h} + \|a_0\| \|v\| + \|a_0\| \|u\|).$$

Hence

$$(3.10) \quad \|u + v\| \leq \alpha_0^{-1}\|f\|_{X_h} + \gamma^{-1}(1 + \alpha_0^{-1}\|a_0\|) \|g\|_{W_h},$$

$$(3.11) \quad \|w\| \leq \gamma^{-1}(1 + \alpha_0^{-1}\|a_0\|) (\|f\|_{X_h} + \gamma^{-1}\|a_0\| \|g\|_{W_h}).$$

For any subspace M of X let P_M denote the orthogonal projection onto it. Note that

$$(3.12) \quad (N_0^\perp \cap V_h) + V_h^\perp = \{u \in X_h : u \perp P_{V_h}(N_0)\},$$

whereas

$$(3.13) \quad N_0^\perp \cap X_h = \{u \in X_h : u \perp P_{X_h}(N_0)\}.$$

Now since $N_0 \subset V$, $\delta(V_h, V) \rightarrow 0$ and V_h is approximating V , it follows that N_0 , $P_{V_h}(N_0)$ and $P_{X_h}(N_0)$ are uniformly close to each other (all the gaps tend to 0). Hence $(N_0^\perp \cap V_h) + V_h^\perp$ is uniformly close to $N_0^\perp \cap X_h$ and therefore we deduce from the above that

$$(3.14) \quad \liminf_h \inf_{W_h \times (N_0^\perp \cap X_h)} \sup_{W_h \times X_h} \Theta_0 > 0.$$

Considering that W and X are orthogonal subspaces of the product space, the theorem follows. \square

We will give some explicit examples of such Galerkin spaces in Section 3.5.

3.2. Spurious electric currents. Suppose for the moment that the hypotheses of Theorem 3.2 is verified, so that for *any* nonzero wavenumber k the associated bilinear form a_k is discrete LSF on $(X_h \times X_h)$. Let I be a compact interval of positive reals (which might very well contain resonant wavenumbers). Since the map $k \mapsto a_k$ is continuous, it maps I to a compact set, and we can apply Corollary 1.17, which we express in terms of sequences:

Suppose we have for each integer n , a wavenumber $k_n \in I$, a discretization parameter h_n and a current $u_n \in X_{h_n}$, such that (h_n) tends to 0 and

$$(3.15) \quad \sup_{u' \in X_{h_n}} \frac{|a_{k_n}(u_n, u')|}{\|u'\|_{-1/2}} = o(\|u_n\|_{-1/2}).$$

Then we have (the constant C stems from the standard duality $\langle \cdot, \cdot \rangle$ between $H_{\text{rot}}^{-1/2}(\Gamma)$ and $H_{\text{div}}^{-1/2}(\Gamma)$ and is of course independent of the wavenumber)

$$(3.16) \quad \|A_{k_n} u_n\|_{H_{\text{rot}}^{-1/2}(\Gamma)} \leq C \sup_{u' \in X} \frac{|a_{k_n}(u_n, u')|}{\|u'\|_{-1/2}} = o(\|u_n\|_{-1/2}).$$

From the uniform continuity of the solution operator for the exterior problem on I , we deduce that the electromagnetic field generated by u_n is negligible compared with $\|u_n\|_{-1/2}$. More precisely if (E_n, H_n) is given by

$$(3.17) \quad E_n = (1 - (1/k_n^2) \text{grad div}) \Phi_{k_n} u_n, \quad H_n = 1/(ik_n Z) \text{curl } \Phi_{k_n} u_n,$$

then for any R ,

$$(3.18) \quad \|E_n\|_{H_{\text{curl}}^0(\Omega_+ \cap B_R)} = o(\|u_n\|_{-1/2}),$$

and a similar estimate holds for H_n . Expressed differently, if we have *bounded* currents producing small *discrete* residual errors, then the near-fields they radiate are also small. The far-field pattern associated with an exterior electromagnetic field depends continuously upon the boundary data, so it is also negligible compared with $\|u_n\|_{-1/2}$.

If one looks at a single wavenumber of interest (for instance a resonant one), then by Remark 1.16 one can exhibit rather explicit estimates as follows: Suppose k is a resonant frequency, put $a = a_k$, and let l be a linear form which is a compatible right-hand side. Let u denote a solution

$$(3.19) \quad \forall v \in X \quad a(u, v) = l(v).$$

Let u_h^b be the best approximation of u in X_h , and suppose we solve the Galerkin equation with an error ϵ_h

$$(3.20) \quad \sup_{v \in Y_h} |a(u_h, v) - l(v)| / \|v\| \leq \epsilon_h.$$

We also have

$$(3.21) \quad \sup_{v \in Y_h} |a(u_h^b, v) - l(v)| / \|v\| \leq \|a\| \|u_h^b - u\|.$$

Hence

$$(3.22) \quad \sup_{v \in Y_h} |a(u_h - u_h^b, v)| / \|v\| \leq \epsilon_h + \|a\| \|u_h^b - u\|.$$

Let N denote the (right) kernel of a . Let δ_h be the order of best approximation of the elements of N by elements of X_h . Then by equation (1.49) we have the estimate

$$(3.23) \quad \inf_{v \in N} \|u_h - u_h^b - v\| \leq C(\epsilon_h + \|u_h^b - u\| + \delta_h \|u_h - u_h^b\|).$$

As will be seen later, for the standard Finite Element approximation, δ_h and $\|u_h^b - u\|$ are of order $h^{3/2}$, and an ϵ_h of this order is also possible. In other words the discrete Galerkin solution can be written as a best approximation, plus an element of the kernel (which does not radiate at all), plus a current of magnitude $h^{3/2}$ less than $\|u_h\|$.

Notice however that for an ϵ_h of order $h^{3/2}$, $\|u_h\|$ can be bounded, but if one asks for too small an ϵ_h (smaller than $h^{3/2}$), then $\|u_h\|$ might be forced to be very big.

3.3. Sufficient conditions in integer exponent Sobolev spaces. The preceding sufficient conditions for the good behavior of the EFIE were formulated in half-integer Sobolev norms. We prove here that these condition hold under some hypotheses formulated in the more familiar integer Sobolev norms.

Recall notation from Section 2.4. Consider the following hypotheses for a family of spaces X_h :

(H0) The spaces X_h are finite dimensional conjugation-stable subspaces of $\mathbf{H}_{\text{div}}^0(\Gamma)$.

(H1) There is $C > 0$ such that for all $u \in \mathbf{H}_{\text{div}}^1(\Gamma)$

$$(3.24) \quad \inf_{u' \in X_h} \|u - u'\|_0 \leq Ch \|u\|_1.$$

(H2) There is $C > 0$ such that for all $u \in X_h$, $\|u\|_0 \leq Ch^{-1} \|u\|_{-1}$.

(H3) There is $C > 0$ such that for all $u \in X_h$ if

$$(3.25) \quad \forall w \in W_h \quad \langle u, w \rangle = 0,$$

then the solution p of

$$(3.26) \quad p \in \mathbf{H}^1(\Gamma)^\bullet \quad \text{and} \quad \Delta p = \text{div } u$$

satisfies

$$(3.27) \quad \|u - \text{grad } p\|_0 \leq Ch \|\text{div } u\|_0.$$

Notice that by (H3) we have the usual Inf-Sup estimate: There is $C > 0$ such that

$$(3.28) \quad \inf_{q \in \text{div } X_h} \sup_{u \in X_h} \frac{|\langle q, \text{div } u \rangle|}{\|q\|_0 \|u\|_0} \geq \frac{1}{C}.$$

Let $\Omega_h : \mathbf{H}_{\text{div}}^0 \rightarrow X_h$ map $u_0 \in \mathbf{H}_{\text{div}}^0$ to $u \in X_h$, the solution of

$$(3.29) \quad \begin{cases} u \in X_h \\ q \in \text{div } X_h \end{cases} \quad \begin{cases} \forall u' \in X_h & \langle u, u' \rangle + \langle q, \text{div } u' \rangle = \langle u_0, u' \rangle, \\ \forall q' \in \text{div } X_h & \langle q', \text{div } u \rangle = \langle q', \text{div } u_0 \rangle. \end{cases}$$

Under the above hypotheses one has an estimate of the form, for $u_0 \in \mathbf{H}_{\text{div}}^1$,

$$(3.30) \quad \|\Omega_h u_0 - u_0\|_0 \leq Ch \|u_0\|_1.$$

Moreover Ω_h maps divergence-free fields to divergence-free fields.

We denote by V the right kernel of c on $W \times X$ and by V_h the one on $W_h \times X_h$.

Theorem 3.3. *Under the above hypotheses (V_h) is quasi-conforming in V . More precisely there is C such that for all h and all $v \in V_h$, if p solves*

$$(3.31) \quad p \in H^1(\Gamma)^\bullet \quad \text{and} \quad \Delta p = \operatorname{div} v$$

and w solves

$$(3.32) \quad w \in W \quad \forall w' \in W \quad d(w, w') = -d(\operatorname{grad} p, w'),$$

then $\operatorname{grad} p + w \in V$ and

$$(3.33) \quad \|v - (\operatorname{grad} p + w)\|_{-1/2} \leq Ch^{1/2} \|v\|_{-1/2}.$$

Proof. Put $v = v_0 + v_1$ with $v_0 \in W_h$ and $v_1 \in X_h$ such that $\forall w \in W_h \langle w, v_1 \rangle = 0$. According to (H3) we have

$$(3.34) \quad \|v_1 - \operatorname{grad} p\|_0 \leq Ch \|u\|_0.$$

So from (H2) we deduce

$$(3.35) \quad \|v_1 - \operatorname{grad} p\|_{-1/2} \leq Ch^{1/2} \|u\|_{-1/2}.$$

We will use a regularity result that we state without proof. Let Λ be the operator on tangent fields that maps any u to a divergence-free w such that for all divergence-free w' we have $d(w, w') = d(u, w')$. Then Λ is continuous $H_T^s(\Gamma) \rightarrow H_T^s(\Gamma)$.¹

The field w is defined by $w = -\Lambda \operatorname{grad} p$. From the H_T^1 continuity of Λ , it follows that

$$(3.36) \quad \|\Omega_h w - w\|_0 \leq Ch \|w\|_1 = Ch |w|_1 \leq Ch |\operatorname{grad} p|_1 \leq Ch \|u\|_0.$$

Hence

$$(3.37) \quad \|\Omega_h w - w\|_{-1/2} \leq Ch^{1/2} \|u\|_{-1/2}.$$

But for all $w' \in W_h$

$$(3.38) \quad d(\Omega_h w - v_0, w') = d(\Omega_h w + v_1, w') = d((\Omega_h w - w) + (v_1 - \operatorname{grad} p), w').$$

The absolute value of this last expression is bounded by

$$(3.39) \quad C(\|\Omega_h w - w\|_{-1/2} + \|v_1 - \operatorname{grad} p\|_{-1/2}) \|w'\|_{-1/2}.$$

Taking w' to be the conjugate of $\Omega_h w - v_0 \in W_h$ gives

$$(3.40) \quad \|\Omega_h w - v_0\|_{-1/2} \leq Ch^{1/2} \|u\|_{-1/2}.$$

Combining equation (3.35) with (3.37) and (3.40) gives the theorem. □

¹This regularity result is comparable to the following more standard one. The pseudo-differential operator A defined by

$$\langle A\varphi, \varphi' \rangle = \iint \frac{1}{|x-y|} \operatorname{rot} \varphi(x) \cdot \operatorname{rot} \varphi'(y) dx dy$$

is an isomorphism of order 1, and the pseudo-differential operator B defined by

$$\langle B\varphi, \varphi' \rangle = \iint \frac{1}{|x-y|} \operatorname{grad} \varphi(x) \cdot \operatorname{rot} \varphi'(y) dx dy$$

is a morphism of order less than 1 (one can show that it is of order 0). The operator $A^{-1}B$ is therefore of order less than 0.

3.4. Convergence rates. Let (X_h) be a family of Galerkin spaces satisfying (H0) and (H1). Let \mathcal{Q}_h be the $H_{\text{div}}^0(\Gamma)$ -orthogonal projection onto X_h . Then we have

$$(3.41) \quad \|\mathcal{Q}_h u\|_0 \leq C\|u\|_0 \quad \text{and} \quad \|u - \mathcal{Q}_h u\|_0 \leq Ch\|u\|_1.$$

The spaces $H_{\text{div}}^s(\Gamma)$ for $0 \leq s \leq 1$ can be obtained by interpolation. Hence interpolation on the operator $\mathcal{I} - \mathcal{Q}_h$, for $0 \leq s \leq 1$, gives

$$(3.42) \quad \|u - \mathcal{Q}_h u\|_0 \leq Ch^s \|u\|_s.$$

Then one uses the regularity of the $H_{\text{div}}^0(\Gamma)$ -inner product (written $(\cdot|\cdot)_0$) on various Sobolev spaces. This technique is the familiar Aubin-Nitsche trick. That $H_{\text{div}}^s(\Gamma)$ and $H_{\text{div}}^{-s}(\Gamma)$ are dual with respect to the $H_{\text{div}}^0(\Gamma)$ -inner product can be deduced from the fact that the operator $I - \text{grad div}$ is an isomorphism $H_{\text{div}}^s(\Gamma) \rightarrow H_{\text{rot}}^{s-1}(\Gamma)$ and that this space, as already mentioned, is the L_T^2 -dual of $H_{\text{div}}^{-s}(\Gamma)$. Both of these facts can be proved using the Helmholtz decomposition and regularity of the Laplacian. For $0 \leq s \leq 1$ we have

$$(3.43) \quad \|u - \mathcal{Q}_h u\|_{-s} \leq C \sup_{v \in H_{\text{div}}^s(\Gamma)} \frac{|(u - \mathcal{Q}_h u|v)_0|}{\|v\|_s}$$

$$(3.44) \quad \leq C \sup_{v \in H_{\text{div}}^s(\Gamma)} \frac{|(u - \mathcal{Q}_h u|v - \mathcal{Q}_h v)_0|}{\|v\|_s}$$

$$(3.45) \quad \leq C\|u - \mathcal{Q}_h u\|_0 \|\mathcal{I} - \mathcal{Q}_h\|_{0,s}.$$

Here $\|\mathcal{I} - \mathcal{Q}_h\|_{0,s}$ is of course the norm of the induced map

$$(3.46) \quad \mathcal{I} - \mathcal{Q}_h : H_{\text{div}}^s(\Gamma) \rightarrow H_{\text{div}}^0(\Gamma).$$

This gives for $0 \leq s, s' \leq 1$

$$(3.47) \quad \|u - \mathcal{Q}_h u\|_{-s} \leq Ch^{s+s'} \|u\|_{s'}.$$

For smooth scatterers and smooth incident waves (such as plane waves) the solution u of the EFIE is known to be smooth, and in particular $u \in H_{\text{div}}^1(\Gamma)$. Since the Inf-Sup condition yields quasi-optimal convergence, we therefore have

Theorem 3.4. *Under the above hypotheses the Galerkin solution u_h of the EFIE satisfies the convergence estimate $\|u - u_h\|_{-1/2} \leq Ch^{3/2}$.*

3.5. Some well-known spaces. Let φ be the orthogonal projection onto Γ , which is defined and smooth on a tubular neighborhood of Γ . Let (\mathcal{T}_h) be a family of triangulations of Γ , where for all h the largest diameter of a triangle of \mathcal{T}_h is h . We will always suppose that (\mathcal{T}_h) has the minimum angle property. Let Γ_h be the affine polyhedron determined by \mathcal{T}_h , considered as a Lipschitz manifold. For small enough h , φ induces Lipschitz-isomorphisms $\Gamma_h \rightarrow \Gamma$, and we denote by Ξ_h the inverse mappings.

Fix a nonzero $m \in \mathbb{N}$. On Γ_h we consider the space $\mathcal{S}^0(\mathcal{T}_h)$ of continuous scalar functions whose restriction to any triangle is P^m (a polynomial of degree m), the space $\mathcal{S}^1(\mathcal{T}_h)$ of Raviart-Thomas H_{div}^0 conforming vector fields of degree m , and the space $\mathcal{S}^2(\mathcal{T}_h)$ of scalar functions whose restriction to any triangle is P^{m-1} .

From these finite element spaces on Γ_h we deduce finite element spaces on Γ by the transport formulas

$$(3.48) \quad \begin{aligned} \mathcal{S}_h^0 &= \{x \mapsto p(\Xi_h(x)) : p \in \mathcal{S}^0(\mathcal{T}_h)\}, \\ \mathcal{S}_h^1 &= \{x \mapsto \text{Jac } \Xi_h(h) D \Xi_h(x)^{-1} u(\Xi_h(x)) : u \in \mathcal{S}^1(\mathcal{T}_h)\}, \\ \mathcal{S}_h^2 &= \{x \mapsto \text{Jac } \Xi_h(x) q(\Xi_h(x)) : q \in \mathcal{S}^2(\mathcal{T}_h)\}. \end{aligned}$$

These transport formulas were chosen to make the following diagram commute. The horizontal arrows are the differential operators rot and div , whereas the vertical ones are the above transport formulas.

$$(3.49) \quad \begin{array}{ccccc} S^0(\mathcal{T}_h) & \rightarrow & S^1(\mathcal{T}_h) & \rightarrow & S^2(\mathcal{T}_h) \\ \downarrow & & \downarrow & & \downarrow \\ S_h^0 & \rightarrow & S_h^1 & \rightarrow & S_h^2 \end{array}$$

If we wish to be precise about the order m , we use superscripts $S_h^{0,m}$, $S_h^{1,m}$ and $S_h^{2,m-1}$.

It is well known that (S_h^1) satisfies (H1). If the mesh is quasi-uniform, then (H2) also holds. The only remaining point is (H3). This property has been found to be very important in the study of eigenvalue problems by mixed formulations; see in particular Boffi et al. [8] and Demkowicz et al. [20]. This is because it implies a discrete compactness result which in the three-dimensional setting is due to Kikuchi [24]. Since this property is equally important to the present problem, we include a short proof of it. It goes without saying that the absence of a boundary on Γ greatly simplifies our task. Ramifications can be found in Boffi [7].

The usual degrees of freedom pertaining to \mathcal{T}_h on Γ_h can be transported to the curved triangles $\Xi_h^{-1}(T)$ for $T \in \mathcal{T}_h$. With these, an interpolation operator Π_h onto S_h^1 can be defined on spaces with slightly stronger regularity than $H_{\text{div}}^0(\Gamma)$; see Brezzi-Fortin [12], p. 125, and Roberts-Thomas [34], p. 549, for two different variants in the planar setting. We will use the following properties of this interpolation operator:

- Π_h is a projector onto S_h^1 defined on some extension of $H_T^1(\Gamma) + S_h^1$;
- Π_h maps divergence-free fields to divergence-free fields;
- $\exists C > 0 \forall u \in H_T^1(\Gamma) \forall h \quad |u - \Pi_h u|_0 \leq Ch|u|_1$.

Theorem 3.5. *For any regular family of triangulations (\mathcal{T}_h) , (S_h^1) satisfies (H3), i.e., with $X_h = S_h^1$, there is $C > 0$ such that for all $u \in X_h$ if*

$$(3.50) \quad \forall w \in W_h \quad \langle u, w \rangle = 0,$$

then the solution p of

$$(3.51) \quad p \in H^1(\Gamma)^\bullet \quad \Delta p = \text{div } u,$$

satisfies

$$(3.52) \quad |u - \text{grad } p|_0 \leq Ch|\text{div } u|_0.$$

Proof. Note that $\text{div}(\text{grad } p - u) = 0$, so

$$(3.53) \quad \text{div}(\Pi_h \text{grad } p - u) = \text{div } \Pi_h(\text{grad } p - u) = 0.$$

Put $\tilde{u} = \Pi_h \text{grad } p$. The approximation property of Π_h and the regularity of the Laplacian yield

$$(3.54) \quad |\tilde{u} - \text{grad } p|_0 \leq Ch|\text{grad } p|_1 \leq Ch|\text{div } u|_0.$$

Moreover for any divergence-free $u' \in X_h$ we have

$$(3.55) \quad \langle \tilde{u} - u, u' \rangle = \langle \tilde{u}, u' \rangle = \langle \tilde{u} - \text{grad } p, u' \rangle.$$

Applying this identity with u' , the conjugate of $\tilde{u} - u$ gives

$$(3.56) \quad |\tilde{u} - u|_0 \leq |\tilde{u} - \text{grad } p|_0 \leq Ch|\text{div } u|_0.$$

Therefore we have

$$(3.57) \quad |u - \text{grad } p|_0 \leq |u - \tilde{u}|_0 + |\tilde{u} - \text{grad } p|_0 \leq Ch |\text{div } u|_0. \quad \square$$

ACKNOWLEDGMENTS

I am indebted to F. Béreux for the important empirical observation that at resonant frequencies, the spurious currents do not pollute the computed far-field pattern. Also this work would not have been possible had I not received the invaluable advice of J.-C. Nédélec.

REFERENCES

- [1] T. Abboud, *Etude mathématique et numérique de quelques problèmes de diffraction d'ondes électromagnétiques*, PhD thesis, Ecole Polytechnique, 1991.
- [2] I. Babuska, *Error bounds for the finite element method*, Numer. Math., Vol. 16, pp. 322-333, 1971. MR **44**:6166
- [3] I. Babuska, A.K. Aziz, *Survey lectures on the mathematical foundations of the finite element method*, pp. 3-359, in A.K. Aziz (ed.), "The mathematical foundations of the finite element method with applications to partial differential equations", Academic Press, New York, 1972. MR **54**:9111
- [4] I. Babuska, J. Osborn, *Eigenvalue problems*, pp. 641-788, in P.G. Ciarlet, J.-L. Lions (eds.), "Handbook of numerical analysis, Vol. II, Finite element methods (Part 1)", North-Holland, 1991.
- [5] A. Bendali, *Numerical analysis of the exterior boundary value problem for the time-harmonic Maxwell equations by a boundary finite element method, Part 1: The continuous problem*, Math. Comp., Vol. 43, No. 167, pp. 29-46, 1984, *Part 2: The discrete problem*, Math. Comp., Vol. 43, No. 167, pp. 47-68, 1984. MR **86i**:65071a, MR **86i**:65071b
- [6] C. Bernardi, C. Canuto, Y. Maday, *Generalized Inf-Sup conditions for the Chebyshev spectral approximation of the Stokes problem*, SIAM J. Numer. Anal., Vol. 25, No. 6, pp. 1237-1271, 1988. MR **90e**:65151
- [7] D. Boffi, *Discrete compactness and Fortin operator for edge elements*, Numer. Math., Vol. 87, No. 2, pp. 229-246, 2000. MR **2001k**:65168
- [8] D. Boffi, F. Brezzi, L. Gastaldi, *On the problem of spurious eigenvalues in the approximation of linear elliptic problems in mixed form*, Math. Comp., Vol. 69, No. 229, pp. 121-140, 1999. MR **2000i**:65175
- [9] A. DeLaBourdonnaye, *Décomposition de $H_{\text{div}}^{-1/2}(\Gamma)$ et nature de l'opérateur de Steklov-Poincaré du problème extérieur de l'électromagnétisme*, C. R. Acad. Sci. Paris Sér. I Math., Vol. 316, No. 4, pp. 369-372, 1993. MR **93k**:78004
- [10] F. Brezzi, *On the existence, uniqueness and approximation of saddle-point problems arising from Lagrangian multipliers*, RAIRO Anal. Numér., Vol. 8, No. R-2, pp. 129-151, 1974. MR **51**:1540
- [11] F. Brezzi, J. Douglas (Jr.), L.D. Marini, *Two families of mixed finite elements for second order elliptic problems*, Numer. Math., Vol. 47, pp. 217-235, 1985. MR **87g**:65133
- [12] F. Brezzi, M. Fortin, *Mixed and hybrid finite element methods*, Springer-Verlag, 1991. MR **92d**:65187
- [13] A. Buffa, M. Costabel, C. Schwab, *Boundary element methods for Maxwell's equations on non-smooth domains*, Numer. Math., Vol. 92, No. 4, pp. 679-710, 2002.
- [14] M. Cessenat, *Mathematical methods in electromagnetism, Linear theory and applications*, World Scientific Publishing Co., River Edge, NJ, 1996. MR **97j**:78001
- [15] S.H. Christiansen, F. Béreux, J.-C. Nédélec, J.-P. Martinaud, *Algorithme de simulation électromagnétique, notamment des performances d'une antenne*, Patent by Thomson-CSF Detexis, Reg. No. 0007456 at INPI, dated June 9, 2000.
- [16] S.H. Christiansen, J.-C. Nédélec, *Des préconditionneurs pour la résolution numérique des équations intégrales de frontière de l'électromagnétisme*, C. R. Acad. Sci. Paris, Sér. I Math., Vol. 331, No. 9, pp. 733-738, 2000. MR **2001i**:78022
- [17] D.L. Colton, R. Kress, *Integral equation methods in scattering theory*, John Wiley & Sons, New York, 1983. MR **85d**:35001

- [18] M. Costabel, *Boundary integral operators on Lipschitz domains: elementary results*, SIAM J. Math. Anal., Vol. 19, No. 3, pp. 613-626, 1988. MR **89h**:35090
- [19] L. Demkowicz, *Asymptotic convergence in finite and boundary element methods, Part 1, Theoretical results*, Comput. Math. Appl., Vol. 27, No. 12, pp. 69-84, 1994, *Part 2, The LBB constant for rigid and elastic scattering problems*, Comput. Math. Appl., Vol. 28, No. 6, pp. 93-109, 1994. MR **95h**:65080, MR **95k**:65106
- [20] L. Demkowicz, P. Monk, C. Schwab, L. Vardapetyan, *Maxwell eigenvalues and discrete compactness in two dimensions*, Comput. Math. Appl., Vol. 40, No. 4-5, pp. 589-605, 2000. MR **2002e**:65166
- [21] G.C. Hsiao, R.E. Kleinman, *Mathematical foundations of error estimates in numerical solutions of integral equations in electromagnetics*, IEEE Trans. Ant. Prop., Vol. 47, No. 3, pp. 316-328, 1997. MR **98a**:78001
- [22] J.D. Jackson, *Classical electrodynamics, Second edition*, John Wiley & Sons, New York-London-Sydney, 1975. MR **55**:9721
- [23] T. Kato, *Perturbation theory for linear operators, Second edition*, Springer-Verlag, Berlin-New York 1976. MR **53**:11389
- [24] F. Kikuchi, *On a discrete compactness property for the Nédélec finite elements*, J. Fac. Sci. Univ. Tokyo, Sect. 1A Math., Vol. 36, pp. 479-490, 1989. MR **91h**:65173
- [25] J.-L. Lions, E. Magenes, *Problèmes aux limites non homogènes et applications, Vol. I, Vol. II*, Dunod, Paris, 1968. MR **40**:512, MR **40**:513
- [26] J.-C. Nédélec, *Curved finite element methods for the solution of singular integral equations on surfaces in \mathbb{R}^3* , Comput. Methods Appl. Mech. Engrg., Vol. 8, No. 1, pp. 61-80, 1976. MR **56**:13741
- [27] J.-C. Nédélec, *Computation of eddy currents on a surface in \mathbb{R}^3 by finite element methods*, SIAM J. Numer. Anal., Vol. 15, No. 3, pp. 580-594, 1978. MR **58**:14409
- [28] J.-C. Nédélec, *Éléments finis mixtes incompressibles pour l'équation de Stokes dans \mathbb{R}^3* , Numer. Math., Vol. 39, pp. 97-112, 1982. MR **83g**:65111
- [29] J.-C. Nédélec, *Acoustic and Electromagnetic Equations, Integral Representations for Harmonic Problems*, Springer-Verlag, 2001. MR **2002c**:35003
- [30] R.A. Nicolaides, *Existence, uniqueness and approximation for generalized saddle point problems*, SIAM J. Numer. Anal., Vol. 19, No. 2, pp. 349-357, 1982. MR **83d**:49023
- [31] L. Paquet, *Problèmes mixtes pour le système de Maxwell*, Ann. Fac. Sci. Toulouse Math., Vol. 4, No. 2, pp. 103-141, 1982. MR **84e**:58075
- [32] S.S.M. Rao, D.R. Wilton, A.W. Glisson, *Electromagnetic scattering by surfaces of arbitrary shape*, IEEE Trans. Ant. Prop. AP-30, pp. 409-418, 1982.
- [33] P.A. Raviart, J.-M. Thomas, *A mixed finite element method for 2nd order elliptic problems*, pp. 292-315, in I. Galligani, E. Magenes (eds.), "Mathematical aspects of the finite element method", Lecture Notes in Math., Vol. 606, Springer-Verlag, Berlin and New York, 1977. MR **58**:3547
- [34] J.E. Roberts, J.-M. Thomas, *Mixed and hybrid methods*, pp. 523-640, in P.G. Ciarlet, J.-L. Lions (eds.), "Handbook of numerical analysis, Vol. II, Finite element methods (Part 1)", North-Holland, 1991.
- [35] W. Rudin, *Functional analysis, Second edition*, McGraw-Hill, New York, 1991. MR **92k**:46001
- [36] A.H. Schatz, *An observation concerning Ritz-Galerkin methods with indefinite bilinear forms*, Math. Comp., Vol. 28, pp. 959-962, 1974. MR **51**:9526
- [37] G. Strang, *Variational crimes in the finite element method*, pp. 689-710, in A.K. Aziz (ed.), "The mathematical foundations of the finite element method with applications to partial differential equations", Academic Press, New York, 1972. MR **54**:1668
- [38] M. Taylor, *Partial differential equations, Vol. I Basic theory, Vol. II Qualitative studies in linear equations*, Springer-Verlag, New York, 1996. MR **98b**:35002b, MR **98b**:35003
- [39] I. Terrasse, *Résolution mathématique et numérique des équations de Maxwell instationnaires par une méthode de potentiels retardés*, PhD thesis, Ecole Polytechnique, 1993.
- [40] W.L. Wendland, *Strongly elliptic boundary integral equations*, pp. 511-561, in A. Iserles, M. Powell (eds.), "The state of the art in numerical analysis (Birmingham 1986)", Inst. Math. Appl. Conf. Ser. New Ser. 9, Oxford Univ. Press, New York, 1987. MR **88m**:65209