

PERFECT PARALLELEPIPEDS EXIST

JORGE F. SAWYER AND CLIFFORD A. REITER

ABSTRACT. There are parallelepipeds with edge lengths, face diagonal lengths and body diagonal lengths that are all positive integers. In particular, there is a parallelepiped with edge lengths 271, 106, 103, minor face diagonal lengths 101, 266, 255, major face diagonal lengths 183, 312, 323, and body diagonal lengths 374, 300, 278, 272. Focused brute force searches give dozens of primitive perfect parallelepipeds. Examples include parallelepipeds with up to two rectangular faces.

1. INTRODUCTION

A famous open problem in Number Theory is whether there exists a perfect cuboid. That is, is there a rectangular box in \mathbb{R}^3 with positive integer edge lengths, face diagonal lengths and body diagonal lengths [2, 3]? In [2] Richard Guy poses the weaker question of whether there exist perfect parallelepipeds in \mathbb{R}^3 . A perfect parallelepiped is a parallelepiped with edge lengths, face diagonal lengths and body diagonal lengths that are all positive integers. Previous attempts at finding perfect parallelepipeds focused on using rational coordinates [1, 6, 8]. Here we show that perfect parallelepipeds exist by giving examples and we describe a technique using necessary conditions within brute force searches that check at the last stage whether proposed perfect parallelepipeds can be realized in \mathbb{R}^3 .

2. THERE IS A PERFECT PARALLELEPIPED

While we will discuss our search strategy in the next section, it is straightforward to exhibit and verify that a perfect parallelepiped exists, which is our main result. We call the shorter diagonal of a parallelogram the minor diagonal and the longer diagonal the major diagonal. These will be the same for a rectangle.

Theorem 1. *There is a perfect parallelepiped with edge lengths 271, 106, 103, minor face diagonal lengths 101, 266, 255, major face diagonal lengths 183, 312, 323, and body diagonal lengths 374, 300, 278, 272.*

Proof. Consider the parallelepiped with edge vectors given by $\vec{u} = \langle 271, 0, 0 \rangle$, $\vec{v} = \langle \frac{9826}{271}, \frac{60\sqrt{202398}}{271}, 0 \rangle$, $\vec{w} = \langle \frac{6647}{271}, \frac{143754}{271}\sqrt{\frac{42}{4819}}, 66\sqrt{\frac{8358}{4819}} \rangle$. Direct computation verifies that $\|\vec{u}\| = 271$, $\|\vec{v}\| = 106$, $\|\vec{w}\| = 103$, $\|\vec{u}-\vec{v}\| = 255$, $\|\vec{u}-\vec{w}\| = 266$, $\|\vec{v}-\vec{w}\| = 101$, $\|\vec{u}+\vec{v}\| = 323$, $\|\vec{u}+\vec{w}\| = 312$, $\|\vec{v}+\vec{w}\| = 183$, $\|\vec{u}+\vec{v}+\vec{w}\| = 374$, $\|\vec{u}+\vec{v}-\vec{w}\| = 300$, $\|\vec{u}-\vec{v}+\vec{w}\| = 278$, and $\|-\vec{u}+\vec{v}+\vec{w}\| = 272$. \square

Received by the editor November 16, 2009 and, in revised form, December 3, 2009.

2010 *Mathematics Subject Classification.* Primary 11D09.

The support of a Lafayette EXCEL grant is appreciated.

©2010 American Mathematical Society
Reverts to public domain 28 years from publication

A *Mathematica*TM script verifying those computations may be found at [7]. Additional examples of perfect parallelepipeds may also be found there. These include parallelepipeds with two rectangular faces. The parallelepiped with edge vectors $\vec{u} = \langle 1120, 0, 0 \rangle$, $\vec{v} = \langle 0, 1035, 0 \rangle$, $\vec{w} = \langle 0, \frac{46548}{115}, \frac{12}{115}\sqrt{49755859} \rangle$ has that form. In particular, it has edge lengths $\|\vec{u}\| = 1120$, $\|\vec{v}\| = 1035$ and $\|\vec{w}\| = 840$. The rectangular face diagonal lengths are $\|\vec{u} + \vec{v}\| = \|\vec{u} - \vec{v}\| = 1525$ and $\|\vec{u} + \vec{w}\| = \|\vec{u} - \vec{w}\| = 1400$ and the other face has diagonal lengths $\|\vec{v} - \vec{w}\| = 969$ and $\|\vec{v} + \vec{w}\| = 1617$. The body diagonals lengths are $\|\vec{u} + \vec{v} + \vec{w}\| = \|\vec{u} - \vec{v} - \vec{w}\| = 1967$ and $\|\vec{u} + \vec{v} - \vec{w}\| = \|\vec{u} - \vec{v} + \vec{w}\| = 1481$.

3. THE SEARCH

First, we observe that the major diagonal of a parallelogram can be expressed in terms of the edges and the minor diagonal. That can be used to facilitate the search for perfect parallelograms; that is, parallelograms with edge and diagonal lengths that are all positive integers.

Lemma 2. *Let x_1, x_2 , and d_{12} be positive integers with $1 \leq x_2 \leq x_1$ and $x_1 - x_2 < d_{12} \leq \sqrt{x_1^2 + x_2^2}$. Then the parallelogram with edge length x_1 and x_2 and minor diagonal length d_{12} is perfect if and only if $2x_1^2 + 2x_2^2 - d_{12}^2$ is a square.*

Proof. Let \vec{u} and \vec{v} be edge vectors for the parallelogram so that $\|\vec{u}\| = x_1$, $\|\vec{v}\| = x_2$, $\|\vec{u} - \vec{v}\| = d_{12}$, then the result follows from observing that $\|\vec{u} + \vec{v}\|^2 = 2\|\vec{u}\|^2 + 2\|\vec{v}\|^2 - \|\vec{u} - \vec{v}\|^2$. \square

The search technique that we used determined by brute force all such x_1, x_2, d_{12} where x_1 was below some bound. Then all non-oblique assemblies of three such perfect parallelograms with matching pairs of edges were considered as possible perfect parallelepipeds. The search was implemented in J [4].

Whether the body diagonals were of integer length was determined by the following lemma. We use d_{ij} to denote the minor diagonal length of the parallelogram with edges i and j . The body diagonal with edge i having negative contribution is denoted m_i , $1 \leq i \leq 3$ and m_4 denotes the length of the body diagonal when all edges contribute positively.

Lemma 3. *Suppose there is a parallelepiped with edge lengths x_1, x_2 , and x_3 and minor face diagonal lengths d_{12}, d_{13} and d_{23} . Then the square of the body diagonal lengths are $m_1^2 = -x_1^2 + x_2^2 + x_3^2 + d_{12}^2 + d_{13}^2 - d_{23}^2$, $m_2^2 = x_1^2 - x_2^2 + x_3^2 + d_{12}^2 - d_{13}^2 + d_{23}^2$, $m_3^2 = x_1^2 + x_2^2 - x_3^2 - d_{12}^2 + d_{13}^2 + d_{23}^2$, and $m_4^2 = 3x_1^2 + 3x_2^2 + 3x_3^2 - d_{12}^2 - d_{13}^2 - d_{23}^2$.*

Proof. Let $\vec{u}, \vec{v}, \vec{w}$ be edge vectors for the parallelepiped such that $\|\vec{u}\| = x_1$, $\|\vec{v}\| = x_2$, $\|\vec{w}\| = x_3$, $\|\vec{u} - \vec{v}\| = d_{12}$, $\|\vec{u} - \vec{w}\| = d_{13}$, $\|\vec{v} - \vec{w}\| = d_{23}$. Note that $2\vec{u} \cdot \vec{v} = x_1^2 + x_2^2 - d_{12}^2$ and likewise for the other dot products. We see that

$$\begin{aligned} \|\vec{u} + \vec{v} + \vec{w}\|^2 &= \|\vec{u}\|^2 + \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\vec{u} \cdot \vec{v} - 2\vec{u} \cdot \vec{w} + 2\vec{v} \cdot \vec{w} \\ &= x_1^2 + x_2^2 + x_3^2 - (x_1^2 + x_2^2 - d_{12}^2) - (x_1^2 + x_3^2 - d_{13}^2) \\ &\quad + (x_2^2 + x_3^2 - d_{23}^2) \\ &= -x_1^2 + x_2^2 + x_3^2 + d_{12}^2 + d_{13}^2 - d_{23}^2 \end{aligned}$$

as desired. The other cases are similar. \square

Our search quickly located triples of perfect parallelograms with matching edge lengths $x_1, x_2,$ and x_3 and minor diagonal lengths d_{12}, d_{13} and d_{23} that also had all four proposed body diagonals m_1, m_2, m_3 and m_4 of positive integer length. For example, the smallest such is given by $x_1 = 115, x_2 = 106, x_3 = 83, d_{12} = 31, d_{13} = 58$ and $d_{23} = 75$. However, these perfect parallelograms cannot be realized as a parallelepiped in \mathbb{R}^3 .

The following lemma gives the final criterion necessary for the assembly to be realizable. We let θ_{ij} denote the angle between edges x_i and x_j in the triangle with sides x_i, x_j and d_{ij} and let c_{ij} denote the cosine of that angle. Note that $c_{ij} = \cos(\theta_{ij}) = \frac{x_i^2 + x_j^2 - d_{ij}^2}{2x_i x_j}$ and by our choice of minor diagonal $0 \leq c_{ij} < 1$.

Lemma 4. *An edge-matched assembly of three perfect parallelograms with edge lengths $x_1, x_2,$ and x_3 and minor diagonal lengths d_{12}, d_{13} and d_{23} can be assembled in \mathbb{R}^3 into a parallelepiped if $c_{12}^2 + c_{13}^2 + c_{23}^2 < 1 + 2c_{12}c_{13}c_{23}$.*

Proof. Let $\vec{u} = x_1\langle 1, 0, 0 \rangle, \vec{v} = x_2\langle c_{12}, \sqrt{1 - c_{12}^2}, 0 \rangle, \rho = \frac{c_{23} - c_{12}c_{13}}{\sqrt{1 - c_{12}^2}\sqrt{1 - c_{13}^2}}$, and $\vec{w} = x_3\langle c_{13}, \rho\sqrt{1 - c_{13}^2}, \sqrt{1 - \rho^2}\sqrt{1 - c_{13}^2} \rangle$. Direct computation shows that the parallelepiped generated by $\vec{u}, \vec{v}, \vec{w}$ realizes the parallelepiped with desired edges and minor diagonals provided that $-1 < \rho < 1$. Note that $\rho = \pm 1$ would yield a degenerate parallelepiped. A *Mathematica*TM script verifying those computations may be found at [7]. The condition that $-1 < \rho < 1$ is equivalent to $\rho^2 < 1$ which is equivalent to $(c_{23} - c_{12}c_{13})^2 < (1 - c_{12}^2)(1 - c_{13}^2)$ and simplifies to the required inequality. □

The above lemma describes realizability using non-oblique assemblies at one vertex. Note that moving along any edge of such a perfect parallelepiped leads to a vertex with two angles becoming non-acute. Thus, configurations with an odd number of oblique angles at each vertex would be distinct from those above and these too exist [7]. At least the first two of those were first found by Randall Rathbun [5].

To give some sense of the number of edge-matched non-oblique configurations checked we offer sample statistics. When checking edges up to 3949 there were about 2×10^{10} non-oblique edge-matched configurations tested. Of those, about 9×10^7 satisfied one of the necessary body diagonal conditions from Lemma 3. About 1.7×10^6 satisfied two; 33403 satisfied three; 414 satisfied all four. Of those, 27 gave realizable perfect parallelepipeds.

We have established that perfect parallelepipeds exist, and some with two rectangular faces exist. The question of whether perfect cuboids exist remains open. Intermediate questions are also open. Is there a perfect parallelepiped with integer volume? Is there a perfect parallelepiped with rational coordinates?

REFERENCES

1. D. D’Argenio, C. Reiter, *Families of nearly perfect parallelepipeds*, JP Jour. Algebra Number Theory & Appl. 9 (2007), 105-111. MR2407809 (2009d:11049)
2. R. Guy, *Unsolved problems in number theory*, Third ed., Springer, Springer-Verlag, 2004. MR2076335 (2005h:11003)
3. J. Leech, *The rational cuboid revisited*, Amer. Math. Monthly. 84 (1977), 518-533. Corrections 85 (1978) 472. MR0447106 (56:5421)
4. Jsoftware, <http://www.jsoftware.com>.
5. Randall Rathbun, Private Communication.

6. C. Reiter, J. Tirrell, *Pursuing the perfect parallelepiped*, JP Jour. Algebra Number Theory & Appl. 6 (2006), 279-274. MR2283937 (2008g:11050)
7. J. Sawyer, C. Reiter, *Auxiliary materials for perfect parallelepipeds exist*, <http://www.lafayette.edu/~reiterc/nt/ppe/index.html>.
8. J. Tirrell, C. Reiter, *Matrix generations of the diophantine solutions to sums of $3 \leq n \leq 9$ squares that are square*, JP Jour. Algebra, Number Theory & Appl. 8 (2007), 69-80. MR2370192 (2008i:11046)

BOX 8681 FARINON CENTER, LAFAYETTE COLLEGE, EASTON, PENNSYLVANIA 18042
E-mail address: sawyerj@lafayette.edu

DEPARTMENT OF MATHEMATICS, LAFAYETTE COLLEGE, EASTON, PENNSYLVANIA 18042
E-mail address: reiterc@lafayette.edu