

S and the integrals of the second member are taken along the arcs $s^{(v)}$ of the spirals (S_n^v) from (2''). The proof, proceeding along the same lines as that of (6), is suppressed here.

REMARK. The explicit values of the elementary integrals (3) are, of course, well known; but we refrain purposely from using them, as they are not needed. It is, indeed, sufficient for our proofs to know that those integrals depend only on the exponents m, n and are independent of ϕ, θ , or r .

UNIVERSITY OF SASKATCHEWAN

ON THE DENSITY THEOREM

A. PAPOULIS

1. **Introduction.** Let F be a set on the plane and x a point of F . With $\{I_n\}$ an arbitrary sequence of intervals¹ containing the point x and with diameter tending to zero, we form the sequence $|F \cdot I_n|/|I_n|$.² It has been shown (see [1] and [2])³ that for almost⁴ all points x of F ,

$$(1) \quad \lim_{I_n} \frac{|F \cdot I_n|}{|I_n|} = 1.$$

If the sequence $\{I_n\}$ of intervals is replaced by a sequence of arbitrary rectangles with sides not necessarily parallel to the axes of coordinates, then the above ceases to be true. H. Busemann and W. Feller (see [1]) have shown that if the direction of some one of the sides of the rectangles $\{I_n\}$ varies within any nonzero angle, then (1) is no longer true for all sets F .

The purpose of the following is to show that even if the direction of the rectangles $\{I_n\}$ converging to the point x is fixed, then (1) is still not true for some sets, provided of course that the fixed direction may vary from point to point.

Received by the editors October 17, 1950.

¹ Rectangles with sides parallel to the coordinate axes.

² The number $|E|$ will mean the two-dimensional Lebesgue-measure of the set E .

³ Numbers in brackets refer to the references at the end of the paper.

⁴ By "almost all points x of a set E " we shall mean all points of E except for a set of measure zero; this will also be indicated by p.p.

2. **Part A.** We shall construct⁵ a set E included in the unit square Q and such that:

(2A) $|E| < 3\epsilon$, where ϵ is an arbitrary number; given $x \in Q$ there exists a sequence $\{I_n\}$ of rectangles containing the point x , with diameter tending to zero, and such that:

$$\limsup_{I_n} \frac{|E \cdot I_n|}{|I_n|} > \frac{1}{3}.$$

This construction will be given in §§3-7.

3. **Definition.** Given a plane set E , let $M(E)$ be the set of points x of the plane with the property that there exists a line-segment s containing x and such that

$$(3) \quad \frac{m_1(E \cdot s)}{m_1(s)} > \frac{1}{2}.$$

If E is a circle of radius R , then clearly $M(E)$ is a concentric circle of radius $3R$.

4. **Lemma.**⁷

$$\text{g.l.b.}_E \frac{|E|}{|M(E)|} = 0$$

as E ranges over all plane sets.

If E is a circle, then $|E|/|M(E)| = 1/9$; if E is a triangle, then $|E|/|M(E)| = 1/13$.

PROOF. It suffices to show that given an ϵ we can construct a set E such that

$$(4) \quad \frac{|E|}{|M(E)|} < \epsilon.$$

Consider the open isosceles triangle $S_0 = ABC$ of base $BC = a$, angle $BAC = \theta$, and altitude h , with an axis $\alpha - \alpha$ parallel to the y -axis. Clearly the points of the trapezoid $T_0 = BCED$, where DE is the line $y = -h$, belong to $M(S_0)$. Draw the lines $y = h - h'$ and $y = h + h'$

⁵ This construction is similar to the one given by H. Busemann and W. Feller in [1].

⁶ The number $m_1(s)$ will mean the linear measure of the set s .

⁷ The above lemma is based on a construction used by O. Perron in giving a simple solution to the Besicovitch-Kakeya problem; it was utilized also by H. Busemann and W. Feller in their treatment of the density theorem.

$$\begin{aligned} T_2 &\subset M(S_2), \\ |S_2| &= |S_0| \{1 + 2(1/2)^2 + 2(1/3)^2\}, \\ |T_2| &> |T_0| \{1 + 1/2 + 1/3\}. \end{aligned}$$

Repeating the above process on each of the four new triangles, whose base is part of BC , with $h' = h/4$, we obtain the open figure S_3 , consisting of S_2 and the eight added triangles, and the trapezoid $T_3 = BCE_3D_3$; as before

$$\begin{aligned} T_3 &\subset M(S_3), \\ |S_3| &= |S_0| \{1 + 2(1/2)^2 + 2(1/3)^2 + 2(1/4)^2\}, \\ |T_3| &> |T_0| \{1 + 1/2 + 1/3 + 1/4\}. \end{aligned}$$

At the n th such step with $h' = h/n$, we obtain the open figure S_n and the trapezoid $T_n = BCE_nD_n$; they satisfy the relation

$$\begin{aligned} T_n &\subset M(S_n), \\ (6) \quad |S_n| &= |S_0| \{1 + 2(1/2)^2 + \cdots + 2(1/n)^2\}, \\ |T_n| &> |T_0| \{1 + 1/2 + \cdots + 1/n\}. \end{aligned}$$

Since $1 + 1/2 + \cdots + 1/n + \cdots$ diverges, $1 + (1/2)^2 + \cdots + (1/n)^2 + \cdots$ converges, and $|T_0| \neq 0$, it follows that for a certain $n = N(\epsilon)$ we shall have

$$(7) \quad \frac{|S_N|}{|S_N + T_N|} < \epsilon,$$

and since $S_N + T_N \subset M(S_N)$, (4) has been established and the lemma is proved.

5. The set $S_N + T_N$ will be denoted by $B(\epsilon; \theta)$, where ϵ and θ are as in the previous section, and the set S_N by $S(\epsilon; \theta)$. Thus

$$(8) \quad \frac{|S(\epsilon; \theta)|}{|B(\epsilon; \theta)|} < \epsilon, \quad B(\epsilon; \theta) \subset M(S(\epsilon; \theta)).$$

REMARK 1. Obviously ϵ and θ can be taken arbitrarily small in the above construction.

6. **Construction of E .** In this section θ will be kept constant and will be omitted in the expressions $S(\epsilon; \theta)$ and $B(\epsilon; \theta)$.

First step. By $[B(\epsilon)]$ we shall mean the class of all sets similar to $B(\epsilon)$ and similarly placed, with diameter smaller than one. For every $x \in Q$, there exists a sequence of sets belonging to $[B(\epsilon)]$, containing the point x and with diameter tending to zero, hence (Vitali covering

theorem) there exists a sequence $\{B(\epsilon)\}$ of disjoint sets belonging to $[B(\epsilon)]$ and covering Q p.p. The sets $S(\epsilon)$ which are parts of $B(\epsilon)$ of the sequence $\{B(\epsilon)\}$ form another sequence $\{S(\epsilon)\}$ whose sum we denote by E_0 ; from (8) and the disjointness of the sets $B(\epsilon)$ of $\{B(\epsilon)\}$ we conclude that

$$(9) \quad |E_0| < \epsilon |Q| = \epsilon.$$

Second step. By $[B(\epsilon/2)]$ we shall mean the class of all sets similar to $B(\epsilon/2)$ and similarly placed with diameter less than $1/2$. As before there exists a sequence $\{B(\epsilon/2)\}$ of disjoint sets belonging to $[B(\epsilon/2)]$ and covering Q p.p.; the sets $S(\epsilon/2)$ which are parts of $B(\epsilon/2)$ of the sequence $\{B(\epsilon/2)\}$ form another sequence $\{S(\epsilon/2)\}$ whose sum we denote by E_1 ; as before,

$$(10) \quad |E_1| < \frac{\epsilon}{2} |Q| = \epsilon/2^k.$$

(k+1)th step. By $[B(\epsilon/2^k)]$ we shall mean the class of all sets similar to $B(\epsilon/2^k)$ and similarly placed, with diameter smaller than $2/2^k$. Again there exists a sequence $\{B(\epsilon/2^k)\}$ of disjoint sets belonging to $[B(\epsilon/2^k)]$ and covering Q p.p., and the sequence $\{S(\epsilon/2^k)\}$ of the $S(\epsilon/2^k)$ sets whose sum we denote by E_k ; as before

$$(11) \quad |E_k| < \epsilon/2^k$$

and so we continue.

Since we repeat the process an enumerable number of times, the set of points of Q which are not covered at least once in this process has a zero measure; hence it can be covered by an open set G of area smaller than ϵ ; with $E = G + \sum_{k=0}^{\infty} E_k$, we have

$$(12) \quad |E| \leq |G| + \sum_{k=0}^{\infty} |E_k| < \epsilon + \sum_{k=0}^{\infty} \frac{\epsilon}{2^k} = 3\epsilon.$$

7. We shall now show that E satisfies the requirements of §2. Suppose $x \in Q$; if x belongs to E , E being open, we can cover x with a circle $C_x \subset E$. If we take any sequence $\{I_n\}$ of intervals included in C_x and with diameter tending to zero, we have $|E \cdot I_n| / |I_n| = 1 > 1/3$; hence (2A) is true. Suppose x is not a point of E ; it suffices to show that given $\delta > 0$ there exists an interval I_k of diameter smaller than δ and such that

$$(13) \quad x \in I_k, \quad \frac{|E \cdot I_k|}{|I_k|} > \frac{1}{3}.$$

Take k such that $1/2^k < \delta$; since $x \notin E$ it follows that $x \notin G$, hence it is covered by one set $B(\epsilon/2^k)$ belonging to the sequence $\{B(\epsilon/2^k)\}$ of the $(k+1)$ th step. From the way these sets have been constructed, it follows that there exists a line segment s such that

$$(14) \quad x \in s, \quad s \subset B\left(\frac{\epsilon}{2^k}\right), \quad \frac{m_1(s \cdot S(\epsilon/2^k))}{m_1(s)} > \frac{1}{2}$$

where $S(\epsilon/2^k)$ is the set contained in $B(\epsilon/2^k)$. Since the maximum diameter of the sets $\{B(\epsilon/2^k)\}$ is smaller than $1/2^k$, and $s \subset B(\epsilon/2^k)$, we conclude that $ds < 1/2^k$.⁸ But $S(\epsilon/2^k) \subset E_k \subset E$, hence also

$$(15) \quad \frac{m_1(s \cdot E)}{m_1(s)} > \frac{1}{2}$$

and since E is open, we can find an interval I_k containing s and satisfying (15); and so our contention is proved.

REMARK 2. Since the orientation of the I_k 's can be taken within the angle θ , and since θ can be chosen arbitrarily small, it follows that the rectangles of the sequence $\{I_k\}$ in (2A) can be chosen with an orientation within an arbitrarily small angle.

8. With $F = Q - E$ we have from (2A) with $x \in F \subset Q$

$$\liminf_{I_n} \frac{|F \cdot I_n|}{|I_n|} < \frac{2}{3}.$$

Hence (1) is not true because now we have a sequence $\{I_n\}$ of rectangles and not of intervals.

9. **Part B.** We shall construct a set E included in the unit square Q and such that:

(2B) $|E| < 5\epsilon$ where ϵ is an arbitrary number; given $x \in Q$ there exists a sequence $\{I_n\}$ of rectangles of "fixed orientation," containing the point x , with diameter tending to zero, and such that

$$\limsup_{I_n} \frac{|E \cdot I_n|}{|I_n|} > \frac{1}{3}.$$

10. **Notations.** Suppose angle $BAC = \theta_1$, angle $EDF = \theta_2$, and γ a given direction; if from A we draw a line parallel to γ and it falls inside BAC , this will be indicated by writing $\gamma \subset \theta_1$. If from A we draw two lines parallel to DE and DF respectively and both fall inside BAC , this will be indicated by writing $\theta_2 \subset \theta_1$. By s_γ we shall mean a line segment parallel to the direction γ .

⁸ ds will mean the diameter of the set s .

11. **Outline of the method.** In the construction of Part A the direction of the line segments satisfying (14), and hence of the corresponding rectangles I_k , cannot be the same for every k for the following reason: for a given $x \in Q$ the segments s satisfying (14) for $k=0$ may have any direction $\gamma \subset \delta^0$; for $k=1$ may have any direction $\gamma \subset \delta^1$, for any k a direction $\gamma \subset \delta^k$; the angle δ^k depends on the location of x inside $B(\epsilon/2^k; \theta)$. It is possible that there is no direction $\gamma \subset \delta^k$ for every k . If we modify our second covering in such a manner that the covering sets $B(\epsilon/2; \theta^1)$ have an angle $\theta^1 \subset \delta^0$, then all the directions which satisfy (14) for $k=1$ must be included in θ^1 , hence in δ^0 , they therefore must satisfy (14) also for $k=0$; these possible directions form a nonzero angle θ^2 which is used for the following step. Thus for each point x we have a sequence of closed angles $\theta \supset \theta^1 \supset \theta^2 \supset \dots$, hence there exists a $\gamma \subset \theta^k$ and a segment s_γ satisfying (14) for every k ; and that is our objective.

12. We now come to our construction.

First step. We begin as in §6; we thus have the sequences $\{B(\epsilon; \theta)\}$, $\{S(\epsilon; \theta)\}$ and the set E_1 with

$$(9) \quad |E_1| < \epsilon.$$

For almost every $x \in Q$ there exists an s_{γ_0} of direction γ_0 and such that

$$(16) \quad x \in s, \quad ds < 1, \quad \frac{m_1(s \cdot E_1)}{m_1(s)} > \frac{1}{2}.$$

Since E_1 is open, it follows that all the possible directions γ obtained by continuous rotation of γ_0 , and for which the above is true, form a nonzero angle $w(x)$ depending on x . Consider the points x of Q such that $w(x) > \alpha$ where α is a given angle; they form a set P_α ; since to almost every x of Q there corresponds a nonzero angle $w(x)$, $\lim P_\alpha$ as α tends to zero must include almost all points of Q . Therefore given an ϵ there exists an α_1 such that

$$(17) \quad |Q - P_{\alpha_1}| < \epsilon.$$

The points of $Q - P_{\alpha_1}$ can be included in an open set G_0 of measure less than ϵ ; with $R_1 = E_1 + G_1$ we have

$$(18) \quad |R_1| < \epsilon + \epsilon = 2\epsilon.$$

Consider all points x of Q such that for every $\gamma \subset \theta$ there exists an s such that (16) is true; they form a set which we denote by $1A_0^1$; clearly all points of E_1 belong to $1A_0^1$ since their corresponding w equals 2π .

In the following, angles will be considered with their orientation; thus when we speak of the set $B(\epsilon; \phi)$ we shall mean a set as in §5 but with its axis α — α not parallel to the y -axis but to the bisector of the angle ϕ .

If we bisect θ , we obtain the angles θ_1^1 and θ_1^2 . We denote by $1A_1^t$ ($t=1, 2$) the set of points of $Q-1A_0^1$ such that for every $\gamma \subset \theta_1^t$ ($t=1, 2$) there exists an s_γ satisfying (16). We continue as follows:

We divide θ into 2^n ($n=2, 3, \dots$) equal angles θ_n^t ($t=1, 2, \dots, 2^n$) and denote by $1A_n^t$ ($t=1, \dots, 2^n$) the set of points of $Q - \{1A_0^1 + 1A_1^1 + 1A_1^2 + \dots + 1A_{n-1}^1 + \dots + 1A_{n-1}^{2^{n-1}}\}$ with the property that for every $\gamma \subset \theta_n^t$ there exists an s_γ satisfying (16). After n_1 such steps, with $\theta/2^{n_1+1} > \alpha_1$, all points of $Q-G_0$ will be included in the sets $1A_h^t$ ($h=0, 1, \dots, n_1; t=1, 2, \dots, 2^{n_1}$) (see (17)).

To avoid three indices let us enumerate the sets $1A_h^t$ and their corresponding angles θ_h^t writing $1A(i)$ and $1\theta(i)$ with $i=1, 2, \dots, N_1$.

*k*th step. Assuming that we have defined $(k-1)A(i)$, $(k-1)\theta(i)$, R_{k-1} , and E_{k-1} , $i=1, 2, \dots, N_{k-1}$, we define $kA(i)$, $k\theta(i)$, R_k , and E_k as follows: we cover p.p. the open kernel of $(k-1)A(i)$ (which has the same measure as $(k-1)A(i)$ as we can easily see) with a sequence $\{B(\epsilon/2^{k-1}; (k-1)\theta(i))\}$ of disjoint sets similar to $B(\epsilon/2^{k-1}; (k-1)\theta(i))$ and similarly placed. The sets S which are parts of the B 's form another sequence $\{S(\epsilon/2^{k-1}; (k-1)\theta(i))\}$ whose sum we denote by E_k ; clearly

$$(19) \quad |E_k| < \frac{\epsilon}{2^{k-1}}.$$

The subdivision of Q in the first step into the sets $1A_h^t$ we now perform on $(k-1)A(i)$ for every i , the only change being that instead of (16) we now must satisfy

$$(20) \quad x \in s, \quad ds < \frac{1}{2^{k-1}}, \quad \frac{m_1(s \cdot E_{k-1})}{m_1(s)} > \frac{1}{2}.$$

We thus obtain the sets $kA(i)_h^t$ and the angles $k\theta(i)_h^t$; we continue this process until we cover all points of the $(k-1)A(i)$ except for a set of measure less than $\epsilon/2^{k-1}$ (see (17)) which we include in an open set G_k of measure less than $\epsilon/2^{k-1}$; and with $R_k = E_k + G_k$ we have

$$(21) \quad |R_k| < \frac{\epsilon}{2^{k-1}} + \frac{\epsilon}{2^{k-1}} = \frac{\epsilon}{2^{k-2}}.$$

The sets $kA(i)_h^t$ and $k\theta(i)_h^t$ we again enumerate for all i , h , and t , and we thus obtain $kA(i)$ and $k\theta(i)$.

We continue similarly. In this process the part of Q which is not covered at least once by the sets $\{B\}$ has a zero measure; we cover it with a set G of area smaller than ϵ ; with $E = G + \sum_{k=1}^{\infty} R_k$, we have $|E| < \epsilon + \sum_{k=1}^{\infty} \epsilon/2^{k-2} = 5\epsilon$.

14. We shall now prove that E satisfies (2B). Suppose $x \in Q$ if $x \in G_k + G$; then we have no problem since x can be covered by a circle included in E .

Suppose then that $x \notin G_k + G$; then $x \in B(\epsilon/2^{k-1}; (k-1)\theta(i))$ for a certain i depending on x (and k), and every k . From the way the different coverings have been performed, it follows that

$$1\theta(i) \supset 2\theta(i) \supset \cdots \supset (k-1)\theta(i) \supset \cdots$$

and since these angles are closed there must exist a direction γ_x included in all of them and such that (20) is true for every k if the segment s is parallel to γ_x ; from that and the openness of E , (2B) follows easily.

The author is indebted to Professor A. S. Besicovitch for his many suggestions.

REFERENCES

1. H. Busemann and W. Feller, *Zur Differentiation der Lebesgueschen Integrale*, Fund. Math. vol. 22.
2. F. Riesz, *Sur les points de densité au sens fort*, Fund. Math. vol. 22.

UNIVERSITY OF PENNSYLVANIA