

THEOREMS ON SEPARABILITY IN RIEMANNIAN n -SPACE

PARRY MOON AND DOMINA EBERLE SPENCER

Despite the importance of separation of variables as a means for solving partial differential equations, no comprehensive study of separability conditions seems to have been made. We limit investigation to the Helmholtz equation

$$(1) \quad \nabla^2\phi + (k)^2\phi = 0$$

and the Laplace equation

$$(2) \quad \nabla^2\phi = 0$$

and consider separability in an n -space with Riemannian metric $g_{ij} = 0$, where $i \neq j$.

DEFINITION I. *If the assumption*

$$(3) \quad \phi = \prod_{i=1}^n U^i(u^i)$$

permits the separation of the partial differential equation into n ordinary differential equations, the equation is said to be simply separable.

DEFINITION II. *If the assumption*

$$(4) \quad \phi = \frac{1}{R(u^1, u^2, \dots, u^n)} \prod_{i=1}^n U^i(u^i)$$

permits the separation of the partial differential equation into n ordinary differential equations, and if $R \neq \text{const.}$, the equation is said to be an R -separable.

1. General case.

THEOREM I. *The necessary and sufficient conditions for simple separability of the Helmholtz equation, in Riemannian n -space with an orthogonal coordinate system and $\phi = \phi(u^1, u^2, \dots, u^n)$, are*

$$(5) \quad \begin{aligned} g_{ii} &= \frac{S}{M_{ii}}, \\ \frac{g^{1/2}}{S} &= \prod_{i=1}^n f_i(u^i), \end{aligned}$$

Presented to the International Congress of Mathematicians, Cambridge, Mass., September 6, 1950; received by the editors November 28, 1951.

where S is the *Stäckel determinant* [1]

$$S = \begin{vmatrix} \Phi_{11}(u^1) & \Phi_{12}(u^1) & \cdots & \Phi_{1n}(u^1) \\ \Phi_{21}(u^2) & \Phi_{22}(u^2) & \cdots & \Phi_{2n}(u^2) \\ \cdots & \cdots & \cdots & \cdots \\ \Phi_{n1}(u^n) & \Phi_{n2}(u^n) & \cdots & \Phi_{nn}(u^n) \end{vmatrix}$$

and M_{ii} is the cofactor of Φ_{ii} .

PROOF. The Helmholtz equation may be written

$$(6) \quad \frac{1}{g^{1/2}} \sum_{i=1}^n \frac{\partial}{\partial u^i} \left(\frac{g^{1/2}}{g_{ii}} \frac{\partial \phi}{\partial u^i} \right) + (k)^2 \phi = 0.$$

Evidently [2] a necessary condition for separation is

$$(7) \quad \frac{g^{1/2}}{g_{ii}} = f_i(u^i) \cdot F_i(u^1, \dots, u^{i-1}, u^{i+1}, \dots, u^n).$$

Substitute Equation (7) in Equation (6) and differentiate with respect to the separation constants $\alpha_1 = (k)^2, \alpha_2, \alpha_3, \dots, \alpha_n$. Then we obtain a set of n simultaneous algebraic equations,

$$(8) \quad \sum_{i=1}^n f_i F_i \Phi_{ij}(u^i) = \delta_j^1 g^{1/2},$$

where

$$\Phi_{ij}(u^i) = - \frac{1}{f_i(u^i)} \frac{\partial}{\partial \alpha_j} \left[\frac{1}{U^i} \frac{d}{du^i} \left(f_i \frac{dU^i}{du^i} \right) \right].$$

For $S \neq 0$, which is the only case of practical significance, the solution is

$$(9) \quad f_i F_i = \frac{g^{1/2}}{S} M_{ii}.$$

Combination of Equations (7) and (9) gives the conditions of Theorem I.

To show that these conditions are sufficient as well as necessary [3; 4], we substitute Equation (5) into Equation (6) and find that separation is always effected, with the separated equations

$$(10) \quad \frac{1}{f_i} \frac{d}{du^i} \left(f_i \frac{dU^i}{du^i} \right) + U^i \sum_{j=1}^n \alpha_j \Phi_{ij} = 0, \quad i = 1, 2, \dots, n.$$

THEOREM II. *The necessary and sufficient conditions for simple*

separability of the Laplace equation, in Riemannian n -space with an orthogonal coordinate system and $\phi = \phi(u^1, u^2, \dots, u^n)$, are

$$(11) \quad \frac{g_{ii}}{g_{jj}} = \frac{M_{j1}}{M_{i1}}, \quad \frac{g^{1/2}}{g_{ii}} = \left[\prod_{i=1}^n f_i(u^i) \right] M_{i1}.$$

PROOF. Procedure is exactly as in the proof of Theorem I. The only difference is that the algebraic equations (8) are now homogeneous:

$$(8a) \quad \sum_{i=1}^n f_i F_i \Phi_i(u^i) = 0.$$

The solution therefore gives only ratios

$$\frac{g_{ii}}{g_{jj}} = \frac{M_{j1}}{M_{i1}},$$

and we have the lighter restrictions of Equation (11) as compared with those of Equation (5). As in the previous proof, the conditions are both necessary and sufficient, and the separated equations are again given by Equation (10).

THEOREM III. *The necessary and sufficient conditions for R -separability of the Helmholtz equation, in Riemannian n -space with an orthogonal coordinate system and $\phi = \phi(u^1, u^2, \dots, u^n)$, are*

$$(12) \quad \sum_{i=1}^n \frac{1}{f_i g_{ii}} \frac{\partial}{\partial u^i} \left(f_i \frac{\partial R}{\partial u^i} \right) + AR = 0, \quad g_{ii} = \frac{S}{M_{i1}},$$

where A is a constant and $R(u^1, u^2, \dots, u^n)$ is determined by

$$(13) \quad \frac{g^{1/2}}{S} = \left[\prod_{i=1}^n f_i(u^i) \right] (R)^2.$$

PROOF. Substitution of Equation (4) into Equation (6) gives

$$(14) \quad \sum_{i=1}^n \frac{1}{U^i} \frac{\partial}{\partial u^i} \left[\frac{g^{1/2}}{g_{ii}} \left(\frac{1}{R} \frac{dU^i}{du^i} - \frac{U^i}{(R)^2} \frac{\partial R}{\partial u^i} \right) \right] + \frac{(k)^2 g^{1/2}}{R} = 0.$$

If terms in $\partial R / \partial u^i$ are to be separated from those in dU^i / du^i , it is necessary that

$$(15) \quad \frac{g^{1/2}}{g_{ii}} = (R)^2 f_i F_i.$$

Substitution of Equation (15) into Equation (14) and multiplication by an arbitrary function $Q(u^1, u^2, \dots, u^n)$ gives

$$(R)^2 Q \sum_{i=1}^n \frac{F_i}{U^i} \frac{d}{du^i} \left(f_i \frac{dU^i}{du^i} \right) + \alpha_1 g^{1/2} = 0,$$

where

$$\alpha_1 = (k)^2 Q - \frac{RQ}{g^{1/2}} \sum_{i=1}^n F_i \frac{\partial}{\partial u^i} \left(f_i \frac{\partial R}{\partial u^i} \right).$$

Thus a necessary condition for separability is

$$(16) \quad \sum_{i=1}^n \frac{1}{f_i g_{ii}} \frac{\partial}{\partial u^i} \left(f_i \frac{\partial R}{\partial u^i} \right) + \left(\frac{\alpha_1}{Q} - (k)^2 \right) R = 0.$$

Differentiation with respect to the separation constants leads to n simultaneous equations

$$(8b) \quad \sum_{i=1}^n f_i F_i \Phi_{i,j}(u^i) = \delta_j^1 \frac{g^{1/2}}{(R)^2 Q},$$

the solution of which ($S \neq 0$) is

$$f_i F_i = \frac{g^{1/2}}{(R)^2 Q} \frac{M_{i1}}{S}.$$

Comparison with Equation (15) shows that

$$(17) \quad g_{ii} = \frac{S}{M_{i1}} Q,$$

and

$$(18) \quad \frac{g^{1/2}}{QS} = \left(\prod_{i=1}^n f_i(u^i) \right) (R)^2.$$

But it is easily shown that $Q = \text{const.}$, which may be taken as unity without loss of generality, and $\alpha_1 = (k)^2 + A$, where A is an arbitrary constant. Thus Equations (16), (17), and (18) assume the form given in Theorem III.

If $k = 0$, we obtain immediately:

THEOREM IV. *The necessary and sufficient condition for R -separability of Laplace's equation, in Riemannian n -space with an orthogonal coordinate system and $\phi = \phi(u^1, u^2, \dots, u^n)$, is*

$$(19) \quad \sum_{i=1}^n \frac{1}{f_i g_{ii}} \frac{\partial}{\partial u^i} \left(f_i \frac{\partial R}{\partial u^i} \right) + \alpha_1 \frac{R}{Q} = 0;$$

where R and Q are defined by the relations

$$(20) \quad g_{ii} = \frac{S}{M_{ii}} Q, \quad \frac{g^{1/2}}{QS} = \left[\prod_{i=1}^n f_i(u^i) \right] (R)^2.$$

2. **Degenerate cases.** In most physical applications, ϕ is not a function of all n coordinates but depends on a smaller number, say m . Without loss of generality, we number the coordinates $1, 2, \dots, m, \dots, n$. Then the Helmholtz equation becomes

$$\frac{1}{g^{1/2}} \sum_1^m \frac{\partial}{\partial u^i} \left(\frac{g^{1/2}}{g_{ii}} \frac{\partial \phi}{\partial u^i} \right) + (k)^2 \phi = 0$$

and the solution for simple separability is

$$\phi = U^1 \cdot U^2 \dots U^m;$$

but g_{ii} and $g^{1/2}$ may still contain all n variables, and

$$(21) \quad \frac{g^{1/2}}{g_{ii}} = f_i(u^i) \cdot F_i(u^1, u^2, \dots, u^{i-1}, u^{i+1}, \dots, u^n).$$

Separability conditions are considerably lightened:

THEOREM V. *The necessary and sufficient conditions for simple separability of the Helmholtz equation, in Riemannian n -space with an orthogonal coordinate system and $\phi = \phi(u^1, u^2, \dots, u^m)$, are*

$$(22) \quad \begin{aligned} g_{ii} &= \frac{S}{M_{ii}}, & i &= 1, 2, \dots, m, \\ \frac{g^{1/2}}{S} &= \left[\prod_{i=1}^m f_i(u^i) \right] \cdot \mathcal{F}(u^{m+1}, \dots, u^n), \end{aligned}$$

where

$$S = \begin{vmatrix} \Phi_{11} & \dots & \Phi_{1m} \\ \cdot & \cdot & \cdot \\ \Phi_{m1} & \dots & \Phi_{mm} \end{vmatrix}.$$

PROOF. Though the space remains n -dimensional, there are only m separation constants. Thus Equation (8) becomes

$$(8c) \quad \sum_{i=1}^m f_i F_i \Phi_{i,j}(u^i) = \delta_j^1 g^{1/2}$$

and the Stäckel matrix contains only m^2 terms. Procedure as in Theorem I results in Theorem V. Similarly, we have for the Laplace equation:

THEOREM VI. *The necessary and sufficient conditions for simple separability of the Laplace equation, in Riemannian n -space with an orthogonal coordinate system and $\phi = \phi(u^1, u^2, \dots, u^m)$, are*

$$(23) \quad \begin{aligned} \frac{g_{ii}}{g_{jj}} &= \frac{M_{j1}}{M_{i1}}, & i, j &= 1, 2, \dots, m, \\ \frac{g^{1/2}}{g_{ii}} &= \left[\prod_{j=1}^m f_j(u^j) \right] \cdot \mathcal{F}(u^{m+1}, \dots, u^n), \end{aligned}$$

where M_{i1} and M_{j1} are cofactors in the $m \times m$ Stäckel matrix of Theorem V.

As a special case, we have $m = 1$. The question of separation does not arise, but a one-dimensional solution is possible only under certain conditions. Evidently the requirements are obtained from Equations (7) and (6):

THEOREM VII. *The necessary and sufficient conditions that the Helmholtz equation have a one-dimensional solution $\phi = \phi(u^i)$, in Riemannian n -space with an orthogonal coordinate system, are*

$$(24) \quad \frac{g^{1/2}}{g_{ii}} = f_j(u^i) \cdot F_j(u^1, \dots, u^{i-1}, u^{i+1}, \dots, u^n), \quad g_{ii} = g_{ii}(u^i).$$

THEOREM VIII. *The necessary and sufficient condition that the Laplace equation have a one-dimensional solution $\phi = \phi(u^i)$, in Riemannian n -space with an orthogonal coordinate system, is that (21) holds.*

3. Euclidean 3-space. Separability conditions in euclidean 3-space are obtained directly from the theorems of §§1 and 2. In a number of special cases, however, the requirements are so greatly simplified that they should be mentioned. In any orthogonal cylindrical coordinate system in 3-space, with $g_{11} = g_{22}$, the Stäckel matrix may be written

$$[S] = \begin{bmatrix} 0 & \Phi_{12} & \Phi_{13} \\ 0 & \Phi_{22} & \Phi_{23} \\ 1 & 0 & 1 \end{bmatrix}.$$

The corresponding matrix for any orthogonal rotational system in 3-space, written with $g_{11} = g_{22}$, is

$$[S] = \begin{bmatrix} \Phi_{11} & \Phi_{12} & \Phi_{13} \\ \Phi_{21} & \Phi_{22} & \Phi_{23} \\ 0 & 0 & 1 \end{bmatrix}.$$

The following theorems are obvious consequences of §§1 and 2:

THEOREM IX. *The Helmholtz and Laplace equations are never R-separable in any orthogonal cylindrical coordinate system in euclidean 3-space.*

THEOREM X. *The necessary and sufficient condition for simple separability of the Helmholtz or the Laplace equation, in euclidean 3-space with an orthogonal cylindrical coordinate system, $g_{11}=g_{22}$, and $\phi = \phi(u^1, u^2, u^3)$, is that the metric coefficient be a separable sum:*

$$(25) \quad g_{11} = - [\Phi_{13}(u^1) + \Phi_{23}(u^2)].$$

THEOREM XI. *The necessary and sufficient conditions for simple separability of the Helmholtz equation, in euclidean 3-space with an orthogonal rotational coordinate system, $g_{11}=g_{22}$, and $\phi = \phi(u^1, u^2, u^3)$, are*

$$(26) \quad g_{11} = \Phi_{11}(u^1) + \Phi_{21}(u^2), \quad \frac{g_{11}}{g_{33}} = - [\Phi_{13}(u^1) + \Phi_{23}(u^2)],$$

$$g_{33} = [f_1(u^1) \cdot f_2(u^2)]^2.$$

THEOREM XII. *The necessary and sufficient conditions for simple separability of the Laplace equation, in euclidean 3-space with an orthogonal rotational coordinate system, $g_{11}=g_{22}$, and $\phi = \phi(u^1, u^2, u^3)$, are*

$$(27) \quad \frac{g_{11}}{g_{33}} = - [\Phi_{13}(u^1) + \Phi_{23}(u^2)], \quad g_{33} = [f_1(u^1) \cdot f_2(u^2)]^2.$$

THEOREM XIII. *Simple separability of the Laplace equation occurs in every orthogonal cylindrical coordinate system in euclidean 3-space with $\phi = \phi(u^1, u^2)$.*

THEOREM XIV. *The necessary and sufficient condition for simple separability of the Helmholtz equation, in euclidean 3-space with orthogonal cylindrical coordinates, $g_{11}=g_{22}$, and $\phi = \phi(u^1, u^2)$, is*

$$(28) \quad g_{11} = \Phi_{11}(u^1) + \Phi_{21}(u^2).$$

THEOREM XV. *The necessary and sufficient conditions for simple separability of the Helmholtz equation, in euclidean 3-space with an orthogonal rotational coordinate system, $g_{11}=g_{22}$, and $\phi = \phi(u^1, u^2)$, are*

$$(29) \quad g_{11} = \Phi_{11}(u^1) + \Phi_{21}(u^2), \quad g_{33} = [f_1(u^1) \cdot f_2(u^2)]^2.$$

THEOREM XVI. *The necessary and sufficient condition for simple separability of the Laplace equation, in euclidean 3-space with an*

orthogonal rotational coordinate system, $g_{11} = g_{22}$, and $\phi = \phi(u^1, u^2)$, is

$$(30) \quad g_{33} = [f_1(u^1) \cdot f_2(u^2)]^2.$$

THEOREM XVII. *The necessary and sufficient conditions for R-separability of the Helmholtz equation, in euclidean 3-space with an orthogonal rotational coordinate system, $g_{11} = g_{22}$ and $\phi = \phi(u^1, u^2)$, are*

$$(31) \quad g_{11} = \Phi_{11}(u^1) + \Phi_{21}(u^2), \quad \frac{g_{11}}{g_{33}} = \Omega_1(u^1) + \Omega_2(u^2),$$

where Ω_1 and Ω_2 are arbitrary functions.

THEOREM XVIII. *The necessary and sufficient condition for R-separability of the Laplace equation, in euclidean 3-space with an orthogonal rotational coordinate system, $g_{11} = g_{22}$, and $\phi = \phi(u^1, u^2)$, is*

$$(32) \quad \frac{g_{11}}{g_{33}} = \Omega_1(u^1) + \Omega_2(u^2).$$

BIBLIOGRAPHY

1. Paul Stäckel, *Über die Integration der Hamilton-Jacobischen Differentialgleichung mittels Separation der Variablen*, Habilitationsschrift, Halle, 1891.
 ———, *Sur une classe de problèmes de dynamique*, C.R. Acad. Sci. Paris vol. 116 (1893) p. 485.
 ———, *Über die Integration der Hamiltonschen Differentialgleichung mittels Separation der Variablen*, Math. Ann. vol. 49 (1897) p. 145.
2. H. P. Robertson, *Bemerkung über separierbare Systeme in der Wellenmechanik*, Math. Ann. vol. 98 (1927) p. 749.
3. L. P. Eisenhart, *Separable systems of Stäckel*, Ann. of Math. vol. 35 (1934) p. 284.
 ———, *Stäckel systems in conformal euclidean space*, Ann. of Math. vol. 36 (1935) p. 57.
4. P. M. Morse and H. Feshbach, *Methods of theoretical physics*, Cambridge, Technology Press, 1946.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY AND
 UNIVERSITY OF CONNECTICUT