

ERROR ESTIMATION IN THE WEINSTEIN METHOD FOR EIGENVALUES¹

H. F. WEINBERGER

1. **Introduction.** We are given the eigenvalues λ_n and eigenvectors u_n of a completely continuous positive operator L in a Hilbert space \mathfrak{H} . The problem is to determine the eigenvalues of the projection L' of L into a subspace \mathfrak{G} .²

The method of Weinstein [1]³ gives upper bounds for these eigenvalues. If (p_1, p_2, \dots) is any complete set of vectors in the space

$$(1) \quad \mathfrak{B} = \mathfrak{H} \ominus \mathfrak{G},$$

the m th intermediate problem is to determine the eigenvalues $\lambda_n^{(m)}$ of the projection of L into the space

$$(2) \quad \mathfrak{H} \ominus \{p_1, \dots, p_m\}.$$

By the minimax principle, we have

$$(3) \quad \lambda_n^{(m)} \geq \lambda'_n,$$

so that the solutions of the m th intermediate problem provide upper bounds for the eigenvalues λ'_n . The method of Weinstein consists of explicitly solving the m th intermediate problem in terms of the known eigenvalues and eigenvectors of L .

It has been shown (Aronszajn and Weinstein [1]; Aronszajn [1, p. 476; 2, pp. 30–35]) that for each fixed n , and for any complete sequence (p_1, p_2, \dots) ,

$$(4) \quad \lim_{m \rightarrow \infty} \lambda_n^{(m)} = \lambda'_n.$$

We are here concerned with the speed of this convergence, that is, with an estimate of the error

$$(5) \quad \lambda_n^{(m)} - \lambda'_n.$$

Presented to the Society, September 4, 1951; received by the editors December 10, 1951.

¹ This work was supported by the Office of Naval Research.

² We make the convention that the projection of an operator into a subspace is restricted to this subspace, so that the domain of L' is \mathfrak{G} .

³ The formulation of the method in Hilbert space as here presented was given for a special operator by Aronszajn and Weinstein [1], and for the general operator by Aronszajn [1; 2].

2. **The error estimate.** The upper bounds $\lambda_n^{(m)}$, and hence the errors (5), depend upon the sequence (p_1, p_2, \dots) . We obtain our error estimate by choosing⁴

$$(6) \quad p_n = \text{projection into } \mathfrak{B} \text{ of } u_n.$$

(p_1, p_2, \dots) is clearly complete in \mathfrak{B} since (u_1, u_2, \dots) is complete in \mathfrak{S} . We shall show that with the choice (6) of the vectors p_n ,

$$(7) \quad \lambda_n^{(m)} - \lambda'_n \leq \lambda_{m+1}.$$

Thus, the known eigenvalue λ_{m+1} is a *uniform* estimate of the error (5). Since L is completely continuous, this error estimate can be made arbitrarily small by choosing m sufficiently large.

In proving the inequality (7), we make use of the following inequality given by N. Aronszajn [1, p. 476, Corollary I']. For any m th intermediate problem,

$$(8) \quad \lambda_n^{(m)} \leq \lambda'_n + \mu,$$

where μ is the largest eigenvalue of the projection of L into $\mathfrak{B} \ominus \{p_1, \dots, p_m\}$, that is,

$$(9) \quad \mu = \max_{p \in \mathfrak{B} \ominus \{p_1, \dots, p_m\}} (Lp, p) / (p, p).$$

For a general sequence $\{p_n\}$, μ is unknown so that (8) does not give an error estimate. However, for the choice (6) of the vectors p_n we can estimate μ . We have, for p in $\mathfrak{B} \ominus \{p_1, \dots, p_m\}$,

$$(10) \quad (p, u_n) = (p, p_n) = 0, \quad n = 1, \dots, m.$$

Then, by the minimax principle,

$$(11) \quad (Lp, p) / (p, p) \leq \lambda_{m+1} \quad \text{for } p \in \mathfrak{B} \ominus \{p_1, \dots, p_m\}$$

and so, by (9),

$$(12) \quad \mu \leq \lambda_{m+1}.$$

Combining (12) with (8) gives the error estimate (7).

3. **Optimum property of the estimate.** The error estimate (7) is uniform with regard to the eigenvalues and moreover depends only on the eigenvalues λ_n of L . We now show that it is the best error estimate for the m th intermediate problem having these two properties.

⁴ It is easily verified that, if L and L' have no common eigenvectors, this sequence is a "suite privilégiée" in the terminology of Weinstein [1].

The explicit construction of the projection in (6) in specific cases will be discussed in another paper.

Suppose μ_m is such an error estimate. Since it depends only on the eigenvalues of L , it must be valid for all projections L' of L . Take for L' the projection into the space

$$(13) \quad \{u_r, u_{r+1}, \dots\}.$$

Obviously, we have for this L'

$$(14) \quad \lambda_1' = \lambda_r.$$

By the minimax principle, we find that for any m th intermediate problem

$$(15) \quad \lambda_1^{(m)} \geq \lambda_{m+1}.$$

Subtracting (14) from this gives

$$(16) \quad \lambda_1^{(m)} - \lambda_1' \geq \lambda_{m+1} - \lambda_r.$$

$$(17) \quad \mu_m \geq \lambda_1^{(m)} - \lambda_1',$$

by definition of μ_m , and hence by (16) and (17),

$$(18) \quad \mu_m \geq \lambda_{m+1} - \lambda_r.$$

But r is arbitrary and $\lambda_r \rightarrow 0$ as $r \rightarrow \infty$. Therefore,

$$(19) \quad \mu_m \geq \lambda_{m+1},$$

which proves the optimum property of λ_{m+1} .

4. Remarks on the Rayleigh-Ritz method. In the Rayleigh-Ritz method as generalized by Aronszajn [1; 2], the eigenvalues and eigenfunctions of the projection L' of L are assumed to be known and the eigenvalues of L are sought. When L' is the projection of L into a finite space, this reduces to the ordinary Rayleigh-Ritz method.

Aronszajn indicated that the same inequality from which (8) is obtained may also lead to an error estimate for the Rayleigh-Ritz method. However, the error term in this case involves the maximum of $(Lp, p)/(p, p)$ in a space outside the domain of L' , so that this estimate cannot be obtained in terms of the eigenvalues of L' .

In fact, no such estimate is possible. For the eigenvalues of L' in no way determine the behavior of L outside the domain of L' .

However, the situation is different if one knows the eigenvalues and eigenvectors not only of the projection L' of L into \mathfrak{G} but also of the extension⁵ L'' of L into a space \mathfrak{S} such that

⁵ That is, L'' is completely continuous and $L = L''$ in $\mathfrak{S} \cap \mathfrak{G}$.

$$(20) \quad \mathfrak{S} \supset \mathfrak{P} = \mathfrak{S} \ominus \mathfrak{G}.$$

We let the eigenvectors and eigenvalues of L'' be u_n'' and λ_n'' . We choose as the n th constraint vector p_n to be relaxed by the generalized Rayleigh-Ritz method the projection of u_n'' into \mathfrak{P} . The generalized Rayleigh-Ritz method enables us to calculate the eigenvalues $\lambda_n^{(m)}$ of the projection of L into $\mathfrak{G} \oplus \{p_1, \dots, p_m\}$ in terms of the eigenvalues and eigenvectors of L' . By the definition of eigenvalues, we find that for our choice of p_n ,

$$(21) \quad \max_{p \in \mathfrak{G} \oplus \{p_1, \dots, p_m\}} (Lp, p)/(p, p) \leq \lambda_{m+1}''.$$

Aronszajn's inequality then gives

$$(22) \quad \lambda_n \leq \lambda_n^{(m)} + \lambda_{m+1}'', \quad n = 1, 2, \dots$$

But the $\lambda_n^{(m)}$ are now lower bounds for the λ_n , so that

$$(23) \quad \lambda_n^{(m)} \leq \lambda_n \leq \lambda_n^{(m)} + \lambda_{m+1}''.$$

Thus, for the special case where the eigenvalue problems of L in both \mathfrak{G} and a space containing \mathfrak{P} are solved, we can obtain the uniform error estimate λ_{m+1}'' for the lower bounds given by the m th intermediate problem of the generalized Rayleigh-Ritz method. This is done by choosing as the n th constraint to be released the projection of the eigenvector u_n'' of L'' . If $\mathfrak{S} \supset \mathfrak{P}$, an alternative to the Weinstein method using projections and the Rayleigh-Ritz method is thus obtained.

BIBLIOGRAPHY

N. Aronszajn

1. *Rayleigh-Ritz and A. Weinstein methods for approximation of eigenvalues. I. Operators in a Hilbert space*, Proc. Nat. Acad. Sci. U.S.A. vol. 34 (1948) pp. 474-480, 594-601.
2. *Studies in eigenvalue problems. The Rayleigh-Ritz and Weinstein methods for approximation of eigenvalues. I. Operators in a Hilbert space*. Technical Report No. 1 (Oklahoma Agricultural and Mechanical College, Stillwater, Oklahoma, 1949).

A. WEINSTEIN

1. *Études des spectres des équations aux dérivées partielles de la théorie des plaques élastiques*, Mémorial des Sciences Mathématiques vol. 88 (1937).

N. Aronszajn and A. Weinstein

1. *On the unified theory of eigenvalues of plates and membranes*, Amer. J. Math. vol. 64 (1942) pp. 623-645.

UNIVERSITY OF MARYLAND