

## BIBLIOGRAPHY

1. H. Seifert and W. Threlfall, *Lehrbuch der Topologie*, Leipzig and Berlin, 1934.
2. A. M. Peiser, *Covering mappings*, Duke Math. J. vol. 10 (1943) pp. 305–307.
3. K. Borsuk, *Sur les rétractes*, Fund. Math. vol. 17 (1931) pp. 152–170.
4. C. W. Saalfrank, *Retraction properties for normal Hausdorff spaces*, Fund. Math. vol. 36 (1949) pp. 93–108.

RUTGERS UNIVERSITY AND  
LAFAYETTE COLLEGE

## DENSE IMBEDDING OF TOPOLOGICAL GROUPS

MORIKUNI GOTO<sup>1</sup>

The present article originated from the problem of determining when a continuous representation of a Lie group into another is open. Important cases of the problem have been discussed by A. Malcev<sup>2</sup> and the author,<sup>3</sup> and some of them have been extended to the case of more general groups by the author and H. Yamabe.<sup>4</sup>

Recently W. T. van Est obtained new results concerning the same problem on Lie groups.<sup>5</sup> Here I shall give an extension of his essential result to a more general case in a simpler way.

Let  $G$  be a locally compact connected group and let  $A(G)$  be the group of all continuous automorphisms of  $G$ . Let  $A(G)$  be topologized by the notion of uniform convergence in the wider sense.

Now let  $I(G)$  be the subgroup of  $A(G)$  composed of all inner automorphisms of  $G$ . We shall call  $G$  a (CA) group<sup>6</sup> if  $I(G)$  is a closed subgroup of  $A(G)$ .

**LEMMA.<sup>7</sup>** *Let  $G$  be a locally compact connected and locally connected group, and  $H$  a locally compact group. If  $\phi$  is a continuous isomorphism which maps  $G$  in an everywhere dense subgroup in  $H$ , then the following propositions hold:*

- (1)  $\phi(G)$  is an invariant subgroup of  $H$ .
- (2) Let  $h$  be an element of  $H$ . Let us consider the automorphism  $\sigma_h(x)$  defined by  $\sigma_h x = \phi^{-1}(h^{-1}\phi(x))h$  for  $x \in G$ . Then  $\sigma_h(x)$  is a continuous

Received by the editors October 6, 1952.

<sup>1</sup> The author's name will be hereafter spelled "Goto" instead of "Gotó" which has been used thus far.

<sup>2</sup> Malcev [4] in the bibliography.

<sup>3</sup> Goto [1].

<sup>4</sup> Goto and Yamabe [3]. See also [2].

<sup>5</sup> van Est [5], where he solved a prize problem (Wiskundig Genootschap Amsterdam, 1950), which had already been established in [1] and generalized in [3], independently of the author.

<sup>6</sup> The notion of a (CA) group is a generalization of van Est's (CA) Lie group.

<sup>7</sup> See [3] and [2].

automorphism of  $G$ .

(3) The correspondence  $h \rightarrow \sigma_h$  defines a continuous homomorphism from  $H$  into  $A(G)$ .

From this lemma we infer the following theorem.

**THEOREM.<sup>8</sup>** *Let  $G$  be a locally compact connected and locally connected (CA) group, and  $H$  a locally compact group. Suppose that the center  $Z$  of  $G$  is compact. If there exists a continuous isomorphism  $\phi$  mapping  $G$  into  $H$ , then the image  $\phi(G)$  is a closed subgroup of  $H$ .*

**PROOF.** Clearly we need consider only the case when  $H$  coincides with the closure  $\text{Cl}(\phi(G))$  of  $\phi(G)$ . In such a case we get a continuous homomorphism  $h \rightarrow \sigma_h$  from  $H$  into  $A(G)$  by virtue of the above lemma, and  $I(G)$  corresponds to  $\phi(G)$  under this homomorphism:  $\sigma\phi(G) = I(G)$ . Now since  $\sigma$  is continuous we get  $\sigma\text{Cl}(\phi(G)) \subseteq \text{Cl}(I(G))$  and thus  $\sigma(H) \subseteq I(G)$ .<sup>9</sup> The above relation means that for every  $h$  in  $H$  there exists an element  $g$  of  $G$  such that

$$\phi(g)^{-1}x\phi(g) = h^{-1}xh \quad \text{for all } x \in \phi(G).$$

Then  $h\phi(g)^{-1}$  commutes with every element of  $\phi(G)$ , whence  $h\phi(g)^{-1}$  is contained in the center  $A$  of  $H$ . Thus we have

$$H = \phi(G)A.$$

Since  $Z$  is compact,  $\phi(Z)$  is also compact, and the relation  $\phi(G) \cap A = \phi(Z)$  is obvious. Hence we have algebraically

$$H/\phi(Z) = \phi(G)/\phi(Z) \times A/\phi(Z)$$

where  $\times$  means the direct product of abstract groups.

Let us now consider the topological direct product group

$$L = G/Z \times A/\phi(Z)$$

of locally compact groups  $G/Z$  and  $A/\phi(Z)$ . It is easy to show that  $L$  can be covered by countable compact sets. Hence the continuous isomorphism from  $L$  onto  $H/\phi(Z)$ , obtained by extending the mapping  $\phi$  and the identity mapping of  $A/\phi(Z)$ , is necessarily open. Therefore  $\phi(G)/\phi(Z)$  is closed in  $H/\phi(Z)$ , whence  $\phi(G)$  is closed in  $H$ ; this completes the proof.

**COROLLARY.<sup>10</sup>** *Let  $G$  be a local Lie group. Then  $G$  always generates a*

<sup>8</sup> This theorem includes the essential theorems in both [3] and [5].

<sup>9</sup> The following argument is entirely the same as one in [3].

<sup>10</sup> Our theorem is superfluous for obtaining this corollary, which can be proved by the theorem of van Est only.

closed subgroup whenever  $G$  is imbedded in a Lie group as a local subgroup if and only if the following two conditions are satisfied:

- (1) The adjoint group of  $G$  is a closed linear group.
- (2) The center of the simply-connected group corresponding to  $G$  is a finite group.

PROOF. Necessity: (2) is already known.<sup>11</sup> Then as the adjoint group of  $G$  is locally isomorphic with  $G$ , it must be closed in the general linear group.

The sufficiency follows readily from the theorem.

#### BIBLIOGRAPHY

1. M. Goto, *Faithful representations of Lie groups I*, *Mathematica Japonicae* vol. 1 (1948).
2. ———, *On local Lie groups in a locally compact group*, *Ann. of Math.* vol. 54 (1951).
3. M. Goto and H. Yamabe, *On continuous isomorphisms of topological groups*, *Nagoya Math. J.* vol. 1 (1950).
4. A. Malcev, *On the theory of Lie groups in the large*, *Rec. Math. (Mat. Sbornik)* vol. 16 (1945).
5. W. T. van Est, *Dense imbeddings of Lie groups*, *Neder. Akad. Wetensch.* vol. 54 (1951).

THE INSTITUTE FOR ADVANCED STUDY

---

<sup>11</sup> See [1] and [5].