

## BANDS OF SEMIGROUPS

A. H. CLIFFORD

In studying a general semigroup  $S$ , a natural thing to do is to decompose  $S$  (if possible) into the class sum of a set  $\{S_\alpha; \alpha \in I\}$  of mutually disjoint subsemigroups  $S_\alpha$  such that (1) each  $S_\alpha$  belongs to some more or less restrictive type  $\mathcal{C}$  of semigroup, and (2) the product  $S_\alpha S_\beta$  of any two of them is wholly contained in a third:  $S_\alpha S_\beta \subseteq S_\gamma$ , for some  $\gamma \in I$  depending upon  $\alpha$  and  $\beta$ . We shall then say that  $S$  is a *band of semigroups of type*  $\mathcal{C}$ . If, for every  $\alpha$  and  $\beta$  in  $I$ ,  $S_\alpha S_\beta$  and  $S_\beta S_\alpha$  are both contained in the same  $S_\gamma$ , then we shall call  $S$  a *semilattice of semigroups of type*  $\mathcal{C}$ .

We shall also be concerned with the following specialization of the notion of band of semigroups. Suppose that  $I$  is the direct product  $J \times K$  of two classes  $J$  and  $K$ . The subsemigroups  $S_\alpha$  are then described by two subscripts:  $S_{i\kappa}$  ( $i \in J, \kappa \in K$ ). Suppose moreover that  $S_{i\kappa} S_{j\lambda} \subseteq S_{i\lambda}$  for all  $i, j \in J$  and all  $\kappa, \lambda \in K$ . We shall then call  $S$  a *matrix of semigroups of type*  $\mathcal{C}$ . The primary purpose of the present paper is to show (Theorem 4) that a band of semigroups of type  $\mathcal{C}$  is a semilattice of semigroups each of which is a matrix of semigroups of type  $\mathcal{C}$ .

The rest of the paper is devoted to giving necessary and sufficient conditions on a semigroup  $S$  that it be a band or a semilattice of (1) simple semigroups, (2) completely simple semigroups, and (3) groups. (Throughout this paper we use the term *simple* to mean *simple without zero*, i.e. a simple semigroup is one containing no proper two-sided ideal whatever.) For (1), we have the elegant condition,  $a \in Sa^2S$  for all  $a \in S$ , due to<sup>1</sup> Olaf Andersen [1]. If a semigroup  $S$  is a class sum of [completely] simple semigroups, it is also a semilattice of [completely] simple semigroups. But a class sum of groups need not be a band of groups, nor need a band of groups be a semilattice of groups; these three categories are characterized by Theorems 6, 7, and 8, respectively.

We note that a semigroup  $S$  is a "band of groups of order one" if and only if each element of  $S$  is idempotent. In this case we call  $S$  simply a "band," and consequently make the definition: a *band* is a semigroup every element of which is idempotent. By the same token, we define a *semilattice* to be a commutative band. A "matrix of groups

---

Presented to the Society, December 30, 1953; received by the editors November 20, 1953.

<sup>1</sup> (Added in proof.) This result has also been obtained recently by R. Croisot [8, p. 369].

of order one" will be called a "rectangular band." Thus we define a *rectangular band* to be the direct product  $J \times K$  of two classes  $J$  and  $K$ , with multiplication defined by

$$(1) \quad (i, \kappa)(j, \lambda) = (i, \lambda) \quad (\text{all } i, j \in J; \kappa, \lambda \in K).$$

The reason for the term is the following. Think of  $J \times K$  as a rectangular array of points, the point  $\alpha = (i, \kappa)$  lying in the  $i$ th row and  $\kappa$ th column. Then  $\alpha = (i, \kappa)$  and  $\beta = (j, \lambda)$  are opposite vertices of a rectangle of which the other two vertices are  $\alpha\beta = (i, \lambda)$  and  $\beta\alpha = (j, \kappa)$ .

**THEOREM 1.** *A semigroup  $S$  is a band of semigroups of type  $\mathcal{T}$  if and only if there exists a band  $I$  and a homomorphism  $\phi$  of  $S$  onto  $I$  such that the inverse image  $\phi^{-1}(\alpha)$  of each element  $\alpha$  of  $I$  is a subsemigroup  $S_\alpha$  of  $S$  of type  $\mathcal{T}$ .*

**PROOF.** Let  $S$  be a band of semigroups  $S_\alpha$  ( $\alpha \in I$ ) of type  $\mathcal{T}$ . Then to each pair of elements  $\alpha, \beta$  of the index class  $I$  corresponds a unique element  $\gamma$  of  $I$  such that  $S_\alpha S_\beta \subseteq S_\gamma$ . Define  $\alpha\beta = \gamma$ . Since  $S_\alpha S_\beta \cdot S_\gamma = S_\alpha \cdot S_\beta S_\gamma$ , and  $S_\alpha S_\alpha \subseteq S_\alpha$ , this multiplication is associative and idempotent, i.e.  $I$  becomes a band thereunder. Each element  $a$  of  $S$  belongs to exactly one  $S_\alpha$ , and, if we define  $\phi(a) = \alpha$ , the mapping  $\phi$  of  $S$  onto  $I$  is clearly a homomorphism with the property stated in the theorem. The converse is evident.

We shall use the expression, " $S$  is a band  $I$  of semigroups  $S_\alpha$  ( $\alpha \in I$ ) of type  $\mathcal{T}$ ," to indicate the situation described in Theorem 1.  $\phi$  will be called the natural homomorphism of  $S$  onto  $I$ .

We shall call a semigroup  $S$  *simple* if  $SaS = S$  for every  $a \in S$ , and *completely simple* if  $S$  is simple and contains a primitive idempotent (cf. Rees [3]). (An idempotent  $e$  is primitive if from  $f^2 = f$  and  $ef = fe = f$  we conclude  $f = e$ .) We shall not have occasion to consider semigroups with a zero element, and we consequently omit the phrase *without zero*. Rees [2] showed (in particular) that every completely simple semigroup is isomorphic with a "matrix semigroup  $S$  over a group  $G$ ," defined as follows. Let  $J$  and  $K$  be any two sets. Let  $S$  be the set of all triples  $(a; i, \kappa)$ , with  $a \in G$ ,  $i \in J$ ,  $\kappa \in K$ , and let the product of two elements of  $S$  be defined by

$$(2) \quad (a; i, \kappa)(b; j, \lambda) = (ap_{\kappa j}b; i, \lambda) \quad (a, b \in G; i, j \in J; \kappa, \lambda \in K),$$

where  $(p_{\kappa j})$  is a fixed  $K \times J$  matrix of elements  $p_{\kappa j}$  of  $G$ . Since Suschkewitsch [4] proved essentially the same theorem for finite  $S$ , we shall refer to this result as the Suschkewitsch-Rees Theorem.

The following theorem is simply a restatement, in the terminology of the present paper, of Theorem 2 of an earlier paper [5].

**THEOREM 2.** *A semigroup is a class sum of groups if and only if it is a semilattice of completely simple semigroups.*

We remark that if a semigroup  $S$  is a class sum of groups, then it is a class sum of mutually disjoint groups. This follows from Theorem 1 of [5], and the evident fact that  $S$  admits relative inverses. A similar remark holds if  $S$  is a class sum of simple or of completely simple semigroups, the former by Andersen's Theorem (stated below) and the latter by Theorem 2.

We next express in our present terminology a result due to McLean [6]. Call a band *nowhere commutative* if  $ab = ba$  implies  $a = b$ .

**THEOREM (McLean).** *Every band is a semilattice of nowhere commutative bands.*

Since a rectangular band is clearly nowhere commutative, the next theorem sharpens that of McLean.

**THEOREM 3.** *Every band is a semilattice of rectangular bands.*

**PROOF.** A band is a class sum of groups (of order one), and hence, by Theorem 2, is a semilattice of completely simple bands. Applying the Suschkewitsch-Rees Theorem to a completely simple band  $B$ , we see that the structure group of  $B$  must have order one, and hence the law (2) of multiplication reduces to (1).

We come now to the main result of the present paper.

**THEOREM 4.** *A band of semigroups of type  $\mathcal{G}$  is a semilattice of semigroups each of which is a matrix of semigroups of type  $\mathcal{G}$ .*

**PROOF.** Let  $S$  be a band  $I$  of semigroups  $S_\alpha$  ( $\alpha \in I$ ) of type  $\mathcal{G}$ . By Theorem 3,  $I$  is itself a semilattice  $P$  of rectangular bands  $I_\tau$  ( $\tau \in P$ ). Let  $\phi$  be the natural homomorphism of  $S$  onto  $I$ , and let  $\psi$  be that of  $I$  onto  $P$ . The mapping  $a \rightarrow \psi(\phi(a))$  is clearly a homomorphism of  $S$  onto  $P$ . The inverse image  $S'_\tau$  of an element  $\tau$  of  $P$  is the class sum of all the  $S_\alpha$  for which  $\psi(\alpha) = \tau$ , i.e. for which  $\alpha \in I_\tau$ . Describing the elements  $\alpha$  of  $I_\tau$  as pairs  $(i, \kappa)$  of elements of index classes  $J_\tau$  and  $K_\tau$ , as above, then we may write  $S_{i\kappa}$  for  $S_\alpha$  and obtain  $S_{i\kappa}S_{j\lambda} \subseteq S_{i\lambda}$  for all  $i, j \in J_\tau$  and all  $\kappa, \lambda \in K_\tau$ . Thus  $S'_\tau$  is a matrix of semigroups  $S_{i\kappa}$ , and the latter (being the  $S_\alpha$ ) are of type  $\mathcal{G}$ .

**THEOREM 5.** *A matrix of [completely] simple semigroups is [completely] simple.*

**PROOF.** Let  $S$  be a matrix of simple semigroups  $S_{i\kappa}$  ( $i \in J, \kappa \in K$ ). Let  $a$  and  $b$  be arbitrary elements of  $S$ ; we are to show that there

exist  $x, y$  in  $S$  such that  $xay = b$ . Suppose  $a \in S_{i\kappa}$  and  $b \in S_{j\lambda}$ . Then  $bab \in S_{j\lambda}$ . Since  $S_{j\lambda}$  is simple, there exist  $u, v$  in  $S_{j\lambda}$  such that  $u \cdot bab \cdot v = b$ . Hence we may take  $x = ub, y = bv$ .

Now suppose that the semigroups  $S_{i\kappa}$  are completely simple. To show that  $S$  is completely simple, it suffices to show that any idempotent element  $e$  of  $S_{i\kappa}$  is primitive in  $S$ , as well as in  $S_{i\kappa}$ . Let  $f$  be an idempotent element of  $S$  such that  $fe = ef = f$ . Suppose  $f \in S_{j\lambda}$ . Then  $fe \in S_{j\lambda}S_{i\kappa} \subseteq S_{j\kappa}$  and  $ef \in S_{i\kappa}S_{j\lambda} \subseteq S_{i\lambda}$ . Hence  $S_{j\kappa}$  and  $S_{i\lambda}$  must be the same as  $S_{j\lambda}$ , i.e.  $i = j$  and  $\kappa = \lambda$ . Thus  $f \in S_{i\kappa}$ , and since  $e$  is primitive in  $S_{i\kappa}$ , we conclude that  $f = e$ .

From Theorem 5 and the Suschkewitsch-Rees Theorem, it follows that a semigroup is a matrix of groups if and only if it is completely simple.

The following is Theorem 9.5 in [1]. (A proof may be constructed for it by suitably modifying the proof of Theorem 2.)<sup>2</sup>

**THEOREM (Croisot-Andersen).** *The following three assertions concerning a semigroup  $S$  are equivalent:*

- (A)  $S$  is a class sum of simple semigroups.
- (B)  $S$  is a semilattice of simple semigroups.
- (C)  $a \in Sa^2S$  for every  $a \in S$ .

Let us call two elements  $a, b$  of a semigroup  $S$   $r$ -associate ( $l$ -associate) if they generate the same right (left) ideal of  $S$ , and  $r \cap l$ -associate if they are both  $r$ - and  $l$ -associate. The following is Theorem 7 in [7]:

**THEOREM (J. A. Green).** *If an element  $a$  of a semigroup  $S$  is  $r \cap l$ -associate with  $a^2$ , then the set of all elements of  $S$  which are  $r \cap l$ -associate with  $a$  is a subgroup of  $S$ .*

**THEOREM 6.** *The following four assertions concerning a semigroup  $S$  are equivalent:*

- (A)  $S$  is a class sum of completely simple semigroups.
- (B)  $S$  is a class sum of groups.
- (C)  $S$  is a semilattice of completely simple semigroups.
- (D)  $a \in Sa^2 \cap a^2S$  for every  $a \in S$ .

**PROOF.** (A) implies (B), since every completely simple semigroup is a class sum of groups. (B) implies (C) by Theorem 2, and evidently (C) implies (A). (B) implies (D), for, if  $a \in S$ , then  $a$  and  $a^2$  both belong to the same subgroup  $G$  of  $S$ , so that  $a \in Ga^2 \cap a^2G \subseteq Sa^2 \cap a^2S$ . (D) implies (B), for, if  $a \in S$ , (D) implies that  $a$  and  $a^2$  are

<sup>2</sup> See footnote 1.

$r \cap l$ -associate, and hence  $a$  belongs to a subgroup of  $S$  by Green's theorem.

**THEOREM 7.** *A semigroup  $S$  is a band of groups if and only if it satisfies the following two conditions:*

- (1)  $a \in Sa^2 \cap a^2S$  for every  $a \in S$ ;
- (2) for every pair of elements  $a, b$  of  $S$ ,  $Sba = Sba^2$  and  $abS = a^2bS$ .

**PROOF.** Let  $S$  be a band of groups. In particular,  $S$  is a class sum of groups, and (1) follows as in the proof of Theorem 6. Let  $a, b \in S$ . From the fact that  $a$  and  $a^2$  belong to the same subgroup of  $S$ , and the assumption that  $S$  is a band of groups, we conclude that  $ba$  and  $ba^2$  belong to the same subgroup of  $S$ , and hence  $Sba = Sba^2$ . Similarly,  $abS = a^2bS$ .

Suppose conversely that (1) and (2) hold in  $S$ . By (1) and Green's Theorem,  $S$  is the class sum of groups  $G_\alpha$  ( $\alpha \in I$ ). Let  $a, b \in S$ . Let  $G_\alpha$  be the subgroup of  $S$  to which  $a$  belongs, let  $e_\alpha$  be the identity element of  $G_\alpha$ , and let  $a^{-1}$  be the inverse of  $a$  in  $G_\alpha$ . Replacing  $b$  by  $a^{-1}b$  in the second part of (2), we conclude  $e_\alpha bS = abS$ . If  $a'$  is any other element of  $G_\alpha$ , we conclude similarly that  $e_\alpha bS = a'bS$ , and hence  $abS = a'bS$ . Evidently  $Sa = Sa'$ , and hence  $Sab = Sa'b$ . Consequently  $ab$  and  $a'b$  are  $r \cap l$ -associate. Now, in a semigroup  $S$  which is the class sum of groups, two elements of  $S$  are  $r \cap l$ -associate if and only if they belong to the same subgroup of  $S$ . Consequently  $ab$  and  $a'b$  belong to the same subgroup of  $S$ . By the left-right dual of this argument,  $ba$  and  $ba'$  also belong to the same subgroup of  $S$ . Thus the relation of belonging to the same subgroup of  $S$  is a congruence relation, whence  $S$  is a band of groups.

**THEOREM 8.<sup>3</sup>** *A semigroup  $S$  is a semilattice of groups if and only if it satisfies the following two conditions:*

- (1)  $a \in Sa^2 \cap a^2S$  for every  $a \in S$ ;
- (2) if  $e$  and  $f$  are idempotent elements of  $S$ , then  $ef = fe$ .

**PROOF.** Assume that  $S$  is a semilattice  $I$  of groups  $G_\alpha$  ( $\alpha \in I$ ). Since  $S$  is in particular a class sum of groups, (1) follows as in the proof of Theorem 6. To show (2), let  $e$  and  $f$  be idempotent elements of  $S$ . Then  $e \in G_\alpha$  and  $f \in G_\beta$  for some  $\alpha, \beta \in I$ . Since  $G_\alpha$  and  $G_\beta$  are groups,  $e$  and  $f$  must be the identity elements thereof:  $e = e_\alpha$ ,  $f = e_\beta$ . Let  $\gamma = \alpha\beta$  ( $= \beta\alpha$  since  $I$  is by hypothesis commutative). Let  $e_\gamma$  be the identity element of  $G_\gamma$ . Since  $e_\alpha e_\beta \in G_\gamma$  we have  $e_\gamma \cdot e_\alpha e_\beta = e_\alpha e_\beta$ .

Now  $e_\gamma e_\alpha \in G_{\gamma\alpha} = G_\gamma$ , so that  $e_\gamma e_\alpha \cdot e_\gamma = e_\gamma e_\alpha$ . Hence

<sup>3</sup> (Added in proof.) This result has also been noted by R. Croisot [8, p. 375].

$$(e_\gamma e_\alpha)^2 = (e_\gamma e_\alpha \cdot e_\gamma) e_\alpha = e_\gamma e_\alpha e_\alpha = e_\gamma e_\alpha.$$

Since  $e_\gamma$  is the only idempotent element of  $G_\gamma$ , we conclude that  $e_\gamma e_\alpha = e_\gamma$ . Similarly,  $e_\gamma e_\beta = e_\gamma$ . Hence

$$e_\alpha e_\beta = e_\gamma e_\alpha e_\beta = e_\gamma e_\beta = e_\gamma.$$

Thus  $e_\alpha e_\beta = e_{\alpha\beta}$  for all  $\alpha, \beta$  in  $I$ , whence  $e_\beta e_\alpha = e_{\beta\alpha} = e_{\alpha\beta} = e_\alpha e_\beta$ .

Conversely, let  $S$  be a semigroup satisfying conditions (1) and (2). By (1) and Green's theorem,  $S$  is a class sum of groups. From Theorem 3 of [5] and condition (2) it then follows that  $S$  is a semilattice of groups; moreover, the theorem cited provides an explicit construction for any such  $S$ .

#### REFERENCES

1. Olaf Andersen, *Ein Bericht über die Struktur abstrakter Halbgruppen*, Thesis Hamburg, 1952, unpublished.
2. D. Rees, *On semi-groups*, Proc. Cambridge Philos. Soc. vol. 36 (1940) pp. 387–400.
3. ———, *Note on semi-groups*, Proc. Cambridge Philos. Soc. vol. 37 (1941) pp. 434–435.
4. A. Suschkewitsch, *Über die endlichen Gruppen ohne das Gesetz der eindeutigen Umkehrbarkeit*, Math. Ann. vol. 99 (1928) pp. 30–50.
5. A. H. Clifford, *Semigroups admitting relative inverses*, Ann. of Math. vol. 42 (1941) pp. 1037–1049.
6. David McLean, *Idempotent semigroups*, Amer. Math. Monthly vol. 61 (1954) pp. 110–113.
7. J. A. Green, *On the structure of semigroups*, Ann. of Math. vol. 54 (1951) pp. 163–172.
8. R. Croisot, *Demi-groupes inversifs et demi-groupes réunions de demi-groupes simples*, Ann. École Norm. (3) vol. 70 (1953) pp. 361–379. (Added in proof.)

THE JOHNS HOPKINS UNIVERSITY