

ON A LOCALLY COMPACT GROUP ACTING ON A MANIFOLD

PAUL S. MOSTERT¹

Introduction. Let G be a locally compact group, admissibly topologized [5], operating as a transitive transformation group on a manifold M . Using the notation of [1], let G_x be the subgroup of G which leaves the point x fixed. Montgomery [1] has proved the following result.

THEOREM (MONTGOMERY). *If G is a connected Lie group which acts transitively on a compact manifold M , and if G_x is connected, then G contains a compact subgroup which acts transitively on M .*

The purpose of this note is to show that the theorem is true when G is any locally compact group satisfying the second axiom of countability. It will follow, by [4], that the corollaries of [1] are true for the more general case.

1. A lemma. Arens [5] has proved that M is homeomorphic to G/G_x . The author [2; 3] has proved that G is a fibre bundle over G/G_x .

LEMMA. *Let G be a connected locally compact group satisfying the second axiom of countability and acting transitively on a manifold M . Then there exists a Lie group G^* acting transitively on M such that G^* is a factor group of G by a compact normal subgroup of G_x .*

PROOF. Let N be a compact normal subgroup of G such that G/N is a Lie group [6], and $N^* = N \cap G_x$. We shall show that $G/N^* = G^*$ is a Lie group which acts transitively on M . As mentioned above, the author [3] has shown that G/N^* is a fibre bundle over G/G_x with fibre G_x/N^* , and hence G/N^* is locally homeomorphic to $(G_x/N^*) \times K$ where K is homeomorphic to a euclidean space. Hence, G/N^* is a manifold. Now G/G_x is homeomorphic and isomorphic to $(G/N)/(G_x/N^*)$ so that by a proper choice of N , $\dim(N/N^*) = 0$. Since G/N^* is a fibre bundle over G/N , G/N^* is locally homeomorphic with $(N/N^*) \times L$ where L is a euclidean space. Hence, N/N^* is discrete. This implies that a subgroup of N^* (which we may assume to be N^* itself) has the property that G/N^* is a Lie group. Define, for $g^* \in G^*$, $g^* = gN^*$, $y \in M$,

Received by the editors February 12, 1954.

¹ This work was done under Contract N7-Onr-434, Task Order III, Navy Department.

$$g^*(y) = gN(y).$$

It is not difficult to show that G^* forms a transformation group on M with this definition. Since $N^* \subset G_x$, G^* is transitive.

2. The theorem. Under the hypothesis of the lemma, and if G_x is moreover connected, then Montgomery's theorem implies that G^* contains a compact subgroup H^* which acts transitively on M for M compact. If p is the natural projection $G \rightarrow G^*$, then $p^{-1}(H^*) = H$ is compact and acts transitively on M . Hence, we have proved the theorem as stated below.

THEOREM. *Let G be a connected locally compact group satisfying the second axiom of countability which acts transitively on a compact manifold. If G_x is connected, then G contains a compact subgroup which acts transitively on M .*

BIBLIOGRAPHY

1. D. Montgomery, *Simply connected homogeneous spaces*, Proc. Amer. Math. Soc. vol. 1 (1950) pp. 467-469.
2. P. S. Mostert, *Local cross sections in locally compact groups*, Proc. Amer. Math. Soc. vol. 4 (1953) pp. 645-649.
3. ———, *Sections in principal fibre spaces*, to appear.
4. ———, *On fibre spaces with 0-dimensional fibres*, Duke Math. J. vol. 21 (1954) pp. 67-74.
5. R. Arens, *Topologies for homeomorphism groups*, Amer. J. Math. vol. 68 (1946) pp. 593-610.
6. H. Yamabe, *A generalization of a theorem of Gleason*, Ann. of Math. vol. 58 (1953) pp. 351-365.

TULANE UNIVERSITY