

GENERALIZED THEOREMS OF DESARGUES FOR n -DIMENSIONAL PROJECTIVE SPACE

P. O. BELL

The purpose of the present note is to generalize to n dimensions the celebrated two-triangle theorem of Desargues and its converse.¹ The generalized theorems can be stated in simple forms which, nevertheless, suffice for each of the large number of special cases of the configurations involved. The terms concurrent and collinear will be used in the sense that n lines are concurrent if there is at least one point lying on all n of them, three points are collinear if there is at least one line containing them. The word simplex will denote an n -tuple of linearly independent points.

Two simplexes $A_1A_2 \cdots A_n, A'_1A'_2 \cdots A'_n$ will be said to be *in perspective from ϕ* if there exists a point ϕ such that $\phi A_1A'_1, \phi A_2A'_2, \cdots$, and $\phi A_nA'_n$ are each collinear triads. Each simplex determines a linear space of $n-1$ dimensions. Let k points of one simplex coincide with the corresponding points of the other simplex ($k \geq 0$). Without loss of generality, the simplexes may be named in a manner such that A_i coincides with A'_i if $i \leq k$. With these definitions a generalization of Desargues' theorem is

THEOREM 1. *If two simplexes $A_1, A_2 \cdots A_n, A'_1A'_2 \cdots A'_n$ are in perspective from a point ϕ , and if A_j does not coincide with A'_j for $k < j \leq n$, there exists a point P_{ij} common to $A_iA_j, A'_iA'_j$ ($k < i < j, i = 1, 2, \cdots, n-1$) which is linearly dependent upon the $j-i$ points $P_i, P_{i+1}, \cdots, P_{j-1}$, in which P_h denotes the point $P_{h, h+1}$.*

For $n=3$, the above stated theorem is obviously the theorem of Desargues.

For the proof of the theorem two cases are to be considered: Case (1). The point ϕ is distinct from each vertex of at least one of the simplexes. Case (2). The point ϕ coincides with a vertex of each simplex.

PROOF. CASE (1). Let ϕ be assumed to be distinct from each of the vertices A_1, A_2, \cdots, A_n . Let a fixed coordinate system be postulated in the join² of the linear spaces determined by the simplexes. The

Received by the editors June 22, 1954 and, in revised form, January 6, 1955.

¹ A statement of Desargues' theorem and of its converse may be found, for example, in Hodge and Pedoe, *Methods of algebraic geometry*, Cambridge University Press, 1947, p. 192.

² The join of two linear spaces is the smallest linear space which contains the two spaces.

property that the simplexes are in perspective from ϕ is that corresponding homogeneous coordinates of ϕ and the points of the simplexes are related by a system of linear equations which, on choosing multipliers of the points A'_i properly, assume the forms

$$\begin{aligned} A'_1 &= A_1, & A'_2 &= A_2, \dots, A'_k &= A_k, \\ A'_i &= \phi + \lambda_i A_i & & & (k < i \leq n). \end{aligned}$$

The points $P_{k+1}, P_{k+2}, \dots, P_{n-1}$ defined by

$$P_i = A'_i - A'_{i+1} = \lambda_i A_i - \lambda_{i+1} A_{i+2} \quad (k < i < n)$$

are points common to the corresponding pairs of lines $A_i A_{i+1}, A'_i A'_{i+1}$. Moreover, a point P_{ij} common to the pair of lines $A_i A_j, A'_i A'_j$ is given by

$$P_{ij} = A'_i - A'_j = \lambda_i A_i - \lambda_j A_j \quad (k < i < j < n).$$

In view of the expressions for the points P_i ($k < i < n$) it follows immediately that

$$(1) \quad P_{ij} = P_i + P_{i+1} + \dots + P_{j-1} \quad (k < i < j \leq n).$$

In particular,

$$P_{k+1 n} = P_{k+1} + P_{k+2} + \dots + P_{n-1}.$$

This completes the proof.

For the case of the theorem in which $k=0$, it is significant that the point P_{ij} common to the lines $A_i A_j, A'_i A'_j$ lies in a linear space of $j-i-1$ dimensions which is a subspace of the linear space S_{n-2} determined by the points P_1, P_2, \dots, P_{n-1} .³

In case $k \neq 0$, A_i coincides with A'_i ($i=1, 2, \dots, k$). The points P_{ij} ($i \leq k$) may therefore be defined by the relations

$$P_{ij} = P_i = A_i \quad (i = 1, 2, \dots, k, j \neq i).$$

Since the points P_{ij} for $i > k$ satisfy the relations (1), any point P_{ij} whatever lies in the linear space of $j-i-1$ dimensions determined by the points $P_i, P_{i+1}, \dots, P_{j-1}$, each point P_{ij} being either the point P_i or a linear combination of these $j-i$ points.

PROOF. CASE (2). The point ϕ coincides with a vertex of each simplex. Let the vertices which coincide with ϕ be A_l and A'_m . The sub-case in which $l=m$ can be disposed of as follows: Since k denotes, as in Case (1), the number of points A_i which coincide with their cor-

³ If the simplexes $A_1 A_2 \dots A_n, A'_1 A'_2 \dots A'_n$ are in distinct linear spaces S_{n-1}, S'_{n-1} it is, of course, obvious that the P 's lie in the intersection space S_{n-2} .

respondents A'_i , the index l will be one of the numbers $1, 2, \dots, k$. Without loss of generality, let the points A_1, A_l , and A'_1, A'_l be re-named by permuting their indices so that the vertices which coincide with ϕ are said to be the points A_1, A'_1 . Therefore, with properly selected multipliers for A_1 and $A'_1, A_1 = A'_1 = \phi$. Moreover, as in Case (1) of the theorem, A_h coincides with A'_h if $h \leq k$. In this case A_h is the common center of two sets of lines $A_h A_j, A'_h A'_j$ ($j \neq h$). Therefore, the points P_{hj} ($h \leq k$) may be defined as follows:

$$(2) \quad P_{hj} = A_h = P_h \quad (h \leq k, j > h).$$

Since A_i, A'_i are collinear with ϕ , when A_i does not coincide with A'_i , they satisfy linear relations which, on choosing multipliers of the points A'_i properly, assume the forms

$$A'_i = \phi + \lambda_i A_i \quad (i > k).$$

The points $P_{k+1}, P_{k+2}, \dots, P_{n-1}$ are therefore defined by

$$(3) \quad P_i = A'_i - A'_{i+1} = \lambda_i A_i - \lambda_{i+1} A_{i+1} \quad (k < i < n).$$

It follows that points common to pairs of lines $A_i A_j, A'_i A'_j$ are defined by

$$(4) \quad P_{ij} = A'_i - A'_j = \lambda_i A_i - \lambda_j A_j \quad (k < i < j < n).$$

In view of (2), (3), and (4), the following relations

$$P_{ij} = P_i + P_{i+1} + \dots + P_{j-1} \quad (k < i < j < n),$$

$$P_{hj} = P_h \quad (h \leq k, j > h)$$

hold. The proof of the theorem in this sub-case is complete.

For the proof of the remaining sub-case of the theorem, in which A_l coincides with A'_m in ϕ ($l \neq m$), it is evident at once that l and m must both be greater than k since $A_i = A'_i$ ($i \leq k$). (The distinguishing feature of this case is that the points of the simplexes which coincide in ϕ are not corresponding elements in the perspectivity.) Without loss of generality the names of the points A_l, A_{k+1} and those of the points A'_m, A'_{k+2} may be interchanged, so that the point A_{k+1} will be said to coincide with the point A'_{k+2} in ϕ .

As in the sub-case of the theorem proved above, points P_{hj} ($h \leq k$) may be defined by equations (2).

Moreover, since on selecting multipliers of A_{k+1} and A'_{k+2} properly, the relations $A_{k+1} = A'_{k+2} = \phi$ hold, it follows from the hypothesis that with a proper selection of multipliers of A'_i , we have

$$(5) \quad A'_i = \phi + \lambda_i A_i, \quad \lambda_{k+2} = 0 \quad (i > k, i \neq k + 1).$$

The points P_{ij} may, therefore, be defined by the relations

$$P_{ij} = A'_i - A'_j = \lambda_i A_i - \lambda_j A_j \quad (i, j > k, i \neq j).$$

On making use of the definitions of the points P_h ($h > k$) it follows that

$$P_{k+1 j} = \lambda_{k+1} A_{k+1} = P_{k+1} \quad (j > k + 1),$$

$$P_{k+2} = -\lambda_{k+3} A_{k+3}, \quad P_{ij} = P_i + P_{i+1} + \dots + P_{j-1} \quad (k + 1 < i < j).$$

The proof of the theorem is now complete.

Involved in the conclusion of Theorem 1 there are $(n - k - 2) \cdot (n - k - 3) / 2$ nontrivial linear relations of the form

$$P_{ij} = P_i + P_{i+1} + \dots + P_{j-1} \quad (k < i < j - 1 \leq n - 1).$$

The direct converse of Theorem 1 would postulate that among the points P_{ij} common to the lines $A_i A_j, A'_i A'_j$ all of these linear relations are satisfied. It is interesting to find, however, that if any particular one of these relations holds, all of them will hold. Consequently, a generalization of Desargues' converse theorem appears in the following form:

THEOREM 2. *Let two simplexes $A_1 A_2 \dots A_n, A'_1 A'_2 \dots A'_n$ be such that for each pair of integers i, j ($i \neq j; i, j = 1, 2, \dots, n$) there exists a corresponding point P_{ij} common to the lines $A_i A_j, A'_i A'_j$. Let k denote the number of points of the simplex $A_1 A_2 \dots A_n$ which coincide with the corresponding points of the simplex $A'_1 A'_2 \dots A'_n$ ($n \geq k \geq 0$). If $k \geq n - 2$, the simplexes are in perspective from a point. If $k < n - 2$, let the points of the simplexes be so named that A_i coincides with A'_i ($i = 1, 2, \dots, k$). If a particular point P_{rs} ($k < r < s - 1 \leq n - 1$) is linearly dependent upon all of the $s - r$ points $P_r, P_{r+1}, \dots, P_{s-1}$, the simplexes are in perspective from a point.*

If $k = n$, the simplexes coincide. They are therefore in perspective from an arbitrary point ϕ .

If $k = n - 1$, the simplexes are in perspective from any point ϕ collinear with A_n, A'_n .

If $k = n - 2$, according to hypothesis a point P_{n-1} common to the lines $A_{n-1} A_n, A'_{n-1} A'_n$ exists. The lines $A_{n-1} A'_{n-1}, A_n A'_n$ are, therefore, in a common plane and consequently concur in a point ϕ from which the simplexes are in the perspective.

If $k < n - 2$, let multipliers of the linearly dependent points $P_{rs}, P_r, P_{r+1}, \dots, P_{s-1}$ be selected such that the linear relation assumes the form

$$(6) \quad P_r + P_{r+1} + \cdots + P_{s-1} + P_{sr} = 0, \quad P_{rs} = - P_{sr}.$$

The points are themselves defined by linear relations

$$(7) \quad \begin{aligned} P_{rs} &= \alpha_{rs}A_r + \beta_{rs}A_s = \alpha'_rA'_r + \beta'_{rs}A'_s, \\ P_i &= \alpha_iA_i + \beta_{i+1}A_{i+1} = \alpha'_iA'_i + \beta'_{i+1}A'_{i+1} \end{aligned}$$

($i = r, r + 1, \dots, s - 1$).

In view of the linear independence of the A 's and of the A' 's, the coefficients of the A 's and of the A' 's which result from substituting the above relations into (6) must be equated to zero. The results follow:

$$(8) \quad \begin{aligned} \alpha_r - \alpha_{rs} &= \beta_s - \beta_{rs} = \alpha_i + \beta_i = 0, \\ \alpha'_r - \alpha'_{rs} &= \beta'_s - \beta'_{rs} = \alpha'_i + \beta'_i = 0 \quad (i = r + 1, \dots, s - 1). \end{aligned}$$

The relations (7) together with (8) imply that

$$\alpha_rA_r - \alpha'_rA'_r = \alpha_{r+1}A_{r+1} - \alpha'_{r+1}A'_{r+1} = \cdots = \beta'_sA'_s - \beta_sA_s.$$

A point ϕ is thus defined which is a point common to each of the lines $A_rA'_r, A_{r+1}A'_{r+1}, \dots, A_sA'_s$.

By means of the following argument the point ϕ will be shown to be a point common to each of the lines $A_{s+1}A'_{s+1}, \dots, A_nA'_n$. Let p denote any one of the integers $s+1, s+2, \dots, n$. According to the hypothesis, points $P_{pr}, P_{p r+1}, \dots, P_{ps}$ exist which are common to the respective pairs of lines $(A_pA_r, A'_pA'_r), (A_pA_{r+1}, A'_pA'_{r+1}), \dots, (A_pA_s, A'_pA'_s)$. These points are, therefore, defined by such linear forms as the following

$$\begin{aligned} P_{pr} &= a_{pr}A_p + b_{pr}A_r = a'_{pr}A'_p + b'_{pr}A'_r, \\ P_{p r+1} &= a_{p r+1}A_p + b_{p r+1}A_{r+1} = a'_{p r+1}A'_p + b'_{p r+1}A'_{r+1}, \\ &\dots \dots \dots \\ P_{ps} &= a_{ps}A_p + b_{ps}A_s = a'_{ps}A'_p + b'_{ps}A'_s. \end{aligned}$$

From these relations the following ones are deduced:

$$\begin{aligned} a_{pr}A_p - a'_{pr}A'_p &= b'_{pr}A'_r - b_{pr}A_r, \\ a_{p r+1}A_p - a'_{p r+1}A'_p &= b'_{p r+1}A'_{r+1} - b_{p r+1}A_{r+1}, \\ &\dots \dots \dots \\ a_{ps}A_p - a'_{ps}A'_p &= b'_{ps}A'_s - b_{ps}A_s. \end{aligned}$$

It follows that the line $A_pA'_p$ intersects each of the lines $A_rA'_r$, $A_{r+1}A'_{r+1}$, \dots , $A_sA'_s$. If $A_pA'_p$ should coincide with one of these lines, it could not coincide with another, because of the linear independence of the points of the simplexes. Therefore in such a case $A_pA'_p$ contains the point ϕ . If $A_pA'_p$ does not coincide with any of these lines, it must, nevertheless, contain the point ϕ . For if $A_pA'_p$ does not contain ϕ , its points of intersection with the above named lines and ϕ are in a common plane. But this latter condition cannot hold because the lines $A_pA'_p$, $A_rA'_r$, $A_{r+1}A'_{r+1}$, \dots , $A_sA'_s$ would then be in a common plane and the linear independence of the points of each simplex would be violated. The proof of the theorem is now complete.

Various corollaries can be stated, among which the following ones are noteworthy:

COROLLARY 1. *Let two simplexes $A_1A_2 \dots A_n$, $A'_1A'_2 \dots A'_n$ be defined in a manner such that corresponding to each pair (i, j) a point P_{ij} common to the lines A_iA_j , $A'_iA'_j$ exists. If there exists a triad (r, s, t) such that (a) A_r, A_s, A_t do not coincide with the corresponding points A'_r, A'_s, A'_t , and (b) P_{rt} is collinear with P_{rs} and P_{st} , the simplexes are in perspective from a point.*

COROLLARY 2. *Let two simplexes $A_1A_2 \dots A_n$, $A'_1A'_2 \dots A'_n$ and points P_{ij} ($i \neq j, i = 1, 2, \dots, n$) be defined as in Corollary 1. Let P_i denote the points $P_{i\ i+1}$ ($i = 1, 2, \dots, n-1$). If the point P_n is linearly dependent upon the $n-1$ points P_1, P_2, \dots, P_{n-1} but not upon a smaller number of these points, the simplexes are in perspective from a point.*

The following theorem provides a geometric interpretation relative to the simplexes of a linear relation among the points P_1, P_2, \dots, P_{n-1} .

THEOREM 3. *Let two simplexes $A_1A_2 \dots A_n$, $A'_1A'_2 \dots A'_n$ be defined in a manner such that corresponding to each pair (i, j) a point P_{ij} common to the lines A_iA_j , $A'_iA'_j$ exists. Let P_i denote the point $P_{i\ i+1}$. If a linear relation exists among the $n-1$ points P_1, P_2, \dots, P_{n-1} , but no such relation exists among a smaller number of these points, corresponding points A_i, A'_i ($i \neq 1, n$) of the two simplexes coincide, and the simplexes are in perspective from a point.*

The case of Theorem 1 which corresponds to $k=0$ may be generalized in another direction as follows:

THEOREM 4. *If two simplexes $A_1A_2 \dots A_n$, $A'_1A'_2 \dots A'_n$ are in*

perspective from a point and if each point A_i is distinct from the corresponding point A'_i , corresponding to each pair (i, j) ($i \neq j$, $i=1, 2, \dots, n$) there exists a point P_{ij} common to A_iA_j , $A'_iA'_j$ such that the points of each set $P_{1j_1}, P_{1j_2}, \dots, P_{nj_n}$, in which j_1j_2, \dots, j_n is a permutation of the set of integers $1, 2, \dots, n$, are linearly dependent.

In a similar manner, the following general theorem incorporates, as special cases, Theorem 3 and Corollary 2 of Theorem 2:

THEOREM 5. *If two simplexes $A_1A_2 \dots A_n$, $A'_1A'_2 \dots A'_n$ are such that corresponding to each pair (i, j) ($i \neq j$, $i=1, 2, \dots, n$) there exists a point P_{ij} common to A_iA_j , $A'_iA'_j$, and if either one of the following conditions holds, the simplexes are in perspective from a point: (a) among the n -points of any set $P_{1j_1}, P_{2j_2}, \dots, P_{nj_n}$, in which j_1, j_2, \dots, j_n is a permutation of the set of integers $1, 2, \dots, n$ such that $i \neq j_i$ ($i=1, 2, \dots, n$), but not among any subset of these points, there exists a linear dependence; (b) among $n-1$ points of such a set $P_{1j_1}, P_{2j_2}, \dots, P_{nj_n}$, but not among a smaller number of these points, there exists a linear dependence. If (a) forms part of the hypothesis, A_j does not coincide with A'_j for any value of j . If (b) forms part of the hypothesis and if P_{hj_h} is the point not involved in the linear dependence, A_i coincides with A'_i in each of the $n-2$ instances in which $i \neq h, j_h$.*

Theorems 3, 4, and 5 may be proved, without serious difficulty, by methods similar to those already applied. Their proofs will therefore be omitted.

If in the hypothesis of Theorem 5 condition (b) is replaced by the condition that a linear dependence exists among $n-2$ points of a set $P_{1j_1}, P_{2j_2}, \dots, P_{nj_n}$, the conditions are insufficient to force the simplexes to be in perspective from a point. To illustrate, let P_h, P_k denote two points which do not appear in a given linear relation among the remaining $n-2$ points of the set P_1, P_2, \dots, P_n . In view of the linear independence of the points of each simplex, it may be easily shown that such a relation places no restriction upon any of the points $A_h, A_{h+1}, A_k, A_{k+1}, A'_h, A'_{h+1}, A'_k, A'_{k+1}$. The existence of the points P_h, P_k having already been assumed, the existence of points ϕ_h, ϕ_k common to the pairs of lines $A_hA'_h, A_{h+1}A'_{h+1}$ and $A_kA'_k, A_{k+1}A'_{k+1}$, respectively, is assured, but the points ϕ_h, ϕ_k do not necessarily coincide. Consequently, under these conditions the simplexes are not, in general, in perspective from a point.

UNIVERSITY OF KANSAS